

Comparing deformation subspaces across anatomical variability: a population finite-element study of the human pelvis

Petr Henyš^{1,2}, Michal Kuchař⁵, Alexander Morávek⁵, Niels Hammer^{2,3,4*}

¹Institute of New Technologies and Applied Informatics, Faculty of Mechatronics, Informatics and Interdisciplinary Studies, Technical University of Liberec, Liberec, Czech Republic.

²Division of Macroscopic and Clinical Anatomy, Gottfried Schatz Research Center, Medical University of Graz, Graz, Austria.

³Department of Orthopedic and Trauma Surgery, University of Leipzig, Leipzig, Germany.

⁴Division of Biomechanics, Fraunhofer Institute for Machine Tools and Forming Technology IWU, Chemnitz and Dresden, Germany.

⁵Department of Anatomy, Faculty of Medicine in Hradec Králové, Charles University, Hradec Králové, Czech Republic.

*Corresponding author(s). E-mail(s): niels.hammer@medunigraz.at;

Abstract

Comparing deformation modes across subject-specific finite-element models requires distinguishing changes in mechanical behaviour from changes in spectral ordering. We investigated this distinction in 278 computed-tomography-derived pelvic models with individual geometry and bone elasticity on a shared reference domain. The 15 lowest linearised stiffness modes were compared using spectral gaps, modal assignment, principal angles, anatomical diameter functionals, and modal strain-energy fractions under standing and internal-ring expansion loads. Modes 1–3 formed a well-separated, order-stable group with label-exchange rates below 0.4%. A recurrent rank 9–10 cluster occurred in 54.7% of subjects. Within this cluster, median anteroposterior inlet coupling assigned to rank 9 decreased from 0.454 to 0.139 mm/mm between subjects without and with label exchange, whereas the corresponding subspace capacity changed from 0.551 to 0.525 mm/mm. Thus, the approximately 69% difference at one rank greatly exceeded the approximately 5% difference at the subspace level. Standing primarily recruited low-rank modes, whereas internal-ring expansion preferentially recruited the 9–10 block. An exploratory first-order material sensitivity analysis and a minimal analytical model contextualised the separation between rank instability and subspace-level deformation capacity. These findings support comparing compact modal subspaces when spectral crowding makes individual ranks unreliable. The framework distinguishes representation changes from changes in anatomical coupling within a linearised pelvic model. Expansion loads do not predict childbirth outcomes.

Keywords: Population biomechanics, Pelvic mechanics, Finite-element modelling, Modal analysis, Anatomical variability, Invariant subspaces

1 Introduction

The human pelvis transfers loads between the trunk and lower limbs while surrounding a canal whose dimensions have anatomical and obstetric significance. Its mechanical behaviour depends on the geometry and material distribution of the bones together with the sacroiliac joints, pubic symphysis, and ligamentous restraints. Accounts of sacroiliac load transfer emphasise this interaction between structural arrangement and stabilising tissues (Sniijders et al. 1993; Vleeming et al. 2012). Variation in pelvic form therefore raises a mechanical question beyond the description of shape: do anatomically different pelves retain comparable deformation pathways, and how are those pathways recruited by different loads? This question is relevant to understanding the relationship between pelvic form, load transmission, and canal deformation (Gruss and Schmitt 2015), even when the available models address only the initial elastic response.

Population modelling is necessary because a representative geometry cannot establish how mechanical behaviour varies among individuals. More generally, average model inputs need not produce an average biomechanical response (Cook and Robertson 2016). Population approaches have consequently become an explicit direction in computational musculoskeletal research (Fernandez et al. 2020). For the pelvis, Arand et al. (2019) used CT-derived statistical modelling to characterise variation in pelvic-ring anatomy. Such models describe the distribution of geometry, but their principal components maximise geometric variance rather than identify mechanically compliant directions. A statistical shape mode and an elastic deformation mode thus answer different questions. Connecting anatomical variability to mechanics requires constitutive information, joint representation, and specified boundary conditions in addition to geometric correspondence.

CT-derived finite-element (FE) studies already provide substantial evidence for the importance of these factors. Salo et al. (2017) compared pelvic strains across gait configurations in a validated cohort of five specimen-specific models. Their study demonstrates a route from individual anatomy to load-dependent strain predictions and documents variability between specimens. At the level of pelvic restraints, Henyš et al. (2022) examined how the sacrospinous and sacrotuberous ligaments influence pelvic kinematics and load redistribution. More recently, Chen et al. (2026) combined statistical shape modelling with musculoskeletal analysis to estimate sacroiliac and pubic joint reaction forces in females during standing and walking tasks. These studies establish relevant precedents for personalised pelvic mechanics. They also motivate a complementary question: which features of the available deformation repertoire remain comparable when both anatomy and the applied load differ?

Spectral analysis offers one way to organise that repertoire. Instead of describing the response to only one prescribed force pattern, a stiffness eigenproblem identifies deformation directions that can be combined to describe responses within a retained modal space. Pelvic modal analysis itself is established. Henyš and Čapek (2019) investigated convergence and experimental validation of a CT-derived composite pelvic-bone model. A more direct antecedent is the study by Henyš et al. (2021), which derived skeletal mechanical metrics from stiffness-matrix spectra and applied them to pelvic bones, relating these descriptors to anthropometric measurements and sex. That work established the value of spectral information beyond geometric measurements alone. It also left a consequential issue for population interpretation: an ordered eigenpair is a numerical descriptor, but its rank does not establish that its deformation pattern has the same anatomical meaning in every subject.

This issue becomes acute in crowded spectral regions. When neighbouring eigenvalues approach each other, changes in model parameters can produce marked changes in eigenvectors and their correspondence to reference patterns. Eigenvalue veering and mode localisation are well-established phenomena in structural mechanics (Pierre 1988; Manconi and Mace 2017). For clustered spectra, Ghosh and Ghanem (2012) explicitly developed an invariant-subspace approach to stochastic eigenproblems, tracking the space spanned by neighbouring modes rather than requiring each eigenvector to remain individually identifiable. Thus, neither the spectral description of pelvic mechanics nor the use of subspaces for clustered eigenvalues is introduced here. The biomechanical question is what this distinction changes when interpreting anatomical deformation across a pelvic population.

For example, one eigenvector may strongly increase an inlet diameter in one subject, while a neighbouring eigenvector carries much of that displacement pattern in another. Comparing the same rank alone could then suggest a loss of inlet-opening capability even though a similar deformation remains available through a different combination of the two modes. Conversely, proximity of eigenvalues does not guarantee that an anatomical capacity is preserved. A clustered subspace can itself change with anatomy, and its coupling to a specific diameter can change even when its dimension is unchanged. A useful comparison must therefore assess spectral separation, correspondence of the spanned spaces, and anatomically defined observables together. Basis invariance removes dependence on the coordinates chosen within one space; preservation across subjects remains a hypothesis to test.

Anatomical observables make this distinction interpretable beyond numerical stability. Here, inlet and outlet diameter changes link displacement fields to recognisable pelvic dimensions. Maximising a diameter change within a modal subspace at a prescribed maximum nodal displacement measures the kinematic capacity available in that space. This quantity differs from compliance per unit force: a pathway can permit a particular diameter change without being strongly excited by a given load. Modal strain-energy decomposition provides the complementary assessment of recruitment under controlled loading. Considering both quantities allows us to ask whether standing

and internal-ring expansion access different parts of the deformation repertoire, rather than inferring load-bearing function from mode shape alone.

Resolving this ambiguity matters whenever modal descriptors are used to compare individuals or groups. Before a difference at one rank is attributed to anatomy, sex, or material distribution, it is necessary to determine whether the compared modes retain correspondence. It also matters for the interpretation of reduced models: retaining a compact deformation space can be more meaningful than selecting individually labelled modes whose identities vary among subjects. These are immediate methodological uses of the present analysis. The broader biological motivation is to determine whether variation in pelvic form accompanies a reorganisation of mechanical pathways or a change in the anatomical deformation they can support. A linear elastic model can address the initial structural component of that question, while leaving physiological adaptation and actual childbirth mechanics unresolved.

Here, we analyse 278 CT-derived pelvic FE models using standardised geometries from BoneDat (Henyš and Kuchař 2025), subject-specific bone elasticity, and a shared reference domain. We combine spectral gaps and modal assignment with principal-angle comparisons, basis-invariant anatomical capacities, and static load decomposition. We test three linked propositions: that well-separated low-rank modes retain correspondence across the cohort; that anatomical capacity within recurrent clustered subspaces varies less with mode-exchange status than coupling assigned to individual ranks; and that standing and internal-ring expansion recruit different spectral regions. Material-sensitivity analyses and a minimal analytical model contextualise these comparisons. The contribution is a population-level connection between spectral correspondence, anatomically defined deformation capacity, and load recruitment in the pelvic ring. The inferred rank groups describe this model family; the expansion loads are controlled mechanical probes rather than simulations of labour.

2 Materials and methods

The Methods below state the computational framework in self-contained form. The core objects are (i) a pullback-assembled linear-elasticity problem on a shared reference domain, (ii) a subject-specific generalised eigenproblem, (iii) a cross-subject pairing on a common mass metric, (iv) a gap-based detector for recurrent clustered regimes, (v) invariant-subspace comparison via principal angles and the Grassmann distance, and (vi) basis-invariant anatomical functionals that supply physically interpretable coordinates inside otherwise ambiguous clustered subspaces. Wherever possible, we use notation that separates three distinct objects: the solver *rank* index j (the j -th smallest eigenvalue for each subject), the paired *reference* label $i = \pi_s^{-1}(j)$ (a cross-subject identifier via inverse assignment), and the *anatomical* functional ℓ (a linear observable defined on the template mesh).

2.1 Cohort and common-reference geometry


We analysed a precomputed set of $N=278$ CT-derived pelvic finite-element models assembled on a common reference domain ($n_F=150$, $n_M=128$; age 16–91 yr, mean 54 ± 19). No new cohort was recruited. The source repository provides standardised pelvic geometries suitable for in silico analysis (Henyš and Kuchař 2025). The reference pelvic domain Ω_{ref} was discretized using a conforming volumetric mesh comprising 166 808 linear four-node tetrahedral elements (Lagrange P_1 , 48 291 nodes, totaling 144 873 displacement degrees of freedom), with conforming interfaces between the bony segments (sacrum, left and right ossa coxae) and the joint cartilage domains (bilateral sacroiliac articular cartilage and the pubic symphysis disc). A nonrigid volumetric registration $\chi_s : \Omega_{\text{ref}} \rightarrow \Omega_s$ with deformation gradient $F_s(X) = \nabla_X \chi_s(X)$ and Jacobian $J_s = \det F_s$ maps the template domain Ω_{ref} onto each subject domain Ω_s . The registration was obtained by radial-basis-function interpolation. The registered geometries share anatomical correspondence, but size variation remains in the archived cohort. The primary analysis uses stored solver-order spectra without the separate post-hoc eigenvalue scale adjustment; isotropic size is examined as a covariate in the allometric analysis. Fig. 1 summarises the anatomical dimensions, ligament representation, and boundary-condition layout used to define the pelvic models analysed here.

2.2 Pullback-assembled linear elasticity on a common reference domain

Let $\Omega_{\text{ref}} \subset \mathbb{R}^3$ be the template domain and Ω_s the subject domain, related by a fixed volumetric registration $x = \chi_s(X)$ with deformation gradient $F_s(X) = \nabla_X \chi_s(X)$ and local Jacobian $J_s(X) = \det F_s$. Writing $\chi_s(X) = X + \varphi_s(X)$, so $F_s = I + \nabla_X \varphi_s$, small-strain linear elasticity is solved on Ω_s but integrated on Ω_{ref} through the standard change of variables (Marsden and Hughes 1983; Bonet and Wood 1997). For trial and test fields $\hat{\mathbf{u}}, \hat{\mathbf{v}} \in V \subset [H^1(\Omega_{\text{ref}})]^3$, the physical strain tensor at $x = \chi_s(X)$ is


$$\begin{aligned} \boldsymbol{\varepsilon}_x(\mathbf{u}) &= \text{sym}(\nabla_x \mathbf{u}) = \text{sym}(\nabla_X \hat{\mathbf{u}} F_s^{-1}), \\ \boldsymbol{\sigma}_x &= C_{\text{phys}}(x) : \boldsymbol{\varepsilon}_x(\mathbf{u}), \end{aligned} \tag{1}$$

A Pelvic finite-element assembly & soft tissues



- Anterior SIJ lig. ■ Posterior SIJ lig.
- Interosseous SIJ lig. ■ Sacrotuberous lig.
- Sacrospinous lig. ■ Pubic ligaments

Habitual Locomotor Loads (Standing)




Bilateral standing

2 × 400 N vertical
Acetabular notches

Unilateral standing

1 × 800 N vertical
Right acetabular notch


B1 Inlet plane (superior view)



Pelvic Inlet Diameters

- Conjugata vera (AP): 123 mm
- Transverse diameter (ML): 131 mm


B2 Outlet plane (inferior view)



Pelvic Outlet Diameters

- Biischiadic diameter (BIS): 103 mm
- Bituberous diameter (BIT): 132 mm

Parturition-Motivated Proxy Loads (Birth Canal Transit Stages)



Inlet ring expansion

±400 N lateral
Iliopectineal inner ring

Tuberosity distraction

±400 N lateral
Ischial tuberosities/spines

AP outlet distraction

±400 N anteroposterior
Pubic crest & SCJ

Fig. 1 Pelvic model and standardised loading. (A) Bony segments, joint cartilage, ligament bundles, and superior sacral restraint. (B) Inlet and outlet diameter definitions on the reference anatomy. (C) Two-leg and one-leg standing loads. (D) Comparative internal-ring expansion, ischial distraction, and anteroposterior outlet distraction loads (LAB₁–LAB₃). Each traction is normalised to the prescribed resultant across subjects. Expansion loads probe linear structural response and do not represent successive simulated stages of labour

and the pullback-assembled elastic bilinear form reads

$$a(\hat{\mathbf{u}}, \hat{\mathbf{v}}) = \int_{\Omega_{\text{ref}}} \left[C_{\text{phys}}(\chi_s(X)) : \text{sym}(\nabla_X \hat{\mathbf{u}} F_s^{-1}) \right] : \text{sym}(\nabla_X \hat{\mathbf{v}} F_s^{-1}) J_s \, dX. \quad (2)$$

Here F_s and J_s arise only from the fixed image registration; no finite-strain mechanics is implied. The mass bilinear form inherits the geometric volume factor,

$$m(\hat{\mathbf{u}}, \hat{\mathbf{v}}) = \int_{\Omega_{\text{ref}}} J_s(X) \hat{\mathbf{u}} \cdot \hat{\mathbf{v}} \, dX, \quad (3)$$

defining an L^2 volume inner product weighted by local volumetric dilation rather than physical dynamic mass density. The resulting generalised eigenvalues $\lambda_k = \phi_k^\top K_s \phi_k / (\phi_k^\top M_s \phi_k)$ are stiffness-spectrum descriptors in a volume metric. With coordinates in mm and modulus in MPa they have units N/mm⁴; their reciprocals have units mm⁴/N and are not point compliances. They are not physical resonance frequencies because M_s does not include mass density. The *unmapped* reference mass matrix M_{ref}^0 is obtained from (3) with $J \equiv 1$ and provides a shared, registration-independent inner product across subjects.

Material model. The physical elasticity tensor $C_{\text{phys}}(x)$ is isotropic and heterogeneous. For bone, the CT-derived hydroxyapatite density $\rho(x)$ is converted to ash density $\rho_{\text{ash}} = (\rho + 0.09)/1.14$ and mapped to a Young's modulus

$E(x)$ using an implemented piecewise approximation motivated by Keller (Keller 1994),

$$E(x) = \begin{cases} 2398 \text{ MPa}, & \rho_{\text{ash}}(x) \leq 0.486, \\ \alpha \rho_{\text{ash}}(x)^\beta, & \rho_{\text{ash}}(x) > 0.486, \end{cases} \quad (4)$$

$$\alpha = 10\,200, \quad \beta = 2.0,$$

with Poisson’s ratio $\nu = 0.3$. Negative input densities are clipped to zero. The law is applied at source points, after which the resulting modulus is interpolated to FE cells using RBF interpolation (smoothing parameter 50, ten neighbours). The constant low-density branch and the exponent 2.0 are modelling choices rather than an exact reproduction of the cited regression. Density–modulus calibration depends on anatomical site (Morgan et al. 2003) and was not performed individually for this cohort. Cartilage (sacroiliac joints and pubic symphysis) is homogeneous linear elastic with $E = 1.0 \text{ MPa}$ and $\nu = 0.45$. These parameters are prescribed effective values. General accounts of bone adaptation (Frost 1987; Huiskes et al. 1987) and pelvic joint mechanics (Snijders et al. 1993; Vleeming et al. 2012; Goode et al. 2008) provide context but do not validate their numerical values.

Ligaments in pullback coordinates. Ligament families are represented as polyline bundles mapped from template nodes X_i to physical points $x_i = \chi_s(X_i)$. For each segment (X_i, X_j) define the physical unit tangent $\mathbf{e} = (x_j - x_i)/\|x_j - x_i\|$ and length $L = \|x_j - x_i\|$. The total bundle stiffness and pretension are distributed across N_{seg} segments, yielding per-segment $k_{\text{seg}} = K_{\text{bundle}}/N_{\text{seg}}$ and $T_{\text{seg}} = T_{\text{bundle}}/N_{\text{seg}}$. Each segment contributes an axial stiffness block

$$W = k_{\text{seg}} (\mathbf{e}\mathbf{e}^\top), \quad K_{ij}^{\text{ax}} = \begin{bmatrix} +W & -W \\ -W & +W \end{bmatrix}, \quad (5)$$

and, in the presence of pretension $T_{\text{seg}} \geq 0$, a geometric (prestress) stiffness

$$W_g = \frac{T_{\text{seg}}}{L} (I - \mathbf{e}\mathbf{e}^\top), \quad K_{ij}^{\text{geo}} = \begin{bmatrix} W_g & -W_g \\ -W_g & W_g \end{bmatrix} \succeq 0, \quad (6)$$

together with paired nodal forces $\pm T_{\text{seg}} \mathbf{e}$ on the right-hand side. The assembled linear stiffness, including prescribed ligament prestress, is

$$K_s = K_{\text{elas},s} + K_{\text{lig},s} + K_s^{\text{geo}}(T) + K_{\Gamma,s}, \quad f = f_{\text{ext}} + f_{\text{pret}}(T), \quad (7)$$

where geometric stiffness $K_s^{\text{geo}}(T)$ incorporates the stabilizing effect of initial ligament pretension T . This is a linear approximation with prescribed prestress; a geometrically nonlinear equilibrated prestressed configuration was not established. The term $K_{\Gamma,s}$ implements Dirichlet penalties on the restrained superior sacral facet Γ_{ref} , $a_\Gamma(\hat{\mathbf{u}}, \hat{\mathbf{v}}) = \gamma \int_{\Gamma_{\text{ref}}} \hat{\mathbf{u}} \cdot \hat{\mathbf{v}} J_{\Gamma,s} dS_X$ with $J_{\Gamma,s} = J_s \|F_s^{-T} N\|$, using a fixed penalty parameter $\gamma = 10^6 \text{ N/mm}^3$ prescribed directly without element-size or modulus scaling. A decade sweep of γ across $[10^5, 10^7] \text{ N/mm}^3$ left the reported cluster classification and modal subspace capacities unchanged.

Force-controlled loading. Neumann terms are pulled back to Γ_{ref} so that physical resultants are invariant under χ_s :

$$\int_{\Gamma_s} \mathbf{t}_{\text{phys}} \cdot \mathbf{v} dS_x = \int_{\Gamma_{\text{ref}}} J_{\Gamma,s}(X) \mathbf{t}_{\text{phys}}(\chi_s(X)) \cdot \hat{\mathbf{v}}(X) dS_X. \quad (8)$$

All load cases in this study are defined by a total force $\mathbf{F}$ on a tagged physical region; the applied traction is $\mathbf{t}_{\text{phys}} = \mathbf{F}/A_{\text{phys}}$ with $A_{\text{phys}} = \int_{\Gamma_{\text{ref}}} J_{\Gamma,s} dS_X$, so that $\int_{\Gamma_s} \mathbf{t}_{\text{phys}} dS_x = \mathbf{F}$ for every subject.

2.3 Standardised load cases

Loads were applied as force-controlled Neumann tractions via (8) so that every subject experienced the same integrated force vector regardless of physical surface area. Two standing cases were considered: symmetric two-leg stance (SP2leg; 400 N per acetabulum, +CC) and one-leg stance (SP1leg; 800 N on the stance acetabulum, +CC). The 400 N per hip and 800 N one-leg magnitudes are prescribed comparative loads. The cited in-vivo hip-contact forces provide context for physiological loading but do not validate these particular traction fields (Bergmann et al. 2001, 2016). Three proxy load cases probed parturition-relevant routing without modelling contact mechanics: LAB₁ applied bilateral mediolateral internal ring expansion forces ($\pm 400 \text{ N}$ outward), LAB₂ applied bilateral ischial-tuberosity distraction ($\pm 400 \text{ N}$), and LAB₃ applied anteroposterior outlet distraction between the pubis insertion ($-AP$, 400 N) and the sacrococcygeal joint ($+AP$, 400 N). These loads act as standardized comparative probes of pelvic structural compliance rather than clinical simulations of labour (Chen and Grimm 2021; Roque de Almeida et al. 2024). Static finite-element solutions solve the linear equilibrium equation $K_s \mathbf{u} = \mathbf{b}$ with

$$\mathbf{b} = \mathbf{f}_{\text{ext}} + \mathbf{f}_{\text{pret}} \quad (9)$$

for the primary total-response analysis, where $\mathbf{f}_{\text{pret}}$ represents baseline ligament pretension (20 N per ligament bundle) and $\mathbf{f}_{\text{ext}}$ represents external load. Primary cohort-wide modal strain-energy fractions (Section 3.7; Table 13) are computed directly from these total displacement fields $\mathbf{u}$. By linearity, total displacement decomposes as $\mathbf{u} = \mathbf{u}_{\text{pret}} + \delta\mathbf{u}_{\text{ext}}$, where $\delta\mathbf{u}_{\text{ext}} = K_s^{-1}\mathbf{f}_{\text{ext}}$ is the incremental external load response. Under external load scaling $\mathbf{f}_{\text{ext}} \rightarrow c\mathbf{f}_{\text{ext}}$, incremental displacements scale proportionally ($\delta\mathbf{u}_{\text{ext}} \rightarrow c\delta\mathbf{u}_{\text{ext}}$), such that modal expansion coefficients $q_i = \phi_i^\top M_s \delta\mathbf{u}_{\text{ext}}$ scale by c , modal energies $E_i = \frac{1}{2}\lambda_i q_i^2$ scale by c^2 , and the normalized modal strain-energy fractions $f_i = E_i / \sum_j E_j$ of the incremental response are strictly scale-invariant. To examine the effect of baseline pretension on routing, we compared archived total-response and incremental-response modal fractions in 18 subjects with complete stored incremental outputs. The top recruited mode agreed for all five loads in all 18 subjects. The maximum individual change in the 9–10 block fraction was 4.24 percentage points, while its mean change under LAB₁ was +3.48 percentage points. The smallest dominant set reaching 80% of retained energy had the same size in all 18 standing cases and in 14 of 18 LAB₁ cases. This subset supports similar dominant routing but does not establish exact invariance across the full cohort.

2.4 Generalised eigenproblem, rigid-mode filtering, and cross-subject pairing

For each subject s we assembled (K_s, M_s) on Ω_{ref} via (7) and (3) and solved the generalised eigenproblem

$$K_s \phi_k = \lambda_k M_s \phi_k, \quad \phi_i^\top M_s \phi_j = \delta_{ij}, \quad (10)$$

using a Krylov–Schur solver (SLEPc) with shift-invert spectral transform and MUMPS direct factorisation, extracting the $N_{\text{ev}} = 15$ lowest non-rigid eigenpairs. Rigid modes (numerically $\lambda < 10^{-8}$) were filtered. For cross-subject comparison we used the mass-weighted modal assurance criterion on the shared reference metric,

$$\text{MAC}_{M_0}(\mathbf{x}, \mathbf{y}) = \frac{|\mathbf{x}^\top M_{\text{ref}}^0 \mathbf{y}|^2}{(\mathbf{x}^\top M_{\text{ref}}^0 \mathbf{x})(\mathbf{y}^\top M_{\text{ref}}^0 \mathbf{y})} \in [0, 1], \quad (11)$$

and solved the Hungarian optimal-assignment problem on $-\text{MAC}_{M_0}$ over the full $N_{\text{ev}} \times N_{\text{ev}}$ matrix (Kuhn 1955), yielding an assignment bijection $\pi_s : \{1, \dots, N_{\text{ev}}\} \rightarrow \{1, \dots, N_{\text{ev}}\}$, where $\pi_s(i_{\text{ref}}) = j_{\text{rank}}$ maps reference identifier i_{ref} to its rank position in subject s , and the inverse permutation $\pi_s^{-1}(j_{\text{rank}}) = i_{\text{ref}}$ assigns reference identity to each solver rank. Entries with $\text{MAC}_{M_0} < 0.05$ were flagged as unpaired. Rank index j (solver order) is required for gap statistics and cluster detection; the inverse assignment $\pi_s^{-1}(j)$ maps each solver mode back to its reference identity and is used only for anatomical interpretation, preserving the distinction between rank order and pathway identity emphasised in the Introduction.

2.5 Spectral gap structure and clustered-regime detection

For each subject with ordered eigenvalues $\lambda_1 \leq \dots \leq \lambda_{N_{\text{ev}}}$, relative consecutive gaps are

$$\text{gap}_{\text{in}}(i) = \frac{\lambda_{i+1} - \lambda_i}{\lambda_i}, \quad i = 1, \dots, N_{\text{ev}} - 1. \quad (12)$$

A clustered regime $[i..j]$ is defined as the maximal contiguous rank block starting at i with internal gaps below threshold, $\text{gap}_{\text{in}}(k) < \varepsilon_{\text{in}}$ for $k = i, \dots, j - 1$. The threshold is set from data as the 25th percentile of the pooled gap_{in} distribution across the cohort, yielding $\varepsilon_{\text{in}} \approx 0.13$; sensitivity to the percentile choice is reported in Section 3.10. External separation is quantified by

$$\text{sep}_{\text{out}}([i..j]) = \min\left(\frac{\lambda_i - \lambda_{i-1}}{\lambda_i}, \frac{\lambda_{j+1} - \lambda_j}{\lambda_j}\right), \quad (13)$$

defined for interior clusters ($i > 1$ and $j < N_{\text{ev}}$) where both adjacent neighbours exist. For boundary blocks reaching modal truncation (such as the 12–15 cluster with $N_{\text{ev}} = 15$), the upper neighbour is uncomputed and two-sided separation is formally undefined. Exact mathematical multiplicity ($\lambda_i = \lambda_{i+1}$) was not observed under the operational numerical tolerance $\text{gap}_{\text{in}} \leq \varepsilon_{\text{deg}} = 10^{-6}$ in any subject. No obvious spatial symmetry was identified that would enforce repeated eigenvalues, although accidental parameter-dependent crossings are not mathematically excluded; nonetheless, the regimes detected by (12) are operational small-gap clusters in which perturbation sensitivity may be elevated (Pierre 1988; Manconi and Mace 2017).

2.6 Invariant subspaces: principal angles, Grassmann distance, and Procrustes alignment

For each clustered rank block of dimension r , the corresponding subject-specific subspace was represented by an M_{ref}^0 -orthonormal basis

$$Q_s \in \mathbb{R}^{n \times r}, \quad Q_s^\top M_{\text{ref}}^0 Q_s = I_r. \quad (14)$$

With a fixed reference basis Q_{ref} , define the overlap matrix

$$S = Q_{\text{ref}}^\top M_{\text{ref}}^0 Q_s \in \mathbb{R}^{r \times r}, \quad S = U \Sigma V^\top, \quad \cos \theta_k = \sigma_k(S), \quad (15)$$

with singular values σ_k yielding the principal angles $\theta_k \in [0, \pi/2]$ between the two subspaces (Björck and Golub 1973). The Grassmann distance is

$$d_G(Q_{\text{ref}}, Q_s) = \|\theta\|_2 = \left(\sum_{k=1}^r \theta_k^2 \right)^{1/2}. \quad (16)$$

When a stable basis representation inside a clustered subspace is required, we apply an orthogonal Procrustes rotation (Schönemann 1966),

$$\tilde{Q}_s = Q_s (V U^\top), \quad (17)$$

which minimises the M_{ref}^0 -weighted distance $\|Q_{\text{ref}} - Q_s R\|_{M_{\text{ref}}^0}$ over orthogonal matrices $R \in O(r)$ while preserving the subspace $\text{span}(Q_s)$.

Subspace comparisons use basis-invariant scalars derived below. Individual-rank couplings remain explicit comparators whose sensitivity to mode exchange is being tested.

Degeneracy vs. near-degenerate mixing. Exact mathematical multiplicity requires $\lambda_i = \lambda_{i+1}$ (operationalized numerically as $\text{gap}_{\text{in}}(k) \leq \varepsilon_{\text{deg}}$ for all k in the block, with $\varepsilon_{\text{deg}} \ll \varepsilon_{\text{in}}$); individual eigenvectors inside a degenerate subspace are then not unique, and any unitary rotation inside $\text{span}(Q_s)$ is an equally valid basis. When $\varepsilon_{\text{deg}} < \text{gap}_{\text{in}}(k) < \varepsilon_{\text{in}}$ the eigenvalues are close but distinct; each eigenvector is unique up to sign, yet small perturbations can rotate the internal basis substantially while the spanned subspace can be less sensitive if its external separation is sufficiently large relative to the perturbation. Neither a small internal gap nor label exchange alone proves preservation of anatomical coupling.

2.7 Canonical anatomical functionals inside clustered subspaces

To interpret clustered subspaces biologically, we use basis-invariant linear functionals defined by anatomical landmark pairs. On the reference template mesh, five key obstetric and locomotor diameters are defined:

- (i) **AP inlet** (s_{AP}): Promontory (X_{prom}) to superior pubic symphysis (X_{pub}), with unit direction $\mathbf{e}_{\text{AP}} = (X_{\text{pub}} - X_{\text{prom}}) / \|X_{\text{pub}} - X_{\text{prom}}\|$;
- (ii) **ML inlet** (s_{ML}): Widest transverse diameter across left and right iliopectineal lines ($X_{\text{left}}, X_{\text{right}}$), with unit direction $\mathbf{e}_{\text{ML}} = (X_{\text{right}} - X_{\text{left}}) / \|X_{\text{right}} - X_{\text{left}}\|$;
- (iii) **Interspinous midpelvic diameter (BIS)** (s_{BIS}): Left to right ischial spine ($X_{\text{sp, left}}, X_{\text{sp, right}}$), with unit direction $\mathbf{e}_{\text{BIS}} = (X_{\text{sp, right}} - X_{\text{sp, left}}) / \|X_{\text{sp, right}} - X_{\text{sp, left}}\|$;
- (iv) **Bituberous outlet** (s_{BIT}): Left to right inferior ischial tuberosity ($X_{\text{tub, left}}, X_{\text{tub, right}}$), with unit direction $\mathbf{e}_{\text{BIT}} = (X_{\text{tub, right}} - X_{\text{tub, left}}) / \|X_{\text{tub, right}} - X_{\text{tub, left}}\|$;
- (v) **AP outlet** (s_{OUTLETAP}): Sacrococcygeal joint (X_{cocc}) to inferior pubic symphysis ($X_{\text{pub, inf}}$), with unit direction $\mathbf{e}_{\text{OUTAP}} = (X_{\text{pub, inf}} - X_{\text{cocc}}) / \|X_{\text{pub, inf}} - X_{\text{cocc}}\|$.

For each subject s , patient-specific landmark coordinates follow the registration mapping $X^s = X + \varphi_s(X)$, and the resulting functional observable for a displacement field $\mathbf{u}$ is

$$s_\ell^{(s)}(\mathbf{u}) = \mathbf{e}_\ell^{(s)\top} (\mathbf{u}(X_{\text{target}}^s) - \mathbf{u}(X_{\text{origin}}^s)) = \mathbf{b}_\ell^{(s)\top} \mathbf{u}, \quad \ell \in \{\text{AP}, \text{ML}, \text{BIS}, \text{BIT}, \text{OUTLETAP}\}, \quad (18)$$

with measurement vectors $\mathbf{b}_\ell^{(s)} \in \mathbb{R}^n$ constructed using inverse-distance-squared weighting over the 32 nearest reference-mesh nodes at each template landmark. These weights remain fixed across subjects; the direction $\mathbf{e}_\ell^{(s)}$ is updated from RBF-mapped landmark positions. BIS measures the midpelvis between the ischial spines, whereas BIT measures the intertuberous outlet. The AP-outlet observable uses the sacrococcygeal landmark and is not a coccygeal-tip diameter.

Basis-invariant max-nodal anatomical capacity. For any individual functional $\mathbf{b}_\ell^{(s)}$ and an orthonormal basis $Q_s \in \mathbb{R}^{n \times r}$ spanning a clustered subspace of dimension r ($Q_s^\top M_{\text{ref}}^0 Q_s = I_r$), every displacement in the subspace is

parameterized by $\mathbf{u} = Q_s \boldsymbol{\alpha}$ with $\boldsymbol{\alpha} \in \mathbb{R}^r$. The maximum anatomical diameter response per 1 mm maximum nodal displacement is defined by the constrained max-nodal capacity:

$$\kappa_{\ell, \infty}(Q_s) = \max_{\boldsymbol{\alpha} \neq \mathbf{0}} \frac{|\mathbf{b}_\ell^{(s)\top} Q_s \boldsymbol{\alpha}|}{\max_a \|(Q_s \boldsymbol{\alpha})_a\|_2} = \max_{\boldsymbol{\alpha}} |\mathbf{b}_\ell^{(s)\top} Q_s \boldsymbol{\alpha}| \quad \text{s.t.} \quad \max_a \|(Q_s \boldsymbol{\alpha})_a\|_2 \leq 1, \quad (19)$$

where a indexes the finite-element mesh nodes and $\|(Q_s \boldsymbol{\alpha})_a\|_2$ denotes the Euclidean displacement norm of node a . Because both the numerator and the max-nodal constraint depend only on the physical displacement field $\mathbf{u} = Q_s \boldsymbol{\alpha}$, this quantity is strictly invariant under arbitrary internal orthogonal basis rotations $\tilde{Q}_s = Q_s R$ with $R \in O(r)$, since $\tilde{Q}_s(R^\top \boldsymbol{\alpha}) = Q_s \boldsymbol{\alpha}$. For $r = 2$, the zero-homogeneity of the objective allows parameterization of $\boldsymbol{\alpha}$ by polar angle $\theta \in [0, \pi)$ with $\boldsymbol{\alpha}(\theta) = (\cos \theta, \sin \theta)^\top$, estimated from 90 angular samples followed by scalar refinement. For $r \geq 3$, we solve the equivalent convex program $\max_{\boldsymbol{\alpha}} \mathbf{c}^\top \boldsymbol{\alpha}$ subject to $\|(Q_s \boldsymbol{\alpha})_a\|_2 \leq 1$ at every node, where $\mathbf{c} = Q_s^\top \mathbf{b}_\ell^{(s)}$; the sign symmetry removes the absolute value. Constraint generation starts with 16 nodes, solves each reduced problem with SLSQP and analytic Jacobians, adds the most violated mesh node, and terminates only when the full-mesh maximum squared norm is at most $1 + 10^{-8}$. The objective is linear and every constraint is convex, so a feasible stationary solution is a global optimum of the finite-dimensional problem up to numerical tolerance. Four independent full-constraint optimisations provide an additional check. The optimal spatial displacement field $\boldsymbol{\psi}_\ell^{(s)} = Q_s \boldsymbol{\alpha}^* / \max_a \|(Q_s \boldsymbol{\alpha}^*)_a\|_2$ satisfies $\max_a \|\boldsymbol{\psi}_{\ell, a}^{(s)}\|_2 = 1.0$ exactly, and the resulting anatomical capacity $\kappa_{\ell, \infty}(Q_s) = |\mathbf{b}_\ell^{(s)\top} \boldsymbol{\psi}_\ell^{(s)}|$ has physical units of millimetres diameter change per millimetre maximum nodal displacement.

Multivariate canonical decomposition. When joint coupling across a pair of measurement vectors $(\mathbf{b}_1^{(s)}, \mathbf{b}_2^{(s)})$ (e.g., AP and ML inlet, or BIS and BIT outlet) is examined, we assemble the covector matrix $B_s = [\mathbf{b}_1^{(s)} \quad \mathbf{b}_2^{(s)}] \in \mathbb{R}^{n \times 2}$ and its subspace projection:

$$C_s = Q_s^\top B_s = [Q_s^\top \mathbf{b}_1^{(s)} \quad Q_s^\top \mathbf{b}_2^{(s)}] \in \mathbb{R}^{r \times 2}, \quad (20)$$

$$C_s = U_s \Sigma_s V_s^\top,$$

where the singular values $\sigma_1 \geq \sigma_2$ describe joint coupling under the coefficient constraint $\|\boldsymbol{\alpha}\|_2 = 1$, equivalent to unit norm in M_{ref}^0 . They do not give maxima under the max-nodal displacement constraint. The canonical displacement modes in physical space are $\boldsymbol{\psi}_k^{(s)} = Q_s U_s[:, k]$ ($k = 1, 2$), with corresponding dual functionals in covector space $\tilde{\mathbf{b}}_k^{(s)} = B_s V_s[:, k] = V_s[0, k] \mathbf{b}_1^{(s)} + V_s[1, k] \mathbf{b}_2^{(s)}$. By construction, their action is exactly equal to the singular values: $\tilde{\mathbf{b}}_k^{(s)\top} \boldsymbol{\psi}_k^{(s)} = \sigma_k$. The singular-direction scores subsequently normalized by maximum nodal displacement are $\hat{s}_k^{(s)} = \sigma_k / \max_a \|\boldsymbol{\psi}_{k, a}^{(s)}\|_2$. Because $V_s \in O(2)$ mixes the two anatomical axes, mode $\boldsymbol{\psi}_1^{(s)}$ is primary-dominant only when $|V_s[0, 0]| \geq |V_s[1, 0]|$; across the pelvic cohort, the first singular mode in the 9–10 subspace is predominantly aligned with the more strongly coupled transverse (ML) axis ($|V_s[1, 0]| > |V_s[0, 0]|$ in 99.6% of subjects). For unambiguous, axis-specific evaluation of anatomical opening without coordinate mixing, the basis-invariant single-functional capacity $\kappa_{\ell, \infty}(Q_s)$ (Eq. (19)) serves as our primary quantitative metric, while joint-SVD scores provide complementary bivariate structure.

2.8 Whole-pelvis deformation decomposition

To distinguish global ring kinematics from joint-localised strains, we decompose the reference displacement fields after removing infinitesimal rigid motion. The fractions below describe the explained squared displacement norm within a specified polynomial dictionary, not elastic energy. Synthetic checks, the reference-mode atlas, and complete profile tables are provided in Section 3.1.

2.8.1 Joint strain and global ring kinematics

The pelvic ring is a closed, doubly-curved structural biomechanical loop composed of three major bony segments (the sacrum and bilateral coxal bones) joined anteriorly at the pubic symphysis and posteriorly at the bilateral sacroiliac joints (SIJs). In continuum mechanics, the standard approach to characterise modal kinematics would be to evaluate the infinitesimal Cauchy strain tensor field,

$$\boldsymbol{\varepsilon}(\mathbf{u}) = \frac{1}{2} (\nabla \mathbf{u} + \nabla \mathbf{u}^\top). \quad (21)$$

In the present quasi-static finite-element formulation, an effective joint modulus of $E_{\text{cartilage}} = 1.0 \text{ MPa}$ is prescribed for the sacroiliac cartilage cleft and pubic symphysis disc. This prescribed modulus represents compliant joint regions within the model, which are three to four orders of magnitude more compliant than cortical and trabecular

bone ($E_{\text{bone}} \sim 5\text{--}18$ GPa). Under the generalised modal formulation, elastic deformation naturally channels into these narrow compliant interfaces. Consequently, direct L^2 -projection of the Cauchy strain field onto low-order polynomial strain templates leaves between 97% and 99% of the strain norm unexplained in the residual across all fifteen computed eigenmodes. The intense local strain spikes in the millimeter-thin joint clefts completely overpower and conceal the systemic, ring-scale kinematics (such as whole-pelvis flaring, torsion, and inlet/outlet dilation).

A dictionary conditioned along only one anatomical axis did not describe coupled three-dimensional ring deformation adequately in the tested example. For complex three-dimensional deformation pathways involving coupled out-of-plane flexure and transverse shearing (such as Mode 2), the tested one-axis dictionary left an 84% residual. This reflects the scope of that descriptive basis and is not evidence that all beam approximations are unphysical. To overcome both limitations, our framework evaluates the overall shape change across the entire three-dimensional pelvic continuum without anatomical partitioning and without imposing artificial 1D beam axes, removing infinitesimal rigid-body motion upfront before decomposing the remaining shape change into four canonical 3D polynomial deformation families.

2.8.2 Continuum inner product and tetrahedral quadrature

Let $\Omega \subset \mathbb{R}^3$ denote the reference pelvic continuum domain with volume $V = \int_{\Omega} dV$. The natural inner product on the displacement space $[L^2(\Omega)]^3$ weighted by volume is

$$\begin{aligned} \langle \mathbf{u}, \mathbf{v} \rangle_V &= \frac{1}{V} \int_{\Omega} \mathbf{u}(\mathbf{x}) \cdot \mathbf{v}(\mathbf{x}) dV, \\ \|\mathbf{u}\|_V^2 &= \langle \mathbf{u}, \mathbf{u} \rangle_V. \end{aligned} \quad (22)$$

The continuous reference domain Ω is discretized into an unstructured tetrahedral mesh $\mathcal{T}_h = \{K_e\}_{e=1}^{N_e}$ using piecewise-linear (P_1) Lagrange continuous finite elements. Within any tetrahedron K , a P_1 displacement field $\mathbf{u}_h(\mathbf{x})$ is an affine vector function of position ($\mathbf{u}_h|_K \in \mathcal{P}_1(K)$). When computing projections against quadratic deformation modes $\mathbf{p}_2 \in \mathcal{P}_2(K)$, the modal cross-product integrand $\mathbf{u}_h \cdot \mathbf{p}_2$ is a cubic polynomial ($\mathcal{P}_3(K)$), and the dictionary Gram matrix integrand $\mathbf{p}_2 \cdot \mathbf{p}_2$ is a quartic polynomial ($\mathcal{P}_4(K)$).

A four-point tetrahedral rule is exact for the squared norm of a P_1 displacement field ($\mathcal{P}_2(K)$), but not generally for the quadratic dictionary products used below. The reference atlas and its accompanying table were regenerated with a positive degree-4 tetrahedral quadrature rule with 14 points per tetrahedron (constructed via Basix). For a rule exact to polynomial degree d ($d = 2$ for four points, $d = 4$ for 14 points), the quadrature points $\mathbf{x}_{K,q}$ and strictly positive weights $w_{K,q} > 0$ satisfy $\sum_{q=1}^{N_{q,\text{tet}}} w_{K,q} = |K|$:

$$\int_K p(\mathbf{x}) dV = \sum_{q=1}^{N_{q,\text{tet}}} w_{K,q} p(\mathbf{x}_{K,q}) \quad \forall p \in \mathcal{P}_d(K). \quad (23)$$

Summing across all N_e tetrahedra gives $N_q = N_{q,\text{tet}} N_e$ discrete sampling points with weights $w_p > 0$ ($N_q = 4N_e$ for the 4-point rule; $N_q = 14N_e$ for the degree-4 rule). Defining normalized quadrature weights $\tilde{w}_p = w_p / \sum_{j=1}^{N_q} w_j$, the continuum inner product (22) is evaluated as:

$$\langle \mathbf{u}, \mathbf{v} \rangle_V \approx \sum_{p=1}^{N_q} \tilde{w}_p \mathbf{u}(\mathbf{x}_p) \cdot \mathbf{v}(\mathbf{x}_p). \quad (24)$$

The degree-4 rule integrates the products of quadratic dictionary functions exactly on each tetrahedron. Because the reference modes are P_1 , it also integrates their cross-products with the dictionary exactly. Figure 2 links the element rule to the whole-domain inner product and to the degrees of the three products used in the decomposition.

2.8.3 Infinitesimal rigid-body motion elimination

Infinitesimal rigid-body displacements in $\mathbb{R}^3$ constitute a 6-dimensional linear Lie algebra subspace $\mathfrak{se}(3)$ consisting of 3 translations and 3 rotations:

$$\begin{aligned} \mathbf{u}_{\text{rigid}}(\mathbf{x}) &= \mathbf{t} + \boldsymbol{\omega} \times (\mathbf{x} - \mathbf{x}_c), \\ \mathbf{t}, \boldsymbol{\omega} &\in \mathbb{R}^3, \end{aligned} \quad (25)$$

where $\mathbf{x}_c$ is the volume-weighted spatial centroid of the pelvic domain,

$$\mathbf{x}_c = \langle \mathbf{x} \rangle_V = \sum_{p=1}^{N_q} \tilde{w}_p \mathbf{x}_p. \quad (26)$$

Volume integration of modal and polynomial fields




Fig. 2 Volume quadrature for the whole-pelvis displacement decomposition. (A) Reference pelvic mesh with actual quadrature positions displayed for every 80th tetrahedron, solely to make the volume sampling visible; the computation uses all 166 808 cells. (B) The positive 14-point degree-4 Basix rule in one reference tetrahedron. Colour and sphere size distinguish its three weight groups (six, four, and four points); weights are positive and sum to the tetrahedral volume. (C) Polynomial degrees of the squared P_1 FE displacement, its cross-product with a quadratic dictionary field, and the quadratic dictionary Gram product. The 14-point rule is exact through degree 4 on each tetrahedron. The illustrations show sampling geometry, not subject-specific deformation.

To ensure scale-free, dimensionally consistent kinematics across subjects, we define the reference gyration radius r_g and non-dimensionalised anatomical coordinates $\tilde{\mathbf{x}} = (\tilde{x}, \tilde{y}, \tilde{z})^\top$:

$$r_g = \sqrt{\langle \|\mathbf{x} - \mathbf{x}_c\|^2 \rangle_V} = \sqrt{\sum_{p=1}^{N_q} \tilde{w}_p \|\mathbf{x}_p - \mathbf{x}_c\|^2}, \quad (27)$$

$$\tilde{\mathbf{x}} = \frac{\mathbf{x} - \mathbf{x}_c}{r_g}.$$

Here the Cartesian coordinate directions correspond to the standard anatomical axes: x is mediolateral (ML), y is anteroposterior (AP), and z is craniocaudal (CC).

In the discrete quadrature representation, each displacement field is a stacked vector $\mathbf{U} \in \mathbb{R}^{3N_q}$ with components $U_{3p-3+i} = \sqrt{\tilde{w}_p} u_i(\mathbf{x}_p)$ for $i \in \{1, 2, 3\}$. The 6 rigid-body design modes $\mathbf{r}_1, \dots, \mathbf{r}_6 \in \mathbb{R}^{3N_q}$ (corresponding to unit translations along x, y, z and unit infinitesimal rotations about x, y, z) form the rigid design matrix $\mathbf{R} \in \mathbb{R}^{3N_q \times 6}$. We compute its thin QR decomposition,

$$\mathbf{R} = \mathbf{Q}_{\text{rigid}} \mathbf{R}_{\text{upper}}, \quad (28)$$

$$\mathbf{Q}_{\text{rigid}} \in \mathbb{R}^{3N_q \times 6}, \quad \mathbf{Q}_{\text{rigid}}^\top \mathbf{Q}_{\text{rigid}} = \mathbf{I}_6.$$

The orthogonal projection operator onto the nonrigid displacement subspace is then:

$$\mathbf{P}_{\text{nonrigid}} = \mathbf{I}_{3N_q} - \mathbf{Q}_{\text{rigid}} \mathbf{Q}_{\text{rigid}}^\top. \quad (29)$$

For any raw modal displacement field $\mathbf{u}$, the nonrigid component is $\mathbf{u}_{\text{nonrigid}} = \mathbf{P}_{\text{nonrigid}} \mathbf{u}$, and the total squared displacement norm partitions orthogonally:

$$\|\mathbf{u}\|_V^2 = \|\mathbf{u}_{\text{rigid}}\|_V^2 + \|\mathbf{u}_{\text{nonrigid}}\|_V^2, \quad (30)$$

$$f_{\text{nonrigid}} = \frac{\|\mathbf{u}_{\text{nonrigid}}\|_V^2}{\|\mathbf{u}\|_V^2}.$$

Crucially, rigid rotation is eliminated at this foundational stage: it is neither retained in the target deformation field nor reported as an explained deformation category. If a displacement field satisfies $\|\mathbf{u}_{\text{nonrigid}}\|_V^2 / \|\mathbf{u}\|_V^2 \leq 10^{-20}$, it is identified as a pure rigid motion and excluded from nonrigid decomposition.

2.8.4 Canonical 3D polynomial dictionary

The vector space of polynomial displacement fields $\mathbf{u}(\tilde{\mathbf{x}})$ of degree at most 2 in three dimensions has complete algebraic dimension $3 \times \binom{3+2}{2} = 30$. Subtracting the 6 degrees of freedom of infinitesimal rigid motion leaves an exact 24-dimensional nonrigid polynomial quotient space. We decompose this space into four canonical physical deformation families spanning all 24 nonrigid degrees of freedom:

- (i) **Axial family** ($d_{\text{axial}} = 6$ degrees of freedom): Uniform normal extension/compression along each anatomical axis, together with quadratic normal strain gradients:

$$\begin{aligned} \mathbf{a}_{x,1} &= (\tilde{x}, 0, 0)^\top, \\ \mathbf{a}_{x,2} &= (\tfrac{1}{2}\tilde{x}^2, 0, 0)^\top, \\ \mathbf{a}_{y,1} &= (0, \tilde{y}, 0)^\top, \\ \mathbf{a}_{y,2} &= (0, \tfrac{1}{2}\tilde{y}^2, 0)^\top, \\ \mathbf{a}_{z,1} &= (0, 0, \tilde{z})^\top, \\ \mathbf{a}_{z,2} &= (0, 0, \tfrac{1}{2}\tilde{z}^2)^\top. \end{aligned} \tag{31}$$

- (ii) **Bending family** ($d_{\text{bending}} = 6$ degrees of freedom): Classical pure Euler-Bernoulli flexural modes. For any pair of distinct coordinate axes $i \neq j \in \{x, y, z\}$, the displacement field

$$\mathbf{b}_{ij} = \tfrac{1}{2}\tilde{x}_i^2 \mathbf{e}_j - \tilde{x}_i \tilde{x}_j \mathbf{e}_i \tag{32}$$

satisfies $\partial_i u_j = \tilde{x}_i$ and $\partial_j u_i = -\tilde{x}_i$. Consequently, the Cauchy shear strain vanishes identically:

$$\varepsilon_{ij}(\mathbf{b}_{ij}) = \tfrac{1}{2}(\partial_j u_i + \partial_i u_j) = \tfrac{1}{2}(-\tilde{x}_i + \tilde{x}_i) \equiv 0. \tag{33}$$

The six ordered pairs (x, y) , (x, z) , (y, x) , (y, z) , (z, x) , (z, y) thus define six independent shear-free flexural modes.

- (iii) **Shear family** ($d_{\text{shear}} = 10$ degrees of freedom): Comprises three uniform transverse shears, six planar quadratic transverse shear gradients, and one triaxial cross-gradient shear field $\mathbf{w}$:

$$\begin{aligned} \mathbf{s}_{xy,1} &= (\tilde{y}, \tilde{x}, 0)^\top, \\ \mathbf{s}_{xy,2} &= (\tilde{x}\tilde{y}, \tfrac{1}{2}\tilde{x}^2, 0)^\top, \\ \mathbf{s}_{yx,2} &= (\tfrac{1}{2}\tilde{y}^2, \tilde{y}\tilde{x}, 0)^\top, \\ \mathbf{s}_{yz,1} &= (0, \tilde{z}, \tilde{y})^\top, \\ \mathbf{s}_{yz,2} &= (0, \tilde{y}\tilde{z}, \tfrac{1}{2}\tilde{y}^2)^\top, \\ \mathbf{s}_{zy,2} &= (0, \tfrac{1}{2}\tilde{z}^2, \tilde{z}\tilde{y})^\top, \\ \mathbf{s}_{zx,1} &= (\tilde{z}, 0, \tilde{x})^\top, \\ \mathbf{s}_{zx,2} &= (\tfrac{1}{2}\tilde{z}^2, 0, \tilde{z}\tilde{x})^\top, \\ \mathbf{s}_{xz,2} &= (\tilde{x}\tilde{z}, 0, \tfrac{1}{2}\tilde{x}^2)^\top, \\ \mathbf{w} &= (\tilde{y}\tilde{z}, \tilde{x}\tilde{z}, \tilde{x}\tilde{y})^\top = \nabla(\tilde{x}\tilde{y}\tilde{z}). \end{aligned} \tag{34}$$

Unlike bending, these modes produce non-zero shear strains $\varepsilon_{ij} \neq 0$. The triaxial mode $\mathbf{w}$ is irrotational ($\nabla \times \mathbf{w} = \mathbf{0}$) and divergence-free ($\nabla \cdot \mathbf{w} = 0$), completing the chosen basis of the 24-dimensional nonrigid polynomial quotient; the complement is not unique.

- (iv) **Torsion family** ($d_{\text{torsion}} = 2$ degrees of freedom): Saint-Venant torsional twist gradients around the three Cartesian axes:

$$\mathbf{t}_x = (0, -\tilde{x}\tilde{z}, \tilde{x}\tilde{y})^\top, \quad \mathbf{t}_y = (\tilde{y}\tilde{z}, 0, -\tilde{x}\tilde{y})^\top, \quad \mathbf{t}_z = (-\tilde{y}\tilde{z}, \tilde{x}\tilde{z}, 0)^\top. \tag{35}$$


Each torsion field exhibits identically zero normal strain ($\varepsilon_{xx} = \varepsilon_{yy} = \varepsilon_{zz} \equiv 0$). Crucially, these three vector fields satisfy the identity:

$$\mathbf{t}_x + \mathbf{t}_y + \mathbf{t}_z \equiv \mathbf{0}. \tag{36}$$


Hence, this selected torsion-like polynomial family has algebraic dimension 2, spanned by $\{\mathbf{t}_x, \mathbf{t}_y\}$.

Representative fields in the 3D polynomial dictionary


A Axial · Normal extension




B Bending · Shear-free flexure



C Shear · Transverse shear



D Torsion · Twist gradient



Ghost: undeformed reference · Colour: normalized displacement magnitude · Each field scaled to 50 mm for display

Fig. 3 Representative fields of the 3D polynomial deformation dictionary. One example per family is evaluated on the reference pelvis: (A) axial extension, $\mathbf{a}_{x,1}$; (B) shear-free bending, $\mathbf{b}_{xy}$; (C) uniform transverse shear, $\mathbf{s}_{xy,1}$; and (D) torsion-like twist, $\mathbf{t}_x$. Each card shows anterior and cranial views. Grey denotes the undeformed geometry; colour indicates displacement magnitude relative to the maximum within that field. The formulas are the raw polynomial dictionary members in dimensionless coordinates, and the rendered fields have their volume-weighted infinitesimal rigid component removed. Each rendered field is independently scaled to 50 mm maximum displacement for visibility; these are illustrative basis fields, not computed FE eigenmodes or predicted load responses.

Figure 3 shows one representative field from each family on the same reference anatomy. The visualisations display the polynomial fields after the rigid-motion projection used in the decomposition; they do not depict separate eigenmodes or unique physical mechanisms.

To assemble the discrete kinematic design matrix, the basis fields in each family $m \in \{\text{axial, bending, shear, torsion}\}$ are sampled at the N_q quadrature points, weighted by $\sqrt{\bar{w}_p}$, projected orthogonal to $\mathbf{Q}_{\text{rigid}}$ via $\mathbf{P}_{\text{nonrigid}}$, and orthonormalised via reduced singular value decomposition (SVD):

$$\begin{aligned} \mathbf{P}_{\text{nonrigid}} \mathbf{A}_m &= \mathbf{Q}_m \Sigma_m \mathbf{V}_m^\top, \\ \mathbf{Q}_m &\in \mathbb{R}^{3N_q \times d_m}, \quad \mathbf{Q}_m^\top \mathbf{Q}_m = \mathbf{I}_{d_m}. \end{aligned} \quad (37)$$

The concatenated design matrix $\mathbf{A} = [\mathbf{Q}_{\text{axial}}, \mathbf{Q}_{\text{bending}}, \mathbf{Q}_{\text{shear}}, \mathbf{Q}_{\text{torsion}}] \in \mathbb{R}^{3N_q \times 24}$ spans the complete 24-dimensional canonical nonrigid deformation subspace.

2.8.5 Numerical conditioning, family non-orthogonality, and the Gram matrix

It is important to note that the four canonical deformation families exhibit intrinsic non-orthogonality and are not mutually orthogonal even on an idealized symmetric domain. For example, on the symmetric unit cube $[-1, 1]^3$ (volume $V = 8$), the unnormalized continuum volume integral between the bending mode $\mathbf{b} = (-xy, \frac{1}{2}x^2, 0)^\top$ and the shear mode $\mathbf{s} = (xy, \frac{1}{2}x^2, 0)^\top$ evaluates analytically to $\int_{[-1,1]^3} \mathbf{b} \cdot \mathbf{s} dV = -22/45 \approx -0.4889$, corresponding to a volume-normalized inner product $\langle \mathbf{b}, \mathbf{s} \rangle_V = \frac{1}{V} \int_{[-1,1]^3} \mathbf{b} \cdot \mathbf{s} dV = -11/180 \approx -0.0611$ (angular cosine $\cos \theta = -11/29 \approx -0.3793$). Both example fields have the same nonzero mean translation $(0, 1/6, 0)^\top$. After subtracting it, their volume-normalised inner product is $-4/45$ and their cosine is $-2/3$. Thus non-orthogonality also holds after the rigid-motion projection used here. This does not mean the two families have a nonzero vector in their intersection.

On the real, geometrically asymmetric pelvic ring, this intrinsic non-orthogonality combines with anatomical geometry. The kinematic design matrix $\mathbf{A}$ has 2-norm condition number $\kappa(\mathbf{A}) \approx 4.58$, and the associated Gram matrix $\mathbf{G} = \mathbf{A}^\top \mathbf{A} \in \mathbb{R}^{24 \times 24}$ has condition number:

$$\kappa(\mathbf{G}) = \frac{\sigma_{\max}(\mathbf{G})}{\sigma_{\min}(\mathbf{G})} = \kappa(\mathbf{A})^2 \approx 20.98. \quad (38)$$

A Gram condition number of ~ 21 indicates that the sampled design columns were linearly independent to numerical precision at the reported geometry; it does not establish unique physical attribution to the families.

2.8.6 Cooperative game-theoretic Shapley value norm allocation

Because the four deformation subspaces exhibit intrinsic non-orthogonality (shared explained norm) and cannot be mutually orthogonal, projecting onto them sequentially (e.g., Gram-Schmidt orthogonalisation) would introduce an arbitrary, order-dependent bias: the first chosen family would absorb all shared variance, artificially starving subsequent families. To obtain an order-independent allocation conditional on this chosen dictionary, we formulate the problem using cooperative game theory.

Let $M = \{\text{axial, bending, shear, torsion}\}$ denote the set of four pattern players ($|M| = 4$). There are $2^4 = 16$ possible coalitions $S \subseteq M$. For any coalition S , we construct the concatenated family design matrix $\mathbf{A}_S \in \mathbb{R}^{3N_q \times \sum_{m \in S} d_m}$ and define the characteristic function $v(S)$ as the fraction of nonrigid squared displacement norm explained by the joint subspace:

$$v(S) = R^2(S) = \frac{\mathbf{u}_{\text{nonrigid}}^\top \mathbf{P}_S \mathbf{u}_{\text{nonrigid}}}{\|\mathbf{u}_{\text{nonrigid}}\|_V^2}, \quad (39)$$

$$\mathbf{P}_S = \mathbf{A}_S (\mathbf{A}_S^\top \mathbf{A}_S)^+ \mathbf{A}_S^\top,$$

with $v(\emptyset) = 0$, where $(\cdot)^+$ denotes the Moore-Penrose pseudoinverse. The Shapley value f_m is uniquely specified by the following axioms for this chosen coalition value function and set of families:

1. **Efficiency:** The sum of allocated values equals the value of the grand coalition:

$$\sum_{m \in M} f_m = v(M) = R^2(M) = 1 - f_{\text{residual}}. \quad (40)$$

2. **Symmetry:** If two families $m, k \in M$ contribute identically to every coalition ($v(S \cup \{m\}) = v(S \cup \{k\})$ for all $S \subseteq M \setminus \{m, k\}$), then $f_m = f_k$.
3. **Dummy player:** If a family m adds zero marginal variance to every coalition ($v(S \cup \{m\}) = v(S)$ for all $S \subseteq M \setminus \{m\}$), then $f_m = 0$.
4. **Additivity:** For any two characteristic value functions v and w defined on the same set of players, the values allocated under the combined sum game $(v + w)(S) = v(S) + w(S)$ satisfy $\phi_m(v + w) = \phi_m(v) + \phi_m(w)$.

The explicit Shapley allocation formula evaluates the marginal contribution of family m averaged over all coalition orders:

$$f_m = \sum_{S \subseteq M \setminus \{m\}} \frac{|S|!(|M| - |S| - 1)!}{|M|!} [v(S \cup \{m\}) - v(S)]. \quad (41)$$

The unexplained shape change is retained as an explicit residual:

$$f_{\text{residual}} = 1 - v(M) = 1 - R^2(\text{axial} \cup \text{bending} \cup \text{shear} \cup \text{torsion}). \quad (42)$$

Because subspace projection operators are monotonic ($V_S \subseteq V_{S \cup \{m\}} \implies v(S \cup \{m\}) \geq v(S)$), every marginal contribution is non-negative, which guarantees $f_m \geq 0$ for all m . Exact closure holds identically:

$$f_{\text{axial}} + f_{\text{bending}} + f_{\text{shear}} + f_{\text{torsion}} + f_{\text{residual}} \equiv 1.0000 \quad (100.0\%). \quad (43)$$

Furthermore, the fractions are strictly scale-invariant ($\mathbf{u} \mapsto c\mathbf{u}$ for $c \neq 0$) and sign-invariant ($\mathbf{u} \mapsto -\mathbf{u}$). For comparative diagnostics, we also compute the standalone coefficient of determination for each isolated family:

$$R_m^2 = v(\{m\}) = \frac{\mathbf{u}_{\text{nonrigid}}^\top \mathbf{P}_m \mathbf{u}_{\text{nonrigid}}}{\|\mathbf{u}_{\text{nonrigid}}\|_V^2}. \quad (44)$$

Due to the intrinsic geometric coupling between families ($\kappa(\mathbf{G}) \approx 20.98$), the sum $\sum_{m \in M} R_m^2$ may moderately exceed the total explained fraction $R^2(M)$; the Shapley values allocate this overlap in an order-independent way conditional on the selected dictionary and anatomical axes.

2.9 Load routing and modal strain-energy fractions

For each load case we solved the static problem $K_s \mathbf{u}_s = \mathbf{b}$ on Ω_{ref} , projected the result onto the M_s -orthonormal eigenbasis Φ ,

$$q = \Phi^\top M_s \mathbf{u}_s, \quad E_i = \frac{1}{2} \lambda_i q_i^2, \quad f_i = \frac{E_i}{\sum_{j=1}^{N_{\text{ev}}} E_j}, \quad (45)$$

and summarised each pathway's contribution by its retained modal strain-energy fraction f_i evaluated across the $N_{\text{ev}} = 15$ computed modes. Block throughput for a clustered regime C is $f_C = \sum_{i \in C} f_i$, and the dominant mode set N_{80} of a load case is the smallest subset of modes (selected in descending order of mean energy contribution) whose cumulative mean energy fractions reach at least 0.80. For cross-load comparison we also report cosine similarity of full 15-mode mean profiles and Jaccard overlap of dominant mode sets.

2.10 First-order eigenvalue sensitivities and analytical uncertainty propagation

To propagate material-parameter uncertainty to the spectrum without repeated finite-element solves, we use classical first-order sensitivities of the generalised eigenproblem (10): for a parameter p_j that affects only the stiffness,

$$\frac{\partial \lambda_m}{\partial p_j} = \frac{\phi_m^\top (\partial K_s / \partial p_j) \phi_m}{\phi_m^\top M_s \phi_m}. \quad (46)$$

For ligament parameters, derivatives of local 3×3 blocks are $\partial W / \partial k_{\text{seg}} = \mathbf{e} \mathbf{e}^\top$ and $\partial W_g / \partial T_{\text{seg}} = (I - \mathbf{e} \mathbf{e}^\top) / L$. They enter the signed 6×6 segment assembly, with factor $1/N_{\text{seg}}$ for differentiation with respect to a bundle parameter. For cartilage moduli, $\partial K_s / \partial E_{\text{region}}$ is the element-wise elastic stiffness integrated over the corresponding joint domain, scaled by the baseline modulus. At source points on the high-density branch of (4), $\partial E / \partial \alpha = \rho_{\text{ash}}^\beta$ and $\partial E / \partial \beta = \alpha \rho_{\text{ash}}^\beta \ln \rho_{\text{ash}}$; the derivatives vanish on the constant branch. We interpolate each source derivative with the same radial-basis-function operator used for the modulus, then integrate it against the modal strain-energy density. This differentiates the implemented map, in which the nonlinear density law precedes interpolation. A finite-difference check at 1669 sampled cells in one subject agreed to relative L^2 errors below 4×10^{-11} for both parameters. Storing the dimensionless Jacobians

$$J_j^{(m)} = \frac{p_j}{\lambda_m} \frac{\partial \lambda_m}{\partial p_j} \quad (47)$$

once per subject lets us compute first-order log-space variance propagation as $\text{Var}(\log \lambda_m) \approx \mathbf{J}_m^\top \Sigma_{\log p} \mathbf{J}_m$ for an assumed covariance $\Sigma_{\log p} = \text{Cov}(\log \mathbf{p})$ on the material parameters. The implemented scenarios use log-space standard deviations (0.25, 0.20, 0.10) for global, type, and side effects of ligament stiffness; (0.35, 0.25, 0.12) for ligament pretension; and (0.35, 0.25, 0.10) for cartilage modulus. The global ligament stiffness–pretension correlation is 0.50. Bone-law parameters α and β have log-space standard deviations 0.30 and 0.20 with correlation -0.40 . These are assumed scenarios, not parameters estimated from the cohort. Eigenvalue CoVs use the lognormal conversion $\sqrt{\exp(v) - 1}$ from propagated log-variance v . For adjacent eigenvalues λ_i, λ_{i+1} , the variance of the gap $\Delta \lambda = \lambda_{i+1} - \lambda_i$ is computed from the combined sensitivity vector $\mathbf{g}_{i,i+1} = \lambda_{i+1} \mathbf{J}_{i+1} - \lambda_i \mathbf{J}_i$ as

$$\text{Var}(\lambda_{i+1} - \lambda_i) \approx \mathbf{g}_{i,i+1}^\top \Sigma_{\log p} \mathbf{g}_{i,i+1} \approx \text{Var}(\lambda_{i+1}) + \text{Var}(\lambda_i) - 2 \text{Cov}(\lambda_i, \lambda_{i+1}), \quad (48)$$

accounting for the substantial positive covariance induced by shared tissue parameters. The gap signal-to-noise ratio compares the nominal gap with its propagated standard deviation (Section 3.9). It measures local susceptibility of modal ordering under the specified parameter covariance. These gap diagnostics do not directly propagate uncertainty in eigenvectors or anatomical capacities and therefore do not establish subspace robustness under large parameter changes. Sex-stratified summaries and combined uncertainty results are reported in Sections 3.6 and 3.10.

2.11 Decomposition of variability sources and robustness analyses

To separate shape and material contributions to spectral variability, we compared the full cohort against a shape-only cohort (χ_s , fixed reference bone modulus) and a material-only cohort (identity deformation, subject-specific bone modulus). Robustness was assessed by (i) random permutation of subject order, (ii) a 500-replicate bootstrap of 222 draws with replacement at the fixed threshold and a separate 2,000-replicate bootstrap of 278 draws

with replacement and threshold re-estimation, and (iii) morphology-axis permutation null tests in which a one-dimensional anatomical proxy μ (e.g. AP inlet diameter) was shuffled 1000 times to assess whether swap-rate localisation along μ exceeded chance (Table 8). Post-hoc allometric regressions of $\log \lambda_k$ on isotropic body-size scale s (with instrumental variables from volume, surface, and mass) and Shapley/LMG variance decomposition into scale, sex, shape, and material contributions are reported in Section 3.10.

2.12 Minimal analytical model for the near-degenerate regime

The cluster/subspace framework admits a compact analytical interpretation via a two-degree-of-freedom model. Let q_1, q_2 denote two generalised rotational coordinates and define a stiffness matrix

$$K_2 = \begin{bmatrix} k_1 & c \\ c & k_2 \end{bmatrix}, \quad \Delta = k_1 - k_2, \quad k_1, k_2 > 0, \quad k_1 k_2 > c^2, \quad (49)$$

with detuning Δ and coupling c . The eigenvalues are

$$\lambda_{\pm} = \frac{k_1 + k_2}{2} \pm \sqrt{\left(\frac{\Delta}{2}\right)^2 + c^2}, \quad \tan(2\theta) = \frac{2c}{\Delta}, \quad (50)$$

and the minimum separation $\lambda_+ - \lambda_- = 2|c|$ is attained at $\Delta = 0$, for constant nonzero c as Δ is varied. With a continuous branch convention, $2\theta = \text{atan2}(2c, \Delta)$; at $\Delta = 0$ the angle is $\pi/4$ for $c > 0$, up to eigenvector sign. When $c = 0$, an exact crossing occurs. The operational criterion

$$|\Delta(\mu)| \leq \tau |c(\mu)| \quad \text{for some morphology parameter } \mu, \quad \tau \in [1.5, 2], \quad (51)$$

separates near-degenerate/mixing from stable-ordering regimes. Adding a third coordinate q_3 with stiffness s and coupling vector $\mathbf{d} = [d_1, d_2]^\top$, dynamic reduction of the 3×3 eigenproblem $(K - \lambda I)\mathbf{q} = \mathbf{0}$ gives the exact non-linear reduced eigenproblem

$$\left(K_2 - \frac{\mathbf{d}\mathbf{d}^\top}{s - \lambda}\right)\mathbf{q}_{12} = \lambda\mathbf{q}_{12}. \quad (52)$$

In the low-frequency limit ($\lambda \ll s$), static condensation yields the linear effective stiffness

$$K_{\text{eff}} = K_2 - \frac{\mathbf{d}\mathbf{d}^\top}{s} = \begin{bmatrix} k_1 - d_1^2/s & c - d_1 d_2/s \\ c - d_1 d_2/s & k_2 - d_2^2/s \end{bmatrix}, \quad (53)$$

showing how coupling to an additional coordinate shifts effective detuning and coupling. Depending on the parameters, the shift can move the system toward or away from avoided crossing. The assumption $\lambda \ll s$ excludes using this condensation for an arbitrarily soft hidden coordinate. In all stochastic numerical simulations (Section 3.11), the full 3×3 eigenproblem is solved directly without static or dynamic truncation. In the n -DOF finite-element setting, the detected clusters correspond to $\{q_1, q_2\}$, the canonical basis (ψ_1, ψ_2) to the rotated eigenvectors, and gap_{in} to $(\lambda_+ - \lambda_-)/\lambda_-$. A stochastic population instantiation of this model, evaluated in Section 3.11, reproduces the label-subspace dissociation, the standing-vs-task routing contrast, and the effect of hidden-pathway variability observed in the full pelvic cohort. The stochastic model used in the figure is specified in Section 2.12.1. Static condensation is only a limiting argument; it is not used for the reported stochastic eigensolves.

2.12.1 Stochastic implementation

This specification defines the stochastic implementation evaluated in Section 3.11. Parameters are dimensionless and illustrative; they were not fitted to the pelvic cohort. The stiffness matrix is

$$K = \begin{bmatrix} 12 + \Delta/2 & c & d_1 \\ c & 12 - \Delta/2 & d_2 \\ d_1 & d_2 & s \end{bmatrix}, \quad M = I. \quad (54)$$

For morphology coordinate μ and nuisance scale η , independent standard-normal draws determine

$$\Delta = 1.6\mu + (0.05 + 0.22\eta)Z_\Delta, \quad (55)$$

$$c = |0.22 + (0.008 + 0.018\eta)Z_c|, \quad (56)$$

$$s = 5.2 \exp[(0.02 + 0.12\eta)Z_s], \quad (57)$$

$$d_1 = 0.28 + 0.42\eta + (0.01 + 0.08\eta)Z_1, \quad (58)$$

$$d_2 = 0.24 + 0.40\eta + (0.01 + 0.08\eta)Z_2. \quad (59)$$

Draws are accepted only when all eigenvalues exceed 10^{-10} and the largest absolute component of the third coordinate occurs in the lowest eigenvector. The low-rank backbone is therefore conditioned by construction, not independently discovered in this example. Up to 200 draws are attempted per subject.

The anatomical observable and task-like load are the unit-normalised vector $(1, 0, 0.2)^\top$; the standing-like load is the unit-normalised vector $(0.10, 0.05, 1)^\top$. For unit-normalised eigenvectors $\mathbf{v}_i$, the observable projection is $(\mathbf{v}_i^\top \mathbf{b})^2$ and the upper-pair score is the sum for $i = 2, 3$. Load fractions are proportional to $(\mathbf{v}_i^\top \mathbf{f})^2 / \lambda_i$. These squared projections differ from the max-nodal anatomical capacity used in the finite-element cohort and serve only as analogous illustrative observables.

The main illustrative cohort contains 240 evenly spaced coordinates in $[-1.10, 1.10]$, perturbed by normal noise of standard deviation 0.035 (random seed 11), sorted, and simulated at $\eta = 0.85$ (seed 19). The nuisance sweep uses nine values in $[0, 1.6]$ and 480 coordinates in $[-0.35, 0.35]$, with seed 31 at each value. Carrier change is the proportion whose larger upper-pair observable projection changes rank relative to $\eta = 0$. Pair projection and task routing are summarised by their medians.

The full 3×3 eigenproblem is solved for every accepted draw. The static-condensation formula in Eq. (53) is a limiting argument valid for $\lambda \ll s$; it is not an approximation used for the upper-pair eigenvalues in this example. The model illustrates rank redistribution under specified parameter variation and does not validate a particular pelvic tissue mechanism.

2.13 Statistical analysis

All statistical analyses were executed using SciPy, statsmodels, and custom Python routines. Continuous metrics across non-normal cohorts were evaluated using non-parametric methods. Two-group comparisons (e.g., non-swapped vs. swapped cluster members, or male vs. female distributions) were conducted using two-sided Mann–Whitney U tests. Effect sizes for Mann–Whitney tests are reported as rank-biserial correlations $r_{rb} = 1 - 2U_1/(n_1n_2)$, where U_1 is computed for the first listed group. With this orientation, positive values indicate larger observations in the second group; the statistic equals Cliff's delta for second versus first group. Paired differences across loading conditions or mode pairs were assessed via two-sided Wilcoxon signed-rank tests. Bivariate associations between continuous anatomical dimensions (such as AP inlet diameter or isotropic scale) and modal metrics (energy fractions, spectral gaps) were quantified using Spearman's rank correlation coefficient ρ_s . The four pre-specified rank-versus-capacity comparisons in each label-exchange table were adjusted with the Benjamini–Hochberg procedure. Other exploratory test families were not uniformly adjusted; their raw p -values should not be treated as confirmatory discoveries. Confidence intervals for median differences and median ratios in the label-exchange comparisons were obtained by percentile bootstrap resampling (5,000 independent within-group resamples; random seed 20260918), using the 2.5th and 97.5th percentiles. Mann–Whitney probabilities concern distributional differences and should not be interpreted as tests specifically of the median difference. Cohort-level cluster stability was evaluated by repeating the detection pipeline across 500 random subcohort draws of 80% sample size performed with replacement ($m = 222$), yielding 95% bootstrap stability intervals. Separate prevalence intervals used 2,000 replicates of 278 draws with replacement and threshold re-estimation (seed 20260918). These are resampling summaries, not independent biological validation.

Differences in subspace orientation across the cohort were quantified using the Grassmann distance $d_G(Q_{\text{ref}}, Q_s) = \|\boldsymbol{\theta}\|_2$. Importantly, d_G measures the intrinsic geometric distance between the linear subspaces $\text{span}(Q_{\text{ref}})$ and $\text{span}(Q_s)$ in the Grassmannian manifold $\text{Gr}(r, n)$, which is strictly invariant to arbitrary unitary rotations of the internal basis within either subspace. Variation in d_G cannot be caused by internal basis re-indexing, although sex-group differences in distance to one reference do not by themselves establish a direction of subspace shift or rule out reference effects.

2.14 Software and computational implementation

Finite-element assembly, eigensolves, and postprocessing were implemented in Python 3.11 with DOLFINx/FEniCSx (≥ 0.10), PETSc, and SLEPc. Reported figures, tables, and numerical values were produced by the study analysis scripts; a public code archive has not yet been established. Results are stored as compressed Zarr arrays for reproducible downstream analysis. The formulation and stochastic implementation are specified above; a public source-code release remains to be prepared.

An OpenAI language model assisted the restructuring and drafting of this manuscript and cover letter from the existing manuscript and analysis outputs. The revised numerical analyses were executed with the documented project scripts and checked against the archived finite-element fields; the authors retain responsibility for the scientific content.

3 Results

3.1 Reference deformation atlas and numerical validation

This analysis provides the reference-mode atlas, synthetic validation checks, and full deformation profiles. The mathematical formulation, 14-point degree-4 quadrature, rigid-motion projection, polynomial dictionary, and Shapley allocation are given in the Methods. Table 1 tabulates the complete 15-mode quantitative profile across all four canonical deformation families, standalone R^2 values, and residual fractions. Table 2 reports the synthetic benchmark validations across idealized symmetric and irregular pelvic domains.

3.1.1 Synthetic benchmark and backwards-identification validation

To verify the accuracy, robustness, and backwards-identifiability of the global deformation decomposition, we executed synthetic validation experiments on both an idealized symmetric domain (cube) and the realistic reference pelvis mesh (Table 2).

On the symmetric domain, pure synthetic fields belonging to the axial, shear, and torsion families are identified with 100.0% fidelity, zero cross-talk, and exactly 0.0% residual. For pure Euler-Bernoulli bending ($\mathbf{u} = (-xy, \frac{1}{2}x^2, 0)$), the displacement field consists of an axial cross-term $-xy$ and a transverse parabolic deflection $\frac{1}{2}x^2$. Because the quadratic shear mode $\mathbf{s}_{xy,2} = (xy, \frac{1}{2}x^2, 0)$ shares the identical transverse parabolic deflection $\frac{1}{2}x^2$, the pure flexural field lies in the joint span $V_{\text{bending}} \oplus V_{\text{shear}}$. The equal-weight $15 \times 15 \times 15$ grid benchmark allocates 77.5% to bending and 22.5% to shear, with residual rounding to zero. These numbers describe the sampled grid and the chosen Shapley game, not a unique physical decomposition of the field.

On the actual reference pelvis mesh, pure canonical polynomial fields are recovered with exactly 0.0% residual (100.0% fitted fraction):

- Pure axial stretch ($x\mathbf{e}_x$): 85.3% axial share, 0.0% residual (100.0% fitted).
- Pure triaxial extension ($x\mathbf{e}_x - \frac{1}{2}y\mathbf{e}_y - \frac{1}{2}z\mathbf{e}_z$): 84.8% axial share, 0.0% residual (100.0% fitted).
- Pure Euler-Bernoulli bending: 74.1% bending share, 0.0% residual (100.0% fitted).
- Pure transverse shear: 82.8% shear share, 0.0% residual (100.0% fitted).
- Pure torsion-like polynomial field: 87.1% torsion share, 0.0% residual (100.0% fitted).

When subjected to pure infinitesimal rigid translations or pure infinitesimal rotations, the nonrigid fraction is detected as $f_{\text{nonrigid}} < 10^{-27}$, correctly triggering the rigid-motion filter and showing negligible contamination in these numerical checks. Finally, adding arbitrary rigid-body translations and rotations to an axial deformation field yields identical Shapley shares (85.3% axial, 7.2% bending, 7.5% shear, 0.0% torsion, 0.0% residual) to within machine precision ($< 10^{-10}$), demonstrating invariance to addition of infinitesimal rigid displacement fields, rather than invariance to arbitrary rotation of anatomical axes.

3.1.2 Reference mode deformation profiles and biomechanical interpretation

Table 1 compiles the quantitative whole-pelvis 3D deformation decomposition for all fifteen reference eigenmodes. The results resolve several key biomechanical features of the pelvic ring:

- **Modes 1 and 2 (low-rank backbone):** For Mode 1, the nonrigid deformation accounts for 3.3% of the total displacement norm (the remainder reflecting whole-pelvis rocking on the sacroiliac/pubis supports). Its nonrigid shape change is predominantly out-of-plane torsional twist (22.7%) and transverse shear (17.7%), with a total fitted fraction of 42.7% (residual 57.3%). For Mode 2, nonrigid shape change (5.5% of total norm) is a coupled sagittal ring flexure and shear (23.6% bending, 23.7% shear, 20.8% axial, 68.6% fitted, residual 31.4%). The selected 3D displacement dictionary explains more than two-thirds of the nonrigid squared displacement norm; the tested one-axis dictionary gave a lower descriptive fit.
- **Modes 4 and 5 (dominant flexural pathways):** Modes 4 and 5 exhibit strong bending dominance, explaining 43.6% and 47.4% of their nonrigid norm respectively (with standalone bending fits R^2_{bending} reaching 47.6% and 62.4%, and total explained fractions of 74.4% and 86.3%). These modes capture lateral and anteroposterior flaring of the iliac wings.
- **The inlet-related 9–10 pair:** Mode 9 is the most highly fitted mode in the entire spectrum (91.8% explained, residual only 8.2%). Its deformation is overwhelmingly axial (60.6% Shapley share; standalone $R^2_{\text{axial}} = 67.7\%$), consistent with Cartesian normal-extension content. Its relation to inlet opening is evaluated separately with the anatomical diameter functional; the polynomial allocation does not measure circumferential strain. In contrast, its near-degenerate partner Mode 10 (74.9% fitted) is dominated by shear (35.2%) and torsion (34.3%), with minimal axial contribution (4.3%; standalone $R^2_{\text{torsion}} = 30.3\%$, $R^2_{\text{shear}} = 32.7\%$, residual 25.1%).



Fig. 4 Whole-pelvis deformation patterns of reference modes. Anterior (left) and cranial (right) views show displacement magnitude from purple to yellow, scaled to a 50 mm maximum; grey geometry is undeformed. Each bar partitions the whole-pelvis nonrigid squared shape norm into fitted axial tension/compression, bending, transverse shear, torsion, and residual. Infinitesimal rigid-body translation and rotation are projected out upfront; rigid rotation is not a reported deformation category. The decomposition operates in full 3D Cartesian space across the entire pelvic ring without imposing a 1D reference axis. Overlapping pattern subspaces are allocated using Shapley values. “Fitted” reports the percentage of nonrigid shape change explained by the four canonical 3D families. All fractions use the complete continuum without bone-wise subdivision and describe kinematic shape change rather than elastic-energy shares.

These reference-mode descriptions illustrate distinct kinematic patterns within the pair. Their rank-specific interpretation should not be transferred unchanged across subjects with label exchange.

Table 1 Reference mode whole-pelvis 3D deformation decomposition.

Mode	Nonrigid	Fitted	Shapley norm partition (%)					Standalone R^2 (%)			
			Axial	Bending	Shear	Torsion	Residual	Axial	Bending	Shear	Torsion
1	3.3	42.7	0.9	1.5	17.7	22.7	57.3	0.4	2.2	15.4	19.0
2	5.5	68.6	20.8	23.6	23.7	0.5	31.4	15.4	16.1	16.9	0.8
3	29.0	76.7	2.8	4.7	41.7	27.5	23.3	5.1	5.1	52.7	37.4
4	24.5	74.4	13.5	43.6	15.8	1.5	25.6	24.6	47.6	14.9	2.2
5	73.0	86.3	4.3	47.4	24.2	10.4	13.7	5.4	62.4	38.7	13.7
6	71.9	82.3	3.1	26.9	24.9	27.3	17.7	2.2	33.0	37.2	35.2
7	69.5	72.1	16.9	7.3	44.3	3.5	27.9	29.4	4.6	55.8	4.1
8	57.0	72.1	22.1	20.3	29.5	0.2	27.9	32.7	32.5	41.3	0.4
9	92.1	91.8	60.6	18.4	12.2	0.5	8.2	67.7	31.6	14.9	0.3
10	95.7	74.9	4.3	1.1	35.2	34.3	25.1	6.2	1.5	32.7	30.3
11	71.4	43.1	5.4	3.9	21.1	12.8	56.9	2.8	0.4	9.5	12.2
12	94.2	62.3	9.6	22.4	27.4	3.0	37.7	13.1	12.7	20.3	2.4
13	98.3	48.4	9.2	19.9	19.1	0.2	51.6	8.3	27.0	25.7	0.0
14	92.0	53.4	1.2	15.9	33.4	3.0	46.6	0.9	5.6	24.9	4.7
15	98.7	55.6	18.4	23.5	13.7	0.0	44.4	16.1	28.2	13.7	0.0

Table 2 Synthetic benchmark and backwards-identification validation.

Domain	Synthetic deformation field	Axial (%)	Bending (%)	Shear (%)	Torsion (%)	Resid. (%)	Outcome
Symmetric cube	Pure axial stretch ($x\mathbf{e}_x$)	100.0	0.0	0.0	0.0	0.0	Exact closure (0.0% residual)
Symmetric cube	Pure bending ($-xy\mathbf{e}_x + \frac{1}{2}x^2\mathbf{e}_y$)	0.0	77.5	22.5	0.0	0.0	Exact closure (0.0% residual)
Symmetric cube	Pure shear ($y\mathbf{e}_x + x\mathbf{e}_y$)	0.0	0.0	100.0	0.0	0.0	Exact closure (0.0% residual)
Symmetric cube	Pure torsion ($yz\mathbf{e}_x - xy\mathbf{e}_z$)	0.0	0.0	0.0	100.0	0.0	Exact closure (0.0% residual)
Pelvis mesh	Pure axial stretch ($x\mathbf{e}_x$)	85.3	7.2	7.5	0.0	0.0	Exact closure (0.0% residual)
Pelvis mesh	Pure triaxial ($x\mathbf{e}_x - \frac{1}{2}y\mathbf{e}_y - \frac{1}{2}z\mathbf{e}_z$)	84.8	11.1	4.1	0.0	0.0	Exact closure (0.0% residual)
Pelvis mesh	Pure bending ($-xy\mathbf{e}_x + \frac{1}{2}x^2\mathbf{e}_y$)	17.1	74.1	8.8	0.0	0.0	Exact closure (0.0% residual)
Pelvis mesh	Pure shear ($y\mathbf{e}_x + x\mathbf{e}_y$)	13.7	1.8	82.8	1.7	0.0	Exact closure (0.0% residual)
Pelvis mesh	Pure torsion ($yz\mathbf{e}_x - xy\mathbf{e}_z$)	5.5	0.5	7.0	87.1	0.0	Exact closure (0.0% residual)
Pelvis mesh	Infinitesimal rigid translation	0.0	0.0	0.0	0.0	0.0	Rigid: rejected ($f_{\text{nonrigid}} < 10^{-20}$)
Pelvis mesh	Infinitesimal rigid rotation	0.0	0.0	0.0	0.0	0.0	Rigid: rejected ($f_{\text{nonrigid}} < 10^{-20}$)
Pelvis mesh	Axial stretch + rigid perturbation	85.3	7.2	7.5	0.0	0.0	Exact closure (0.0% residual)

3.2 Population variability separates an order-locked backbone from recurrent clustered regimes

The cohort of $N=278$ subjects resolves the spectrum into two regimes: an order-locked low-rank backbone and recurrent higher-rank clustered blocks (Fig. 5, Table 3). The reference deformation atlas is provided in Fig. 4. Modes 1–3 are rarely reordered across the cohort (swap rate $< 0.4\%$), whereas higher ranks contain small consecutive gaps and frequent label exchange. Using a data-driven cluster threshold $\varepsilon_{\text{in}} \approx 0.131$ (25th percentile of pooled relative consecutive gaps; see Methods), we detect nine near-degenerate rank clusters above a 3% prevalence floor. The most prevalent cluster is rank modes 9–10 (54.7% of subjects), followed by rank modes 12–15 (41.7%) and rank modes 5–6 (33.8%). No eigenvalue pair met the operational numerical multiplicity criterion ($\text{gap}_{\text{in}} \leq 10^{-6}$). No obvious spatial symmetry was identified that would enforce repeated eigenvalues, although accidental degeneracy is not mathematically excluded.

Anatomical variability does not destabilise the spectrum uniformly. Label fragility is concentrated in a small number of compact families, while low-rank order is preserved. For the dominant 9–10 and 12–15 clusters, median relative internal gaps are small enough that modest perturbations can reverse local ordering. The 9–10 block has a larger median external separation than internal gap, whereas the 12–15 regime is left-separated but right-truncated by the eigensolve; the 9–10 pair can therefore be assessed as an interior subspace, whereas conclusions about isolation of 12–15 remain limited. The result identifies spectral regions in which rank-based comparison is more or less reliable under the present model assumptions.

Permutation statistics reinforce this interpretation. Modes 12 and 13 exhibit the highest swap rates, and mode 15 also shows a sizeable unpaired fraction under strict cross-subject pairing, whereas the backbone is essentially fixed in its ordering. For cross-subject modal comparison, the consequence is concrete: while one-to-one

Table 3 Recurrent clustered rank blocks above the 3% prevalence floor. External separation is unavailable (—) when the block reaches the upper limit of 15 computed modes; such blocks are not established to be isolated on both sides

Cluster	Dim	n	Prev.	Med. $\text{gap}_{\text{in},\text{min}}$	Med. sep_{out}	Type
Rank modes 9–10	2	152	54.7%	0.08114	0.4926	near-degenerate
Rank modes 12–15	4	116	41.7%	0.03153	—	near-degenerate
Rank modes 5–6	2	94	33.8%	0.08737	0.2345	near-degenerate
Rank modes 12–13	2	78	28.1%	0.03839	0.1941	near-degenerate
Rank modes 14–15	2	70	25.2%	0.05116	—	near-degenerate
Rank modes 12–14	3	45	16.2%	0.04109	0.1855	near-degenerate
Rank modes 4–5	2	35	12.6%	0.09521	0.2185	near-degenerate
Rank modes 13–15	3	29	10.4%	0.04465	—	near-degenerate
Rank modes 7–8	2	21	7.6%	0.1051	0.3676	near-degenerate

modal comparison is well-conditioned for the low-rank backbone, higher-rank comparisons must accommodate mode mixing and label exchange through subspace representations.

A variability decomposition into shape-only and material-only datasets (Fig. 5C) shows that the shape-only dataset has the larger coefficient of variation for modes 1–8, whereas bone-density heterogeneity contributes with increasing relative weight in modes 9–15. Shape changes alter *how* the spatial density heterogeneity enters the stiffness operator: shape dominates lower modal variance while material heterogeneity contributes increasingly at higher ranks. The shape-only and material-only coefficients of variation are not additive variance shares, nor do they isolate interactions or establish a causal mechanism for cluster formation.

These prevalence estimates are not driven by a few outliers or by the order in which subjects are processed. The 500 bootstrap draws of 222 subjects with replacement gave 95% percentile ranges of 48.2–61.7% for cluster 9–10 and 35.6–48.6% for cluster 12–15 (Table 9). A separate 2,000-draw full-cohort bootstrap with threshold re-estimation gave 49.6–59.4% and 36.7–46.0%, respectively. Random permutation of subject order leaves cluster prevalence and swap statistics unchanged to machine precision. The central spectral hierarchy is therefore a cohort-level feature rather than a post-processing artefact. Additional cluster-mapping, swap, and robustness diagnostics are provided in Section 3.3 and Section 3.6.




Fig. 5 Population variability partitions the stiffness spectrum into an order-locked backbone and clustered higher ranks. (A) Consecutive relative spectral gaps $\text{gap}_{\text{in}}(i) = (\lambda_{i+1} - \lambda_i)/\lambda_i$ across the cohort ($N=278$); the dashed line marks the data-driven cluster threshold $\epsilon_{\text{in}} \approx 0.131$ (25th percentile). (B) Subject fraction exhibiting label exchange (blue) and unpaired pairing failures (grey; $\text{MAC}_{M_0} < 0.05$) per reference label. (C) Eigenvalue variability (coefficient of variation, CoV) across the combined cohort (blue), shape-only cohort (orange), and material-only cohort (green).

3.3 Spectral cluster architecture

This analysis expands the bookkeeping behind the recurrent clustered regimes, including reference-mode occupancy, swap frequencies, and morphology-axis diagnostics. The cohort-level cluster list is given in Table 3. Table 4 records the mapping between recurrent rank clusters and their dominant reference-mode identities. Table 5 reports per-mode swap rates, Table 6 gives the adjacent rank-pair diagnostics, Table 7 summarises the exchange-regime classification, and Table 8 reports the morphology-axis permutation null test used to localise swap behaviour.

Table 4 Rank-cluster to reference-mode mapping.

rank_cluster	dim	n_subjects	prevalence	paired_all_rate	ref_modes_mode	ref_modes_mode_rate	ref_modes_top3
Rank modes 9–10	2	152	54.7%	1	9–10	1	9–10 (100%)
Rank modes 12–15	4	116	41.7%	0.7414	12–15	1	12–15 (100%)
Rank modes 5–6	2	94	33.8%	1	5–6	0.8298	5–6 (83%); 4,6 (17%)
Rank modes 12–13	2	78	28.1%	0.9872	13–14	0.4805	13–14 (48%); 12–13 (44%); 12,15 (6%)
Rank modes 14–15	2	70	25.2%	0.8571	12,15	0.45	12,15 (45%); 14–15 (38%); 13–14 (15%)
Rank modes 12–14	3	45	16.2%	0.9333	12–14	0.8571	12–14 (86%); 12,13,15 (10%); 13–15 (5%)
Rank modes 4–5	2	35	12.6%	1	4–5	1	4–5 (100%)
Rank modes 13–15	3	29	10.4%	0.8966	12–14	0.5	12–14 (50%); 13–15 (31%); 12,14,15 (15%)
Rank modes 7–8	2	21	7.6%	1	7–8	1	7–8 (100%)

Table 5 Per-mode swap rates.

Mode	Swap rate	Fail rate	Non-id. rate	<i>n</i> swapped	<i>n</i> failed
1	0	0	0	0	0
2	0	0	0	0	0
3	0.003597	0	0.003597	1	0
4	0.2698	0	0.2698	75	0
5	0.3777	0	0.3777	105	0
6	0.1835	0	0.1835	51	0
7	0.01079	0	0.01079	3	0
8	0.003597	0	0.003597	1	0
9	0.2698	0	0.2698	75	0
10	0.2698	0	0.2698	75	0
11	0.003597	0	0.003597	1	0
12	0.6151	0.007194	0.6223	171	2
13	0.6295	0	0.6295	175	0
14	0.5072	0.01439	0.5216	141	4
15	0.3813	0.1978	0.5791	106	55

Table 6 Adjacent rank-pair swap diagnostics.

cluster	cluster_key	rank_pair	ref_pair_mode	swap_rate	invalid_rate	match_rate	median_gap_in	near_deg_rate	true_deg_rate	peak_rate	active_frac	cv_swap	verdict
Rank modes 9–10	modes_9_10	9 $\rightleftharpoons$ 10	9 $\rightleftharpoons$ 10	0.3750	0.0000	1.0000	0.1206	0.5468	0.0000	1.0000	0.2050	0.6157	veering
Rank modes 12–15	modes_12_15	12 $\rightleftharpoons$ 13	12 $\rightleftharpoons$ 13	0.4154	0.0345	0.5603	0.0516	0.8597	0.0000	1.0000	0.3094	0.7588	veering
Rank modes 12–15	modes_12_15	13 $\rightleftharpoons$ 14	13 $\rightleftharpoons$ 14	0.1429	0.0603	0.3017	0.0802	0.7014	0.0000	1.0000	0.0324	2.0521	veering
Rank modes 12–15	modes_12_15	14 $\rightleftharpoons$ 15	14 $\rightleftharpoons$ 15	0.1818	0.2241	0.3793	0.0721	0.7734	0.0000	1.0000	0.0935	1.6757	veering
Rank modes 5–6	modes_5_6	5 $\rightleftharpoons$ 6	5 $\rightleftharpoons$ 6	0.1282	0.0000	0.8298	0.1661	0.3633	0.0000	1.0000	0.0288	1.6080	veering
Rank modes 12–13	modes_12_13	12 $\rightleftharpoons$ 13	13 $\rightleftharpoons$ 14	0.0811	0.0128	0.4744	0.0516	0.8597	0.0000	1.0000	0.0432	2.9996	veering
Rank modes 14–15	modes_14_15	14 $\rightleftharpoons$ 15	12 $\rightleftharpoons$ 15	0.6296	0.1429	0.3857	0.0721	0.7734	0.0000	1.0000	0.3921	0.8933	veering

Table 7 Heuristic classification of observed permutations. Exchange labels do not establish a continuous eigenvalue-veering trajectory.

cluster	cluster_key	dim	n_subjects	prevalence	median_gap_in	gap_type	swap_verdict	final_verdict	ref_modes_mode	ref_modes_mode_rate	paired_all_rate	near_swap_rate	fail_rate	within_block_rate	transposition_rate	cyclic_2_rate	self_rate	n_veering_gains
Rank modes 9–10	modes_9_10	2	152	54.7%	0.0811	near-degenerate	VEERING (local character exchange)	VEERING (local character exchange)	9–10	1.0000	1.0000	0.3750	0.0000	1.0000	0.3750	0.0000	1	1
Rank modes 12–15	modes_12_15	4	116	41.7%	0.0315	near-degenerate	VEERING (local character exchange)	MIXING (multi-mode; cycle 2 common)	12–15	1.0000	0.7414	0.5431	0.2066	0.7414	0.3021	0.3072	3	3
Rank modes 5–6	modes_5_6	2	94	33.8%	0.0874	near-degenerate	VEERING (local character exchange)	VEERING (local character exchange)	5–6	0.8298	1.0000	0.1064	0.0000	0.8298	0.1282	0.0000	1	1
Rank modes 12–13	modes_12_13	2	78	28.1%	0.0384	near-degenerate	VEERING (local character exchange)	VEERING (local character exchange)	13–14	0.4805	0.9872	0.0385	0.0128	0.4744	0.0811	0.0000	1	1
Rank modes 14–15	modes_14_15	2	70	25.2%	0.0312	near-degenerate	VEERING (local character exchange)	VEERING (local character exchange)	12,15	0.4500	0.8571	0.1429	0.1429	0.3857	0.3704	0.0000	1	1

Table 8 Morphology-axis permutation null test.

Cluster	Rank pair	Peak _{obs}	<i>p</i> (peak)	Active _{obs}	<i>p</i> (active)	CV _{obs}	<i>p</i> (CV)	Significant
Rank modes 9–10	9 $\rightleftharpoons$ 10	1	0.2547	0.2162	0.8202	0.6042	0.02697	NO
Rank modes 12–15	12 $\rightleftharpoons$ 13	1	0.4496	0.283	0.7223	0.5553	0.4046	NO
Rank modes 12–15	13 $\rightleftharpoons$ 14	0.25	0.8891	0.5	0.3437	1	0.4336	NO
Rank modes 12–15	14 $\rightleftharpoons$ 15	0.5	0.7443	0.09302	0.8492	0.9727	0.4256	NO
Rank modes 5–6	5 $\rightleftharpoons$ 6	0.5	0.9001	0.1267	0.4535	1.048	0.7263	NO
Rank modes 12–13	12 $\rightleftharpoons$ 13	0.25	0.8192	0.4375	0.2438	1.146	0.5455	NO
Rank modes 14–15	14 $\rightleftharpoons$ 15	0.75	0.6563	1	0.5504	0	1	NO

3.4 Anatomical coupling distinguishes recurrent subspaces

Once individual labels become unstable, the next question is whether the corresponding small recurrent subspaces remain mechanically distinct. Figure 6 shows that several do, when interpreted using invariant-subspace descriptors rather than individual eigenvector labels. Two basis-invariant outputs were evaluated. First, Grassmann distance from a reference subject quantifies how much a subspace rotates across the cohort. Second, basis-invariant anatomical capacities for anatomical diameter functionals (AP and ML inlet; interspinous midpelvic and intertuberos outlet) provide functionally interpretable coordinates inside each clustered block (Methods).

The rank 9–10 subspace has median AP-inlet capacity 0.54 mm/mm and ML-inlet capacity 1.05 mm/mm. The 5–6 block also accommodates outlet deformation, with interspinous capacity 0.95 mm/mm, bituberos capacity 1.48 mm/mm, and modelled AP-outlet capacity 0.50 mm/mm. These anatomical responses are not exclusive to




Fig. 6 Basis-invariant anatomical capacities across recurrent clustered modal subspaces. Basis-invariant anatomical capacities (maximum anatomical diameter response per 1 mm maximum nodal displacement) for the five most prevalent rank blocks across the cohort ($N=278$): (A) anteroposterior (AP) inlet, (B) mediolateral (ML) inlet, (C) interspinous midpelvic diameter, and (D) bituberous outlet. Boxes indicate median and interquartile range (IQR); whiskers extend to $1.5 \times \text{IQR}$.

one block: the corresponding 9–10 capacities are 1.20, 1.55, and 1.28 mm/mm, respectively. Thus, describing 5–6 as outlet-related does not imply that it has the largest outlet capacity. Its relevance also depends on selective recruitment under the applied loads.

The four-dimensional 12–15 block illustrates a third regime. Its capacities were recomputed using the convex constrained optimisation described in Methods. It has high prevalence, small internal gaps, and a richer pattern of within-block mixing than a simple two-mode exchange. Its anatomical capacity structure is more diffuse than in 9–10 or 5–6, limiting interpretation in terms of one anatomical observable. Furthermore, because spectral eigensolves were truncated at $N_{\text{ev}} = 15$, external separation of this upper block on the right cannot be determined without higher eigenvalues; we therefore treat 12–15 as a diffuse higher-rank regime bounded by modal truncation. Not every recurrent cluster corresponds to one clean anatomical function. The distinction between a recurrent spectral block and a specific anatomical response therefore remains necessary.

Subspace orientation in the ambient displacement space (quantified by the Grassmann distance d_G , which is strictly invariant to internal basis rotations) also varies systematically with sex. The largest rank-biserial contrast in distance from the selected reference was observed for 9–10. Distance to one reference does not alone establish the direction or independence of a sex-associated shift. This qualifies how clustered behaviour is distributed across the population without altering the central point: recurrent clustered subspaces retain distinct mechanical meanings across sexes despite modest differences in their ambient subspace orientation.

3.5 Subspace capacity varies less than individual-rank coupling

The empirical question is whether anatomical capacity differs less between exchange groups when evaluated over a subspace than at an individual rank. This hypothesis was tested for the most prevalent cluster, rank modes 9–10 (Fig. 7).




Fig. 7 Inlet coupling redistributes between rank labels within the stable 9–10 subspace. Anteroposterior (AP) inlet coupling for rank mode 9 (A), rank mode 10 (B), and the basis-invariant single-functional AP capacity of the two-dimensional 9–10 subspace (C), stratified by 9 $\rightleftharpoons$ 10 label-exchange status. Points represent individual cluster-member subjects ($n=152$; no-exchange $n=95$, exchange $n=57$; male in blue, $n=90$; female in yellow, $n=62$). Horizontal lines indicate medians; boxes indicate interquartile range; whiskers indicate $1.5 \times \text{IQR}$. Coupling is expressed as diameter change per millimetre of maximum nodal displacement.

Among the 152 cluster members, 95 showed no exchange and 57 showed exchange. Median AP-inlet coupling at rank 9 decreased from 0.454 to 0.139 mm/mm (difference -0.315; 95% confidence interval [-0.360, -0.262]; $p = 2.7 \times 10^{-23}$), whereas rank-10 coupling increased from 0.163 to 0.436 mm/mm ($p = 1.7 \times 10^{-22}$). Thus, the anatomical response redistributed between neighbouring ranks.

The corresponding subspace capacity changed from 0.551 to 0.525 mm/mm (difference -0.026; 95% confidence interval [-0.084, 0.015]; rank-biserial correlation -0.21). This is approximately a 5% median decrease, compared with approximately 69% at rank 9. The difference is smaller, but capacity is not identical between groups: the Mann–Whitney test gives $p = 0.030$ (Benjamini–Hochberg-adjusted $q = 0.040$). The test assesses distributions, whereas the bootstrap interval concerns the median difference; these summaries need not give identical significance conclusions.

Across all 278 subjects, irrespective of cluster membership, subspace capacity differed by -0.006 mm/mm (95% confidence interval [-0.048, 0.019]; $p = 0.106$), while rank-level coupling again redistributed strongly. This supports relative stability of the subspace descriptor, without establishing equivalence. Full estimates, effect sizes, joint-SVD comparisons, and adjusted probabilities are provided in Tables 11–12.

Load routing reinforces the same conclusion (Fig. 11). The 9–10 exchange accompanies a large change in rank-level coupling and a smaller change in subspace capacity. Subspace rotation and the internal 9–10 gap vary across subjects, AP coupling shifts modestly with swap status, and the throughput of the 9–10 block under LAB₁–LAB₃ depends on morphology in a load-specific way. For LAB₁, block throughput decreases with larger AP inlet diameter ($\rho_s = -0.48$, $p = 4.3 \times 10^{-17}$), whereas LAB₂ and LAB₃ show weaker and differently signed trends. These patterns support the interpretation that inlet-related mechanics remains associated with the 9–10 subspace even as anatomical variability redistributes its basis representation and modulates its load-specific amplitude.

3.6 Cohort robustness and sex effects

The analysis reports exploratory sex-group comparisons where they help describe variation of the recurrent clustered subspaces, especially the 9–10 inlet-related pair. This analysis provides the broader sex-stratified summary and the bootstrap stability evidence that the dominant clusters are stable cohort-level features rather than products of a few outlying subjects. Table 9 gives the bootstrap stability estimates (500 replicates of 80% cohort size drawn with replacement, $m = 222$) for the dominant clusters and gap statistics. Fig. 8 summarises the sex-stratified spectral structure, and Table 10 reports the accompanying sex-stratified statistical tests for the spectral and subspace metrics.

Table 9 Cohort bootstrap stability under random 80% fractional resamples ($m = 222, 500$ replicates with replacement).

Metric	Target	Observed	Boot. mean	Boot. std	95
prevalence	Rank modes 9–10	0.547	0.545	0.035	[0.482, 0.617]
prevalence	Rank modes 12–15	0.417	0.417	0.033	[0.356, 0.486]
prevalence	Rank modes 5–6	0.338	0.337	0.031	[0.279, 0.401]
prevalence	Rank modes 12–13	0.281	0.280	0.029	[0.225, 0.340]
prevalence	Rank modes 14–15	0.252	0.252	0.028	[0.198, 0.311]
swap_rate	Mode 4	0.270	0.271	0.030	[0.216, 0.329]
swap_rate	Mode 5	0.378	0.378	0.033	[0.315, 0.444]
swap_rate	Mode 6	0.183	0.182	0.026	[0.135, 0.239]
swap_rate	Mode 7	0.011	0.011	0.007	[0.000, 0.027]
swap_rate	Mode 9	0.270	0.271	0.028	[0.218, 0.331]
swap_rate	Mode 10	0.270	0.271	0.028	[0.218, 0.331]
swap_rate	Mode 12	0.594	0.595	0.033	[0.523, 0.658]
swap_rate	Mode 13	0.622	0.622	0.033	[0.554, 0.689]
swap_rate	Mode 14	0.468	0.470	0.033	[0.405, 0.545]
swap_rate	Mode 15	0.453	0.455	0.035	[0.390, 0.523]
median_gap	Pair 1–2	1.609	1.603	0.038	[1.521, 1.665]
median_gap	Pair 2–3	1.183	1.189	0.019	[1.159, 1.221]
median_gap	Pair 3–4	0.484	0.483	0.020	[0.448, 0.526]
median_gap	Pair 4–5	0.271	0.271	0.011	[0.247, 0.295]
median_gap	Pair 5–6	0.166	0.166	0.008	[0.148, 0.183]
median_gap	Pair 6–7	0.526	0.527	0.013	[0.502, 0.556]
median_gap	Pair 7–8	0.287	0.287	0.011	[0.268, 0.313]
median_gap	Pair 8–9	0.913	0.911	0.014	[0.882, 0.938]
median_gap	Pair 9–10	0.121	0.121	0.007	[0.109, 0.135]
median_gap	Pair 10–11	1.144	1.147	0.021	[1.109, 1.189]
median_gap	Pair 11–12	0.705	0.705	0.014	[0.672, 0.732]
median_gap	Pair 12–13	0.052	0.052	0.004	[0.046, 0.060]
median_gap	Pair 13–14	0.080	0.080	0.005	[0.072, 0.091]
median_gap	Pair 14–15	0.072	0.071	0.006	[0.062, 0.082]

Table 10 Exploratory sex-stratified tests. Probabilities are unadjusted. Subspace distances use cluster-member subjects; other rows use the full cohort. The permutation-rate row reports a Fisher test, whose statistic is an odds ratio rather than Mann–Whitney U .

Metric	Male med.	Female med.	U/OR	p	$r_{rb}/\log OR$
gap_in (9–10)	0.104	0.1534	6384	1.489e-06	0.335
gap_in (12–13)	0.04925	0.0552	9182	0.5321	0.04354
gap_in (13–14)	0.09191	0.07643	1.063e+04	0.1233	-0.1073
gap_in (14–15)	0.06002	0.08448	7631	0.003216	0.2051
gap_in (5–6)	0.1669	0.166	9677	0.9088	-0.008021
gap_in (4–5)	0.2627	0.2768	9299	0.6529	0.03135
gap_in (7–8)	0.3604	0.245	1.412e+04	1.311e-11	-0.471
swap_count_per_subject	4	3	1.352e+04	2.824e-09	-0.4079
permutation_rate	0.8984	0.9	0.9829	1	-0.01724
d.G (modes_9_10)	0.2605	0.3996	377	1.5e-19	0.8649
d.G (modes_12_15)	1.455	0.7866	2848	6.629e-11	-0.7059
d.G (modes_5_6)	0.8472	0.572	1793	1.766e-07	-0.6263
d.G (modes_12_13)	0.5299	0.3729	1120	0.0001815	-0.4973
d.G (modes_14_15)	0.5736	0.9485	270	6.922e-05	0.5559

Table 11 Cluster members: rank and subspace comparisons. No exchange: $n = 95$; exchange: $n = 57$. Couplings and differences are in mm/mm. Difference is exchange minus no exchange; intervals use 5,000 percentile bootstrap resamples. p is the two-sided Mann–Whitney probability, q its Benjamini–Hochberg adjustment across the four metrics in this table, and r_{rb} the rank-biserial effect. The joint-SVD score is distinct from single-functional capacity.

Metric	No exchange	Exchange	Difference [95% CI]	r_{rb}	p	q
Rank 9	0.454	0.139	-0.315 [-0.360, -0.262]	-0.97	2.66×10^{-23}	1.07×10^{-22}
Rank 10	0.163	0.436	0.273 [0.225, 0.335]	0.95	1.70×10^{-22}	3.41×10^{-22}
$\kappa_{AP, \infty}$	0.551	0.525	-0.026 [-0.084, 0.015]	-0.21	0.030	0.040
Joint-SVD AP	0.077	0.075	-0.002 [-0.020, 0.012]	-0.02	0.840	0.840

Table 12 Full cohort, irrespective of cluster membership: rank and subspace comparisons. No exchange: $n = 204$; exchange: $n = 74$. Couplings and differences are in mm/mm. Difference is exchange minus no exchange; intervals use 5,000 percentile bootstrap resamples. p is the two-sided Mann–Whitney probability, q its Benjamini–Hochberg adjustment across the four metrics in this table, and r_{rb} the rank-biserial effect. The joint-SVD score is distinct from single-functional capacity.

Metric	No exchange	Exchange	Difference [95% CI]	r_{rb}	p	q
Rank 9	0.458	0.139	-0.319 [-0.355, -0.284]	-0.95	9.73×10^{-34}	1.95×10^{-33}
Rank 10	0.140	0.441	0.301 [0.259, 0.354]	0.96	4.26×10^{-34}	1.70×10^{-33}
$\kappa_{AP, \infty}$	0.532	0.526	-0.006 [-0.048, 0.019]	-0.13	0.106	0.142
Joint-SVD AP	0.081	0.076	-0.005 [-0.016, 0.011]	0.02	0.780	0.780




Fig. 8 Sex-stratified spectral summary

3.7 Standing and proxy loads distinguish backbone-dominated routing from higher-rank recruitment

Load routing reflects the same separation between conserved baseline support and selective higher-rank recruitment (Fig. 9, Table 13). Standing is dominated by the gap-protected low-rank backbone. Symmetric two-leg standing (SP2leg) is concentrated in mode 2 alone, which carries about 84.5% of the retained modal strain energy; single-leg standing (SP1leg) distributes energy across the full rotational triad (modes 1–3), reaching a top-3 cumulative share of about 92.0%. Under these two prescribed standing loads, a few low-rank modes account for most of the retained modal energy.




Fig. 9 Standing and proxy childbirth loads recruit distinct modal regimes. (A) Heatmap of mean retained modal strain-energy share across the 15 lowest solver ranks for standing (SP2leg, SP1leg) and proxy childbirth loads (LAB₁–LAB₃). (B) Partition of retained modal strain energy across functional blocks: low-rank backbone (modes 1–3; blue), mid-rank regime (modes 4–8; orange), inlet cluster (modes 9–10; green), and higher-rank reserve (modes 11–15; purple). Standing concentrates in the backbone, whereas proxy loads recruit higher ranks in a task-specific pattern.

The proxy load family recruits higher-rank modes. LAB₁, bilateral mediolateral internal ring expansion (± 400 N outward), preferentially engages the rank 9–10 inlet-related subspace, which carries a mean energy share of about 0.51 and is led by the inlet-opening mode. LAB₂, bilateral ischial-tuberosity distraction, reroutes energy into the mid-rank block at modes 4–8, where the 5–6 outlet-related pair becomes especially important. LAB₃, anteroposterior outlet distraction, is hybrid, retaining sizeable low-rank contributions while also recruiting mid-rank and 9–10 pathways. Different loading geometries recruit different parts of the higher-rank spectrum, while continuing to use the backbone as common support.

Among the recurrent clustered regimes, 9–10 remains the clearest case because it combines high cohort prevalence, clear inlet-related interpretation, direct label-versus-subspace evidence (Fig. 7), and substantial recruitment under LAB₁. The 5–6 pair remains an outlet-related secondary contrast under LAB₂, whereas 12–15 is common but behaves, under the present load family, as a mixed higher-rank reserve rather than a single dominant pathway. Primary cohort routing is evaluated on the total static response. Archived incremental outputs for 18 subjects showed unchanged top-mode identity across all five loads, but individual 9–10 block fractions shifted by up to 4.24 percentage points and dominant-set size changed in four LAB₁ cases (Section 2.3). Additional load-specificity and sex-stratified summaries are reported in Section 3.6 and Section 3.8.

These are linear comparative proxy loads, not simulations of labour. They do not model fetal contact, tissue remodelling, hormone-dependent softening, or nonlinear late-stage birth mechanics. Their value is comparative: they reveal which deformation patterns are preferentially recruited under different loading geometries, not whether the model can predict clinical childbirth outcomes.

Table 13 Dominant mode sets for the standing and proxy loading cases. N_{80} denotes the smallest set of modes required to reach 80% cumulative mean energy fraction.

Load	Top mode	Top mode share (%)	Top-3 modes	Top-3 share (%)	N_{80}	Modes in 80% set	Cumulative share at N_{80} (%)
SP2leg	2 (ML rotation)	84.49	2,4,5	93.79	1	2	84.49
SP1leg	1 (CC rotation)	39.89	1,2,3	92.04	3	1,2,3	92.04
LAB ₁	9 (inlet opening)	33.49	9,10,8	66.12	5	9,10,8,4,3	81.48
LAB ₂	5	25.15	5,6,4	74.77	4	5,6,4,9	85.8
LAB ₃	2 (ML rotation)	45.81	2,5,4	68.09	5	2,5,4,9,6	85.8

3.8 Load-routing specificity

The study uses standing and proxy childbirth loads to show the contrast between a robust low-rank backbone and selectively recruited higher-rank subspaces. This analysis provides comprehensive load-specificity diagnostics, including cluster-level function-locking, dominant-mode sets, and pairwise overlap of full routing profiles.

Fig. 10 visualises the load–function locking map across the prevalent clusters. Table 14 summarises the phase-specific coupling synthesis, Table 13 summarises the dominant mode sets, and Table 15 reports the pairwise overlap metrics for the full load-routing profiles.




Fig. 10 Load-function coupling heatmap across prevalent clusters

Table 14 Exploratory load–coupling associations among cluster members. AP/ML and BIS/BIT scores are earlier paired-SVD descriptors; Outlet AP uses an L^2 -optimal direction followed by max-nodal normalisation. These differ from the max-nodal capacity in the main comparison. FLI selects the strongest Spearman association among five axes and is descriptive. The energy column is the within-cluster median.

Cluster	Regime	Best-predictor coupling axis for FLI (AP/ML inlet, BIS/BIT outlet, or Outlet AP scalar)	FLI ρ	Load case	$\bar{f}_C$	$r_{rb}(d_G)$
Rank modes 9–10	Veering	BIS outlet	0.5153	LAB ₁	0.5291	0.8649
Rank modes 9–10	Veering	Outlet AP scalar	0.339	LAB ₂	0.1578	0.8649
Rank modes 9–10	Veering	Outlet AP scalar	0.4354	LAB ₃	0.1407	0.8649
Rank modes 12–15	Mixing (4D)	AP inlet	0.3442	LAB ₁	0.07568	-0.7059
Rank modes 12–15	Mixing (4D)	BIT outlet	0.8222	LAB ₂	0.01016	-0.7059
Rank modes 12–15	Mixing (4D)	ML inlet	-0.6724	LAB ₃	0.06857	-0.7059
Rank modes 5–6	Weak exchange	ML inlet	0.4989	LAB ₁	0.02502	-0.6263
Rank modes 5–6	Weak exchange	BIS outlet	0.7738	LAB ₂	0.5917	-0.6263
Rank modes 5–6	Weak exchange	BIS outlet	0.6931	LAB ₃	0.217	-0.6263
Rank modes 12–13	Weak exchange	AP inlet	0.4367	LAB ₁	0.06863	-0.4973
Rank modes 12–13	Weak exchange	BIS outlet	0.7197	LAB ₂	0.0009939	-0.4973
Rank modes 12–13	Weak exchange	Outlet AP scalar	0.7232	LAB ₃	0.01162	-0.4973
Rank modes 14–15	Mixing	ML inlet	0.3535	LAB ₁	0.02817	0.5559
Rank modes 14–15	Mixing	BIS outlet	0.8028	LAB ₂	0.01237	0.5559
Rank modes 14–15	Mixing	BIS outlet	-0.3534	LAB ₃	0.008386	0.5559

Table 15 Pairwise overlap of load-routing profiles.

Load A	Load B	Jaccard overlap (80% sets)	Cosine similarity (full profile)	Shared modes (80% sets)
SP2leg	SP1leg	0.3333	0.6668	2
SP2leg	LAB ₁	0	0.1202	
SP2leg	LAB ₂	0	0.1343	
SP2leg	LAB ₃	0.2	0.9367	2
SP1leg	LAB ₁	0.1429	0.1233	3
SP1leg	LAB ₂	0	0.1051	
SP1leg	LAB ₃	0.1429	0.6326	2
LAB ₁	LAB ₂	0.2857	0.48	4,9
LAB ₁	LAB ₃	0.25	0.3613	4,9
LAB ₂	LAB ₃	0.8	0.4456	4,5,6,9




Fig. 11 Morphology and loading modulate recruitment of the 9–10 modal block. (A–C) Modal strain-energy fraction carried by the 9–10 block as a function of AP inlet diameter for LAB₁ (bilateral internal ring expansion), LAB₂ (ischial distraction), and LAB₃ (outlet distraction). Blue dots and dashed lines indicate males; yellow dots and dashed lines indicate females. (D) Basis-invariant single-functional AP-inlet capacity of the 9–10 subspace stratified jointly by sex and label-exchange status (M/no, F/no, M/yes, F/yes). Boxes show median and IQR; whiskers extend to $1.5 \times \text{IQR}$.

3.9 First-order material sensitivity differentiates spectral regions

First-order propagation with derivatives of the implemented material map predicts substantial variation in absolute eigenvalues under the assumed covariance; shared-parameter covariance reduces the calculated uncertainty of adjacent differences (Fig. 12; Table 17). Mean gap signal-to-noise ratios exceed 2 for ranks 1–2 and 2–3 in both sexes, whereas the 9–10 mean is 0.997 in females and 0.947 in males. Under this assumed local perturbation model, the internal 9–10 ordering is therefore more susceptible than the low-rank gaps. The calculation concerns eigenvalues and gaps; it does not independently verify preservation of subspace orientation or anatomical capacity under large material changes.




Fig. 12 Analytical propagation of material-parameter uncertainty to eigenvalues and spectral gaps. (A) Total coefficient of variation (CoV, %) of eigenvalues $\lambda_1, \dots, \lambda_{15}$ under hierarchical material-parameter covariance, stratified by sex (blue: male; yellow: female). (B) Relative gap signal-to-noise ratio (spectral gap divided by propagated standard deviation) between adjacent solver ranks, incorporating shared-parameter covariance $\text{Cov}(\lambda_i, \lambda_{i+1})$. The shaded region highlights the 9–10 rank transition. Triangles mark 10th-percentile values across subjects.

3.10 Model-boundary and sensitivity analyses

These analyses bound the main claims rather than extend them. Allometry describes associations between the archived spectra and isotropic size; it is not a recovery of information proven to have been removed by geometric normalisation. The corrected material-uncertainty calculation illustrates how assumed soft-tissue and density-law scatter could affect eigenvalues and spectral gaps to first order. Fig. 13 shows the post-hoc allometric recovery analysis, and Table 16 summarises the corresponding scaling exponents and variance decomposition. Fig. 14 presents the threshold sensitivity, reference sensitivity, isotropic scale vs solver gap, and 15-mode displacement reconstruction error (tabulated in Table 18), alongside Table 17 which tabulates the combined first-order uncertainty propagation used to support the robustness discussion in the article.




Fig. 13 Allometry dashboard

Table 16 Allometric scaling exponents and variance decomposition per eigenvalue mode. β_{scale} is the IV-2SLS log-log coefficient of eigenvalue on isotropic scale, with HC-robust 95 % CIs and FDR-corrected p -values. $\Delta\lambda$ per IQR gives the percent eigenvalue change for an interquartile range shift in log-scale. The last four columns show LMG (Shapley) shares of explained variance divided by full-model R^2 ; they sum to approximately 100% and are not percentages of total outcome variance. These are descriptive associations, not identified causal effects.

Mode	β_{scale}	95% CI	p_{FDR}	$\Delta\lambda$ IQR (%)	R^2 scale (%)	R^2 shape (%)	R^2 material (%)	R^2 sex (%)
1	-1.38	[-1.80, -0.96]	4.7e-10	-10.1	17.4	77.5	1.1	3.9
2	-1.83	[-2.14, -1.53]	8.4e-26	-13.1	24.2	60.6	5.8	9.4
3	-1.68	[-1.99, -1.36]	2.0e-21	-12.1	26.2	68.0	3.1	2.7
4	-2.39	[-2.78, -2.00]	7.7e-27	-16.8	33.9	50.8	10.9	4.4
5	-2.07	[-2.49, -1.64]	1.1e-18	-14.7	30.4	56.8	3.6	9.3
6	-2.14	[-2.48, -1.80]	3.4e-28	-15.1	40.7	48.8	3.5	7.1
7	-2.35	[-2.60, -2.10]	5.7e-49	-16.5	54.8	33.1	4.3	7.8
8	-2.33	[-2.63, -2.02]	2.4e-37	-16.4	40.2	45.3	10.4	4.1
9	-1.89	[-2.16, -1.61]	1.9e-31	-13.5	28.5	43.8	18.3	9.4
10	-2.14	[-2.41, -1.87]	7.7e-39	-15.1	41.5	37.4	17.4	3.7
11	-2.09	[-2.43, -1.75]	6.3e-27	-14.8	27.0	39.1	25.4	8.5
12	-1.38	[-1.81, -0.95]	9.4e-10	-10.1	13.5	58.1	27.1	1.3
13	-1.54	[-2.02, -1.07]	8.2e-10	-11.2	14.1	47.0	37.4	1.5
14	-1.64	[-2.07, -1.21]	1.5e-12	-11.8	18.5	52.2	27.4	1.9
15	-1.39	[-1.77, -1.00]	1.8e-11	-10.1	37.4	43.8	14.8	3.9




Fig. 14 Robustness and reconstruction dashboard

Table 17 Population-level material uncertainty propagation stratified by sex. CoV is the total coefficient of variation (ligament + cartilage + bone). Lig/Cart/Bone columns show each group's fraction of total variance. The bone contribution uses derivatives of the implemented density-to-modulus map. Gap signal-to-noise ratios are displayed in Fig. 12, not in this table.

Mode	Female ($N = 150$)				Male ($N = 128$)			
	CoV (%)	Lig (%)	Cart (%)	Bone (%)	CoV (%)	Lig (%)	Cart (%)	Bone (%)
1	46.0	5	94	0	46.5	7	93	0
2	46.5	4	93	2	46.7	5	93	2
3	42.5	5	94	1	43.3	6	93	1
4	46.0	6	90	4	45.1	8	90	2
5	44.2	8	89	3	44.2	9	89	2
6	44.9	9	89	2	44.7	10	88	2
7	45.9	16	83	1	45.5	15	84	1
8	47.3	7	90	3	46.9	6	91	3
9	45.2	4	88	7	43.9	4	90	5
10	43.7	2	90	8	44.1	5	90	6
11	44.4	1	87	11	43.7	3	89	9
12	45.6	1	86	14	45.3	2	85	13
13	46.1	1	84	15	46.0	1	83	16
14	45.2	1	86	13	45.1	2	88	10
15	45.3	2	88	9	45.0	3	89	8

Table 18 Relative displacement reconstruction error for 15 modes. Errors are relative displacement norms, expressed as fractions rather than percentages. These are displacement diagnostics and do not establish strain-energy or local stress accuracy.

Load	Median relative error	95th percentile	Maximum
SP2leg	0.002	0.003	0.005
SP1leg	0.002	0.003	0.004
LAB1	0.032	0.050	0.066
LAB2	0.035	0.050	0.058
LAB3	0.036	0.056	0.074




Fig. 15 Minimal analytical model illustrates label exchange within a stable subspace. A stochastic cohort of the analytical model with morphology parameter μ and hidden-pathway scale η . **(A)** Spectrum: a separated backbone mode and a nearby eigenvalue pair undergoing near-degeneracy around $\mu \approx 0$. **(B)** Functional projection: coupling shifts between individual rank labels across the transition zone while total coupling within the pair remains high. **(C)** Load routing: standing-like load concentrates in the backbone, whereas task-like load recruits the clustered block with a variable between-label split. **(D)** Effect of hidden-pathway variability scale η : label-exchange rate increases with η , whereas subspace coupling and task routing remain stable. Model derivations and simulation parameters are detailed in Sections 2.12–2.12.1.

3.11 A three-coordinate model illustrates label–subspace dissociation

The constructed analytical example illustrates how a separated low-rank mode and a coupled higher pair can produce contrasting rank and subspace behaviour (Fig. 15). As detuning varies, the dominant projection of the anatomical observable transfers between the upper eigenvectors, whereas their summed projection remains high. Standing-like and task-like loads recruit different spectral regions. Increasing the nuisance scale changes the dominant carrier more strongly than the upper-pair projection or task routing.

The example conditions accepted draws on a retained low-rank backbone. Its observable is a squared projection, rather than the max-nodal capacity used in the pelvic analysis. It therefore illustrates a spectral mechanism without independently establishing either pelvic rank stability or tissue-specific compensation. The cohort permutation test did not detect exchange localisation along AP inlet diameter alone (Table 8; all tested cases $p > 0.05$). A single anatomical coordinate is consequently not supported as an explanation of the observed exchange.

4 Discussion

4.1 The comparative unit depends on spectral separation

The principal finding is that anatomical coupling can be much less sensitive to cross-subject mode exchange when evaluated over a small subspace than at one fixed eigenvalue rank. Within the recurrent 9–10 cluster, median rank-9 AP-inlet coupling decreases from 0.454 to 0.139 mm/mm between non-exchange and exchange

groups, approximately 69%, whereas the corresponding subspace capacity decreases from 0.551 to 0.525 mm/mm, approximately 5%. This difference matters for population biomechanics: interpreting rank 9 alone as a conserved inlet-opening pathway would overstate the change in that block’s kinematic capacity.

The result combines three distinct measurements. Modal assignment identifies changes in correspondence to a reference; principal angles quantify changes in subspace orientation; and anatomical functionals evaluate specific diameter responses. Basis invariance alone guarantees independence from the chosen coordinates within a fixed subspace, not stability across anatomically different subjects. Cross-subject preservation is therefore an empirical question, addressed here by the distribution of anatomical capacities. The observed differences support relative stability, rather than exact conservation or a demonstrated equivalence between exchange groups.

This interpretation applies selectively across the spectrum. Modes 1–3 retain their ordering in almost all subjects, making individual-mode comparison comparatively well conditioned. Higher ranks contain recurrent small-gap blocks. The 9–10 block is the clearest interior example; the 12–15 block remains incompletely characterised because its upper spectral neighbour was not computed. The analysis consequently supports choosing comparison units using both internal gaps and external separation, rather than grouping all higher modes indiscriminately.

4.2 Anatomical capacity and load recruitment answer different questions

Anatomical capacity estimates the largest diameter change within a subspace at a specified displacement normalisation; the two-dimensional estimates were spot-checked against a constrained solver. It does not measure deformation per unit force and does not predict how strongly that subspace is recruited by a particular load. Static modal energy provides the complementary load-dependent quantity. Standing mainly recruits low ranks, whereas internal-ring expansion increases recruitment of the 9–10 block. The 5–6 pair is relevant to ischial distraction despite not having the largest outlet capacity among the compared blocks.

These findings provide a mechanical description of the initial deformation repertoire of a constrained pelvic ring. They may help formulate hypotheses about how load transfer and canal deformation coexist, a longstanding question in pelvic biomechanics (Gruss and Schmitt 2015). They do not demonstrate evolutionary optimisation, physiological compensation, or birth-canal expansion during actual labour. In particular, the mixed-sex cohort and its mean age of approximately 54 years limit direct inference to pregnant or primiparous populations. The minimal analytical example illustrates the spectral mechanism and was constructed to retain the backbone; it is not an independently validated physiological model.

4.3 Implications for population finite-element studies

The present analysis extends the pelvic eigenmetric approach of Henyš et al. (2021) from spectral descriptors toward the correspondence of anatomically meaningful deformation spaces across subjects. Its distinction from load-specific cohort analyses, such as the strain comparisons of Salo et al. (2017), is the separate evaluation of available kinematic capacity and load recruitment. The numerical rationale for following clustered subspaces is established (Ghosh and Ghanem 2012); the evidence added here is the magnitude of the discrepancy between rank-level and subspace-level anatomical comparisons in a pelvic population. This finding does not invalidate earlier spectral descriptors, but identifies a correspondence check needed when assigning anatomical meaning to a fixed rank.

The practical contribution is a sequence of comparisons that could be evaluated in other anatomically variable structures: identify crowded spectral regions, verify their external separation, compare their subspaces in a shared metric, and evaluate anatomically meaningful observables together with load-dependent recruitment. This extends established subject-specific finite-element modelling by addressing ambiguity in the interpretation of population modal comparisons (Fernandez et al. 2020). It does not require a new constitutive formulation to reveal a consequential limitation of rank-based interpretation.

The exchange groups are defined from eigenvectors also used to evaluate anatomical coupling. The rank-level contrast is consequently descriptive, not an independent causal effect of exchange. A maximum over two modes is at least as large as the capacity of either included mode; that inequality alone does not guarantee a smaller difference between groups.

The generalisable component is the analysis strategy. The observed rank partition depends on the pelvic geometry, restraints, constitutive assumptions, and retained spectral range. Applying the approach elsewhere requires new anatomical functionals and a fresh assessment of gaps and truncation. Likewise, low-dimensional representations useful for describing global ring kinematics need not resolve local strains or stresses adequately. The polynomial deformation atlas is a descriptive check, with explicit residuals, rather than a validation of physiological deformation pathways.

4.4 Limitations and validation priorities

The models assume small strains, isotropic bone elasticity, homogeneous joint cartilage, and prescribed ligament properties. The superior sacral restraint and chosen load locations constrain the available deformation repertoire. Subject-specific ligament laxity, pelvic floor support, nonlinear contact, and time-dependent tissue behaviour are not represented. These omissions limit extrapolation to large deformations, altered joint conditions, and childbirth (Vleeming et al. 2012; Chen and Grimm 2021).

The superior sacral restraint uses a fixed penalty parameter ($\gamma = 10^6 \text{ N/mm}^3$); the claimed decade sensitivity sweep could not be verified from retained outputs.

The corrected first-order propagation indicates possible local sensitivity of eigenvalues and gaps under the specified covariance. It does not replace re-solving the full model under broad parameter changes, nor does it verify robustness of eigenvectors and capacities in that setting. The reduced analytical example likewise cannot supply evidence about tissue-specific variability that was not measured in the cohort. Future work should test subspace capacities under explicit soft-tissue perturbations and alternative boundary conditions.

Energy fractions refer to the 15 retained modes. Their sum is unity by normalisation and is not evidence that those modes recover the complete static energy. Displacement reconstruction errors are reported separately, but displacement accuracy does not establish accuracy of local stress or strain. An archived expanded-spectrum check on three subjects reproduced the first 15 eigenvalues with maximum relative discrepancy 3.4×10^{-11} . In these subjects, energy capture by 15 modes relative to the full assembled quadratic energy was 99.3–99.5% for standing and 76.5–94.9% for expansion loads. This does not establish cohort-wide truncation adequacy or the upper separation of the 12–15 block.

Finally, the present evidence establishes computational behaviour within the model family. Experimental measurements of ring displacement under comparable loading would be needed to validate anatomical coupling and recruitment. Robustness to cohort resampling and selected modelling choices strengthens the computational comparison, while leaving that physiological validation open.

5 Conclusions

Across 278 pelvic finite-element models, individual low-rank modes retain stable correspondence, whereas higher-rank anatomical coupling can redistribute between neighbouring eigenvalue ranks. For the recurrent 9–10 block, subspace-level inlet capacity varies substantially less with label-exchange status than single-rank coupling. Comparing compact subspaces therefore helps distinguish changes in modal representation from changes in modelled anatomical deformation capacity. The conclusion concerns population comparisons within a linearised model and motivates targeted experimental and modelling tests.

Statements and Declarations

Funding

This research was supported by the Czech Science Foundation, Grant No. 24-10862S.

Competing interests

The authors declare no competing interests.

Ethics approval and consent to participate

This study used previously published anonymised imaging-derived data and precomputed models; no new imaging or interventions were performed. According to the BoneDat descriptor, the source study was approved by the Ethics Committee of the University Hospital Hradec Kralove (reference 202411IO3P), with waiver of informed consent (Henyš and Kuchař 2025).

Consent for publication

Not applicable; no identifiable individual participant information is presented.

Data availability

Summary results supporting this study are included in this article. The source anatomical resource is described in BoneDat (Henyš and Kuchař 2025); access to underlying clinical imaging is governed by the source study. A public archive of the additional subject-level derived outputs has not yet been established.

Code availability

The study uses custom Python code for model assembly, spectral analysis, and postprocessing. A public archival release of this code has not yet been established.

Author contributions

P.H., M.K., and N.H. conceptualised the study. P.H. and M.K. developed the computational models and performed the simulations. P.H., M.K., and A.M. analysed the data. P.H., M.K., A.M., and N.H. interpreted the results and contributed to manuscript writing.

References

- Roque de Almeida P, Alves Campice de Oliveira I, de Oliveira Campos J, et al (2024) Modeling of the biomechanical behavior and growth of the human uterus during pregnancy. *Journal of Biomechanics* 174:112268. <https://doi.org/10.1016/j.jbiomech.2024.112268>
- Arand C, Wagner D, Richards RG, et al (2019) 3D statistical model of the pelvic ring—a CT-based statistical evaluation of anatomical variation. *Journal of Anatomy* 234(3):376–383. <https://doi.org/10.1111/joa.12928>
- Bergmann G, Deuretzbacher F, Heller M, et al (2001) Hip contact forces and gait patterns from routine activities. *Journal of Biomechanics* 34(7):859–871. [https://doi.org/10.1016/S0021-9290\(01\)00040-9](https://doi.org/10.1016/S0021-9290(01)00040-9)
- Bergmann G, Bender A, Dymke J, et al (2016) Standardized loads acting in hip implants. *PLOS ONE* 11(5):e0155612. <https://doi.org/10.1371/journal.pone.0155612>
- Björck Å, Golub GH (1973) Numerical methods for computing angles between linear subspaces. *Mathematics of Computation* 27(123):579–594. <https://doi.org/10.1090/S0025-5718-1973-0348991-3>
- Bonet J, Wood RD (1997) *Nonlinear Continuum Mechanics for Finite Element Analysis*, 1st edn. Cambridge University Press
- Chen J, Hume PA, Wyatt HE, et al (2026) Development of a pelvic musculoskeletal model based on statistical shape modelling for estimating sacroiliac and pubic joint reaction forces in females. *Biomechanics and Modeling in Mechanobiology* 25:91. <https://doi.org/10.1007/s10237-026-02110-5>
- Chen S, Grimm MJ (2021) Childbirth computational models: Characteristics and applications. *Journal of Biomechanical Engineering* 143(5):050801. <https://doi.org/10.1115/1.4049226>
- Cook DD, Robertson DJ (2016) The generic modeling fallacy: Average biomechanical models often produce non-average results! *Journal of Biomechanics* 49(15):3609–3615. <https://doi.org/10.1016/j.jbiomech.2016.10.004>
- Fernandez J, Dickinson A, Hunter P (2020) Population based approaches to computational musculoskeletal modelling. *Biomechanics and Modeling in Mechanobiology* 19:1165–1168. <https://doi.org/10.1007/s10237-020-01364-x>
- Frost HM (1987) Bone 'mass' and the 'mechanostat': A proposal. *The Anatomical Record* 219(1):1–9. <https://doi.org/10.1002/ar.1092190104>
- Ghosh D, Ghanem R (2012) An invariant subspace-based approach to the random eigenvalue problem of systems with clustered spectrum. *International Journal for Numerical Methods in Engineering* 91(4):378–396. <https://doi.org/10.1002/nme.4276>
- Goode A, Hegedus EJ, Sizer P (2008) Three-dimensional movements of the sacroiliac joint: A systematic review of the literature and assessment of clinical utility. *Journal of Manual & Manipulative Therapy* 16(1):25–38. <https://doi.org/10.1179/106698108790818639>
- Gruss LT, Schmitt D (2015) The evolution of the human pelvis: changing adaptations to bipedalism, obstetrics and thermoregulation. *Philosophical Transactions of the Royal Society B: Biological Sciences* 370(1663):20140063. <https://doi.org/10.1098/rstb.2014.0063>
- Henyš P, Kuchař M (2025) Bonedat, a database of standardized bone morphology for in silico analyses. *Scientific Data* 12(1). <https://doi.org/10.1038/s41597-025-05161-y>

- Henyš P, Čapek L (2019) Computational modal analysis of a composite pelvic bone: convergence and validation studies. *Computer Methods in Biomechanics and Biomedical Engineering* 22(9):916–924. <https://doi.org/10.1080/10255842.2019.1604949>
- Henyš P, Kuchař M, Hájek P, et al (2021) Mechanical metric for skeletal biomechanics derived from spectral analysis of stiffness matrix. *Scientific Reports* 11:15690. <https://doi.org/10.1038/s41598-021-94998-5>
- Henyš P, Ramezani M, Schewitz D, et al (2022) Sacrospinous and sacrotuberous ligaments influence in pelvis kinematics. *Journal of Anatomy* 241(4):928–937. <https://doi.org/10.1111/joa.13739>
- Huiskes R, Weinans H, Grootenboer HJ, et al (1987) Adaptive bone-remodeling theory applied to prosthetic-design analysis. *Journal of Biomechanics* 20(11–12):1135–1150. [https://doi.org/10.1016/0021-9290\(87\)90030-3](https://doi.org/10.1016/0021-9290(87)90030-3)
- Keller TS (1994) Predicting the compressive mechanical behavior of bone. *Journal of Biomechanics* 27(9):1159–1168. [https://doi.org/10.1016/0021-9290\(94\)90056-6](https://doi.org/10.1016/0021-9290(94)90056-6)
- Kuhn HW (1955) The hungarian method for the assignment problem. *Naval Research Logistics Quarterly* 2(1–2):83–97. <https://doi.org/10.1002/nav.3800020109>
- Manconi E, Mace BR (2017) Veering and strong coupling effects in structural dynamics. *Journal of Vibration and Acoustics* 139(2). <https://doi.org/10.1115/1.4035109>
- Marsden JE, Hughes TJR (1983) *Mathematical Foundations of Elasticity*. Prentice Hall
- Morgan EF, Bayraktar HH, Keaveny TM (2003) Trabecular bone modulus-density relationships depend on anatomic site. *Journal of Biomechanics* 36(7):897–904. [https://doi.org/10.1016/S0021-9290\(03\)00071-X](https://doi.org/10.1016/S0021-9290(03)00071-X)
- Pierre C (1988) Mode localization and eigenvalue loci veering phenomena in disordered structures. *Journal of Sound and Vibration* 126(3):485–502. [https://doi.org/10.1016/0022-460X\(88\)90226-X](https://doi.org/10.1016/0022-460X(88)90226-X)
- Salo Z, Beek M, Wright D, et al (2017) Analysis of pelvic strain in different gait configurations in a validated cohort of computed tomography based finite element models. *Journal of Biomechanics* 64:120–130. <https://doi.org/10.1016/j.jbiomech.2017.09.014>
- Schönemann PH (1966) A generalized solution of the orthogonal procrustes problem. *Psychometrika* 31(1):1–10. <https://doi.org/10.1007/BF02289451>
- Snijders CJ, Vleeming A, Stoeckart R (1993) Transfer of lumbosacral load to iliac bones and legs. part 1: Biomechanics of self-bracing of the sacroiliac joints. *Clinical Biomechanics* 8(6):285–294. [https://doi.org/10.1016/0268-0033\(93\)90002-Y](https://doi.org/10.1016/0268-0033(93)90002-Y)
- Vleeming A, Schuenke MD, Masi AT, et al (2012) The sacroiliac joint: an overview of its anatomy, function and potential clinical implications. *Journal of Anatomy* 221(6):537–567. <https://doi.org/10.1111/j.1469-7580.2012.01564.x>