TETRODOTOXIN RESISTANCE IN GARTER SNAKES: AN EVOLUTIONARY RESPONSE OF PREDATORS TO DANGEROUS PREY

EDMUND D. BRODIE, III 940 East 57th Street, Department of Ecology and Evolution, University of Chicago, Chicago, IL 60637

AND

EDMUND D. BRODIE, JR.

Research article

THE EVOLUTIONARY RESPONSE OF PREDATORS TO DANGEROUS PREY: HOTSPOTS AND COLDSPOTS IN THE GEOGRAPHIC MOSAIC OF COEVOLUTION BETWEEN GARTER SNAKES AND NEWTS @

Edmund D. Brodie, Jr., B. J. Ridenhour, E. D. III Brodie

Evolution, Volume 56, Issue 10, 1 October 2002, Pages 2067-2082, https://doi.org/10.1111 /i.0014-3820.2002.tb00132.x

Published: 01 October 2002 Article history v Research Article

JOURNAL ARTICLE

Evolutionary Biology

The skin microbiome facilitates adaptive tetrodotoxin production in poisonous newts

Patric M Vaelli, Kevin R Theis, Janet E Williams, Lauren A O'Connell, James A Foster, Heather L Eisthen 🖼

Journal of Animal Ecology

RESEARCH ARTICLE | 6 Free Access

The geographic mosaic in parallel: Matching patterns of newt tetrodotoxin

snake resistance in multiple predator-prey pairs



Individual fluctuations in toxin levels affect breeding site fidelity in a chemically defended amphibian

Gary M. Bucciarelli , David B. Green, H. Bradley Shaffer and Lee B. Kats

Published: 25 May 2016 https://doi.org/10.1098/rspb.2016.0468

Chemical Ecology of the North American Newt Genera Taricha and Notophthalmus

Charles T. Hanifin . Yuta Kudo & Mari Yotsu-Yamashita

Chapter | First (

Published: March 11, 2008 • https://doi.org/10.1371/journal.pbio.0060060





