

# System drift and speciation

Joshua S. Schiffman<sup>†</sup>     Peter L. Ralph<sup>†‡</sup>

<sup>†</sup>Molecular and Computational Biology, University of Southern California, Los Angeles, California 90089, U.S.A.

<sup>‡</sup>Departments of Mathematics and Biology & The Institute for Ecology and Evolution, University of Oregon, Eugene, Oregon 97403, U.S.A.

jsschiff@usc.edu     plr@uoregon.edu

August 16, 2018

## Abstract

Even if a species' phenotype does not change over evolutionary time, the underlying mechanism may change, as distinct molecular pathways can realize identical phenotypes. Here we use linear system theory to explore the consequences of this idea, describing how a gene network underlying a conserved phenotype evolves, as the genetic drift of small changes to these molecular pathways cause a population to explore the set of mechanisms with identical phenotypes. To do this, we model an organism's internal state as a linear system of differential equations for which the environment provides input and the phenotype is the output, in which context there exists an exact characterization of the set of all mechanisms that give the same input–output relationship. This characterization implies that selectively neutral directions in genotype space should be common and that the evolutionary exploration of these distinct but equivalent mechanisms can lead to the reproductive incompatibility of independently evolving populations. This evolutionary exploration, or *system drift*, is expected to proceed at a rate proportional to the amount of intrapopulation genetic variation divided by the effective population size ( $N_e$ ). At biologically reasonable parameter values this could lead to substantial interpopulation incompatibility, and thus speciation, in fewer than  $N_e$  generations. This model also naturally predicts Haldane's rule, thus providing another possible explanation of why heterogametic hybrids tend to be disrupted more often than homogametes during the early stages of speciation.

## Introduction

It is an overarching goal of many biological subdisciplines to attain a general understanding of the function and evolution of the complex molecular machinery that translates an organism's genome into the characteristics on which natural selection acts, the phenotype. For example, there is a growing body of data on the evolutionary histories and molecular characterizations of particular gene regulatory networks [Jaeger, 2011, Davidson and Erwin, 2006, Israel et al., 2016], as well as thoughtful verbal and conceptual models [True and Haag, 2001, Weiss and Fullerton, 2000, Edelman and Gally, 2001, Pavlicev and Wagner, 2012]. Mathematical models of both particular regulatory networks and the evolution of such systems in general can provide guidance where intuition fails, and thus has the potential to discover general principles in the organization of biological systems as well as provide concrete numerical predictions [Servedio et al., 2014]. There is a substantial amount of work studying the evolution of gene regulatory networks, in frameworks both abstract [Wagner, 1994, 1996, Siegal and Bergman, 2002, Bergman and Siegal, 2003, Draghi and Whitlock, 2015] and empirically inspired [Mjolsness et al., 1991, Jaeger et al., 2004, Kozlov et al., 2015, Crombach et al., 2016, Wotton et al., 2015, Chertkova et al., 2017].

It is well known that in many contexts mathematical models can fundamentally be *nonidentifiable* and/or *indistinguishable* – meaning that there can be uncertainty about an inferred model's parameters or even its claims about causal structure, despite access to complete and perfect data [Bellman and Åström, 1970, Grewal and Glover, 1976, Walter et al., 1984]. Models with different parameter schemes, or even different mechanics can make equally accurate predictions, but still not actually reflect the internal dynamics of the

system being modelled. In control theory, where electrical circuits and mechanical systems are often the focus, it is understood that there can be an infinite number of “realizations”, or ways to reverse engineer the dynamics of a “black box”, even if all possible input and output experiments are performed [Kalman, 1963, Anderson et al., 1966, Zadeh and Deoser, 1976]. The inherent nonidentifiability of chemical reaction networks is sometimes referred to as “the fundamental dogma of chemical kinetics” [Craciun and Pantea, 2008]. In computer science, this is framed as the relationship among processes that *simulate* one another [Van der Schaft, 2004]. Finally, the field of *inverse problems* studies those cases in which, despite the existence of a theoretical one-to-one mapping between a model and behavior, tiny amounts of noise make inference problems nonidentifiable in practice [Petrov and Sizikov, 2005].

Nonidentifiability is a major barrier to mechanistic understanding of real systems, but viewed from another angle, this concept can provide a starting point for thinking about externally equivalent systems – systems that evolution can explore, so long as the parameters and structures can be realized biologically. These functional symmetries manifest in convergent and parallel evolution, as well as *developmental system drift*: the observation that macroscopically identical phenotypes in even very closely related species can in fact be divergent at the molecular and sequence level [Kimura, 1981, True and Haag, 2001, Tanay et al., 2005, Tsong et al., 2006, Hare et al., 2008, Lavoie et al., 2010, Vierstra et al., 2014, Matsui et al., 2015, Dalal et al., 2016, Dalal and Johnson, 2017].

The main purpose of this paper is to explore this general idea in the concrete framework of linear systems theory, thought of as modeling gene regulatory networks. First, we apply results from system theory which give an analytical description of the set of all linear gene network architectures that yield identical phenotypes. This suggests that, quite generally, there are many directions in which a gene network can drift without changing the phenotype. Since these phenotypically equivalent gene networks are not necessarily compatible with one another, system drift may result in reproductive incompatibility between isolated populations, even in the absence of any sort of adaptive, or environmental change.

It is not a new observation that there is often more than one way to do the same thing, or that speciation can be the result of (nearly) neutral processes. The potential for speciation has been analyzed in models of traits under stabilizing selection determined additively by alleles at many loci [Wright, 1935, Barton, 1986, 1989, 2001], in related fitness landscape models [Fraïsse et al., 2016], and for pairs of traits that must match but whose value is unconstrained [Sved, 1981]. It has also been shown that population structure can allow long-term stable coexistence of incompatible genotypes encoding identical phenotypes [Phillips, 1996]. However, previous simulations of system drift in regulatory sequences [Tulchinsky et al., 2014] and a regulatory cascade [Porter and Johnson, 2002] found rapid speciation under directional selection but only equivocal support for speciation under models of purely neutral drift. The rate at which hybrid incompatibility accumulates due to genetic drift creating segregation variance between isolated populations is fairly well understood [Slatkin and Lande, 1994, Rosas et al., 2010, Chevin et al., 2014], but model assumptions can strongly affect predictions, including whether variation is due to rare or common alleles [Slatkin and Lande, 1994], and the shape of the fitness landscape [Fraïsse et al., 2016]. Our main aim is to provide a concrete framework that can provide natural predictions of these model parameters across a general class of models. Furthermore, tools from system theory allow analytical predictions to be made for large populations with complex phenotypes that would be inaccessible to population simulations.

## Results

We use a model of gene regulatory networks that describes the temporal dynamics of a collection of  $n$  coregulating molecules – such as transcription factors – as well as external or environmental inputs. We write  $\kappa(t)$  for the vector of  $n$  molecular concentrations at time  $t$ . The vector of  $m$  “inputs” determined exogenously to the system is denoted  $u(t)$ , and the vector of  $\ell$  “outputs” is denoted  $\phi(t)$ . The output is merely a linear function of the internal state:  $\phi_i(t) = \sum_j C_{ij} \kappa_j(t)$  for some matrix  $C$ . Since  $\phi$  is what natural selection acts on, we refer to it as the *phenotype* (meaning the “visible” aspects of the organism), and in contrast refer to  $\kappa$  as the *kryptotype*, as it is “hidden” from direct selection. Although  $\phi$  may depend on all entries of  $\kappa$ , it is usually of lower dimension than  $\kappa$ , and we tend to think of it as the subset of

molecules relevant for survival. The dynamics are determined by the matrix of regulatory coefficients,  $A$ , a time-varying vector of inputs  $u(t)$ , and a matrix  $B$  that encodes the effect of each entry of  $u$  on the elements of the kryptotype. The rate at which the  $i^{\text{th}}$  concentration changes is a weighted sum of the concentrations as well as the input:

$$\begin{aligned}\dot{\kappa}(t) &= A\kappa(t) + Bu(t) \\ \phi(t) &= C\kappa(t).\end{aligned}\tag{1}$$

Furthermore, we always assume that  $\kappa(0) = 0$ , so that the kryptotype measures deviations from initial concentrations. Here  $A$  can be any  $n \times n$  matrix,  $B$  any  $n \times m$ , and  $C$  any  $\ell \times n$  dimensional matrix, with usually  $\ell$  and  $m$  less than  $n$ . We think of the system as the triple  $(A, B, C)$ , which translates (time-varying)  $m$ -dimensional input  $u(t)$  into the  $\ell$ -dimensional output  $\phi(t)$ . Under quite general assumptions, we can write the phenotype as

$$\phi(t) = Ce^{At}\kappa(0) + \int_0^t Ce^{A(t-s)}Bu(s)ds,\tag{2}$$

which is a convolution of the input  $u(t)$  with the system's *impulse response*, which we denote as  $h(t) := Ce^{At}B$ .

In terms of gene regulatory networks,  $A_{ij}$  determines how the  $j^{\text{th}}$  transcription factor regulates the  $i^{\text{th}}$  transcription factor. If  $A_{ij} > 0$ , then  $\kappa_j$  upregulates  $\kappa_i$ , while if  $A_{ij} < 0$ , then  $\kappa_j$  downregulates  $\kappa_i$ . The  $i^{\text{th}}$  row of  $A$  is therefore determined by genetic features such as the strength of  $j$ -binding sites in the promoter of gene  $i$ , factors affecting chromatin accessibility near gene  $i$ , or basal transcription machinery activity. The form of  $B$  determines how the environment influences transcription factor expression levels, and  $C$  might determine the rate of production of downstream enzymes. To demonstrate this approach, we apply it to construct a simple gene network in Example 1 below.

**Example 1** (An oscillator). *For illustration, we consider an extremely simplified model of oscillating gene transcription, as for instance is found in cell cycle control or the circadian rhythm. There are two genes, whose transcript concentrations are given by  $\kappa_1(t)$  and  $\kappa_2(t)$ , and gene-2 upregulates gene-1, while gene-1 downregulates gene-2 with equal strength. Only the dynamics of gene-1 are consequential to the oscillator (perhaps the amount of gene-1 activates another downstream gene network). Lastly, both genes are equally upregulated by an exogenous signal. The dynamics of the system are described by*

$$\begin{aligned}\dot{\kappa}_1(t) &= \kappa_2(t) + u(t) \\ \dot{\kappa}_2(t) &= -\kappa_1(t) + u(t) \\ \phi(t) &= \kappa_1(t).\end{aligned}$$

*In matrix form the system regulatory coefficients are given as,  $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ , and  $C = \begin{bmatrix} 1 & 0 \end{bmatrix}$ . Suppose the input is an impulse at time zero (a delta function), and so its phenotype is equal to its impulse response,*

$$\phi(t) = h(t) = \sin t + \cos t.$$

*The system and its dynamics are referred to in Figure 1. We return to the evolution of such a system below.*




Figure 1: (Left) Diagram of the gene network in Example 1, and (right) plot of the phenotype  $\phi(t)$  against time  $t$ .

## Equivalent gene networks

As reviewed above, some systems with identical phenotypes are known to differ, sometimes substantially, at the molecular level; systems with identical phenotypes do not necessarily have identical kryptotypes. How many different mechanisms perform the same function?

Two systems are equivalent if they produce the same phenotype given the same input, i.e., have the same input–output relationship. We say that the systems defined by  $(A, B, C)$  and  $(\bar{A}, \bar{B}, \bar{C})$  are **phenotypically equivalent** if their impulse response functions are the same:  $h(t) = \bar{h}(t)$  for all  $t \geq 0$ . This implies that for any acceptable input  $u(t)$ , if  $(\kappa_u(t), \phi_u(t))$  and  $(\bar{\kappa}_u(t), \bar{\phi}_u(t))$  are the solutions to equation (1) of these two systems, respectively, then

$$\phi_u(t) = \bar{\phi}_u(t) \quad \text{for all } t \geq 0.$$

In other words, phenotypically equivalent systems respond identically for *any* input.

One way to find other systems phenotypically equivalent to a given one is by change of coordinates: if  $V$  is an invertible matrix, then the systems  $(A, B, C)$  and  $(VAV^{-1}, VB, CV^{-1})$  are phenotypically equivalent because their impulse response functions are equal:

$$\begin{aligned} h(t) &= Ce^{At}B = CV^{-1}Ve^{At}V^{-1}VB \\ &= CV^{-1}e^{VAV^{-1}t}VB = \bar{C}e^{\bar{A}t}\bar{B} = \bar{h}(t). \end{aligned} \tag{3}$$

However, not all phenotypically equivalent systems are of this form: systems can have identical impulse responses without being coordinate changes of each other. In fact, systems with identical impulse responses can involve interactions between different numbers of molecules, and thus have kryptotypes in different dimensions altogether.

This implies that most systems have at least  $n^2$  degrees of freedom, where recall  $n$  is the number of components of the kryptotype vector. This is because for an arbitrary  $n \times n$  matrix  $Z$ , taking  $V$  to be the identity matrix plus a small perturbation in the direction of  $Z$  above implies that moving  $A$  in the direction of  $ZA - AZ$  while also moving  $B$  in the direction of  $ZB$  and  $C$  in the direction of  $-CZ$  will leave the phenotype unchanged to second order in the size of the perturbation. If the columns of  $B$  and the rows of  $C$  are not all eigenvectors of  $A$ , then any such  $Z$  will result in a different system.

It turns out that in general, there are more degrees of freedom, except if the system is *minimal* – meaning, informally, that it uses the smallest possible number of components to achieve the desired dynamics. Results in system theory show that any system can be realized in a particular minimal dimension (the dimension of the kryptotype,  $n_{\min}$ ), and that any two phenotypically equivalent systems of dimension  $n_{\min}$  are related by a change of coordinates. Since gene networks can grow or shrink following gene duplications and deletions, these additional degrees of freedom can apply, in principle, to any system.

Even if the system is not minimal, results from systems theory explicitly describe the set of all phenotypically equivalent systems. We refer to  $\mathcal{N}(A_0, B_0, C_0)$  as the set of all systems phenotypically equivalent to the system defined by  $(A_0, B_0, C_0)$ :

$$\mathcal{N}(A_0, B_0, C_0) = \{(A, B, C) : Ce^{At}B = C_0e^{A_0t}B_0 \text{ for } t \geq 0\}. \quad (4)$$

These systems need not have the same kryptotypic dimension  $n$ , but must have the same input and output dimensions ( $\ell$  and  $m$ , respectively).

The Kalman decomposition, which we now describe informally, elegantly characterizes this set [Kalman, 1963, Kalman et al., 1969, Anderson et al., 1966]. To motivate this, first note that the input  $u(t)$  only directly pushes the system in certain directions (those lying in the span of the columns of  $B$ ). As a result, different combinations of input can move the system in any direction that lies in what is known as the *reachable subspace*. Analogously, we can only observe motion of the system in certain directions (those lying in the span of the rows of  $C$ ), and so can only infer motion in what is known as the *observable subspace*. The Kalman decomposition then classifies each direction in kryptotype space as either reachable or unreachable, and as either observable or unobservable. Only the components that are both reachable and observable determine the system's phenotype – that is, components that both respond to an input and produce an observable output.

Concretely, the **Kalman decomposition** of a system  $(A, B, C)$  gives a change of basis  $P$  such that the transformed system  $(PAP^{-1}, PB, CP^{-1})$  has the following form:

$$PAP^{-1} = \begin{bmatrix} A_{r\bar{o}} & A_{r\bar{o},ro} & A_{r\bar{o},\bar{r}\bar{o}} & A_{r\bar{o},\bar{r}o} \\ 0 & A_{ro} & 0 & A_{ro,\bar{r}o} \\ 0 & 0 & A_{\bar{r}\bar{o}} & A_{\bar{r}\bar{o},\bar{r}o} \\ 0 & 0 & 0 & A_{\bar{r}o} \end{bmatrix},$$

and

$$PB = \begin{bmatrix} B_{r\bar{o}} \\ B_{ro} \\ 0 \\ 0 \end{bmatrix} \quad (CP^{-1})^T = \begin{bmatrix} 0 \\ C_{ro}^T \\ 0 \\ C_{\bar{r}o}^T \end{bmatrix}.$$

The impulse response of the system is given by

$$h(t) = C_{ro}e^{A_{ro}t}B_{ro},$$

and therefore, the system is phenotypically equivalent to the *minimal* system  $(A_{ro}, B_{ro}, C_{ro})$ .

This decomposition is unique up to a change of basis that preserves the block structure. In particular, the minimal subsystem obtained by the Kalman decomposition is unique up to a change of coordinates. This implies that there is no equivalent system with a smaller number of kryptotypic dimensions than the dimension of the minimal system. It is remarkable that the gene regulatory network architecture to achieve a given input–output map is never unique – both the change of basis used to obtain the decomposition and, once in this form, all submatrices other than  $A_{ro}$ ,  $B_{ro}$ , and  $C_{ro}$  can be changed without affecting the phenotype, and so represent degrees of freedom. (However, some of these subspaces may affect how the system deals with noise.)

*Note on implementation:* The *reachable subspace*, which we denote by  $\mathcal{R}$ , is defined to be the closure of  $\text{span}(B)$  under applying  $A$ , and the *unobservable subspace*, denoted  $\bar{\mathcal{O}}$ , is the largest  $A$ -invariant subspace contained in the null space of  $C$ . The four subspaces,  $r\bar{o}$ ,  $ro$ ,  $\bar{r}\bar{o}$ , and  $\bar{r}o$  are defined from these by intersections and orthogonal complements –  $ro$  refers to the both *reachable and observable* subspace, while  $\bar{r}\bar{o}$  refers to the *unreachable and unobservable* subspace, and similarly for  $\bar{r}o$  and  $r\bar{o}$ .

For the remainder of the paper, we interpret  $\mathcal{N}$  as the neutral set in the fitness landscape, along which a large population will drift under environmental and selective stasis. Even if the phenotype is constrained and

remains constant through evolutionary time, the molecular mechanism underpinning it is not constrained and likely will not be conserved.

Finally, note that if  $B$  and  $C$  are held constant – i.e., if the relationships between environment, kryptotype, and phenotype do not change – there are *still* usually degrees of freedom. The following example 2 gives the set of minimal systems equivalent to the oscillator of Example 1, that all share common  $B$  and  $C$  matrices. The oscillator can also be equivalently realized by a three-gene (or larger) network, and will have even more evolutionary degrees of freedom available, as in Figure 3.

**Example 2** (All equivalent rewirings of the oscillator). *The oscillator of example 1 is minimal, and so any equivalent system is a change of coordinates by an invertible matrix  $V$ . If we further require  $B$  and  $C$  to be invariant then we need  $VB = B$  and  $CV = C$ . Therefore the following one-parameter family  $(A(\tau), B, C)$  describes the set of all two-gene systems phenotypically equivalent to the oscillator:*

$$A(\tau) = \frac{-1}{\tau + 1} \begin{bmatrix} -1 & -\tau & -1 \\ 2\tau(\tau + 1) + 1 & \tau & \end{bmatrix} \text{ for } \tau \neq -1.$$

The resulting set of systems are depicted in Figure 2.




Figure 2: (Left)  $A(\tau)$ , the set of all phenotype-equivalent cell cycle control networks. (Right) Gene-1 dynamics (blue) for both systems  $A(0)$  and  $A(-2)$  are identical, however,  $A(0)$  gene-2 dynamics (orange) differ from  $A(-2)$  (green).




Figure 3: A possible non-minimal three-gene oscillator, phenotypically equivalent to  $A(\tau)$ , the systems in Examples 1 and 2.

**Sexual reproduction and recombination** Parents with phenotypically equivalent yet differently wired gene networks may produce offspring with dramatically different phenotypes. If the phenotypes are significantly divergent then the offspring may be inviable or otherwise dysfunctional, despite both parents being well adapted. If this is consistent for the entire population, we would consider them to be separate species, in accord with the biological species concept [Mayr, 2000].

First, we must specify how sexual reproduction acts on these systems. Suppose that each of a diploid organisms' two genomes encodes a set of system coefficients. We assume that a diploid which has inherited systems  $(A', B', C')$  and  $(A'', B'', C'')$  from its two parents has phenotype determined by the system that averages these two,  $((A' + A'')/2, (B' + B'')/2, (C' + C'')/2)$ .

Each genome an organism inherits is generated by meiosis, in which both of its diploid parents recombine their two genomes, and so an  $F_1$  offspring carries one system copy from each parent, and an  $F_2$  is an offspring of two independently formed  $F_1$ s. If the parents are from distinct populations, these are simply first- and second-generation hybrids, respectively.

Exactly how the coefficients (i.e., entries of  $A$ ,  $B$  and  $C$ ) of a haploid system inherited by an offspring from her diploid parent are determined by the parent's two systems depends on the genetic basis of any variation in the coefficients. Thanks to the randomness of meiotic segregation, the result is random to the extent that each parent is heterozygous for alleles that affect the coefficients. Since the  $i^{\text{th}}$  row of  $A$  summarizes how each gene regulates gene  $i$ , and hence is determined by the promoter region of gene  $i$ , the elements of a row of  $A$  tend to be inherited together, which will create covariance between entries of the same row. It is, however, a quite general observation that the variation seen among recombinant systems is proportional to the difference between the two parental systems.

Offspring formed from two phenotypically identical systems do not necessarily exhibit the same phenotype as both of its parents – in other words  $\mathcal{N}$ , the set of all systems phenotypically equivalent to a given one, is not, in general, closed under averaging or recombination. If sexual recombination among systems drawn from  $\mathcal{N}$  yields systems with divergent phenotypes, populations containing significant diversity in  $\mathcal{N}$  can carry genetic load, and isolated populations may fail to produce hybrids with viable phenotypes.





Figure 4: A conceptual figure of the fitness consequences of hybridization: axes represent system coefficients (i.e., entries of  $A$ ); the line of optimal system coefficients is down in black; dotted lines give phenotypic distances to the optimum. A pair of parental populations are shown in black, along the optimum; a hypothetical population of  $F_1$ s are shown in red, and the distribution of one type of  $F_2$  is shown in purple (other types of  $F_2$  are not shown; some would be a similar distance to the other side of the optimal set). The distribution of  $F_2$  hybrids is appropriate for mixed homozygotes if both traits have a simple, one-locus genetic basis, but there is variation within each population at that locus.

## Hybrid incompatibility

Two parents with the optimal phenotype can produce offspring whose phenotype is suboptimal if the parents have different underlying systems. How quickly do hybrid phenotypes break down as genetic distance between parents increases? We will quantify how far a system’s phenotype is from optimal using a weighted difference between impulse response functions. Suppose that  $\rho(t)$  is a nonnegative, smooth, square-integrable weighting function,  $h_0(t)$  is the *optimal* impulse response function and define the “distance to optimum” of another impulse response function to be

$$D(h) = \left( \int_0^\infty \rho(t) \|h(t) - h_0(t)\|^2 dt \right)^{1/2}. \quad (5)$$

Consider reproduction between a parent with system  $(A, B, C)$  and another displaced by distance  $\epsilon$  in the direction  $(X, Y, Z)$ , i.e., having system  $(A + \epsilon X, B + \epsilon Y, C + \epsilon Z)$ . We assume both are “perfectly adapted” systems, i.e., having impulse response function  $h_0(t)$ , and their offspring has impulse response function  $h_\epsilon(t)$ . A Taylor expansion of  $D(h_\epsilon)$  in  $\epsilon$  shows that the phenotype of an  $F_1$  hybrid between these two is at distance proportional to  $\epsilon^2$  from optimal, while  $F_2$  hybrids are at distance proportional to  $\epsilon$ . This is because an  $F_1$  hybrid has one copy of each parental system, and therefore lies directly between the parental systems (see Figure 4) – the parents both lie in  $\mathcal{N}$ , which is the valley defined by  $D$ , and so their midpoint only differs from optimal due to curvature of  $\mathcal{N}$ . In contrast, an  $F_2$  hybrid may be homozygous for one parental type in some coefficients and homozygous for the other parental type in others; this means that each coefficient of an  $F_2$  may be equal to either one of the parents, or intermediate between the two; this means that possible  $F_2$  systems may be as far from the optimal set,  $\mathcal{N}$ , as the distance between the parents. The precise rate at which the phenotype of a hybrid diverges depends on the geometry of the optimal set  $\mathcal{N}$  relative to segregating genetic variation.



242 **Example 3** (Hybrid incompatibility: misregulation due to system drift). *Offspring of two equivalent systems*  
 243 *from Example 2 can easily fail to oscillate. For instance, the  $F_1$  offspring between homozygous parents at*  
 244  *$\tau = 0$  and  $\tau = -2$  has phenotype  $\phi_{F_1}(t) = e^t$ , rather than  $\phi(t) = \sin t + \cos t$ . However, the coefficients*  
 245 *of these two parental systems differ substantially, probably more than would be observed between diverging*  
 246 *populations. In figure 5 we compare the phenotypes for  $F_1$  and  $F_2$  hybrids between more similar parents,*  
 247 *and see increasingly divergent phenotypes as the difference between the parental systems increases. (In this*  
 248 *example, the coefficients of  $A(\epsilon)$  differ from those of  $A(0)$  by an average factor of  $1 + \epsilon/2$ ; such small*  
 249 *differences could plausibly be caused by changes to promoter sequences.) This divergence is quantified in*  
 250 *Figure 6, which shows that mean distance to optimum phenotype of the  $F_1$  and  $F_2$  hybrid offspring between*  
 251  *$A(0)$  and  $A(\epsilon)$  increases with  $\epsilon^2$  and  $\epsilon$ , respectively.*



Figure 5: **(left)** Phenotypes of  $F_1$  hybrids between an  $A(0)$  parent and, top-to-bottom, an  $A(1/100)$ , an  $A(1/10)$ , and  $A(1/2)$  parent; parental coefficients differ by around 0.5%, 5%, and 25% respectively. Parental phenotypes ( $\sin t + \cos t$ ) are shown in solid black, and hybrid phenotypes in dashed blue. **(right)** Phenotypes of all possible  $F_2$  hybrids between the same set of parents, with parental phenotype again in black. Different colored lines correspond to different  $F_2$  hybrids; note that some completely fail to oscillate.



Figure 6: Mean hybrid phenotypic distance from optimum computed with equation (5), using  $\rho(t) = \exp(-t/4\pi)$  for  $F_1$  (black) and  $F_2$  (blue) hybrids between  $A(0)$  and  $A(\epsilon)$  parent oscillators. Genetic distance is computed as  $\left(\sum_{ij}(A_{ij}(0) - A_{ij}(\epsilon))^2\right)^{1/2}$ .

**Haldane's rule** This model naturally predicts Haldane's rule, the observation that if only one hybrid sex is sterile or inviable it is likely the heterogametic sex (e.g., the male in XY sex determination systems) [Haldane, 1922, Orr, 1997]. For example, consider an XY species with a two-gene network where the first gene resides on an autosome and the second gene on the X chromosome. A male whose pair of haplotypes is  $(\begin{bmatrix} A_1 & A_2 \\ X_1 & X_2 \end{bmatrix}, \begin{bmatrix} A_1 & A_2 \\ X_1 & X_2 \end{bmatrix})$  has phenotype determined by  $A = \begin{bmatrix} A_1 & A_2 \\ X_1 & X_2 \end{bmatrix}$ , if dosage compensation upregulates heterogametes by a factor of two relative to homogametes (as with *Drosophila*), while a female homozygous for the haplotype  $\begin{bmatrix} \bar{A}_1 & \bar{A}_2 \\ \bar{X}_1 & \bar{X}_2 \end{bmatrix}$ , has phenotype determined by  $A = \begin{bmatrix} \bar{A}_1 & \bar{A}_2 \\ \bar{X}_1 & \bar{X}_2 \end{bmatrix}$ . An  $F_1$  male offspring of these two will have its phenotype determined by  $\begin{bmatrix} (A_1 + \bar{A}_1)/2 & (A_2 + \bar{A}_2)/2 \\ \bar{X}_1 & \bar{X}_2 \end{bmatrix}$ . If both genes resided on the autosomes, this system would only be possible in an  $F_2$  cross. More generally, if the regulatory coefficients for a system are shared between the sex and one or more autosomal chromosomes,  $F_1$  males are effectively equivalent to purely autosomal-system  $F_2$  hybrids, and recall that  $F_2$ s are significantly less fit on average than  $F_1$ s (see Figure 6). Although many alleles will be dominant if the phenotype–fitness relationship is convex, the underlying mechanism does not depend on the *dominance theory* Turelli and Orr [1995] to explain Haldane's rule: instead, it derives from the nature of segregation variance.

## The speed of speciation

XXX Talk about how drift goes like variance over  $N_e$ .

## Genetic variation in empirical regulatory systems

What is known about the key quantity above, the amount of heritable variation in real regulatory networks? The coefficient  $A_{ij}$  from the system (1) measures how much the rate of net production of  $i$  changes per change in concentration of  $j$ . It is generally thought that regulatory sequence change contributes much more to inter- and intraspecific variation than does coding sequence change affecting molecular structure [Schmidt

et al., 2010]. In the context of transcription factor networks this may be affected not only by the binding strength of molecule  $j$  to the promoter region of gene  $i$  but also the effects of other transcription factors (e.g., cooperativity) and local chromatin accessibility [Stefflova et al., 2013]. For this reason, the mutational target size for variation in  $A_{ij}$  may be much larger than the dozens of base pairs typically implicated in the handful of binding sites for transcription factor  $j$  of a typical promoter region, and single variants may affect many entries of  $\mathcal{N}$  simultaneously.

Variation in binding site occupancy may overestimate variation in  $A$ , since it does not capture buffering effects (if for instance only one site of many needs to be occupied for transcription to begin), and variation in expression level measures changes in steady-state concentration (our  $\kappa_i$ ) rather than the *rate* of change. Nonetheless, these measures likely give us an idea of the scale of variability. It has been shown that between human individuals, there is differential occupancy in 7.5% of binding sites of a transcription factor (p65) [Kasowski et al. 2010]. It has also been inferred that cis-regulatory variation accounts for around 2–6% of expression variation in human blood-derived primary cells [Verlaan et al. 2009], and that human population variation explained about 3% of expression variation [Lappalainen et al. 2013]. Allele-specific expression is indicative of standing genetic *cis*-regulatory variation; allele-specific expression in 7.2–8.5% of transcripts of a flycatcher species has been observed [Wang et al. 2017], as well as allele-specific expression in 23.4% of genes studied in a baboon species [Tung et al. 2015]. Taken together, this suggests that variation in the entries of  $A$  may be on the order of at least a few percent between individuals of a population – doubtless varying substantially between species and between genes.

## Discussion

In this paper, we use tools from linear system theory and quantitative genetics to study the evolution of a mechanistic model of the genotype-phenotype map, in which the phenotype is subject to stabilizing selection. In so doing, we provide an explicit model of phenogenetic drift [Weiss and Fullerton, 2000] and developmental system drift [True and Haag, 2001]. In this context, the Kalman decomposition [Kalman, 1963] gives an analytical description of all phenotypically equivalent gene networks. It also implies that nearly all systems are nonidentifiable, and that, in general, there exist axes of genetic variation unconstrained by natural selection. The independent movement of separated populations along these axes by genetic drift can lead to a significant reduction in hybrid viability, and thus precipitate speciation, at a speed dependent on the effective population size and the amount of genetic variation. In this model, at biologically reasonable parameter values, system drift is a significant – and possibly rapid – driver of speciation. This may be surprising because hybrid inviability appears as a consequence of recombining different, yet functionally equivalent, mechanisms, and since species are often defined by their unique adaptations or morphologies.

Consistent with empirical observation of hybrid breakdown (e.g., Plötner et al. 2017), we see that the fitnesses of  $F_2$  hybrids drop at a much faster rate than those of  $F_1$ s. Another natural consequence of the model is Haldane’s rule, that if only one  $F_1$  hybrid sex is inviable or sterile it is likely to be the heterogametic sex. This occurs because if the genes underlying a regulatory network are distributed among both autosomes and the sex chromosome, then heterogametic  $F_1$ s show variation (and fitnesses) similar to that seen in  $F_2$  hybrids.

Is there evidence that this is actually occurring? System drift and network rewiring has been inferred across the tree of life [Wotton et al., 2015, Crombach et al., 2016, Dalal and Johnson, 2017, Johnson, 2017, Ali et al., 2017], and there is often significant regulatory variation segregating within populations. Transcription in hybrids between closely related species with conserved transcriptional patterns can also be divergent [Haerty and Singh, 2006, Maheshwari and Barbash, 2012, Coolon et al., 2014, Michalak and Noor, 2004, Mack and Nachman, 2016], and hybrid incompatibilities have been attributed to cryptic molecular divergence underlying conserved body plans [Gavin-Smyth and Matute, 2013]. Furthermore, in cryptic species complexes (e.g., sun skinks [Barley et al., 2013]), genetically distinct species may be nearly morphologically indistinguishable.

**The origin of species not by means of natural selection?** As classically formulated, the Dobzhansky-Muller model of hybrid incompatibility is agnostic to the relative importance of neutral versus selective genetic substitutions [Coyne and Orr, 1998], and plausible mechanisms have been proposed whereby Dobzhansky-Muller incompatibilities could originate under neutral genetic drift [Lynch and Force, 2000] or stabilizing selection [Fierst and Hansen, 2009]. The same holds for the “pathway model” [Lindtke and Buerkle, 2015], which is closer to the situation here. However, previous authors have argued that neutral processes are likely too slow to be a significant driver of speciation [Nei et al., 1983, Seehausen et al., 2014]. This has led some to conclude that hybrid incompatibility is typically a byproduct of positive selection [Orr et al., 2004, Schluter, 2009] or a consequence of genetic conflict [Presgraves, 2010, Crespi and Nosil, 2013], two processes that typically act much more rapidly than genetic drift. However, our calculations suggest that even under strictly neutral processes, hybrid fitness breaks down as a function of genetic distance rapidly enough to play a substantial role in species formation across the tree of life. This is consistent with broad patterns such as the relationship between molecular divergence and genetic isolation seen by Roux et al. [2016], and the clocklike speciation rates observed by Hedges et al. [2015].

Neutral processes are certainly not the only drivers of speciation. All of these forces – adaptive shifts, conflict, and network drift – are plausible drivers of speciation, and may even interact. Many of our observations carry over to models of directional selection – for instance, rapid drift along the set of equivalent systems could be driven by adaptation in a different, pleiotropically coupled system. Or, reinforcement due to local adaptation might provide a selective pressure that speeds up system drift. Furthermore, while the fitness consequences of incompatibility in any one given network may be small, the cumulative impact of system drift across the many different networks an organism relies on may be substantial. It remains to be seen how the relative strengths of these forces compare.

**Nonlinearity and model assumptions** Of course, real regulatory networks are not linear dynamical systems. Most notably, physiological limits put upper bounds on expression levels, implying saturating response curves. It remains to be seen how well these results carry over into real systems, but the fact that most nonlinear systems can be locally approximated by a linear one suggests our qualitative results may hold more generally. Furthermore, nonidentifiability (which implies the existence of neutral directions) is often found in practice in moderately complex models of biological systems [Gutenkunst et al., 2007, Piazza et al., 2008].

Finally, despite our model’s precise separation of phenotype and kryptotype, this relationship in nature may be far more complicated as aspects of the kryptotype may be less “hidden” than we currently assume. For instance, attributes excluded from the phenotype as modelled here, ignore the potential energy costs associated with excessively large (non-minimal) kryptotypes, as well as the relationship between a specific network architecture and robustness to mutational, transcriptional, or environmental noise. More precise modeling will require better mechanistic understanding not only of biological systems, but also the nature of selective pressures and genetic variation in these systems.

## Acknowledgements

We would like to thank Sergey Nuzhdin, Stevan Arnold, Michael Turelli, Patrick Phillips, Erik Lundgren and Hossein Asgharian for valuable discussion. We would also like to thank Nick Barton, Sarah Signor, Todd Parsons, Joachim Hermisson for very helpful comments on the manuscript. Work on this project was supported by funds from the Sloan Foundation and the NSF (under DBI-1262645) to PR.

## References

Sammi Ali, Sarah Signor, Konstantin Kozlov, and Sergey Nuzhdin. Quantitative variation and evolution of spatially explicit morphogen expression in *Drosophila*. *bioRxiv*, page 175711, 2017. [11](#)

364 BDO Anderson, RW Newcomb, RE Kalman, and DC Youla. Equivalence of linear time-invariant dynamical  
365 systems. *Journal of the Franklin Institute*, 281(5):371–378, 1966. 2, 5

366 Anthony J Barley, Jordan White, Arvin C Diesmos, and Rafe M Brown. The challenge of species delimitation  
367 at the extremes: diversification without morphological change in philippine sun skinks. *Evolution*, 67(12):  
368 3556–3572, 2013. 11

369 N H Barton. The maintenance of polygenic variation through a balance between mutation and stabilizing  
370 selection. *Genet Res*, 47(3):209–216, June 1986. doi: 10.1017/S0016672300023156. URL [https://www.  
371 ncbi.nlm.nih.gov/pubmed/3744046](https://www.ncbi.nlm.nih.gov/pubmed/3744046). 2

372 Nicholas H. Barton. The divergence of a polygenic system subject to stabilizing selection, mutation and  
373 drift. *Genetics Research*, 54(1):59–78, 1989. 2

374 Nicholas H. Barton. The role of hybridization in evolution. *Molecular Ecology*, 10(3):551–568, 2001. 2

375 Richard Ernest Bellman and Karl Johan Åström. On structural identifiability. *Mathematical biosciences*, 7  
376 (3-4):329–339, 1970. 1

377 Aviv Bergman and Mark L Siegal. Evolutionary capacitance as a general feature of complex gene networks.  
378 *Nature*, 424(6948):549–552, 2003. 1

379 Aleksandra A. Chertkova, Joshua S. Schiffman, Sergey V. Nuzhdin, Konstantin N. Kozlov, Maria G. Sam-  
380 sonova, and Vitaly V. Gursky. In silico evolution of the Drosophila gap gene regulatory sequence un-  
381 der elevated mutational pressure. *BMC Evolutionary Biology*, 17(1):4, 2017. ISSN 1471-2148. doi:  
382 10.1186/s12862-016-0866-y. URL <http://dx.doi.org/10.1186/s12862-016-0866-y>. 1

383 Luis-Miguel Chevin, Guillaume Decorzent, and Thomas Lenormand. Niche dimensionality and the genetics  
384 of ecological speciation. *Evolution*, 68(5):1244–1256, 2014. 2

385 Joseph D Coolon, C Joel McManus, Kraig R Stevenson, Brenton R Graveley, and Patricia J Wittkopp.  
386 Tempo and mode of regulatory evolution in Drosophila. *Genome research*, 24(5):797–808, 2014. 11

387 Jerry A Coyne and H Allen Orr. The evolutionary genetics of speciation. *Philosophical Transactions of the  
388 Royal Society of London B: Biological Sciences*, 353(1366):287–305, 1998. 12

389 Gheorghe Craciun and Casian Pantea. Identifiability of chemical reaction networks. *Journal of Mathematical  
390 Chemistry*, 44(1):244–259, 2008. 2

391 Bernard Crespi and Patrik Nosil. Conflictual speciation: species formation via genomic conflict. *Trends in  
392 Ecology & Evolution*, 28(1):48–57, 2013. 12

393 Anton Crombach, Karl R Wotton, Eva Jiménez-Guri, and Johannes Jaeger. Gap gene regulatory dynamics  
394 evolve along a genotype network. *Molecular biology and evolution*, 33(5):1293–1307, 2016. 1, 11

395 Chiraj K Dalal and Alexander D Johnson. How transcription circuits explore alternative architectures while  
396 maintaining overall circuit output. *Genes & Development*, 31(14):1397–1405, 2017. 2, 11

397 Chiraj K Dalal, Ignacio A Zuleta, Kaitlin F Mitchell, David R Andes, Hana El-Samad, and Alexander D  
398 Johnson. Transcriptional rewiring over evolutionary timescales changes quantitative and qualitative prop-  
399 erties of gene expression. *eLife*, 5:e18981, 2016. 2

400 Eric H Davidson and Douglas H Erwin. Gene regulatory networks and the evolution of animal body plans.  
401 *Science*, 311(5762):796–800, 2006. 1

402 Jeremy Draghi and Michael Whitlock. Robustness to noise in gene expression evolves despite epistatic  
403 constraints in a model of gene networks. *Evolution*, 69(9):2345–2358, 2015. 1

Gerald M Edelman and Joseph A Gally. Degeneracy and complexity in biological systems. *Proceedings of the National Academy of Sciences*, 98(24):13763–13768, 2001. 1

Janna L Fierst and Thomas F Hansen. Genetic architecture and postzygotic reproductive isolation: evolution of Bateson-Dobzhansky-Muller incompatibilities in a polygenic model. *Evolution*, 2009. 12

C Fraïsse, P A Gunnarsson, D Roze, N Bierne, and J J Welch. The genetics of speciation: Insights from Fisher’s geometric model. *Evolution*, 70(7):1450–1464, 07 2016. doi: 10.1111/evo.12968. URL <https://www.ncbi.nlm.nih.gov/pubmed/27252049>. 2

Jackie Gavin-Smyth and Daniel R Matute. Embryonic lethality leads to hybrid male inviability in hybrids between *Drosophila melanogaster* and *D. santomea*. *Ecology and Evolution*, 3(6):1580–1589, 2013. 11

M Grewal and K Glover. Identifiability of linear and nonlinear dynamical systems. *IEEE Transactions on Automatic Control*, 21(6):833–837, Dec 1976. doi: 10.1109/TAC.1976.1101375. 1

Ryan N Gutenkunst, Joshua J Waterfall, Fergal P Casey, Kevin S Brown, Christopher R Myers, and James P Sethna. Universally sloppy parameter sensitivities in systems biology models. *PLoS Computational Biology*, 3(10):e189, 2007. 12

Wilfried Haerty and Rama S Singh. Gene regulation divergence is a major contributor to the evolution of Dobzhansky–Muller incompatibilities between species of *Drosophila*. *Molecular Biology and Evolution*, 23(9):1707–1714, 2006. 11

JBS Haldane. Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics*, 12(2):101–109, 1922. 10

Emily E Hare, Brant K Peterson, Venky N Iyer, Rudolf Meier, and Michael B Eisen. Sepsid even-skipped enhancers are functionally conserved in *Drosophila* despite lack of sequence conservation. *PLoS Genetics*, 4(6):e1000106, 2008. 2

S Blair Hedges, Julie Marin, Michael Suleski, Madeline Paymer, and Sudhir Kumar. Tree of life reveals clock-like speciation and diversification. *Molecular Biology and Evolution*, 32(4):835–845, 2015. 12

Jennifer W Israel, Megan L Martik, Maria Byrne, Elizabeth C Raff, Rudolf A Raff, David R McClay, and Gregory A Wray. Comparative developmental transcriptomics reveals rewiring of a highly conserved gene regulatory network during a major life history switch in the sea urchin genus *Heliocidaris*. *PLoS Biology*, 14(3):e1002391, 2016. 1

Johannes Jaeger. The gap gene network. *Cellular and Molecular Life Sciences*, 68(2):243–274, 2011. 1

Johannes Jaeger, Svetlana Surkova, Maxim Blagov, Hilde Janssens, David Kosman, Konstantin N Kozlov, Ekaterina Myasnikova, Carlos E Vanario-Alonso, Maria Samsonova, David H Sharp, et al. Dynamic control of positional information in the early *Drosophila* embryo. *Nature*, 430(6997):368–371, 2004. 1

Alexander D Johnson. The rewiring of transcription circuits in evolution. *Current Opinion in Genetics & Development*, 47:121–127, 2017. 11

Rudolf E. Kalman. Mathematical description of linear dynamical systems. *J. SIAM Control*, 1963. 2, 5, 11

Rudolf E. Kalman, Peter L. Falb, and Michael A. Arbib. *Topics in mathematical system theory*. McGraw-Hill, New York, 1969. ISBN 0754321069. 5

M. Kasowski, F. Grubert, C. Heffelfinger, M. Hariharan, A. Asabere, S. M. Waszak, L. Habegger, J. Rozowsky, M. Shi, A. E. Urban, M. Y. Hong, K. J. Karczewski, W. Huber, S. M. Weissman, M. B. Gerstein, J. O. Korb, and M. Snyder. Variation in transcription factor binding among humans. *Science*, 328(5975):232–235, April 2010. 11



- Motoo Kimura. Possibility of extensive neutral evolution under stabilizing selection with special reference to nonrandom usage of synonymous codons. *Proceedings of the National Academy of Sciences*, 78(9):5773–5777, 1981. 2
- Konstantin Kozlov, Vitaly V Gursky, Ivan V Kulakovskiy, Arina Dymova, and Maria Samsonova. Analysis of functional importance of binding sites in the *Drosophila* gap gene network model. *BMC Genomics*, 2015. 1
- Tuuli Lappalainen, Michael Sammeth, Marc R Friedländer, Peter AC ’t Hoen, Jean Monlong, Manuel A Rivas, Mar Gonzalez-Porta, Natalja Kurbatova, Thasso Griebel, Pedro G Ferreira, et al. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature*, 501(7468):506–511, 2013. 11
- Hugo Lavoie, Hervé Hogues, Jaideep Mallick, Adnane Sellam, André Nantel, and Malcolm Whiteway. Evolutionary tinkering with conserved components of a transcriptional regulatory network. *PLoS Biology*, 8(3):e1000329, 2010. 2
- Dorothea Lindtke and C Alex Buerkle. The genetic architecture of hybrid incompatibilities and their effect on barriers to introgression in secondary contact. *Evolution*, 69(8):1987–2004, 2015. 12
- Michael Lynch and Allan G Force. The origin of interspecific genomic incompatibility via gene duplication. *The American Naturalist*, 156(6):590–605, 2000. 12
- Katya L Mack and Michael W Nachman. Gene regulation and speciation. *Trends in Genetics*, 2016. 11
- Shamoni Maheshwari and Daniel A Barbash. Cis-by-trans regulatory divergence causes the asymmetric lethal effects of an ancestral hybrid incompatibility gene. *PLoS Genetics*, 8(3):e1002597, 2012. 11
- Takeshi Matsui, Robert Linder, Joann Phan, Fabian Seidl, and Ian M Ehrenreich. Regulatory rewiring in a cross causes extensive genetic heterogeneity. *Genetics*, 201(2):769–777, 2015. 2
- Ernst Mayr. The biological species concept. *Species concepts and phylogenetic theory: a debate*. Columbia University Press, New York, pages 17–29, 2000. 7
- Pawel Michalak and Mohamed AF Noor. Association of misexpression with sterility in hybrids of *Drosophila simulans* and *D. mauritiana*. *Journal of Molecular Evolution*, 59(2):277–282, 2004. 11
- Eric Mjolsness, David H Sharp, and John Reinitz. A connectionist model of development. *Journal of Theoretical Biology*, 152(4):429–453, 1991. 1
- Masatoshi Nei, Takeo Maruyama, and Chung-I Wu. Models of evolution of reproductive isolation. *Genetics*, 103(3):557–579, 1983. 12
- H Allen Orr. Haldane’s rule. *Annual Review of Ecology and Systematics*, 28(1):195–218, 1997. 10
- H Allen Orr, John P Masly, and Daven C Presgraves. Speciation genes. *Current Opinion in Genetics & Development*, 14(6):675–679, 2004. 12
- Mihaela Pavlicev and Günter P Wagner. A model of developmental evolution: selection, pleiotropy and compensation. *Trends in Ecology & Evolution*, 27(6):316–322, 2012. 1
- Y.P. Petrov and V.S. Sizikov. *Well-posed, ill-posed, and intermediate problems with applications*, volume 49. Walter de Gruyter, 2005. 2
- Patrick C Phillips. Maintenance of polygenic variation via a migration–selection balance under uniform selection. *Evolution*, 50(3):1334–1339, 1996. 2
- Matthew Piazza, Xiao-Jiang Feng, Joshua D Rabinowitz, and Herschel Rabitz. Diverse metabolic model parameters generate similar methionine cycle dynamics. *Journal of Theoretical Biology*, 251(4):628–639, 2008. 12



- Björn Plötner, Markus Nurmi, Axel Fischer, Mutsumi Watanabe, Korbinian Schneeberger, Svante Holm, Neha Vaid, Mark Aurel Schöttler, Dirk Walther, Rainer Hoefgen, et al. Chlorosis caused by two recessively interacting genes reveals a role of RNA helicase in hybrid breakdown in *Arabidopsis thaliana*. *The Plant Journal*, 2017. 11
- Adam H Porter and Norman A Johnson. Speciation despite gene flow when developmental pathways evolve. *Evolution*, 56(11):2103–2111, 2002. 2
- Daven C Presgraves. The molecular evolutionary basis of species formation. *Nature Reviews Genetics*, 11(3):175–180, 2010. 12
- Ulises Rosas, Nick H. Barton, Lucy Copsey, Pierre Barbier de Reuille, and Enrico Coen. Cryptic variation between species and the basis of hybrid performance. *PLOS Biology*, 8(7):1–12, 07 2010. doi: 10.1371/journal.pbio.1000429. URL <https://doi.org/10.1371/journal.pbio.1000429>. 2
- Camille Roux, Christelle Fraisse, Jonathan Romiguier, Yoann Anciaux, Nicolas Galtier, and Nicolas Bierne. Shedding light on the grey zone of speciation along a continuum of genomic divergence. *PLoS Biology*, 14(12):e2000234, 2016. 12
- Dolph Schluter. Evidence for ecological speciation and its alternative. *Science*, 323(5915):737–741, 2009. 12
- D Schmidt, M D Wilson, B Ballester, P C Schwalie, G D Brown, A Marshall, C Kutter, S Watt, C P Martinez-Jimenez, S Mackay, I Talianidis, P Flicek, and D T Odom. Five-vertebrate ChIP-seq reveals the evolutionary dynamics of transcription factor binding. *Science*, 328(5981):1036–1040, May 2010. doi: 10.1126/science.1186176. URL <https://www.ncbi.nlm.nih.gov/pubmed/20378774>. 10
- Ole Seehausen, Roger K Butlin, Irene Keller, Catherine E Wagner, Janette W Boughman, Paul A Hohenlohe, Catherine L Peichel, Glenn-Peter Saetre, Claudia Bank, Åke Brännström, et al. Genomics and the origin of species. *Nature Reviews Genetics*, 15(3):176–192, 2014. 12
- Maria R Servedio, Yaniv Brandvain, Sumit Dhole, Courtney L Fitzpatrick, Emma E Goldberg, Caitlin A Stern, Jeremy Van Cleve, and D Justin Yeh. Not just a theory – the utility of mathematical models in evolutionary biology. *PLoS Biology*, 12(12):e1002017, 2014. 1
- Mark L Siegal and Aviv Bergman. Waddington’s canalization revisited: developmental stability and evolution. *Proceedings of the National Academy of Sciences*, 99(16):10528–10532, 2002. 1
- Montgomery Slatkin and Russell Lande. Segregation variance after hybridization of isolated populations. *Genetics Research*, 64(1):51–56, 1994. 2
- K. Stefflova, D. Thybert, M. D. Wilson, I. Streeter, J. Aleksic, P. Karagianni, A. Brazma, D. J. Adams, I. Talianidis, J. C. Marioni, P. Flicek, and D. T. Odom. Cooperativity and rapid evolution of cobound transcription factors in closely related mammals. *Cell*, 154(3):530–540, August 2013. 11
- J A Sved. A two-sex polygenic model for the evolution of premating isolation. ii. computer simulation of experimental selection procedures. *Genetics*, 97(1):217–235, January 1981. URL <https://www.ncbi.nlm.nih.gov/pubmed/17249074>. 2
- Amos Tanay, Aviv Regev, and Ron Shamir. Conservation and evolvability in regulatory networks: the evolution of ribosomal regulation in yeast. *Proceedings of the National Academy of Sciences of the United States of America*, 102(20):7203–7208, 2005. 2
- John R True and Eric S Haag. Developmental system drift and flexibility in evolutionary trajectories. *Evolution & Development*, 3(2):109–119, 2001. 1, 2, 11
- Annie E Tsong, Brian B Tuch, Hao Li, and Alexander D Johnson. Evolution of alternative transcriptional circuits with identical logic. *Nature*, 443(7110):415–420, 2006. 2

- Alexander Y Tulchinsky, Norman A Johnson, Ward B Watt, and Adam H Porter. Hybrid incompatibility arises in a sequence-based bioenergetic model of transcription factor binding. *Genetics*, 198(3):1155–1166, 2014. 2
- Jenny Tung, Xiang Zhou, Susan C Alberts, Matthew Stephens, and Yoav Gilad. The genetic architecture of gene expression levels in wild baboons. *eLife*, 4:e04729, 2015. 11
- Michael Turelli and H Allen Orr. The dominance theory of Haldane’s rule. *Genetics*, 140(1):389–402, 1995. 10
- AJ Van der Schaft. Equivalence of dynamical systems by bisimulation. *IEEE transactions on automatic control*, 49(12):2160–2172, 2004. 2
- Dominique J Verlaan, Bing Ge, Elin Grundberg, Rose Hoberman, Kevin CL Lam, Vonda Koka, Joana Dias, Scott Gurd, Nicolas W Martin, Hans Mallmin, et al. Targeted screening of cis-regulatory variation in human haplotypes. *Genome research*, 19(1):118–127, 2009. 11
- Jeff Vierstra, Eric Rynes, Richard Sandstrom, Miaohua Zhang, Theresa Canfield, R Scott Hansen, Sandra Stehling-Sun, Peter J Sabo, Rachel Byron, Richard Humbert, et al. Mouse regulatory DNA landscapes reveal global principles of cis-regulatory evolution. *Science*, 346(6212):1007–1012, 2014. 2
- Andreas Wagner. Evolution of gene networks by gene duplications: a mathematical model and its implications on genome organization. *Proceedings of the National Academy of Sciences*, 91(10):4387–4391, 1994. 1
- Andreas Wagner. Does evolutionary plasticity evolve? *Evolution*, pages 1008–1023, 1996. 1
- Eric Walter, Yves Lecourtier, and John Happel. On the structural output distinguishability of parametric models, and its relations with structural identifiability. *IEEE Transactions on Automatic Control*, 29(1):56–57, 1984. 1
- Mi Wang, Severin Uebbing, and Hans Ellegren. Bayesian inference of allele-specific gene expression indicates abundant cis-regulatory variation in natural flycatcher populations. *Genome Biology and Evolution*, 9(5):1266–1279, 2017. 11
- Kenneth M Weiss and Stephanie M Fullerton. Phenogenetic drift and the evolution of genotype–phenotype relationships. *Theoretical Population Biology*, 57(3):187–195, 2000. 1, 11
- Karl R Wotton, Eva Jiménez-Guri, Anton Crombach, Hilde Janssens, Anna Alcaine-Colet, Steffen Lemke, Urs Schmidt-Ott, and Johannes Jaeger. Quantitative system drift compensates for altered maternal inputs to the gap gene network of the scuttle fly *Megaselia abdita*. *eLife*, 4:e04785, 2015. 1, 11
- Sewall Wright. Evolution in populations in approximate equilibrium. *Journal of Genetics*, 30(2):257, 1935. URL <http://link.springer.com/content/pdf/10.1007/BF02982240.pdf>. 2
- Lotfi A Zadeh and Charles A Deoser. *Linear system theory*. Robert E. Krieger Publishing Company Huntington, 1976. 2

Resubmission Cover Letter  
*Evolution*

Joshua Schiffman  
*and* Peter Ralph  
August 16, 2018

**To the Editor(s) –**

We are pleased to submit a revision of our manuscript,  
**Sincerely,**

**Joshua Schiffman and Peter Ralph**

## Reviewer AE:

I have received the evaluations of two reviewers, and I have read the paper myself. First, sorry for the long time it took to review your manuscript; it was reviewed on the timescale of mathematicians (for Reviewer2), but both reviewers provided thorough evaluations of your work (which in my opinion is better than a quick but superficial review).

Your manuscript is quite long already, and the reviewers' suggestions of modifications and clarification, which need to be implemented, will make the manuscript even longer. I however share the reviewers' opinion that the manuscript's two parts are only loosely related. My suggestion is therefore to publish them separately. Although connecting the systems biology and the quantitative genetics parts is indeed an exciting endeavor, this is not really achieved in the current version of the manuscript, and it would be more profitable to first better describe each part separately. Regarding the connexion between the two parts, please also pay particular attention to R2's first specific comment about dimensionality.

This may be because I am more familiar with Fisher's geometric model than systems biology models, but it seems to me that the first part is more novel, and should therefore deserve your attention first, should you follow my suggestion of publishing the two parts separately.

---

**(AE.1) Format:** *please add line numbers to your manuscript; it is straightforward with the lineno package in LaTeX. Please also do not increase text width too much, as this decreases legibility (LaTeX's default settings already optimize the number of characters per line). Finally, be careful about not confusing citep and citet (a lot of missing parentheses around citations in the second part).*

Reply:

---

**(AE.2) all figures:** *please ensure that all axes legends and labels are big enough and not distorted (same of arrow labels on the diagrams). Please also make sure that the lines are thick enough to be visible.*

Reply:

---

**(AE.3) Above eq (2):** *"Under quite general assumptions": quickly mention them?*

Reply:

---

**(AE.4) eq (2):** *Do you need the first term on the rhs given that you assume that  $\kappa(0) = 0$ ?*

Reply:

---

**(AE.5) Example 1:** *"and so its phenotype" whose phenotype? (the subject of the sentence is "the input")*

Reply:

---

**(AE.6) p5, Note on implementation** *"the closure of  $\text{span}(B)$ " may not be understandable by most of Evolution's readers; please explain what this means.*

Reply:

---

**(AE.7) Figures 1-3:** *In figures 1 and 3, there are two types of arrow heads (+ and - effects), but not in Figure 2, probably because the sign of the effect depends on the value of  $\tau$ . Maybe use a different type of arrow heads for interactions that may change signs, to avoid the confusion with positive effects?*

Reply:

---

**(AE.8) p7, Hybrid incompatibility:** *Please provide a rationale for choosing a particular weight  $\rho$ .*

Reply:

---

**(AE.9) p11, 2nd paragraph** *Try to better relate these different studies to yours (e.g., what you add, what they do and you do not). In addition to R1's suggestions, I add a few more below.*

Reply:

---

**(AE.10) p12 Hybridization:** *What is  $\mathcal{X}$ ?*

Reply:

---

**(AE.11) p15** *"The importance of including neutral directions in these models, which is not usually done" – Do not line of isofitness correspond to neutral directions, and aren't these neutral directions already included in those models?*

Reply:

---

**(AE.12) Additional references:** Weinreich, D. M. and Knies, J. L. (2013), *FISHER'S GEOMETRIC MODEL OF ADAPTATION MEETS THE FUNCTIONAL SYNTHESIS: DATA ON PAIRWISE EPISTASIS FOR FITNESS YIELDS INSIGHTS INTO THE SHAPE AND SIZE OF PHENOTYPE SPACE*. *Evolution*, 67: 2957-2972. doi:10.1111/evo.12156

François Blanquart and Thomas Bataillon *Epistasis and the Structure of Fitness Landscapes: Are Experimental Fitness Landscapes Compatible with Fisher's Geometric Model?* *GENETICS* Early online April 6, 2016

Reply:

## Reviewer 1:

The ms combines two parts that are only loosely linked: one "systems biology" part and a "popgen" part. The combination of both parts is appealing, primarily because we lack such combined approaches and it is not easy to come up with an adequate framework. Below, I will discuss both parts in turn.

General remark: line numbers are very helpful for reviewing.

**(I)** The first (systems biology) part makes use of the concept of "nonidentifiability" of chemical reaction networks and develops these concepts in the context of a genotype to phenotype map. In particular, this map is analyzed for the case where it can be modeled as a linear system.

---

**(1.1)** *First note that the concept of "nonidentifiability" is closely related to themes that have been discussed in evolutionary biology for a long time, under names like "mutational robustness", "network neutrality", "canalization", "redundancy" etc. In particular, if you characterize your model by "many distinct (and mutationally connected) molecular pathways can realize identical phenotypes", note that this essentially describes what has been called a "neutral network" and studied under this name in many articles. Also concrete developmental networks have been studied in this context (eg, von Dassow et al 2000, Nature 406:188–192). Discussion of this literature - and more importantly discussion of terms and notions used here relative to terms that have been used elsewhere - is largely missing. I won't be able to summarize all this*

here, but an older review is the Evolution Perspective piece “Evolution and detection of genetic robustness” (de Visser et al 2003). You can work backward and forward from there.

Reply:

---

(1.2) *Related to this: I find it non-intuitive to start the introduction with the notion of “nonidentifiability”, which is a consequence rather than a cause, the cause being what you later call “phenotypic equivalence” and what had been discussed under other names in other articles.*

Reply:

---

(1.3) *Still related: You give long lists of references in the introduction, but do not provide the reader with any information about specific contributions. A bit more would be helpful (starting with True and Haag coined the term “developmental system drift”, etc).*

Reply:

---

(1.4) 4. *Your ms does not simply assume that there is a large “neutral network” underlying a phenotype, but suggests a mechanistic underpinning to create this neutral space. Since much of the appeal of the first part of the paper is connected to this fact, I’d like to see some more in-depth discussion. (suggestions to follow)*

Reply:

---

(1.5) **(mechanistic models)** *you should probably mention somewhere that mechanistic models of neutral spaces exist in the (quite different) context of RNA folding (papers by Hofacker, Fontana, etal)*

Reply:

---

(1.6) **(mechanistic models)** *For development, Andreas Wagner once suggested a discrete version of a linear model (iterated matrix multiplication). Variants of this have later been used by others (eg Draghi and GP Wagner, I think). How is your model related to these approaches?*

Reply:

---

(1.7) **(mechanistic models)** *You model development as a linear system and this assumption is essential for all further steps (the explicit solution of the dynamics and the Kalman decomposition). For me, the only justification of a linear model (other than mathematical convenience) is local approximation. Do you agree? This will be important for the second part of the ms, see below.*

Reply:

---

(1.8) **(mechanistic models)** *You gain a lot of “neutrality” by the assumption that the dimension of the “kryptotype” is larger than the dimension of the trait. If the trait is the expression of a gene at the top of some pathway, this is certainly true. However, if we include further traits that are affected by the same genes (i.e., pleiotropy) this is no longer clear. The space of “all traits under selection” of an organism is awfully high-dimensional and pleiotropy is wide-spread (even if we do not believe in “omnigenic” models). It is not clear to me from what kind of data we could learn more about these dimensions and I do not expect an answer in your ms, but the issue deserves more discussion. Currently, “pleiotropy” is not even mentioned in the ms.*

Reply:

---

**(1.9) Introduction:** *“Genotypes encoding identical phenotypes can even persist stably within a species” - if there is population structure, I suppose.*

**Reply:**

---

**(1.10) Introduction:** *“It is not a new observations that there is often more than one way to do the same thing, and that this may lead to speciation” - Isn’t this what people would call a neutral Dobzhansky-Muller incompatibility?*

**Reply:**

---

**(1.11) Results:** *(Eq. 1) development as a linear system: has this been done before (probably yes?) - If so, references?*

**Reply:**

---

**(1.12) Results:** *“Of course, neither of these are necessarily true for real systems” - delete “necessarily”?*

**Reply:**

---

**(1.13) Eq. 3:** *What does a coordinate change mean biologically? In our JC, we discussed for a while whether this is just a different parametrization of the exact same biological thing (like transformation to principle components). I think I now understand that it is more, but this should be explained better.*

**Reply:**

---

**(1.14)** *“Since gene networks can grow or shrink following gene duplications and deletions, these additional degrees of freedom can apply in principle to any system.” - are there examples of equivalent gene networks with a different size due to gene duplication or loss?*

**Reply:**

---

**(1.15) Kalman decomposition:** *Define the sub-matrices directly when first used.*

**Reply:**

---

**(1.16) Figure 2:** *Colors are hardly visible*

**Reply:**

(II) The popgen part. This part models the “system drift”, which is nothing else but neutral drift of a population along a high-fitness ridge in an epistatic landscape (conceptual figure 4), right? If two populations in allopatry drift in different directions, this can lead to hybrid incompatibilities, which are uncovered upon secondary contact. Fitness is modeled by the weighted distance from the optimal impulse response function (eq 5). This is a natural assumption.

**Reply:**

---

**(1.17)** *Your analysis of the fitness loss in F1 and F2 hybrids rests on a local Taylor expansion of the fitness landscape. This is adequate given that the underlying linear network is also only locally valid (see above). However, you then apply this to a discussion of hybrid incompatibility (Haldane’s rule etc). We are thus interested in \*very large\* fitness costs for hybrids. This does not seem to be compatible: to get hybrid incompatibility or sterility, you need to have epsilon sufficiently large (below equation 5). But then  $\epsilon^2$  is no*



longer smaller than epsilon. In other words: it seems to me that for the discussion of hybrid incompatibilities you apply your model of the first part beyond the local range where it is valid.

Reply:

---

(1.18) For system drift, you assume that “Selection will tend to restrain this motion, but movement along the optimal set  $N$  is unconstrained, and so we expect the population mean to drift along the optimal set like a particle diffusing.” I see two major problems with this view. Both lead to slower divergence and therefore run against your conclusions. (further points to follow)

Reply:

---

(1.19) (drift along a ridge) In the presence of epistasis, evolution on a neutral network (a high-fitness ridge) is *not* due to drift alone, but also affected by (weak second order) selection in favor of mutational robustness / genetic canalization. In contrast to what you write, diffusion on the set of network coefficients corresponding to the optimal phenotype is not unbiased - even if the optimal phenotypes do indeed all have the same fitness. Instead, selection drives the population to “thicker” parts of the network where the mean fitness of the population (including a cloud of mutants) is higher than on a narrow ridge. This is the basis of the evolution of robustness/canalization. It is possible to account for this effect, see Hermisson et al 2003. Amer. Nat. 161, 708-734; Alvarez-Castro et al, TPB 75 (2009) 109-122, or also Rice, 1998, Evolution 52, 647-656; van Nimwegen, et al 1999, PNAS 96:9716-9720. Note that epistasis is necessary for the neutral evolution of incompatibilities, which is what you are aiming for.

Reply:

---

(1.20) (drift along a ridge) a second problem results from the fact that evolution on a high-fitness ridge often requires coordinated changes at many loci. Take the oscillator system that you use as an example: simultaneous changes at two genes are required to maintain the phenotype. This leads to a phenomenon called “adaptive inertia” (see Baatz and Wagner, TPB 51, 49-66 and Alvarez-Castro et al. TPB 75, 109-122), which effectively slows down the movement along the ridge considerably. This problem applies, in particular, in a “house-of-cards” mutation regime when where rarely two mutations occur together on the same haplotype. In small populations, it typically requires that populations drift through shallow fitness valleys. While this is possible, it slows down the process.

Reply:

---

(1.21) (drift along a ridge) The relevance of both effects could be studied by simulations in a simple example (eg the oscillator that is used as an illustration in the ms anyway).

Reply:

---

(1.22) If speciation due to accumulation of incompatibilities does not occur in allopatry, but under (even weak) gene flow, some degree of positive selection is always needed (Bank et al, Genetics 191, 845-863 for a proof). This also means that even weak gene flow will counteract the process described in the ms.

Reply:

---

(1.23) Population isolates and genetic load: Isn't this exactly “founder effect speciation”?

Reply:

## Reviewer 2:

The present paper is divided into two parts. In the first part, the authors propose a framework in which distinct genetic architectures can produce the same phenotype. (In the fitness landscape terminology, this would correspond to the existence of an evolutionary ridge in a fitness landscape.) The starting point is a nice analogy with linear systems theory. The authors highlight the fact that in general, two distinct linear (differential) systems can respond identically for any input; i.e., different systems (genotypes) can always produce the same output (phenotype) given the same input (environment). Further, the set of equivalent systems can be nicely characterized through Kalman decomposition, thus providing a nice characterization of level sets in the underlying fitness landscape. All along the first part of the paper, this analogy is well illustrated using a simple (yet quite nice) example of oscillating gene transcription. I especially like Fig. 5 where it is shown that even if  $F_1$  hybrids have a phenotypic response close to their parents,  $F_2$  hybrids can behave in a drastically different manner.

In the second part of the paper, motivated by the previous results, the authors explore a general quantitative genetics model in which populations can drift stochastically near a set of equivalent and optimal systems. Since the optimal set (or evolutionary ridge) is not closed under averaging or recombination, two isolated populations can drift apart and accumulate enough genetic differences so that they do not produce any viable offspring. Using some heuristics, several expressions are derived to quantify the accumulation of genetic incompatibilities.

Overall, I think the paper is well written. The framework developed in the first part of the paper is very interesting and that the analogy with system theory is quite enlightening. I am somehow less convinced by the second part. First, I find the arguments a bit sketchy (see below) and not so easy to follow. Secondly, it is not entirely clear to me what is the main contribution of this part compared to previous works. It seems to me that the main result is somehow contained in the fact that the variance (or “segregation variance”) of an  $F_2$  population is given by

$$\sigma_S^2 + 4\omega\sigma_N^2 T/N_e$$

which was already derived Slatkin and Lande according to the authors (except for the explicit expression of  $\omega$ , but again I am a little bit confused by the arguments derived in the appendix).

In summary, I think this paper could be a nice contribution to Evolution. However, I am also convinced that the second part of the paper would require more work, or more explicit reference to previous works (equation, section etc.) if the authors do not want to re-derive already existing formula.

---

**(2.1) end of p6:** “we assume ...  $((A + A')/2, (B + B')/2, (C + C')/2)$ ” . It is claimed in p4 (right before the last paragraph) that two kryptotypes need not have the same dimension. In this case, the previous sum does not make sense, right?

**Reply:**

---

**(2.2) p7: fig 4.** “The distribution of  $F_2$  ... homozygotes”. I do not really grasp the meaning of this sentence. More importantly, I find the purple cloud of dots ( $F_2$  population) quite confusing. It seems to me that the purple populations should be a cloud of points concentrated around a point sitting on the red dotted line, i.e., a Gaussian distribution with the mean at the average of the two parental populations. According to the authors, the distribution is bi-modal with a peak below the optimal set (as displayed on the figure) and another distinct peak sitting on the other side of the optimal. In fact, if I understand the computations of the second part correctly, the red and purple means should coincide, but the red variance should simply be greater.

Reply:

---

(2.3) p7: end of the page. Why do you need  $\rho$  to be square integrable?

Reply:

---

(2.4) p8: “A Taylor expansion of  $D(h_\epsilon)$  .. ”. It is not clear to me at all. Could you provide some extra explanation (e.g., in the appendix)?

Reply:

---

(2.5) p9. Fig 5. In the left panel, it seems to me that the two parents are homozygotes since there is a single  $F_1$  possible offspring (dashed blue curve). It would be worth being more explicit. For the right panels, could you be more explicit on the number of curves. If I understand correctly, there are 16 possibilities due to recombination (2 per entries of the matrix) since  $F_1$  individuals are heterozygotes. Is that correct? Finally, the labels on the y axis are not easy to read.

Reply:

---

(2.6) p11. Paragraph system drift. “move a random distance  $\sigma$ ”. What is  $\sigma$ ? I think it should be  $\sigma_N$  (the std deviation in the direction of the evolutionary ridge) to be consistent with the assumption that the population drifts along the optimal set. I believe that this is what is assumed thereafter. Also, the sentence “It therefore seems .... as cloud of points of width  $\sigma$ ” is not very accurate, since the covariance matrix is not the identity.

Reply:

---

(2.7) p 11. Approximating the optimal set  $\mathcal{N}$  by a quadratic surface should only be accurate if we look at the genetic divergence at small time scales. This should be at least mentioned.

Reply:

---

(2.8) end of p11.  $1/(\frac{d}{du}D(x+uz))$  should be evaluated at  $u=0$ .

Reply:

---

(2.9) p12. Third paragraph  $\sqrt{4\omega T/N_e} \sim \gamma/\sigma_N$ . I guess the underlying assumption here is that  $\sigma_S \ll \sqrt{4\omega\sigma_N^2 T/N_e}$ ?

Reply:

---

(2.10) p12. before eq. 7.  $\mu = c_\mu \gamma T/N_e$ . Why is  $\mu$  proportional to  $\gamma$ ?

Reply:

---

(2.11) Fig 7. I have one important issue with this figure (and the assumptions of the underlying quantitative genetic model). If one wants to be consistent with the assumption that parental populations drift along the evolutionary ridge, I think one would need to assume that selection is strong enough to constraint the mean of the population on the surface. This would presumably require that  $\sigma S/\gamma \ll 1$ . First, I think this assumption (or something alternative to that) should be made explicit in the text. Secondly, it seems to me that this assumption is not satisfied for panel A and C: under the range of parameters proposed by the authors, the parental populations could easily drift away from the optimal set, and in particular, the heuristics derived in the main text would not be satisfied.

**Reply:**

---

**(2.12)** *Finally, it would be worth mentioning several old and recent works relating genetic drift to speciation : Yamagushi and Iwasa, several articles by Gavrilets et al. (I think several citations are missing here), and Mirò Pina and Schertzer.*

**Reply:**