

Assignment

Pettugadi Pranav
CS21BTECH11063

CONTENTS

1	Uniform Random Numbers	1
2	Central Limit Theorem	2
3	From Uniform to Other	4
4	Triangular Distribution	5
5	Guasssian to Other	6

Abstract—This manual provides solutions to the Assignment of Random Numbers

1 UNIFORM RANDOM NUMBERS

Let U be a uniform random variable between 0 and 1.

- 1.1 Generate 10^6 samples of U using a C program and save into a file called uni.dat .

Solution: Download the following files and execute the C program.

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/1-1.c>
we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/cfunc.h>

and compile and execute above files using following commands

```
$ gcc 1-1.c
$ ./a.out
```

- 1.2 Load the uni.dat file into python and plot the empirical CDF of U using the samples in uni.dat. The CDF is defined as

$$F_U(x) = \Pr(U \leq x) \quad (1)$$

Solution: The following code plots Fig. 1.2

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/1-2.py>

to get above fig compile above code using following commands

```
$ python3 1-2.py
```




Fig. 1.2: The CDF of U

- 1.3 Find a theoretical expression for $F_U(x)$.

Solution:

The CDF of U is given by

$$F_U(x) = \Pr(U \leq x) = \int_{-\infty}^x p_U(u) du \quad (2)$$

We now have three cases:

- a) $x < 0$: $p_X(x) = 0$, and hence $F_U(x) = 0$.
b) $0 \leq x < 1$: Here,

$$F_U(x) = \int_0^x du = x \quad (3)$$

- c) for $x > 1$ $F_U(x) = 1$

Therefore,

$$F_U(x) = \begin{cases} 0 & x < 0 \\ x & 0 \leq x < 1 \\ 1 & x \geq 1 \end{cases} \quad (4)$$

1.4 The mean of U is defined as

$$E[U] = \frac{1}{N} \sum_{i=1}^N U_i \quad (5)$$

and its variance as

$$\text{var}[U] = E[U - E[U]]^2 \quad (6)$$

Write a C program to find the mean and variance of U .

Solution: Download the following files

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/1-4.c>
 we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/cfunc.h>

execute them using following files

```
$ gcc 1-4.c
$ ./a.out
```

The value of mean is 0.500007
 the value of variance is 0.083301

1.5 Verify your result theoretically given that

$$E[U^k] = \int_{-\infty}^{\infty} x^k dF_U(x) \quad (7)$$

Solution: we know that

$$\text{var}[U] = E[U - E[U]]^2 \quad (8)$$

$$= E[U^2] - (E[U])^2 \quad (9)$$

$$E[U^2] = \int_{-\infty}^{\infty} x^2 dF_U(x) \quad (10)$$

$$= \int_{-\infty}^{\infty} x^2 p_U(x) dx \quad (11)$$

$$= \int_0^1 x^2 dx = \frac{1}{3} \quad (12)$$

and

$$E[U] = \int_{-\infty}^{\infty} x dF_U(x) \quad (13)$$

$$= \int_{-\infty}^{\infty} x p_U(x) dx \quad (14)$$

$$= \int_0^1 x dx = \frac{1}{2} \quad (15)$$

we know mean = $E[U] = 0.5$

$$\text{var}[U] = E[U - E[U]]^2 \quad (16)$$

$$= E[U^2] - (E[U])^2 = \frac{1}{3} - \frac{1}{4} = \frac{1}{12} \quad (17)$$

$$(18)$$

2 CENTRAL LIMIT THEOREM

2.1 Generate 10^6 samples of the random variable

$$X = \sum_{i=1}^{12} U_i - 6 \quad (19)$$

using a C program, where $U_i, i = 1, 2, \dots, 12$ are a set of independent uniform random variables between 0 and 1 and save in a file called gau.dat

Solution: download following files

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/2-1.c>
 we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/cfunc.h>

and execute them using below commands

```
$ gcc 2-1.c
$ ./a.out
```

2.2 Load gau.dat in python and plot the empirical CDF of X using the samples in gau.dat. What properties does a CDF have?

Solution: The CDF of X is plotted in Fig. 2.2 by downloading below file

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/2-2.py>

and executed using below commands

```
$ python3 2-2.py
```

the properties of CDF are

- 1)The CDF is non decreasing function
- 2)It is symmetric about one point

2.3 Load gau.dat in python and plot the empirical PDF of X using the samples in gau.dat. The PDF of X is defined as

$$p_X(x) = \frac{d}{dx} F_X(x) \quad (20)$$




Fig. 2.2: The CDF of X

What properties does the PDF have?

Solution: The PDF of X is plotted in Fig. 2.3 by downloading following codes

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/2-3.py>

and executed by using following commands

```
$ python3 2-3.py
```




Fig. 2.3: The PDF of X

some properties are

1) area under curve is 1

2) it is symmetric about $x = \mu$

3) it is increasing for $x < \mu$ and decreasing for $x > \mu$

2.4 Find the mean and variance of X by writing a C program.

Solution: to find mean and variance download following files

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/2-4.c>
we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/cfunc.h>

and execute them using following commands

```
$ gcc 2-4.c  
$ ./a.out
```

the value of mean is 0.000326

the value of variance is 1.000906

2.5 Given that

$$p_X(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right), -\infty < x < \infty, \quad (21)$$

repeat the above exercise theoretically.

Solution:

1) CDF is calculated as follows :

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx \quad (22)$$

$$= \frac{1}{\sqrt{2\pi}} \times \sqrt{2\pi} \quad (23)$$

$$= 1 \quad (24)$$

MEAN can be calculated as follows :

$$\int_{-\infty}^{\infty} x \times \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx \quad (25)$$

$$= 0 \quad + \quad (26)$$

2) mean is

$$E(x) = \int_{-\infty}^{\infty} x p_X(x) dx \quad (27)$$

$$= \int_{-\infty}^{\infty} x \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx \quad (28)$$

$$= 0 \quad (29)$$

$$(30)$$

$$E(x)=0$$

3)VARIANCE can be calculated as follows :
 $var[x] = E[X^2] - E[X]^2$
 $var[X] = E[X^2]$ The variance is given by

$$E[X^2] = \int_{-\infty}^{\infty} x^2 \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx = 0 \quad (31)$$

$$= \int_{-\infty}^{\infty} x \times x \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) dx \quad (32)$$

$$= x \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \Big|_{-\infty}^{\infty} \quad (33)$$

$$+ \int_{-\infty}^{\infty} 1 \times \int \frac{x}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \quad (34)$$

$$= x \times 0 + \int_{-\infty}^{\infty} 1 \times \int \frac{x}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) \quad (35)$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{x^2}{2}\right) \quad (36)$$

$$= 1 \quad (37)$$

$$(38)$$

$\therefore var[X] = 1$

3 FROM UNIFORM TO OTHER

3.1 Generate samples of

$$V = -2 \ln(1 - U) \quad (39)$$

and plot its CDF.

Solution: download the following files

we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/3-1.c>
 we get <https://github.com/pettugadipranav/Randomvar/blob/main/codes/cfunc.h>

and execute them using following commands

```
$ gcc 3-1.c
$ ./a.out
```

The CDF of V is plotted in Fig. 3.1 using following codes

<https://github.com/pettugadipranav/Randomvar/blob/main/codes/3-1.py>

and execute them by following commands

```
$ python3 3-1.py
```

3.2 Find a theoretical expression for $F_V(x)$.

Solution: for $Y = G(X)$ then $F_Y(y) =$




Fig. 3.1: The CDF of V

$$F_X(G^{-1}(y))$$

$$V = G(U)$$

$$U = G^{-1}(V) \quad (40)$$

$$= 1 - e^{-\frac{V}{2}} \quad (41)$$

then

$$F_V(v) = F_U(G^{-1}(v)) \quad (42)$$

$$= F_U(1 - e^{-\frac{v}{2}}) \quad (43)$$

$$\Rightarrow F_V(v) = \begin{cases} 0 & x < 0 \\ 1 - e^{-\frac{v}{2}} & x \geq 0 \end{cases} \quad (44)$$

Solution: Note that the function

$$v = f(u) = -2 \ln(1 - u) \quad (45)$$

is monotonically increasing in $[0, 1]$ and $v \in \mathbb{R}^+$. Hence, it is invertible and the inverse function is given by

$$u = f^{-1}(v) = 1 - \exp\left(-\frac{v}{2}\right) \quad (46)$$

Therefore, from the monotonicity of v , and using (4),

$$F_V(v) = F_U\left(1 - \exp\left(-\frac{v}{2}\right)\right) \quad (47)$$

$$\Rightarrow F_V(v) = \begin{cases} 0 & v < 0 \\ 1 - \exp\left(-\frac{v}{2}\right) & v \geq 0 \end{cases} \quad (48)$$

4 TRIANGULAR DISTRIBUTION

4.1 Generate

$$T = U_1 + U_2 \quad (49)$$

Solution: Download the following files and execute the C program.

```
$ https://github.com/pettugadipranav/
  Randomvar/blob/main/codes/4-1.c
$ https://github.com/pettugadipranav/
  Randomvar/blob/main/codes/cfunc.h
```

Execute the above C program files using the following commands

```
$ gcc 4-1.c
$ ./a.out
```

4.2 Find the CDF of T .

Solution: The CDF of T is plotted in Fig. 4.2 using the code below




Fig. 4.2: The CDF of T

```
$ https://github.com/pettugadipranav/
  Randomvar/blob/main/codes/4-2.py
```

Download the above files and execute the following commands to produce Fig.4.2

```
$ python3 4-2.py
```

4.3 Find the PDF of T .

Solution: The PDF of T is plotted in Fig. 4.2 using the code below

```
$ https://github.com/pettugadipranav/
  Randomvar/blob/main/codes/4-3.py
```

Download the above files and execute the following commands to produce Fig.4.2

```
$ python3 4-3.py
```




Fig. 4.3: The PDF of T

4.4 Find the Theoretical Expression for the PDF and CDF of T

Solution:

$$T = U_1 + U_2 \quad (50)$$

$$\Rightarrow p_T(t) = \int_{-\infty}^t p_{U1}(x)p_{U2}(y)dx \quad (51)$$

$$\text{As, } p_{U1}(x) = p_{U1}(y) = p_U(u) \quad (52)$$

$$\Rightarrow p_T(t) = \int_{-\infty}^t p_U(u)p_U(t-u)du \quad (53)$$

a) Theoretical PDF

i) For $0 < t \leq 1$

$$p_T(t) = \int_0^t p_U(t-u)du \quad (54)$$

$$\Rightarrow p_T(t) = \int_0^t du = t \quad (55)$$

ii) For $1 < t \leq 2$

$$p_T(t) = \int_0^1 p_U(t-u)du \quad (56)$$

$$\Rightarrow p_T(t) = \int_{t-1}^1 du = 2 - t \quad (57)$$

$$\therefore P_T(t) = \begin{cases} 0 & t < 0 \\ t & 0 \leq t \leq 1 \\ 2 - t & 1 < t \leq 2 \\ 0 & t > 2 \end{cases}$$

b) Theoretical CDF

$$F_T(t) = \int_{-\infty}^t p_T(u) du \quad (58)$$

$$\therefore F_T(t) = \begin{cases} 0 & t < 0 \\ \frac{t^2}{2} & 0 \leq t \leq 1 \\ 2t - 1 - \frac{t^2}{2} & 1 < t \leq 2 \\ 1 & t > 2 \end{cases}$$

4.5 Verify your results through a plot

Solution: The above theoretical results are verified through the plots in Fig 4.2 and Fig 4.3

5 GUASSIAN TO OTHER

5.1 Generate equiprobable $X \in \{1, -1\}$. **Solution:** Download the following files and execute the C program.

```
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/5-1.c
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/cfunc.h
```

Download the above files and execute the following commands

```
$ gcc 5-1.c
$ ./a.out
```

5.2 Generate

$$Y = AX + N, \quad (59)$$

Where $A = 5$ dB, and $N \sim \mathcal{N}(0, 1)$.

Solution:

```
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/5-2.c
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/cfunc.h
```

Download the above files and execute the following commands

```
$ gcc 5-2.c
$ ./a.out
```

5.3 Plot Y using a scatter plot.

Solution: The CDF of V is plotted in Fig. 5.3 using the code below

```
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/5-3.py
```

Download the above files and execute the following commands to produce Fig.5.3

```
$ python3 5-3.py
```




Fig. 5.3: The Scatter Plot

5.4 Guess how to estimate X from Y .

Solution:

- a) If $Y < 0$ then probably $X = -1$
- b) If $Y > 0$ then probably $X = 1$

5.5 Find

$$P_{e|0} = \Pr(\hat{X} = -1 | X = 1) \quad (60)$$

and

$$P_{e|1} = \Pr(\hat{X} = 1 | X = -1) \quad (61)$$

Solution:

```
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/5-5.c
$ https://github.com/pettugadipranav/
Randomvar/blob/main/codes/cfunc.h
```

Download the above files and execute the following commands to get the result

```
$ gcc 5-5.c
$ ./a.out
```

$$P_{e|0} = 0.499033 \quad (62)$$

$$P_{e|1} = 0.500138 \quad (63)$$

5.6 Find P_e assuming that X has equiprobable symbols.

Solution:

$$P_e = P(X = 1)P_{e|0} + P(X = -1)P_{e|1} \quad (64)$$

Since X is equiprobable

$$P(X = 1) = P(X = -1) = 0.5 \quad (65)$$

$$\Rightarrow P_e = \frac{P_{e|0} + P_{e|1}}{2} \quad (66)$$

$$\Rightarrow \boxed{P_e = 0.499585} \quad (67)$$

5.7 Verify by plotting the theoretical P_e with respect to A from 0 to 10 dB.

Solution :

$$P_{e|0} = \Pr(\hat{X} = -1|X = 1) \quad (68)$$

$$P_{e|0} = \Pr(AX + N < 0|X = 1) \quad (69)$$

$$P_{e|0} = \Pr(N < -A) \quad (70)$$

$$P_{e|0} = \int_{-\infty}^{-A} \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (71)$$

$$P_{e|0} = \int_A^{\infty} \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \quad (72)$$

$$P_{e|0} = Q_N(A) \quad (73)$$

$$\text{Similarly, } P_{e|1} = Q_N(A) \quad (74)$$

\$ <https://github.com/pettugadipranav/Randomvar/blob/main/codes/5-7.py>

Download the above files and execute the following commands to produce Fig.5.7

\$ python3 5-7.py




Fig. 5.7: $P_e(A)$ with semilog-y axis

5.8 Now, consider a threshold δ while estimating X from Y . Find the value of δ that maximizes the theoretical P_e .

Solution: To estimate X from Y , we now consider the following:

$$X = \begin{cases} 1, & Y > \delta \\ -1, & Y < \delta \end{cases} \quad (75)$$

Therefore,

$$P_{e|0} = \Pr(\hat{X} = -1|X = 1) \quad (76)$$

$$= \Pr(AX + N < \delta|X = 1) \quad (77)$$

$$\Rightarrow P_{e|0} = \Pr(N < \delta - A) \quad (78)$$

$$= \int_{-\infty}^{\delta-A} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \quad (79)$$

$$= \int_{A-\delta}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx \quad (80)$$

$$\Rightarrow P_{e|0} = Q_N(A - \delta) \quad (81)$$

Similarly,

$$P_{e|1} = Q_N(A + \delta) \quad (82)$$

$$P_e = P_{e|0} \Pr(X = 1) + P_{e|1} \Pr(X = -1) \quad (83)$$

$$= \frac{Q_N(A - \delta) + Q_N(A + \delta)}{2} \quad (84)$$

Differentiating the above equation wrt δ :

$$0 = \frac{d}{d\delta} \left(\frac{Q_N(A - \delta) + Q_N(A + \delta)}{2} \right) \quad (85)$$

$$= \frac{1}{2} \left(\frac{1}{\sqrt{2\pi}} e^{-\frac{(\delta-A)^2}{2}} - \frac{1}{\sqrt{2\pi}} e^{-\frac{(A+\delta)^2}{2}} \right) \quad (86)$$

$$\Rightarrow (\delta - A)^2 = (A + \delta)^2 \quad (87)$$

$$\Rightarrow \boxed{\delta = 0} \quad (88)$$

5.9 Repeat the above exercise when

$$p_X(0) = p \quad (89)$$

Solution: Using Eq. (83), we have:

$$P_e = P_{e|0}p + P_{e|1}(1 - p) \quad (90)$$

$$= pQ_N(A - \delta) + (1 - p)Q_N(A + \delta) \quad (91)$$

Differentiating as before, we get:

$$0 = p \frac{1}{\sqrt{2\pi}} e^{-\frac{(\delta-A)^2}{2}} - (1-p) \frac{1}{\sqrt{2\pi}} e^{-\frac{(A+\delta)^2}{2}} \quad (92)$$

$$e^{\frac{(\delta+A)^2 - ((\delta-A))^2}{2}} = \frac{1-p}{p} \quad (93)$$

$$\implies \boxed{\delta = 0} \quad (94)$$

5.10 Repeat the above exercise using the MAP criterion.

Solution: Assume that

$$\Pr(X = -1) = p \quad (95)$$

$$\Pr(X = 1) = (1-p) \quad (96)$$

From Total Probability Theorem, we have:

$$\begin{aligned} p_Y(y) &= p_{Y|X=-1}(y|-1) \Pr(X = -1) \\ &\quad + p_{Y|X=1}(y|1) \Pr(X = 1) \end{aligned} \quad (97)$$

$$p_Y(y) = p \times p_{(-A+N)}(y) \quad (98)$$

$$+ (1-p) \times p_{(A+N)}(y) \quad (99)$$

$$p_Y(y) = p \frac{e^{-\frac{(y+A)^2}{2}}}{\sqrt{2\pi}} + (1-p) \frac{e^{-\frac{(y-A)^2}{2}}}{\sqrt{2\pi}} \quad (100)$$

To use the MAP criterion, we must find $p_{X|Y}(x|y)$. To do this, we use the Theorem of Conditional Probability:

$$p_{X|Y}(x|y) = \frac{p_{Y|X}(y|x) \times p_X(x)}{p_Y(y)} \quad (101)$$

When $X = 1$, we have:

$$p_{X|Y}(1|y) = \frac{p_{Y|X}(y|1) \times p_X(1)}{p_Y(y)} \quad (102)$$

$$\begin{aligned} &= \frac{(1-p) \frac{e^{-\frac{(y-A)^2}{2}}}{\sqrt{2\pi}}}{p \frac{e^{-\frac{(y+A)^2}{2}}}{\sqrt{2\pi}} + (1-p) \frac{e^{-\frac{(y-A)^2}{2}}}{\sqrt{2\pi}}} \end{aligned} \quad (103)$$

$$= \frac{(1-p) e^{2yA}}{p + (1-p) e^{2yA}} \quad (104)$$

Similarly, when $X = -1$, we get:

$$p_{X|Y}(-1|y) = \frac{p}{p + (1-p) e^{2yA}} \quad (105)$$

Therefore, when $p_{X|Y}(1|y) > p_{X|Y}(-1|y)$, we

have:

$$\frac{(1-p) e^{2yA}}{p + (1-p) e^{2yA}} > \frac{p}{p + (1-p) e^{2yA}} \quad (106)$$

$$e^{2yA} > \frac{p}{(1-p)} \quad (107)$$

$$\implies y > \frac{1}{2A} \ln \frac{p}{(1-p)} \quad (108)$$

Therefore, when Eq. (108), we can assert that $X = 1$, and $X = -1$ otherwise. Now, consider when $p = \frac{1}{2}$. We have:

$$y > \frac{1}{2A} \ln \frac{p}{(1-p)} \quad (109)$$

$$= \frac{1}{2A} \ln 1 \quad (110)$$

$$= 0 \quad (111)$$

Therefore, when $y > 0$, we choose $X = 1$, and we choose $X = -1$ otherwise.