Class 2 · Approximating the wave function: basis sets

From LCAO to def2-QZVP: what basis functions are, how to read their names, and what they cost

Table of contents

1 Molecular orbitals from basis functions 2
1.1 The LCAO expansion 2
1.2 The variational principle and the CBS limit 3
2 Slater-type and Gaussian-type functions 3
2.1 Slater-type orbitals 3
2.2 Gaussian-type orbitals 3
2.3 The Gaussian product theorem 4
2.4 Contracted Gaussians: STO-3G 4
2.5 Cartesian and spherical d functions 5
3 Building better basis sets 6
3.1 Split valence: letting orbitals change size 6
3.2 Polarization functions: letting orbitals change shape 6
3.3 Diffuse functions: the far tail 7
4 Pople basis sets 7
5 The modern families used in this course 8
5.1 Karlsruhe def2 basis sets 8
5.2 Dunning correlation-consistent basis sets 8
5.3 Jensen polarization-consistent basis sets 9
5.4 Counting basis functions 9
6 Convergence, cost and basis-set errors 10
6.1 More functions, better results 10
6.2 Extrapolation to the CBS limit 10
6.3 The price: cost grows steeply with K 11
6.4 Two basis-set errors: BSIE and BSSE 11
7 Key points 12
8 Study questions 12
9 Further reading 13
Bibliography 13

Learning goals

After studying this chapter you should be able to:

- explain the LCAO expansion and why, by the variational principle, a bigger basis gives a lower (or equal) energy;
- compare Slater-type and Gaussian-type functions and explain why almost every program uses contracted Gaussians;
- explain what split valence (ζ), polarization and diffuse functions add, and when each matters;
- decode basis-set names such as STO-3G, 6-31G*, 6-311+G**, def2-TZVP, cc-pVQZ and aug-cc-pVTZ;
- count the basis functions of a small molecule from the shell composition of each atom;
- describe the complete-basis-set (CBS) limit, how the cost grows with the number of basis functions, and what the basis-set incompleteness error (BSIE) and basis-set superposition error (BSSE) are.

Review from P-Chem 2

- The chemical bond in H_2^+ and H_2 (chapter 12): the LCAO idea in its simplest form.
- The water molecule with a minimal basis and Hartree–Fock calculation for water (§13.1–13.2): seven basis functions, the overlap matrix and the integrals.
- The Hartree–Fock method (§11.3.1): orbitals expanded in basis functions.
- The variational method (§10.2.1).

1 Molecular orbitals from basis functions

1.1 The LCAO expansion

Class 1 ended with the first of the two approximations of every calculation: the orbitals are unknown functions of three coordinates, and a computer cannot store an arbitrary function. The solution is to write each molecular orbital ϕ_i as a **linear combination** of a finite number K of known functions χ_μ, the **basis functions**:

$$\phi_i(\mathbf{r}) = \sum_{\mu=1}^{K} c_{\mu i}\, \chi_\mu(\mathbf{r}). \tag{1}$$

The basis functions are fixed in advance; only the **coefficients** $c_{\mu i}$ are optimized. In P-Chem 2 you met this as LCAO, the *linear combination of atomic orbitals*: for H_2^+, one 1s function on each proton; for water, a 1s, 2s and three 2p functions on oxygen plus one 1s on each hydrogen, seven functions in total. Modern basis functions are not atomic orbitals in the strict sense (they are not eigenfunctions of any atom), but they are still **centered on the atoms** and look like atomic orbitals, so the name has stuck.

With Equation 1, the hard problem of finding unknown *functions* becomes the much easier problem of finding unknown *numbers*: a $K \times K$ matrix of coefficients. In Hartree–Fock and Kohn–Sham DFT this leads to a matrix eigenvalue equation, the Roothaan–Hall equation $\mathbf{FC} = \mathbf{SC}\boldsymbol{\varepsilon}$, which you will meet in Class 4. Here $\mathbf{S}$ is the **overlap matrix**, $S_{\mu\nu} = \int \chi_\mu^* \chi_\nu \, d\mathbf{r}$. Basis functions on different atoms overlap, so $\mathbf{S}$ is not the identity matrix; §13.2 of the P-Chem 2 textbook shows it for water.

K basis functions give exactly K molecular orbitals. Of these, $N/2$ are occupied in a closed-shell molecule with N electrons; the rest are **virtual** (unoccupied) orbitals. The number of basis functions K is printed in every Q-Chem output and is one of the four numbers you record in Experiment 1.

1.2 The variational principle and the CBS limit

If the set $\{\chi_\mu\}$ were **complete**, any function could be expanded exactly in it, and Equation 1 would be no approximation at all. A complete set is infinite, so any real basis is a **truncation**. The orbitals can only take the shapes that the chosen functions allow.

The variational principle of Class 1 tells us what happens as we enlarge the basis. For Hartree–Fock and for Kohn–Sham DFT with a given functional, the energy is minimized with respect to the coefficients. Adding basis functions enlarges the space of allowed orbitals, so the minimum can only go down or stay the same:

$$E(\text{basis}_1) \geq E(\text{basis}_2) \quad \text{if basis}_2 \supseteq \text{basis}_1.$$

As the basis grows toward completeness, the energy converges from above to a limit, the **complete-basis-set (CBS) limit**. The difference between the energy in a finite basis and the CBS energy (with the same method) is the **basis-set incompleteness error** (BSIE). It is the error that Experiment 1 is designed to measure.

> **i** Strictly nested or not?
>
> The inequality is guaranteed only if the bigger basis *contains* the smaller one. Real basis-set families are re-optimized at each size, so def2-TZVP does not literally contain def2-SVP. In practice the total energy still decreases monotonically along a well-designed family, which is what matters for us. Reaction energies, being differences, do not have to converge monotonically, and Experiment 1 asks you to look at exactly this.

2 Slater-type and Gaussian-type functions

What should the basis functions look like? Two choices dominate the history of quantum chemistry.

2.1 Slater-type orbitals

Slater-type orbitals (STOs) mimic the hydrogen-atom solutions:

$$\chi^{\text{STO}}(r,\theta,\varphi) = N\, r^{n-1} e^{-\zeta r}\, Y_\ell^m(\theta,\varphi). \tag{2}$$

The **exponent** ζ (zeta) sets the size: a large ζ gives a tight function close to the nucleus; a small ζ gives a diffuse one. STOs have the right physics. They have a **cusp** at the nucleus (a finite slope at $r = 0$, as every exact wave function must [1]) and an **exponential tail** at large r, like real bound states (Class 1).

Their weakness is computational. A molecular calculation needs huge numbers of integrals over basis functions, the worst being the **two-electron integrals**

$$(\mu\nu|\lambda\sigma) = \iint \frac{\chi_\mu(\mathbf{r}_1)\chi_\nu(\mathbf{r}_1)\,\chi_\lambda(\mathbf{r}_2)\chi_\sigma(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|}\, d\mathbf{r}_1 d\mathbf{r}_2, \tag{3}$$

in which the four functions may sit on four different atoms. For STOs these multicenter integrals have no closed form and must be computed numerically, which is slow.

2.2 Gaussian-type orbitals

In 1950, Boys proposed using **Gaussian-type orbitals** (GTOs) instead [2]:

$$\chi^{\text{GTO}}(x,y,z) = N\, x^a y^b z^c\, e^{-\alpha r^2}, \qquad \ell = a + b + c. \tag{4}$$

The sum $\ell = a + b + c$ sets the angular momentum: $\ell = 0$ is an s function, $\ell = 1$ a p function (x, y or z in front), $\ell = 2$ a d function, and so on. The exponent α plays the role of ζ.

GTOs have the *wrong* physics in two places, visible in Figure 2:

- **No cusp.** A Gaussian is flat at the nucleus ($d\chi/dr = 0$ at $r = 0$), and its value there is too small.
- **Wrong tail.** $e^{-\alpha r^2}$ decays much faster than $e^{-\zeta r}$. Far from the nucleus, a single Gaussian underestimates the wave function by orders of magnitude.

2.3 The Gaussian product theorem

Why use the wrong functions? Because of one beautiful property. The product of two Gaussians centered on two different points $\mathbf{A}$ and $\mathbf{B}$ is a single Gaussian centered at a point $\mathbf{P}$ on the line between them:

$$e^{-\alpha|\mathbf{r}-\mathbf{A}|^2}\, e^{-\beta|\mathbf{r}-\mathbf{B}|^2} = K_{AB}\, e^{-p|\mathbf{r}-\mathbf{P}|^2}, \qquad p = \alpha + \beta, \quad \mathbf{P} = \frac{\alpha\mathbf{A} + \beta\mathbf{B}}{\alpha + \beta}, \quad K_{AB} = e^{-\frac{\alpha\beta}{\alpha+\beta}|\mathbf{A}-\mathbf{B}|^2}. \tag{5}$$

This is the **Gaussian product theorem** (Figure 1). It collapses every pair of functions in Equation 3 to a single center, so a four-center integral becomes a two-center one, which has an analytic formula. Gaussian integrals are so cheap that it pays to use several Gaussians to imitate one STO. This trade, more functions but much faster integrals, is the reason virtually every molecular quantum-chemistry program, Q-Chem included, uses Gaussian basis functions.




Figure 1: The Gaussian product theorem in one dimension. The product (black) of a Gaussian centered at A (exponent 1.0) and one centered at B (exponent 0.6) is exactly a single Gaussian centered at P, with exponent 1.6 and height reduced by the constant K_{AB} of Equation 5.

2.4 Contracted Gaussians: STO-3G

A **contracted Gaussian** is a fixed linear combination of several **primitive** Gaussians:

$$\chi_\mu(\mathbf{r}) = \sum_{k=1}^{L} d_{k\mu}\, g_k(\alpha_k, \mathbf{r}). \tag{6}$$

The exponents α_k and the **contraction coefficients** $d_{k\mu}$ are chosen once, when the basis set is designed, and never change during a calculation. The program optimizes only the coefficients $c_{\mu i}$ of Equation 1, one per *contracted* function. Contraction gives functions with a better shape without increasing K.

The simplest example is **STO-3G** [3]: each Slater-type orbital is replaced by a contraction of three primitive Gaussians, fitted to reproduce it as closely as possible. For the hydrogen 1s function, which in STO-3G mimics a Slater function with $\zeta = 1.24$ (slightly tighter than the free-atom value of 1.0, because a hydrogen in a molecule holds its electron more tightly), the published primitives are:

Table 1: STO-3G contraction for the hydrogen 1s function (coefficients for normalized primitives).

k	exponent α_k (bohr^{-2})	coefficient d_k
1	3.42525091	0.15432897
2	0.62391373	0.53532814
3	0.16885540	0.44463454

The tight primitive ($\alpha \approx 3.4$) builds up the peak near the nucleus; the diffuse one ($\alpha \approx 0.17$) supplies the tail. Figure 2 compares the result with the Slater function it imitates and with the best *single* Gaussian (the one with maximum overlap with the STO, $\alpha = 0.417$). The single Gaussian has an overlap of 0.978 with the STO; the three-Gaussian contraction reaches 0.9998. Even so, STO-3G still misses the cusp, and its tail still falls too fast beyond about 5 bohr.




Figure 2: The hydrogen 1s Slater function with $\zeta = 1.24$ (black), the best single Gaussian (dashed red) and the STO-3G contraction of three Gaussians (blue), using the exponents and coefficients of Table 1. (a) Near the nucleus, the Slater function has a cusp; Gaussians are flat at $r = 0$. The dotted curves are the three weighted primitives that add up to the STO-3G function. (b) On a logarithmic scale, the Slater function decays as a straight line ($e^{-\zeta r}$); Gaussians bend down ($e^{-\alpha r^2}$).

The shell composition of a basis set is often written as (primitives)/[contracted functions]. For STO-3G on oxygen, (6s3p)/[2s1p]: six s primitives contracted to two s functions (1s and 2s), and three p primitives contracted to one set of p functions. A "set" of p functions means p_x, p_y and p_z, so [2s1p] is $2 + 3 = 5$ basis functions.

2.5 Cartesian and spherical d functions

Equation 4 with $\ell = 2$ gives six **Cartesian** d functions: x^2, y^2, z^2, xy, xz, yz (times the Gaussian). But a true d shell has only five components. The extra one is the combination $x^2 + y^2 + z^2 = r^2$, which is spherical: it behaves like an s function. **Spherical** (or *pure*) d functions keep only the five real combinations with $\ell = 2$ (d_{z^2}, $d_{x^2-y^2}$, d_{xy}, d_{xz}, d_{yz}). The same happens for higher ℓ:

Table 2: Number of functions per shell. Cartesian: $(\ell+1)(\ell+2)/2$; spherical: $2\ell+1$.

Shell	Cartesian components	Spherical (pure) components
s	1	1
p	3	3
d	6 ("6d")	5 ("5d")
f	10	7
g	15	9

The choice changes K, the energy (slightly) and the cost. Each basis set was designed with one convention in mind: the Pople sets such as 6-31G* were defined with six Cartesian d functions, while the def2 and correlation-consistent sets were designed for spherical functions. Programs have defaults that you can override, so **always check the number of basis functions in your output** against what you expect. This is one of the checks you will do in Experiment 1.

3 Building better basis sets

A minimal basis such as STO-3G has a fundamental limitation: each occupied atomic orbital gets exactly one function, with a fixed size and shape. In a molecule, orbitals need to change in three ways, and each one is addressed by a different kind of extra function.

3.1 Split valence: letting orbitals change size

The core orbitals (1s on C, N, O; 1s2s2p on Cl) hardly change when atoms form bonds. The **valence** orbitals change a lot: they expand or contract depending on the bonding partner and on the charge. A **split-valence** basis gives each valence orbital two or more functions with different exponents, a tight one and a loose one. The variational optimization then chooses the right mixture, effectively tuning the size of the orbital in each direction and in each molecule.

The number of functions per valence orbital is called **ζ** ("zeta", from the STO exponent):

- **minimal** (single-ζ): one function per occupied orbital (STO-3G);
- **double-ζ** (DZ): two functions per valence orbital;
- **triple-ζ** (TZ): three;
- **quadruple-ζ** (QZ): four; and so on.

In the Dunning families, ζ is labeled by the **cardinal number** n = 2 (D), 3 (T), 4 (Q), 5, 6.

Worked example: what does split valence buy?

For oxygen, a minimal basis has 1s, 2s, 2p: $1+1+3=5$ functions. A split-valence double-ζ basis keeps one core 1s function but doubles the valence: $1+2+2\times3=9$ functions (written [3s2p]). Water goes from $5+2\times1=7$ functions (minimal) to $9+2\times2=13$ (double-ζ valence on both O and H, no polarization). Splitting the valence nearly doubles the size of the basis, before adding anything else.

3.2 Polarization functions: letting orbitals change shape

An s function is spherical; a p function points along one axis. When an atom sits in a molecule, the electric field of its neighbors distorts its electron cloud: the orbitals need to bend and shift off-center. The way to allow this is to add functions with **higher angular momentum** than any occupied orbital of the free atom: p functions

on hydrogen, d functions on C, N, O and Cl, f functions on top of those, and so on. Mixing a little p into an s function shifts its center of charge; mixing d into p bends it. These are **polarization functions**.

Polarization functions are essential for almost every property: bond angles (NH_3 is too flat without them), reaction energies, dipole moments and, very importantly for this lab, the weak interactions between molecules. In larger basis sets, they come in several layers: def2-TZVP has 2d1f on oxygen; cc-pVQZ has 3d2f1g.

3.3 Diffuse functions: the far tail

Diffuse functions are extra functions with very small exponents, so they extend far from the nucleus. They are needed whenever electrons spend time far out:

- **anions**, whose extra electron is weakly bound and spread out;
- **weak intermolecular interactions** (hydrogen bonds, dispersion), which depend on the overlap of the outer tails of two molecules;
- **Rydberg excited states** and properties such as polarizabilities.

Diffuse functions are expensive and can cause numerical problems (near-linear dependence of the basis) in larger molecules. Use them when the chemistry needs them.

> **i** Three questions, three kinds of functions
>
> Size ⟶ more ζ (split valence). Shape ⟶ polarization (higher ℓ). Reach ⟶ diffuse (small α). A good basis set is **balanced**: it improves all three together rather than piling up one kind.

4 Pople basis sets

The basis sets developed by John Pople and co-workers in the 1970s and 1980s are the most widely known, and you will see their names in thousands of papers. Their names are a code for how the functions are contracted.

Table 3: Reading a Pople basis-set name. The numbers before "G" describe contractions: the digit before the hyphen is the core, the digits after it are the valence functions.

Name	Meaning
STO-3G	minimal; each STO fitted by 3 Gaussians [3]
3-21G	core: 3 primitives in 1 function; valence split in two: 2 primitives + 1 primitive [4]
6-31G	core: 6 primitives; valence: 3 + 1 (double-ζ) [5], [6]
6-311G	core: 6; valence: 3 + 1 + 1 (triple-ζ) [7]
6-31G* = 6-31G(d)	6-31G plus one set of d polarization functions on heavy (non-hydrogen) atoms [8], [9]
6-31G** = 6-31G(d,p)	as above, plus p functions on hydrogen
6-31+G*	plus diffuse s and p functions on heavy atoms
6-31++G**	plus diffuse s functions on hydrogen too
6-311G**	triple-ζ valence, d on heavy atoms, p on H

Worked example: counting functions in 6-31G*

Water. Oxygen in 6-31G: one core 1s function, two 2s functions and two sets of 2p functions, [3s2p] = $3 + 6 = 9$. The star adds one set of d functions: 6 if Cartesian (as 6-31G* was defined), 5 if spherical. Hydrogen in 6-31G has two s functions and no polarization in 6-31G* (only the second star adds p on H). Total: $9 + 6 + 2 \times 2 = 19$ (Cartesian d) or 18 (spherical d).

HCl. Chlorine in 6-31G: one 1s, then the 2s2p shell (one s + one p set), then the valence 3s3p split in two (two s + two p sets): [4s3p] = $4 + 9 = 13$; plus d: 6 or 5. Hydrogen: 2. Total: $13 + 6 + 2 = 21$ (Cartesian) or 20 (spherical). For comparison, STO-3G gives HCl only 10 functions: [3s2p] = 9 on Cl, 1 on H.

Pople sets were designed for Hartree–Fock in the era of slow computers, and they were extraordinarily successful. Their limitations are well known today: the s and p functions of each valence shell share exponents (to save time, which no longer matters), polarization is added in an ad hoc way, and the sequence 6-31G* ⟶ 6-311G** is **not designed to converge systematically** toward the CBS limit. You should recognize them, and you will meet 6-31G* again in Experiment 6, where its famous partnership with B3LYP gets a closer look. For new work, the families in the next section are better choices.

5 The modern families used in this course

5.1 Karlsruhe def2 basis sets

The **def2** basis sets of Weigend and Ahlrichs [10] (the "Ahlrichs" or "Karlsruhe" family) were designed to give consistent quality across almost the whole periodic table, H to Rn. The names describe the valence quality:

- **def2-SVP**: split valence (double-ζ) plus polarization;
- **def2-TZVP**: triple-ζ valence plus polarization;
- **def2-QZVP**: quadruple-ζ valence plus polarization.

Variants with an extra P (def2-TZVPP, def2-QZVPP) add more polarization, and variants ending in D add diffuse functions. "def2" stands for the second generation of the Karlsruhe *default* sets. For elements beyond krypton, def2 replaces the inner-core electrons with an **effective core potential** (ECP): a fixed potential that represents the core and includes the scalar-relativistic effects that matter for heavy atoms. Only the outer electrons are treated explicitly. You will use this in Experiments 8 and 10 for ruthenium and palladium.

The def2 family has been found to be among the most cost-effective choices for DFT [11], which is one reason it is the default basis of most experiments in this lab.

5.2 Dunning correlation-consistent basis sets

The **correlation-consistent** basis sets of Dunning and co-workers are written **cc-pVnZ**: *correlation-consistent polarized valence n-zeta*, with n = D, T, Q, 5, 6 [12], [13]. They were built for correlated wave-function methods (MP2, CCSD(T)). The design rule is that functions are added in groups that each recover a similar amount of **correlation energy**. Going from cc-pVDZ to cc-pVTZ adds one more s, p and d function *and* the first f function on oxygen, because these contribute comparably. The result is a family that converges **systematically** toward the CBS limit, so results at several n can be extrapolated (Section 6.2).

The prefix **aug-** adds one diffuse function of *each* angular momentum present [14]: aug-cc-pVTZ is the standard choice for anions and weak interactions with correlated methods. (Other prefixes, such as cc-pCVnZ for core correlation, exist but are not used here.)

5.3 Jensen polarization-consistent basis sets

Jensen's **pc-n** basis sets (pc-0, pc-1, pc-2, pc-3, pc-4) apply the same systematic idea, but are optimized for Hartree–Fock and DFT instead of correlated methods [15]. pc-1 is double-ζ quality. You will use pc-1 in Experiment 6 to compare double-ζ basis sets from different families.

5.4 Counting basis functions

The size of every basis set follows from the shell composition of each atom. With spherical (pure) functions, each shell contributes $2\ell + 1$ functions: s = 1, p = 3, d = 5, f = 7, g = 9.

Table 4: Number of basis functions for water, from the published shell compositions (Basis Set Exchange). Spherical (5d, 7f, 9g) functions unless noted.

Basis set	ζ	H	O	K for H_2O
STO-3G	1	[1s] = 1	[2s1p] = 5	7
6-31G* (5d)	2	[2s] = 2	[3s2p1d] = 14	18 (19 with 6d)
6-311G** (5d)	3	[3s1p] = 6	[4s3p1d] = 18	30 (31 with 6d)
def2-SVP	2	[2s1p] = 5	[3s2p1d] = 14	24
def2-TZVP	3	[3s1p] = 6	[5s3p2d1f] = 31	43
def2-QZVP	4	[4s3p2d1f] = 30	[7s4p3d2f1g] = 57	117
cc-pVDZ	2	[2s1p] = 5	[3s2p1d] = 14	24
cc-pVTZ	3	[3s2p1d] = 14	[4s3p2d1f] = 30	58
cc-pVQZ	4	[4s3p2d1f] = 30	[5s4p3d2f1g] = 55	115
aug-cc-pVTZ	3	[4s3p2d] = 23	[5s4p3d2f] = 46	92
pc-1	2	[2s1p] = 5	[3s2p1d] = 14	24

Worked example: def2-TZVP versus cc-pVTZ for water

def2-TZVP. Oxygen [5s3p2d1f]: $5 \times 1 + 3 \times 3 + 2 \times 5 + 1 \times 7 = 5 + 9 + 10 + 7 = 31$. Hydrogen [3s1p]: $3 + 3 = 6$. Water: $31 + 2 \times 6 = 43$.

cc-pVTZ. Oxygen [4s3p2d1f]: $4 + 9 + 10 + 7 = 30$. Hydrogen [3s2p1d]: $3 + 6 + 5 = 14$. Water: $30 + 2 \times 14 = 58$.

Both are "triple-ζ", but they distribute the functions differently. cc-pVTZ gives hydrogen much more polarization (2p1d instead of 1p), because correlated methods need it. def2-TZVP puts more s functions on oxygen. Same label, different size: the name tells you the *quality class*, not the cost.

Notice three things in Table 4. First, double-ζ sets from different families can have the same size (24 for def2-SVP, cc-pVDZ and pc-1 here) yet different exponents and different results. Second, the size grows quickly with ζ: about a factor of 2 from double- to triple-ζ, and another factor of 2–3 to quadruple-ζ. Third, hydrogen atoms are cheap at double-ζ but not at quadruple-ζ (30 functions each).

6 Convergence, cost and basis-set errors

6.1 More functions, better results

The central message of this chapter fits in one sentence: **in a systematic family, more functions give results that converge toward the CBS limit.** Figure 3 (a) sketches the typical behavior of the total energy. The first step, double- to triple-ζ, is large; each further step is smaller.

For Hartree–Fock and DFT, the total energy converges *fast*, roughly exponentially with the cardinal number [16]:

$$E(n) = E_{\mathrm{CBS}} + A\, e^{-\alpha n}. \tag{7}$$

The **correlation energy** of wave-function methods converges much more slowly, as an inverse power [17], [18]:

$$E_{\mathrm{corr}}(n) = E_{\mathrm{corr,CBS}} + A\, n^{-3}. \tag{8}$$

The reason is physical. Near the point where two electrons meet, the exact wave function has a cusp in the inter-electron distance r_{12}, and smooth orbital products converge to it only slowly. HF and DFT have no such cusp to describe, so they converge faster.




Figure 3: (a) **Schematic** convergence of a variational energy with the cardinal number n of a systematic basis-set family, toward the CBS limit (Equation 7; the curve is illustrative, not data). (b) Real numbers of basis functions K for water in the def2 and cc-pVnZ families (Table 4), with the formal K^4 cost relative to the double-ζ basis written above each bar.

6.2 Extrapolation to the CBS limit

Because the convergence is systematic, two or three results at consecutive n can be fit to Equation 7 or Equation 8 to estimate the CBS limit without ever computing it. For the n^{-3} form, two points $m < n$ give

$$E_{\mathrm{CBS}} = \frac{n^3 E_n - m^3 E_m}{n^3 - m^3}. \tag{9}$$

Worked example: two-point extrapolation

Suppose (hypothetical numbers) the MP2 correlation energy of a molecule is $-0.2950\ E_\mathrm{h}$ with cc-pVTZ ($m = 3$) and $-0.3080\ E_\mathrm{h}$ with cc-pVQZ ($n = 4$). Then

$$E_{\mathrm{corr,CBS}} = \frac{64(-0.3080) - 27(-0.2950)}{64 - 27} = \frac{-19.712 + 7.965}{37} = -0.3175\ E_\mathrm{h}.$$

The extrapolated value lies $0.0095\ E_\mathrm{h}$ (6 kcal/mol) below the QZ result: the correlation energy is far from converged even at quadruple-ζ. This is why reference values such as those you compare with in this lab are usually CCSD(T)/CBS estimates.

The optional Colab notebook of Experiment 1 lets you try an extrapolation on the HCl dimer yourself.

6.3 The price: cost grows steeply with K

Better results cost more. The number of two-electron integrals (Equation 3) grows as K^4: with permutational symmetry there are about $K^4/8$ unique ones. For water, that is about 41,000 with def2-SVP ($K = 24$), 430,000 with def2-TZVP (43), and 23 million with def2-QZVP (117). Going from double- to quadruple-ζ multiplies the formal cost by about 565 (Figure 3 b).

This K^4 is the *formal* scaling of Hartree–Fock and hybrid DFT. In practice, programs do better: integrals between functions far apart are negligible and are skipped (screening), and approximations such as density fitting reduce the work. For small molecules, fixed overheads (starting the program, reading files) also blur the picture. The *measured* exponent is therefore lower than 4, and it depends on the method and the size of the system. In Experiment 1 you will measure it yourself from your own wall times.

! The 10-minute limit is a basis-set limit

On the free IQmol server every job must finish in 10 minutes. For the molecules in this lab, the basis set is usually what decides whether a job fits. A one-character typo, such as def2-QZVP instead of def2-SVP, can multiply the cost by hundreds and get your job killed.

6.4 Two basis-set errors: BSIE and BSSE

- **Basis-set incompleteness error (BSIE).** The difference between a result in a finite basis and the CBS limit, with the same method. It affects every calculation. It shrinks as the basis grows, and it is what Experiment 1 measures for a reaction energy.
- **Basis-set superposition error (BSSE).** A specific artifact of incomplete basis sets in systems made of fragments, such as a dimer. In the dimer, each monomer can "borrow" the basis functions centered on its partner to improve its own description, lowering its energy. The isolated monomers have no partner functions to borrow. The dimer is therefore artificially stabilized, and the interaction energy is too large in magnitude. BSSE also vanishes in the CBS limit. The standard correction, the counterpoise method of Boys and Bernardi [19], recomputes each monomer in the full dimer basis. We return to BSSE in Classes 8 and 10.

Practical rules of thumb

- Double-ζ (def2-SVP, cc-pVDZ) is fine for quick tests and geometries of large systems, but rarely converged for energies.
- Triple-ζ (def2-TZVP) is the usual compromise between accuracy and cost for DFT, and it is the default basis of Experiments 3–5 and 9.
- Quadruple-ζ is close to the CBS limit for HF and DFT, and expensive.
- Add diffuse functions for anions and when weak interactions matter most.
- Report the basis set exactly, including stars, plus signs and prefixes: "6-31G" and "6-31G*" are different basis sets.

7 Key points

- Molecular orbitals are expanded in a finite set of K atom-centered basis functions, $\phi_i = \sum_\mu c_{\mu i}\chi_\mu$. Only the coefficients are optimized.
- For HF and DFT, a bigger basis gives a lower (or equal) energy; the limit is the complete-basis-set (CBS) limit, and the difference is the basis-set incompleteness error.
- Slater functions have the correct cusp and tail but expensive integrals; Gaussians have neither, but the Gaussian product theorem makes their integrals fast. Contracted Gaussians combine several primitives to imitate one Slater function (STO-3G).
- Cartesian d shells have 6 components and spherical ones 5 (f: 10 vs 7). The convention changes K; check it in the output.
- Split valence (more ζ) lets orbitals change size; polarization (higher ℓ) lets them change shape; diffuse functions (small α) extend their reach.
- Pople names encode contractions (6-31G*, 6-311+G**). Pople sets are historical and not designed to converge systematically.
- def2-SVP/TZVP/QZVP cover almost the whole periodic table, with ECPs beyond Kr. cc-pVnZ (n = D, T, Q, 5) converges systematically for correlated methods; aug- adds diffuse functions. pc-n is optimized for DFT.
- The number of basis functions follows from the shell composition: for water, 24 (def2-SVP, cc-pVDZ), 43 (def2-TZVP), 58 (cc-pVTZ), 117 (def2-QZVP), 115 (cc-pVQZ).
- The formal cost of HF and hybrid DFT grows as K^4; the measured exponent is lower. HF/DFT energies converge roughly exponentially with n; correlation energies as n^{-3}.
- BSSE makes complexes artificially too stable in small basis sets; it vanishes at the CBS limit.

8 Study questions

1. Explain in your own words why the energy cannot increase when functions are added to a basis set (for HF or DFT). Why is this guarantee weaker when comparing def2-SVP and def2-TZVP, and why does it not apply to reaction energies?
2. List two physical shortcomings of Gaussian functions compared with Slater functions, and the one mathematical property that makes Gaussians the standard anyway.
3. Using Equation 5, find the center P and the exponent p of the product of an s Gaussian with $\alpha = 2.0$ at $x = 0$ and one with $\beta = 0.5$ at $x = 1.0$ bohr. Which original Gaussian is P closer to, and why?
4. What is the difference between a primitive and a contracted Gaussian? In STO-3G, how many primitives and how many contracted functions does a hydrogen atom have? How many coefficients $c_{\mu i}$ does the SCF optimize for an H atom?
5. Decode each name: 3-21G, 6-31+G**, 6-311++G(d,p), def2-TZVPP, aug-cc-pVQZ. For each, say whether it has polarization functions on H, and whether it has diffuse functions.

6. Count the basis functions of HCl in def2-SVP from the shell compositions H [2s1p] and Cl [4s3p1d], with spherical functions. Then predict K for the HCl dimer. (Check your answer against your first Experiment 1 output.)
7. Count the basis functions of methane (CH_4) in cc-pVDZ, cc-pVTZ and def2-TZVP. Carbon has the same shell composition as oxygen in Table 4.
8. A water calculation reports 25 basis functions with 6-31G**. Another program reports 24 for the same basis. Explain the difference.
9. If the time of a calculation grew exactly as K^4, how much longer would a water job take with def2-QZVP than with def2-TZVP? Give two reasons why the real ratio is smaller.
10. Why does BSSE make the binding energy of a dimer too large in a small basis set, rather than too small? Why does it disappear at the CBS limit?

9 Further reading

- F. Jensen, "Atomic orbital basis sets" [20], and B. Nagy and F. Jensen, "Basis sets in quantum chemistry" [21]: modern, readable reviews of everything in this chapter.
- A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry* [22], §3.6: the classic derivation of STO-nG and the minimal-basis H_2 calculation.
- T. Helgaker, P. Jørgensen and J. Olsen, *Molecular Electronic-Structure Theory* [23], chapters 6 and 8: Gaussian functions, basis-set design and convergence in depth.
- F. Weigend and R. Ahlrichs [10]: the original def2 paper, with tables of the shell compositions for every element.
- The Basis Set Exchange lists the exponents and coefficients of nearly every published basis set; you can look up any of the sets in Table 4 there.

Bibliography

[1] T. Kato, "On the eigenfunctions of many-particle systems in quantum mechanics," *Commun. Pure Appl. Math.*, vol. 10, pp. 151–177, 1957, doi: 10.1002/cpa.3160100201.

[2] S. F. Boys, "Electronic wave functions. I. A general method of calculation for the stationary states of any molecular system," *Proc. R. Soc. Lond. A*, vol. 200, pp. 542–554, 1950, doi: 10.1098/rspa.1950.0036.

[3] W. J. Hehre, R. F. Stewart, and J. A. Pople, "Self-consistent molecular-orbital methods. I. Use of Gaussian expansions of Slater-type atomic orbitals," *J. Chem. Phys.*, vol. 51, pp. 2657–2664, 1969, doi: 10.1063/1.1672392.

[4] J. S. Binkley, J. A. Pople, and W. J. Hehre, "Self-consistent molecular orbital methods. 21. Small split-valence basis sets for first-row elements," *J. Am. Chem. Soc.*, vol. 102, pp. 939–947, 1980, doi: 10.1021/ja00523a008.

[5] R. Ditchfield, W. J. Hehre, and J. A. J. Pople, *Chem. Phys.*, vol. 54, pp. 724–728, 1971.

[6] W. J. Hehre, R. Ditchfield, and J. A. J. Pople, *Chem. Phys.*, vol. 56, pp. 2257–2261, 1972.

[7] R. Krishnan, J. S. Binkley, R. Seeger, and J. A. Pople, "Self-consistent molecular orbital methods. XX. A basis set for correlated wave functions," *J. Chem. Phys.*, vol. 72, pp. 650–654, 1980, doi: 10.1063/1.438955.

[8] P. C. Hariharan and J. A. Pople, *Theor. Chim. Acta*, vol. 28, pp. 213–222, 1973.

[9] M. M. Francl *et al.*, *Chem. Phys.*, vol. 77, pp. 3654–3665, 1982.

[10] F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, vol. 7, pp. 3297–3305, 2005.

[11] J. Witte, J. B. Neaton, and M. J. Head-Gordon, *Chem. Phys.*, vol. 144, p. 194306, 2016.

[12] T. H. J. Dunning, *Chem. Phys.*, vol. 90, pp. 1007–1023, 1989.

[13] D. E. Woon and T. H. J. Dunning, *Chem. Phys.*, vol. 98, pp. 1358–1371, 1993.

[14] R. A. Kendall, J. Dunning T. H., and R. J. Harrison, "Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions," *J. Chem. Phys.*, vol. 96, pp. 6796–6806, 1992, doi: 10.1063/1.462569.

[15] F. J. Jensen, *Chem. Phys.*, vol. 115, pp. 9113–9125, 2001.

[16] D. Feller, "Application of systematic sequences of wave functions to the water dimer," *J. Chem. Phys.*, vol. 96, pp. 6104–6114, 1992, doi: 10.1063/1.462652.

[17] T. Helgaker, W. Klopper, H. Koch, and J. Noga, "Basis-set convergence of correlated calculations on water," *J. Chem. Phys.*, vol. 106, pp. 9639–9646, 1997, doi: 10.1063/1.473863.

[18] A. Halkier *et al.*, "Basis-set convergence in correlated calculations on Ne, N_2, and H_2O", *Chem. Phys. Lett.*, vol. 286, pp. 243–252, 1998, doi: 10.1016/S0009-2614(98)00111-0.

[19] S. F. Boys and F. Bernardi, "The calculation of small molecular interactions by the differences of separate total energies. Some procedures with reduced errors," *Mol. Phys.*, vol. 19, pp. 553–566, 1970, doi: 10.1080/00268977000101561.

[20] F. Jensen, *WIREs Comp. Mol. Sci.*, vol. 3, pp. 273–295, 2013.

[21] B. Nagy and F. Jensen, "Basis Sets in Quantum Chemistry," in *Reviews in Computational Chemistry*, A. L. Parrill and K. B. Lipkowitz, Eds., Hoboken, NJ: John Wiley & Sons, 2017, pp. 93–149.

[22] A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory*. Mineola, NY: Dover, 1996.

[23] T. Helgaker, P. Jørgensen, and J. Olsen, *Molecular Electronic-Structure Theory*. Chichester: Wiley, 2000.