

Math 4317 (Prof. Swiech, S'18): HW #4

Peter Williams

4/15/2018

Section 20

A. Prove that if f is defined for $x \geq 0$ by $f(x) = \sqrt{x}$, then f is continuous at every point of its domain.

For $f(x) = \sqrt{x}$, $\mathcal{D}(f) = \{x \in \mathbb{R} : x \geq 0\}$, let $a \in \mathcal{D}(f)$.

When $a = 0$, $|f(x) - f(a)| = |\sqrt{x} - 0| = \sqrt{x} < \varepsilon$. If we let $\delta(\varepsilon) = \varepsilon^2$, when $x < \varepsilon^2$, $|f(x)| < \varepsilon$.

When $a \neq 0$, $|f(x) - f(a)| = |\sqrt{x} - \sqrt{a}| = \frac{|\sqrt{x} - \sqrt{a}|}{|\sqrt{x} + \sqrt{a}|} |\sqrt{x} + \sqrt{a}| = \frac{|x - a|}{|\sqrt{x} + \sqrt{a}|} < \frac{|x - a|}{\sqrt{a}} < \varepsilon \implies$ when $|x - a| < \varepsilon\sqrt{a}$, then, $|f(x) - f(a)| < \varepsilon$, thus we can choose $\delta(\varepsilon) = \varepsilon\sqrt{a} \implies f$ is continuous at every point in its domain.

B. Show that a “polynomial function”; that is, a function f with the form $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, $x \in \mathbb{R}$ is continuous at every point of \mathbb{R} .

Relying on the properties of algebraic combinations of continuous functions, we construct f as a combination of continuous functions to show its continuity. Considering the last term of the polynomial function, denoted here, $f_0(x) = a_0$, $f_0(x)$ is a continuous, constant function, since, for any $a \in \mathbb{R}$ we have $|f_0(x) - f_0(a)| = |a_0 - a_0| < \varepsilon = \delta(\varepsilon)$, $\varepsilon > 0$. We consider the second to last term of f , $a_1 x$, as a constant, a_1 multiplied by the identity function, denoted, $f_1(x) = x$. Since $f_1(x) = x$, for any real number $a \in \mathbb{R}$, we have $|f_1(x) - f_1(a)| = |x - a| < \varepsilon = \delta(\varepsilon)$, $\varepsilon > 0 \implies a_1 f_1(x) = a_1 x$ is continuous.

Relying on the continuity of $f_1(x) = x$ multiplied by any constant, we can construct higher order terms of f through repeated multiplication of $f_1(x)$, e.g. $a_2 \cdot f_1(x) \cdot f_1(x) = a_2 x^2$ and $a_n \prod_{j=1}^n f_1(x) = a_n \cdot f_1(x) \cdot f_1(x) \cdot \dots \cdot f_1(x) = a_n x^n$, and so on, where each term constructed $a_n x^n$ is continuous on \mathbb{R} since it is constructed via algebraic combinations of continuous functions $\implies f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, is continuous at every point $x \in \mathbb{R}$.

E. Let f be the function on $\mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x$, x irrational, $f(x) = 1 - x$, x rational. Show that f is continuous at $x = \frac{1}{2}$ and discontinuous elsewhere.

Considering the point $a = \frac{1}{2}$, we have $f(a) = \frac{1}{2}$, and $|f(x) - f(a)| = |1 - x - \frac{1}{2}| = |\frac{1}{2} - x| = |x - a| < \varepsilon = \delta(\varepsilon)$. So if $|f(x) - f(a)| < \varepsilon = \delta(\varepsilon) > 0 \implies |x - a| < \delta(\varepsilon)$, and then we have f continuous at the point $a = \frac{1}{2}$. For the case $a \neq \frac{1}{2}$, a irrational, take a sequence $X = (x_n)$ of rational numbers converging to a . Since the sequence $(f(x_n))$ converges to $1 - a$, and we have $f(a) = a$, f is not continuous at irrational points by the Discontinuity Criterion. For the case $a \neq \frac{1}{2}$, a rational, take a sequence $Y = (Y_n)$ of irrational numbers converging to a , the sequence $(f(y_n))$ converges to a , but $f(a) = 1 - a$, which equation is only satisfied when $a = \frac{1}{2}$, thus f is not continuous for rational numbers at any point other than $\frac{1}{2}$.

F. Let f be continuous on $\mathbb{R} \rightarrow \mathbb{R}$. Show that if $f(x) = 0$ for rational x , then $f(x) = 0$ for all $x \in \mathbb{R}$.

Every real point, $x \in \mathbb{R}$ is the limit of a sequence of rational numbers. If f is continuous \implies for a sequence of rational numbers $X = (x_n) \rightarrow x$, we have $(f(x_n)) = 0$, for all $n \in \mathbb{N}$. Since f is continuous at each rational point $x \in \mathbb{R}$, we can find $|f(x_n) - f(x)| < \varepsilon$, $\varepsilon > 0$, and $|x_n - a| < \delta(\varepsilon) \implies (f(x_n)) \rightarrow f(x) = 0, \forall x \in \mathbb{R}$.


I. Using the results of the preceding exercise, show that the function g , defined on $\mathbb{R} \rightarrow \mathbb{R}$ by $g(x) = x \sin(\frac{1}{x})$, $x \neq 0$, $g(x) = 0$, $x = 0$ is continuous at every point. Sketch a graph of this function.

For the case $a = 0$, we have $|g(x) - g(a)| = |x \sin \frac{1}{x} - 0| = |x| |\sin \frac{1}{x}| \leq |x| \cdot 1 < \varepsilon$, $\varepsilon > 0$, since $-1 \leq \sin \frac{1}{x} \leq 1$. So when $|g(x) - g(0)| < \varepsilon = \delta(\varepsilon)$, we then have $|x| = |x - 0| < \delta(\varepsilon) \implies g$ continuous at 0.

For the case $a \neq 0$, we have $|g(x) - g(a)| = |x \sin \frac{1}{x} - a \sin \frac{1}{a}| = |x \sin \frac{1}{x} - a \sin \frac{1}{a} - a \sin \frac{1}{x} + a \sin \frac{1}{x}| = |(x - a)(\sin \frac{1}{x}) + a(\sin \frac{1}{x} - \sin \frac{1}{a})| \leq |x - a| |\sin \frac{1}{x}| + |a| |\sin \frac{1}{x} - \sin \frac{1}{a}|$, by Triangle Inequality. Since both $|\sin \frac{1}{x}| \leq 1$ and $|\sin \frac{1}{x} - \sin \frac{1}{a}| \leq 1$, we have $|x - a| |\sin \frac{1}{x}| + |a| |\sin \frac{1}{x} - \sin \frac{1}{a}| \leq |x - a| \cdot 1 + |a| \cdot 1 = |x - a| + |a| < \varepsilon$.

It then follows that if $\delta(\varepsilon) = \varepsilon - |a|$, i.e. $\varepsilon > \delta(\varepsilon) + |a|$, when $|g(x) - g(a)| < \varepsilon$, then $|x - a| < \delta(\varepsilon) \implies g$ continuous at every point in \mathbb{R} .

Sketch of function below:



N. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ satisfy the relation $g(x + y) = g(x)g(y)$, $x, y \in \mathbb{R}$. Show that if g is continuous at $x = 0$, then g is continuous at every point. Also if $g(a) = 0$ for some $a \in \mathbb{R}$, then $g(x) = 0$ for all $x \in \mathbb{R}$.

If g is continuous at $x = 0 \implies g(x + y) = g(y) = g(0) \cdot g(y)$. This implies also that $g(0)g(y) = g(y) \implies g(0)g(y) - g(y) = 0 = g(y)(g(0) - 1) = 0 \implies g(0) = 1$, or that $g(0) = 0$.

If $g(0) = 0 \implies -g(y) = 0 = g(y)$. In this case then $g(y) = 0, \forall y \in \mathbb{R} \implies g(x) = 0, \forall x \in \mathbb{R}$.

On the other hand if $g(0) = 1, \implies g(0) \cdot g(y) = g(y)$ continuous for every point $y \in \mathbb{R}$.

Section 21

I. Let g be a linear function from $\mathbb{R}^p \rightarrow \mathbb{R}^q$. Show that g is one-one and only if $g(x) = 0$ implies that $x = 0$. Since g is linear \implies for $x, y \in \mathbb{R}^p$, $g(x + y) = g(x) + g(y)$. Then if $g(x) = 0 \implies g(x + y) = 0 + g(y) = g(y) \implies g(x + y) = g(y) \implies g(x + y) = g(0 + y) = g(y)$ which implies $x = 0$. If we assume that g is one-one, then for any $g(x) = g(y) \implies x = y$. So in the case $g(x) = 0$, and $g(x + y) = g(x) + g(y) = 0 + g(y)$. Since $g(x) + g(y) = g(y) \implies g(y) - g(x) = g(y) \implies x + y = x - y$, which is satisfied when $x = 0$.

J. If h is a one-one linear function from $\mathbb{R}^p \rightarrow \mathbb{R}^p$, show that the inverse function h^{-1} is a linear function from $\mathbb{R}^p \rightarrow \mathbb{R}^p$.

Since h is one-one \implies if $h(x_1) = h(x_2)$, $x_1 = x_2$, $x_1, x_2 \in \mathbb{R}^p$. Extending the linear case, we have if $h(ax + by) = h(ax_1 + by_1) = ah(x) + bh(y) = ah(x_1) + bh(y_1)$ then $ax_1 + by_1 = ax + by$. By definition $h^{-1} = \{ax + by : h(ax + by) \in \mathbb{R}^p\} = \{ax : h(ax) \in \mathbb{R}^p\} + \{by : h(by) \in \mathbb{R}^p\}$. This implies $h^{-1}(ax + by) = h^{-1}(h(ax)) + h^{-1}(h(by)) \implies h^{-1}$ is linear, and $h^{-1} : \mathbb{R}^p \rightarrow \mathbb{R}^p$, since $h^{-1}(h(ax)) + h^{-1}(h(by)) = ax + by \in \mathbb{R}^p$ by construction.

K. Show that the sum and the composition of two linear functions are linear functions.

By definition a function is linear if $f(ax + by) = af(x) + bf(y)$, $a, b \in \mathbb{R}$, $x, y \in \mathbb{R}^p$.

For the sum of two linear functions we then have $(f + g)(ax + by) = f(ax + by) + g(ax + by) = af(x) + bf(y) + ag(x) + bg(y) = a(f(x) + g(x)) + b(f(y) + g(y)) = a(f + g)(x) + b(f + g)(y) \implies$ linearity. For the composition of two linear functions we have $f \circ g(ax + by) = f(g(ax + by)) = f(ag(x) + bg(y)) = af(g(x)) + bf(g(y)) = a(f \circ g)(x) + b(f \circ g)(y) \implies$ composition of two linear functions is linear.

L. If f is a linear map on $\mathbb{R}^p \rightarrow \mathbb{R}^q$, define $\|f\|_{pq} = \sup\{\|f(x)\| : x \in \mathbb{R}^p, \|x\| \leq 1\}$. Show that the mapping $f \rightarrow \|f\|_{pq}$ defines a norm on the vector space $\mathcal{L}(\mathbb{R}^p, \mathbb{R}^q)$ of all linear functions on $\mathbb{R}^p \rightarrow \mathbb{R}^q$. Show that $\|f(x)\| \leq \|f\|_{pq}\|x\|$ for all $x \in \mathbb{R}^p$.

We have $x = (x_1, x_2, \dots, x_p) \in \mathbb{R}^p$, $f(x) = y = (y_1, y_2, \dots, y_q) \in \mathbb{R}^q$, and matrix $A_{q \times p} = (c_{ij})$, $1 \leq i \leq q$, $1 \leq j \leq p$, with

$$y_1 = c_{11}x_1 + c_{12}x_2 + \dots + c_{1p}x_p$$

...

$$y_q = c_{q1}x_1 + c_{q2}x_2 + \dots + c_{qp}x_p$$

We then have $\|f(x)\| = \|(y_1, \dots, y_q)\| = \sqrt{y_1^2 + \dots + y_q^2}$. To show $\|f\|_{qp} = \sup\{\|f(x)\| : x \in \mathbb{R}^p, \|x\| \leq 1\}$ is a norm in $\mathcal{L}(\mathbb{R}^p, \mathbb{R}^q)$, we have (i) $\|f\|_{pq} \geq 0$, $x \in \mathbb{R}^p$? Since each element in $\|f(x)\| = \sqrt{y_1^2 + \dots + y_q^2}$, $y_j^2 \geq 0$, $\forall j = 1, \dots, q \implies \sup\{\|f(x)\|\} \geq 0 \forall x \in \mathbb{R}^p$ since by definition, $\sup\{\|f(x)\|\} \geq \|f(x)\| \forall x \in \mathbb{R}^p \implies \|f\|_{pq} \geq 0$.

(ii) $\|f\|_{pq} = 0 \iff f(x) = 0$? Since $\|f(x)\| = \|y\| = \sqrt{y_1^2 + \dots + y_q^2} = 0 \implies$ each $y_j^2 = 0, \forall j = 1, \dots, q$

(iii) $\sup\|af(x)\| = |a|\sup\|f(x)\| = |a|\|f\|_{qp}$, $a \in \mathbb{R}$? We have $\|af(x)\| = \|ay\| = \sqrt{a^2y_1^2 + \dots + a^2y_q^2} = \sqrt{a^2}\|y\| = |a|\|y\|$, and $|a| > 0 \implies \sup\{\|af(x)\|\} = \sup\{|a|\|f(x)\|\} = |a|\sup\{\|f(x)\|\}$.

(iv) $\sup\{\|f(x+x')\|\} \leq \sup\|f(x)\| + \sup\|f(x')\|$, $x, x' \in \mathbb{R}^p$? Since f is linear $\|f(x+x')\| = \|f(x) + f(x')\| \leq \|f(x)\| + \|f(x')\|$, $\forall x, x' \in \mathbb{R}^p$ by Triangle Inequality, then $\sup\{\|f(x) + f(x')\|\} \leq \sup\{\|f(x)\|\} + \sup\{\|f(x')\|\}$. This implies $\|f\|_{qp}$ is a norm.

To show $\|f(x)\| \leq \|f\|_{pq}\|x\|$, we use the earlier notation for a linear map, $f(x) = Ax$, where, $A_{q \times p} = (c_{ij})$. Thus $\|f(x)\| = \|Ax\| \leq \|A\|\|x\|$ as shown in (21.5). This implies $\sup\{\|f(x)\| : x \in \mathbb{R}^p, \|x\| \leq 1\} = \sup\{\|Ax\|\} \leq \sup\{\|A\|\|x\|\}$ which is achieved when x is the max value in its domain, i.e. $\|x\| = 1$. This implies $\sup\{\|Ax\|\|x\|\} = \sup\{\|f(x)\|\|x\|\} = \sup\{\|f(x)\|\} \cdot 1$. This implies $\|f(x)\| \leq \sup\{\|f(x)\| : x \in \mathbb{R}^p, \|x\| \leq 1\}\|x\| \forall x \in \mathbb{R}^p$.

Section 22

B. Let $H : \mathbb{R} \rightarrow \mathbb{R}$ be defined by, $h(x) = 1, 0 \leq x \leq 1$. $h(x) = 0$, otherwise. Exhibit an open set G such that $h^{-1}(G)$ is not open in \mathbb{R} , and a closed set F , such that $h^{-1}(F)$ is not closed in \mathbb{R} .

If we take $G = (0, 2)$, and open set, $h^{-1}(G) = \{x \in \mathcal{D}(f) : h(x) \in G\} = [0, 1]$, a closed set. If we take $F = [-2, 0]$, a closed set, the inverse image, $h^{-1}(F) = \{x \in \mathcal{D}(f) : h(x) \in F\}$ is the union of two open sets $(-\infty, 0) \cup (1, +\infty)$ which is open.

C. If f is bounded and continuous on $\mathbb{R}^p \rightarrow \mathbb{R}$ and if $f(x_0) > 0$, show that f is strictly positive on some neighborhood of x_0 . Does the same conclusion hold if f is merely continuous at x_0 ?

f is bounded and continuous which implies $0 < f(x_0) < M$, for some $M > 0$. Since f is continuous, for each point $a \in \mathcal{D}(f)$, there is a neighborhood V of $f(a)$ and a neighborhood $U(a) \cap D$ such that if $f(a) \in V \implies a \in U(a)$. Since $f(a) > 0 \implies$ we can take a neighborhood V of $f(a)$ that is also strictly positive, i.e. $V = \{y \in \mathbb{R} : 0 < y < M\}$. If f is not bounded the same argument can be made with $V = \{y \in \mathbb{R} : y > 0\}$.

F. A subset $D \subseteq \mathbb{R}^p$ is disconnected if and only if there exists a continuous function $f : D \rightarrow \mathbb{R}$ such that $f(D) = \{0, 1\}$.

$\rightarrow D$ disconnected implies there exists two open sets B, C such that $B \cap D$ and $C \cap D$ are disjoint and $(B \cap D) \cup (C \cap D) = D$. We can then construct a function f on D , $f(x) = 1, x \in (B \cap D)$, $f(x) = 0, x \in (C \cap D)$. \leftarrow Let $f : D \rightarrow \mathbb{R}$ be such that $f(D) = \{0, 1\} \implies$ the inverse image $f^{-1}(\{0, 1\}) = \{x \in D \subseteq f(x) \in \{0, 1\}\}$ could consist of two disjoint open sets such for f on D , $f(x) = 1, x \in (B \cap D)$, $f(x) = 0, x \in (C \cap D)$, where $D = (B \cap D) \cup (C \cap D) \subseteq \mathcal{D}(f) \implies$ there exists a continuous function $f : D \rightarrow \mathbb{R}$ such that $f(D) = \{0, 1\}$.

H. Let f, g_1, g_2 be related by the formulas in the preceding exercise. Show that from the continuity of g_1 and g_2 at $t = 0$ one cannot prove the continuity of f at $(0, 0)$.

Considering g_1, g_2 which are valid restrictions of the domain of f , given $x = (x_1, x_2) \in \mathbb{R}^2$, we can construct $f(x) = 0, x_1 \cdot x_2 = 0, f(x) = 1, x_1 \cdot x_2 \neq 0$. With this f we have $\lim_{x \rightarrow (0,0)} f(x) \neq 0$, and $f((0,0)) = 0 \implies$ discontinuity for f at $(0, 0)$. Therefore continuity for g_1, g_2 on restrictions of $\mathcal{D}(f)$ does not imply continuity of f .

K. Give an example of a bounded and continuous function g on $\mathbb{R} \rightarrow \mathbb{R}$ which does not take on either of the numbers $\sup\{g(x) : x \in \mathbb{R}\}$ or $\inf\{g(x) : x \in \mathbb{R}\}$

If we take $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x, x \in (0, 1) \subseteq \mathbb{R}$, the function is bounded above by 1, below by 0, and continuous on $(0, 1)$, but $f(x) \neq 1 = \sup\{f(x) : x \in (0, 1)\}$, and $f(x) \neq 0 = \inf\{f(x) : x \in (0, 1)\}$ for any x in interval $(0, 1)$.

O. Let f be a continuous function on $\mathbb{R} \rightarrow \mathbb{R}$ which is strictly increasing (in the sense that if $x' < x''$ then $f(x') < f(x'')$). Prove that f is injective and that its inverse function is continuous and strictly increasing.

For points $x, a, b \in \mathcal{D}(f)$, by f be strictly increasing, we have $a > b \implies f(a) > f(b), a = b \implies f(a) = f(b)$ and $a < b \implies f(a) < f(b)$. If we take point x to be $a < x < b$, we can define two neighborhoods $(a, b) \subseteq \mathcal{D}(f)$, and $(f(a), f(b)) \subseteq \mathcal{R}(f)$, such that $x \in (a, b)$, and $f(x) \in (f(a), f(b))$. This implies f^{-1} is continuous, and since $f^{-1}(f(a)) = a > f^{-1}(f(b)) = b$ if $f(a) > f(b)$, implies f^{-1} is strictly increasing. Also since, $f(a) = f(b) \implies a = b$, f is injective.

Section 23

A. Examine each of the functions in Example 20.5 and either show that the function is uniformly continuous on its domain or not.

(a) The constant function, $\mathcal{D}(f) \subseteq \mathbb{R}, f(x) = c, \forall x \in \mathcal{D}(f)$, where c is a real number.

Let $\varepsilon > 0$, we have $|f(x) - f(y)| = |0 - 0| = 0 < \varepsilon, \forall x, y \in \mathcal{D}(f)$. Regardless of the choice of $\delta(\varepsilon)$, we have $|f(x) - f(y)| < \varepsilon$ whenever $|x - y| < \delta(\varepsilon) \implies$ uniform continuity.

(b) The identity function $f(x) = x, x \in \mathbb{R}$.

For all $x, y \in \mathbb{R}$, we have $|f(x) - f(y)| < \varepsilon, \varepsilon > 0$. Choose $\delta(\varepsilon) = \varepsilon$. Then whenever $|f(x) - f(y)| = |x - y| < \varepsilon = \delta(\varepsilon)$ we have $|x - y| < \delta(\varepsilon) \implies$ uniform continuity.

(c) $f(x) = x^2, x \in \mathbb{R}$

(d) $f(x) = \frac{1}{x}, x \in \{x \in \mathbb{R} : x \neq 0\}$

(e) $f(x) = 0, x \geq 0, f(x) = 1, x > 1$

(f) $f(x) = 1, x$ rational, $f(x) = 0, x$ irrational

(g) $\mathcal{D}(f) = \{x \in \mathbb{R} : x > 0\}, f(x) = 0, x$ irrational, $x > 0$. For rational numbers of the form $\frac{m}{n}$, with $m, n \in \mathbb{N}$ that have no common factor but 1, $f(\frac{m}{n}) = \frac{1}{n}$.

(h) $\mathcal{D}(f) = \mathbb{R}^2, f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f(x, y) = (2x + y, x - 3y)$.

(i) $\mathcal{D}(f) = \mathbb{R}^2, f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f(x, y) = (x^2 + y^2, 2xy)$.

C. If B is bounded in \mathbb{R}^p and $f : B \rightarrow \mathbb{R}^p$ is uniformly continuous, show that f is bounded on B . Show that this conclusion fails if B is not bounded in \mathbb{R}^p .

D. Show that $f(x) = \frac{1}{1+x^2}$ for $x \in \mathbb{R}$ is uniformly continuous.

Take $\varepsilon > 0$, for $x, y \in \mathbb{R}$, we have $|f(x) - f(y)| = |\frac{1}{1+x^2} - \frac{1}{1+y^2}| = |\frac{(1+y^2)}{(1+x^2)(1+y^2)} - \frac{(1+x^2)}{(1+y^2)(1+x^2)}| = |\frac{y^2 - x^2}{(1+y^2)(1+x^2)}| = |x+y||x-y| \frac{1}{(1+x^2)(1+y^2)} \leq (|\frac{x}{(1+x^2)(1+y^2)}| + |\frac{y}{(1+x^2)(1+y^2)}|)|x-y| \leq (|\frac{y}{(1+y^2)}| + |\frac{x}{(1+x^2)}|)|x-y|$. Since $\forall x \in \mathbb{R}$, we have $|\frac{x}{1+x^2}| < 1$, we have $(|\frac{y}{(1+y^2)}| + |\frac{x}{(1+x^2)}|)|x-y| < 2|x-y| = 2\delta(\varepsilon) \implies$ if we choose $\delta(\varepsilon) = \frac{\varepsilon}{2}$ whenever $|x - y| < \frac{\varepsilon}{2}$ we have $|f(x) - f(y)| < \varepsilon$, for all $x, y \in \mathbb{R}$

F. Show that $f(x) = \frac{1}{x^2}$, $\mathcal{D}(f) = \{x \in \mathbb{R} : x > 0\}$ is not uniformly continuous on its domain.

If we take $\delta(\varepsilon) = \varepsilon/2$, and $\varepsilon = 1$, and consider points in a subset of $\mathcal{D}(f)$, namely $x, y \in (0, 1)$, and then take $y = \frac{x}{2} \in (0, 1)$, we have we have $|x - y| = |x/2| < 1/2 \implies |f(x) - f(y)| = |\frac{1}{x^2} - \frac{4}{x^2}| = \frac{3}{x^2} < 1$ for but since for all $x, y \in (0, 1)$, $\frac{3}{x^2} > 1$, $f(x)$ is not uniformly continuous on its domain.

G. A function $g : \mathbb{R} \rightarrow \mathbb{R}^p$ is periodic if there exists a snumber $p > 0$ such that $g(x + p) = g(x)$ for all $x \in \mathbb{R}$. Show that a continuous periodic function is bounded and uniformly continuous on \mathbb{R} .

H. Let f be defined on $D \subseteq \mathbb{R}^p$ to \mathbb{R}^q , and suppose that f is uniformly continuous on D . If (x_n) is a Cauchy sequence in D , show that $(f(x_n))$ is a Cauchy sequence in \mathbb{R}^q .

Section 24

B. Give an example of a sequence of continuous functions which converges to a continuous function but where the convergence is not uniform.

D. Let (f_n) be a sequence of continuous functions on $D \subseteq \mathbb{R}^p$ to \mathbb{R}^q such that (f_n) converges uniformly to f on D , and let (x_n) be a sequence of elements in D which converges to $x \in D$. Does it follow that $(f_n(x_n))$ converges to $f(x)$?

E. Consider the sequences (f_n) defined on $D = \{x \in \mathbb{R} : x \geq 0\}$ to \mathbb{R} by the following formulas:

(b) $\frac{x^n}{1+x^n}$,

(c) $\frac{x^n}{n+x^n}$,

(d) $\frac{x^{2n}}{1+x^n}$,

(e) $\frac{x^n}{1+x^{2n}}$

J. Prove the following theorem of G. Polya. If for each $n \in \mathbb{N}$ the function f_n on $I \rightarrow \mathbb{R}$ is monotone increasing and if $f(x) = \lim(f_n(x))$ is continuous on I , then the convergence is uniform on I . (Observe that it is not assumed that f_n is continuous.)

N. If $f_3(x) = x^3$ for $x \in \mathcal{I}$, calculate the n^{th} Bernstein polynomial for f_3 . Show directly that this sequence of polynomials converges uniformly to f_3 on \mathcal{I} .

S. Show that the Weierstrass Approximation Theorem fails for bounded open intervals.

Section 26

N. If $K \subseteq \mathbb{R}^p$ is compact and (f_n) is a sequence of continuous functions on K to \mathbb{R}^q which is uniformly convergent on K , show that the family $\{f_n\}$ is uniformly equicontinuous on K in the sense of Definition 26.6.

O. Let \mathcal{F} be a bounded and uniformly equicontinuous collection of functions on $D \subseteq \mathbb{R}^p$ to \mathbb{R} and let f be defined on $D \rightarrow \mathbb{R}$ by $f = \sup\{f(x) : f \in \mathcal{F}\}$. Show that f is continuous on $D \rightarrow \mathbb{R}$.

Q. Consider the following sequences of functions which show that the Arzela-Ascoli Theorem 26.7 may fail if the various hypotheses are dropped.

(a) $f_n(x) = x + n$, $x \in [0, 1]$;

(b) $f_n(x) = x^n$, $x \in [0, 1]$;

(c) $f_n(x) = \frac{1}{1+(x-n)^2}$, $x \in [0, +\infty)$.