

# Statistical Inference

## Introduction to Linear Regression

Behnam Bahrak  
Spring 2020

1 of 20



## Multiple Linear Regression

➤ Predicting birth weight of babies from a **variety** of variables:

	bwt	gestation	parity	age	height	weight	smoke
1	120	284	0	27	62	100	0
2	113	282	0	33	64	135	0
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
1236	117	297	0	38	65	129	0

$$y \sim x_1 + x_2 + x_3 + x_4 + x_5 + x_6$$



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

2 of 20



## Multiple Predictors

weights of books



	weight (g)	volume (cm <sup>3</sup> )	cover
1	800	885	hb
2	950	1016	hb
3	1050	1125	hb
4	350	239	hb
5	750	701	hb
6	600	641	hb
7	1075	1228	hb
8	250	412	pb
9	700	953	pb
10	650	929	pb
11	975	1492	pb
12	350	419	pb
13	950	1010	pb
14	425	595	pb
15	725	1034	pb



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

3 of 20

## Hardcover vs. Paperback

- Can you identify a trend in the relationship between volume and weight of hardcover and paperback books?

Paperbacks generally weigh less than hardcover books.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

4 of 20

# Multiple Linear Regression in R

```
R
# load data
> library(DAAG)
> data(allbacks)

# fit model
> book_mlr = lm(weight ~ volume + cover, data = allbacks)
> summary(book_mlr)

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  197.96284   59.19274   3.344 0.005841 **
volume        0.71795    0.06153  11.669 6.6e-08 ***
cover:pb     -184.04727   40.49420  -4.545 0.000672 ***

Residual standard error: 78.2 on 12 degrees of freedom
Multiple R-squared:  0.9275, Adjusted R-squared:  0.9154
F-statistic: 76.73 on 2 and 12 DF, p-value: 1.455e-07
```



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

5 of 20

## Example

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover:pb}$$

- For hardcover books: plug in **0** for cover:

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 0 \\ &= 197.96 + 0.72 \text{ volume}\end{aligned}$$

- For paperback books: plug in **1** for cover:

$$\begin{aligned}\widehat{weight} &= 197.96 + 0.72 \text{ volume} - 184.05 \times 1 \\ &= 13.91 + 0.72 \text{ volume}\end{aligned}$$




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

6 of 20

## Example



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 7 of 20 &gt;

## Interpreting the regression parameters: slope

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover:pb}$$

- Slope of **volume**: All else held constant, for each 1 cm<sup>3</sup> increase in volume the model predicts the books to be heavier on average by 0.72 grams.
- Slope of **cover**: All else held constant, the model predicts that paperback books weigh 184.05 grams lower than hardcover books, on average.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 8 of 20 &gt;

## Interpreting the regression parameters: intercept

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover:pb}$$

**Intercept:** Hardcover books with no volume are expected on average to weigh 198 grams.

- Meaningless in context, serves to adjust the height of the line.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

9 of 20

## Prediction

- Predict the weight of a paperback book that is 600  $cm^3$  in volume.

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	197.96	59.19	3.34	0.01
volume	0.72	0.06	11.67	0.00
cover:pb	-184.05	40.49	-4.55	0.00

$$\widehat{weight} = 197.96 + 0.72 \text{ volume} - 184.05 \text{ cover:pb}$$

$$197.96 + 0.72 \times 600 - 184.05 \times 1 = 445.91 \text{ grams}$$




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

10 of 20

## Interaction Variables

- Model assumes hardcover and paperback books have the same slope for the relationship between their volume and weight.
- If this isn't reasonable, then we would include an interaction variable in the model (beyond the scope of this course).




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

11 of 20

## Adjusted $R^2$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

12 of 20

## Adjusted $R^2$

R

```
# fit model
> pov_slr = lm(poverty ~ female_house, data = states)
> summary(pov_slr)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	3.3094	1.8970	1.745	0.0873 .
female_house	0.6911	0.1599	4.322	7.53e-05 ***

Residual standard error: 2.664 on 49 degrees of freedom  
 Multiple R-squared: 0.276, Adjusted R-squared: 0.2613  
 F-statistic: 18.68 on 1 and 49 DF, p-value: 7.534e-05




Statistical Inference

 Behnam Bahrak  
 bahrak@ut.ac.ir

13 of 20

## Predicting poverty from % female householder



Linear model:	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	3.31	1.90	1.74	0.09
female_house	0.69	0.16	4.32	0.00



Statistical Inference

 Behnam Bahrak  
 bahrak@ut.ac.ir

14 of 20

## Another look at $R^2$

ANOVA:	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female.house	1	132.57	132.57	18.68	0.00
Residuals	49	347.68	7.10		
Total	50	480.25			

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{132.57}{480.25} = 0.28$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

15 of 20

## Predicting poverty from % female householder + % white

R

```
> pov_mlr = lm(poverty ~ female_house + white, data = states)
> summary(pov_mlr)
```

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	-2.58	5.78	-0.45	0.66
female_house	0.89	0.24	3.67	0.00
white	0.04	0.04	1.08	0.29

R

```
> anova(pov_mlr)
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female_house	1	132.57	132.57	18.74	0.00
white	1	8.21	8.21	1.16	0.29
Residuals	48	339.47	7.07		
Total	50	480.25			

$$R^2 = \frac{132.57 + 8.21}{480.25} = 0.29$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

16 of 20



## Adjusted $R^2$

$$\text{adjusted } R^2: \quad R_{adj}^2 = 1 - \left( \frac{SSE}{SST} \times \frac{n-1}{n-k-1} \right)$$

$k$  : number of predictors



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

17 of 20

## Example

- Calculate adjusted  $R^2$  for the multiple linear regression model predicting % living in poverty from % female householders and % white. Remember  $n = 51$  (50 states + DC).

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
female.house	1	132.57	132.57	18.74	0.00
white	1	8.21	8.21	1.16	0.29
Residuals	48	339.47	7.07		
Total	50	480.25			

$$R_{adj}^2 = 1 - \left( \frac{SSE}{SST} \times \frac{n-1}{n-k-1} \right) = 1 - \left( \frac{339.47}{480.25} \times \frac{51-1}{51-2-1} \right) = 0.26$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

18 of 20

## $R^2$ vs. adjusted $R^2$

	$R^2$	adjusted $R^2$
Model 1 (poverty vs. female_house)	0.28	0.26
Model 2 (poverty vs. female_house + white)	0.29	0.26

- When **any** variable is added to the model  $R^2$  increases.
- But if the added variable doesn't really provide any new information, or is completely unrelated, adjusted  $R^2$  does not increase.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

19 of 20

## Properties of adjusted $R^2$

$$R_{adj}^2 = 1 - \left( \frac{SSE}{SST} \times \frac{n-1}{n-k-1} \right)$$

- $k$  is never negative  $\rightarrow$  (adjusted  $R^2$ )  $< R^2$
- Adjusted  $R^2$  applies a penalty for the number of predictors included in the model
- We choose models with higher adjusted  $R^2$  over others



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

20 of 20