

# Statistical Inference


## Introduction to Linear Regression

Behnam Bahrak  
Spring 2020

1 of 27

## Prediction


- Using the linear model to predict the value of the response variable for a given value of the explanatory variable is called **prediction**
- Plug in the value of  $x$  in the linear model equation



## Example

- According to the following linear model, what is the predicted % living in poverty in states where the HS graduation rate is 82%?  $\widehat{\% \text{ in poverty}} = 64.68 - 0.62 \% \text{ HS grad}$

$$\begin{aligned}\% \text{ in poverty} &= 64.68 - 0.62 \times 82 \\ &= 13.84 \%\end{aligned}$$




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 3 of 27 &gt;

## Extrapolation

- Applying a model estimate to values outside of the realm of the original data is called **extrapolation**
- Sometimes the intercept might be an extrapolation




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 4 of 27 &gt;

## Example

- According to the following linear model, what is the predicted % living in poverty in states where the HS graduation rate is 20%?



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

5 of 27

## Will All Americans Become Overweight or Obese? Estimating the Progression and Cost of the US Obesity Epidemic

Youfa Wang<sup>1</sup>, May A. Beydoun<sup>1</sup>, Lan Liang<sup>2</sup>, Benjamin Caballero<sup>1</sup> and Shiriki K. Kumanyika<sup>3</sup>

We projected future prevalence and BMI distribution based on national survey data (National Health and Nutrition Examination Study) collected between 1970s and 2004. Future obesity-related health-care costs for adults were estimated using projected prevalence, Census population projections, and published national estimates of per capita excess health-care costs of obesity/overweight. The objective was to illustrate potential burden of obesity prevalence and health-care costs of obesity and overweight in the United States that would occur if current trends continue. Overweight and obesity prevalence have increased steadily among all US population groups, but with notable differences between groups in annual increase rates. The increase (percentage points) in obesity and overweight in adults was faster than in children (0.77 vs. 0.46–0.49), and in women than in men (0.91 vs. 0.65). If these trends continue, by 2030, 86.3% adults will be overweight or obese; and 51.1%, obese. Black women (96.9%) and Mexican-American men (91.1%) would be the most affected. By 2048, all American adults would become overweight or obese, while black women will reach that state by 2034. In children, the prevalence of overweight (BMI  $\geq$  95th percentile, 30%) will nearly double by 2030. Total health-care costs attributable to obesity/overweight would double every decade to 860.7–956.9 billion US dollars by 2030, accounting for 16–18% of total US health-care costs. We continue to move away from the Healthy People 2010 objectives. Timely, dramatic, and effective development and implementation of corrective programs/policies are needed to avoid the otherwise inevitable health and societal consequences implied by our projections.




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

6 of 27

## Linear Regression




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 7 of 27 &gt;

## Extrapolation




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

&lt; 8 of 27 &gt;

## More Realistic Curve



How not to be wrong: The power of mathematical thinking  
By Jordan Ellenberg



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

9 of 27

## Conditions for linear regression

- **Linearity**
  - relationship between the explanatory and the response variable should be linear
- **Nearly normal residuals**
  - residuals should be nearly normally distributed
- **Constant variability**
  - variability of points around the least squares line should be roughly constant



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

10 of 27


## (1) Linearity

- Relationship between the explanatory and the response variable should be linear
- Methods for fitting a model to non-linear relationships exist
- Check using a scatterplot of the data, or a residuals plot

Scatterplot:



Residuals plot:



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir


11 of 27

## Anatomy of a residuals plot

\* RI

% HS grad = 81%  
% in poverty = 10.3 %

$$\begin{aligned}\%pov &= 64.68 - 0.62 \times 81 \\ &= 14.46\% \\ e &= 10.3 - 14.46 \\ &= -4.16\%\end{aligned}$$



■ DC

% HS grad = 86%  
% in poverty = 16.8 %

$$\begin{aligned}\%pov &= 64.68 - 0.62 \times 86 \\ &= 11.36\% \\ e &= 16.8 - 11.36 \\ &= 5.44\%\end{aligned}$$




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

12 of 27

## (2) Nearly Normal Residuals

- Residuals should be nearly normally distributed, centered at 0
- May not be satisfied if there are unusual observations that don't follow the trend of the rest of the data
- Check using a histogram or normal probability plot of residuals




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

13 of 27

## (3) Constant Variability

- Variability of points around the least squares line should be roughly constant
- Implies that the variability of residuals around the 0 line should be roughly constant as well
- Also called **homoscedasticity**
- Check using a residuals plot



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

14 of 27

# $R^2$

- Strength of the fit of a linear model is most commonly evaluated using  $R^2$
- Calculated as the square of the correlation coefficient
- Tells us **what percent of variability in the response variable is explained by the model**
- The remainder of the variability is explained by variables not included in the model
- Always between 0 and 1

$$R^2 = \frac{(\text{Cov}(x, y))^2}{\sigma_x^2 \sigma_y^2}$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

15 of 27

## Example

- Which of the following is the correct interpretation of the  $R^2$  for this model for predicting % living in poverty from % HS graduation rate? ( $R^2 = 0.5625$ )
  - 56.25% of the time % HS graduates predict % living in poverty correctly.
  - 43.75% of the variability in the % of residents living in poverty among the states is explained by the model.
  - 56.25% of the variability in the % of HS graduates among the states is explained by the model.
  - ☒ 56.25% of the variability in the % of residents living in poverty among the states is explained by the model.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

16 of 27




## Example

- The  $R^2$  for the relationship displayed in the scatterplot is 92.16%. What is the correlation coefficient?

$$|R| = \sqrt{R^2} = \sqrt{0.9216} = 0.96$$

$$\Rightarrow R = -0.96$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

17 of 27

## Regression with categorical explanatory variable

Poverty vs. Region

explanatory variable: region

0: east  
1: west

$$\widehat{poverty} = 11.17 + 0.38 \text{ region} : \text{west}$$

for eastern states  
plug in 0 for  $x$

for western states  
plug in 1 for  $x$

$$\widehat{poverty} = 11.17 + 0.38 \times 0 = 11.17 \quad \widehat{poverty} = 11.17 + 0.38 \times 1 = 11.55$$

reference level:

In regression models, with explanatory categorical variables, we always code one of the levels of that categorical variable to be what we call the reference level. This is the level that we plug in zero for.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

18 of 27

## Slope and Intercept

$$\widehat{poverty} = 11.17 + 0.38 \text{ region : west}$$

- **Intercept:** The model predicts an 11.17% average poverty percentage in eastern states.
- This is the value we get if we plug in 0 for the explanatory variable
- **Slope:** The model predicts that the average poverty percentage in western states is 0.38% higher than in the eastern states.



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

19 of 27

## Example

- Next, we use a new region variable (`region4`) with four levels: *northeast*, *midwest*, *west*, *south*. Write the linear regression model based on the regression output below.

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4:midwest	0.03	1.15	0.02	0.98
region4:west	1.79	1.13	1.59	0.12
region4:south	4.16	1.07	3.87	0.00

$$\% \text{ in poverty} = 9.50 + 0.03 \text{ reg4:mw} + 1.79 \text{ reg4:w} + 4.16 \text{ reg4:s}$$



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

20 of 27

## Example

- Calculate the predicted poverty rate for western states.

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	9.50	0.87	10.94	0.00
region4:midwest	0.03	1.15	0.02	0.98
region4:west	1.79	1.13	1.59	0.12
region4:south	4.16	1.07	3.87	0.00

$$\begin{aligned}
 \% \text{ in poverty} &= 9.50 + 0.03 \text{ reg4:mw} + 1.79 \text{ reg4:w} + 4.16 \text{ reg4:s} \\
 &= 9.50 + 0 + 1.79 + 0 \\
 &= 11.29
 \end{aligned}$$



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

21 of 27

## Outliers in Regression

- How does the outlier influence the least squares line?

Without the outlier there is no relationship between  $x$  and  $y$ .



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

22 of 27

## Types of Outliers

- **Outliers** are points that fall away from the cloud of points
- Outliers that fall horizontally away from the center of the cloud but don't influence the slope of the regression line are called **leverage points**
- Outliers that actually influence the slope of the regression line are called **influential points**
  - usually high leverage points
  - to determine if a point is influential, visualize the regression line with and without the point, and ask: *Does the slope of the line change considerably?*



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

23 of 27

## Example

What type of outlier is this?

leverage point



Statistical Inference


Behnam Bahrak  
bahrak@ut.ac.ir

24 of 27

## Example

What type of outlier is this?

influential point




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

25 of 27

## Influential Points

- Light intensity and surface temperature (logged) of 47 stars in the star cluster CYG OB1




Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

26 of 27

## Example

- True or false? Influential points always reduce  $R^2$ .



Statistical Inference

Behnam Bahrak  
bahrak@ut.ac.ir

27 of 27