

Statistical Inference

Logistic Regression

Behnam Bahrak
Spring 2020

1 of 20 ➤

Regression so far ...

- At this point we have covered:
 - Simple linear regression
 - Relationship between numerical response and a numerical or categorical predictor
 - Multiple regression
 - Relationship between numerical response and multiple numerical and/or categorical predictors
- What we haven't seen is what to do when the predictors are weird (nonlinear, complicated dependence structure, etc.) or when the response is weird (categorical, count data, etc.)



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

◀ 2 of 20 ▶

Odds

- Odds are another way of quantifying the probability of an event, commonly used in gambling (and logistic regression).
- For some event E ,

$$\text{odds}(E) = \frac{P(E)}{P(E^c)} = \frac{P(E)}{1 - P(E)}$$

- Similarly, if we are told the odds of E are x to y then:

$$\text{odds}(E) = \frac{x}{y} = \frac{x/(x+y)}{y/(x+y)}$$

- Which implies:

$$P(E) = \frac{x}{x+y}, \quad P(E^c) = \frac{y}{x+y}$$



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

3 of 20

Example - Donner Party

- In 1846 the Donner and Reed families left Springfield, Illinois, for California by covered wagon.
- In July, the Donner Party reached Fort Bridger, Wyoming. There its leaders decided to attempt a new and untested route to the Sacramento Valley.
- Having reached its full size of 87 people and 20 wagons, the party was delayed by a difficult crossing of the Wasatch Range.
- The group became stranded in the eastern Sierra Nevada mountains when the region was hit by heavy snows in late October.
- By the time the last survivor was rescued on April 21, 1847, 40 of the 87 members had died from famine and exposure to extreme cold.



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

4 of 20

Example - Donner Party - Data

	Age	Sex	Status
1	23.00	Male	Died
2	40.00	Female	Survived
3	40.00	Male	Survived
4	30.00	Male	Died
5	28.00	Male	Died
⋮	⋮	⋮	⋮
43	23.00	Male	Survived
44	24.00	Male	Died
45	25.00	Female	Survived



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir


5 of 20

Example - Donner Party

Status vs. Gender:

	Male	Female
Died	20	5
Survived	10	10

Status vs. Age:



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

6 of 20

Example - Donner Party

- It seems clear that both age and gender have an effect on someone's survival, how do we come up with a model that will let us explore this relationship?
- Even if we set Died to 0 and Survived to 1, this isn't something we can transform our way out of - we need something more.
- One way to think about the problem - we can treat Survived and Died as successes and failures arising from a binomial distribution where the probability of a success is given by a transformation of a linear model of the predictors.



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

< 7 of 20 >

Generalized Linear Models

- It turns out that this is a very general way of addressing this type of problem in regression, and the resulting models are called generalized linear models (GLMs).
- Logistic regression is just one example of this type of model.
- All generalized linear models have the following three characteristics:
 1. A probability distribution describing the outcome variable
 2. A linear model
 3. A link function that relates the linear model to the parameter of the outcome distribution

$$\eta = \beta_0 + \beta_1 X_1 + \dots + \beta_n X_n$$

$$g(p) = \eta, \quad p = g^{-1}(\eta)$$



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

< 8 of 20 >

توزیع برنولی $\leftarrow p = \frac{1}{1+e^{-x}} \quad x = \beta_0 + \beta_1 x_1 + \dots + \beta_n x_n$

$g(p) = \log\left(\frac{p}{1-p}\right) = \eta = \beta_0 + \beta_1 x_1 + \dots + \beta_n x_n$

Logistic Regression

- Logistic regression is a GLM used to model a binary categorical variable using numerical and categorical predictors.
- We assume a Bernoulli distribution produced the outcome variable and we therefore want to model p the probability of success for a given set of predictors.
- To finish specifying the Logistic model we just need to establish a reasonable link function that connects η to p . There are a variety of options but the most commonly used is the logit function.

Logit function:

$$\text{logit}(p) = \log\left(\frac{p}{1-p}\right), \quad \text{for } 0 \leq p \leq 1$$



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

9 of 20

Properties of the Logit

- The logit function takes a value between 0 and 1 and maps it to a value between $-\infty$ and ∞ .

Inverse logit (logistic) function

$$g^{-1}(x) = \frac{e^x}{1+e^x} = \frac{1}{1+e^{-x}}$$

- The inverse logit function takes a value between $-\infty$ and ∞ and maps it to a value between 0 and 1.
- This formulation also has some use when it comes to interpreting the model as logit can be interpreted as the log odds of a success.



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

10 of 20

The logistic regression model

- The three GLM criteria give us:

$$y_i \sim \text{Bernoulli}(p_i)$$

$$\eta = \beta_0 + \beta_1 X_1 + \dots + \beta_n X_n$$

$$\text{logit}(p) = \eta$$

- From which we arrive at:

$$p_i = \frac{e^{\beta_0 + \beta_1 X_{1,i} + \dots + \beta_n X_{n,i}}}{1 + e^{\beta_0 + \beta_1 X_{1,i} + \dots + \beta_n X_{n,i}}}$$



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

11 of 20

Example - Donner Party - Model

- In R we fit a GLM in the same way as a linear model except using `glm` instead of `lm` and we must also specify the type of GLM to fit using the `family` argument.

R

```
> summary(glm(Status ~ Age, data=donner, family=binomial))
```

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

12 of 20

Example - Donner Party - Prediction

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a newborn (Age = 0):

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 0 \Rightarrow \frac{p}{1-p} = e^{1.8185} = 6.16 \Rightarrow p = 0.86$$



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

13 of 20

Example - Donner Party - Prediction

Model:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times \text{Age}$$

Odds / Probability of survival for a 25 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 25 \Rightarrow \frac{p}{1-p} = e^{0.156} = 1.17 \Rightarrow p = 0.539$$

Odds / Probability of survival for a 50 year old:

$$\log\left(\frac{p}{1-p}\right) = 1.8185 - 0.0665 \times 50 \Rightarrow \frac{p}{1-p} = e^{-1.5065} = 0.222 \Rightarrow p = 0.181$$




Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

14 of 20

Example - Donner Party - Prediction



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

15 of 20

Example - Donner Party - Interpretation

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.8185	0.9994	1.82	0.0688
Age	-0.0665	0.0322	-2.06	0.0391

- Simple interpretation is only possible in terms of log odds and log odds ratios for intercept and slope terms.
- **Intercept:** The log odds of survival for a party member with an age of 0. From this we can calculate the odds or probability, but additional calculations are necessary.
- **Slope:** For a unit increase in age (being 1 year older) how much will the log odds ratio change, not particularly intuitive. More often then not we care only about sign and relative magnitude.

(شخصی) \log \rightarrow
 نسبت احتمال زنده ماندن
 نسبت احتمال زنده ماندن
 نسبت احتمال زنده ماندن
 نسبت احتمال زنده ماندن



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

16 of 20

Example - Interpretation of Slope

$$\log\left(\frac{p_1}{1-p_1}\right) = 1.8185 - 0.0665(x+1)$$

$$\log\left(\frac{p_2}{1-p_2}\right) = 1.8185 - 0.0665x$$

$$\log\left(\frac{p_1}{1-p_1}\right) - \log\left(\frac{p_2}{1-p_2}\right) = -0.0665$$

$$\frac{p_1/(1-p_1)}{p_2/(1-p_2)} = e^{-0.0665} = 0.94$$

نسبت پتانسیل میسر =



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

< 17 of 20 >

وقتی چند متغیر داریم همه در ثابت در نظر می گیریم و آن صد نفر صاف در یک واحد کم از یاد داریم بنسبت می رسد

Example: Donner Party - Age and Gender

R

```
> summary(glm(Status ~ Age + Sex, family = binomial, data = donner))
```

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.6331	1.1102	1.47	0.1413
Age	-0.0782	0.0373	-2.10	0.0359
Sex:Female	1.5973	0.7555	2.11	0.0345

Gender slope: When the other predictors are held constant this is the log odds ratio between the given level (Female) and the reference level (Male).



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

< 18 of 20 >

Example: Donner Party - Gender Models

- Just like MLR we can plug in gender to arrive at two status vs. age models for men and women respectively.

General model:

$$\log\left(\frac{p_1}{1-p_1}\right) = 1.63312 - 0.07820 \times \text{Age} + 1.59729 \times \text{Sex}$$

Male model:

$$\log\left(\frac{p_1}{1-p_1}\right) = 1.63312 - 0.07820 \times \text{Age} + 1.59729 \times 0 = 1.63312 - 0.07820 \times \text{Age}$$

Female model:

$$\log\left(\frac{p_1}{1-p_1}\right) = 1.63312 - 0.07820 \times \text{Age} + 1.59729 \times 1 = 3.23041 - 0.07820 \times \text{Age}$$




Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

19 of 20

Example: Donner Party - Gender Models



Statistical Inference

Behnam Bahrak
bahrak@ut.ac.ir

20 of 20