A Simple Orthogonal 22-Gon with Exactly One Non-Illuminating Pair

Peyton Wang
Francis Parker High School

Daniel Wang
Citizen Health

Working draft v0.4 — 10 November 2026
Not submitted or peer reviewed

Abstract

We give exact coordinates for a strictly simple orthogonal polygon with 22 maximal sides and two distinct interior points that cannot be joined by any finite billiard trajectory avoiding polygon vertices. The room is assembled from three reflected copies of a five-square staircase. Folding a hypothetical trajectory onto the staircase reduces the problem to an elementary lattice-parity obstruction. Four smooth boundary points of the assembled room project to staircase corners; we show explicitly that reflection at these exceptional points produces only trajectory reversal and cannot evade the obstruction. The construction uses neither interior mirror partitions nor boundary self-contacts. We then prove that this is the only unordered pair of distinct real interior points that do not illuminate one another. The additional argument uses two rational cylinder decompositions, certified by explicit finite permutations, and an elementary integrality calculation.

1 Introduction

In the polygonal illumination problem, light rays travel in straight lines and reflect specularly from the sides of a polygonal room. A ray that reaches a vertex is terminated. Two interior points are said to *illuminate each other* if some finite trajectory with these reflection rules connects them.

Lattice obstructions give polygonal examples with mutually non-illuminating points [1]. General results on illumination in rational polygons appear in [2]. For this particular polygon, we determine the entire relation on distinct interior points by combining a folding obstruction with two explicit cylinder decompositions.

"Strictly simple" means that the boundary is a Jordan polygonal curve: it has no repeated vertices, crossed edges, or self-contacts. Only the exterior boundary reflects light; the dashed segments in the construction are open passages, not mirrors. Sections 2–5 give an elementary proof of darkness. Section 6 proves illumination for every other pair, using a four-copy translation unfolding defined there.

Theorem 1 (Main result)**.** *The polygon Q constructed below is strictly simple and orthogonal, with exactly 22 maximal sides and no holes or interior mirror partitions. Among all unordered pairs of distinct points in* $\mathrm{int}(Q)$*, its only non-illuminating pair is* $\{(0,0),(2,2)\}$.

2 The construction

Let P be the union of five closed unit squares with lower-left corners

$$(-1,-1), \quad (0,-1), \quad (0,-2), \quad (1,-2), \quad (1,-3). \tag{1}$$

Mark two smooth points of its boundary,

$$a = (0, 0), \qquad b = (2, -2). \tag{2}$$


Take the following isometric copies of P:

$$P_0 = P, \qquad P_1 = \{(x, -y) : (x, y) \in P\}, \qquad P_2 = \{(4 - x, -y) : (x, y) \in P\}. \tag{3}$$

Their interiors are pairwise disjoint. The first two meet exactly along $H = [-1, 1] \times \{0\}$, and the latter two meet exactly along $V = \{2\} \times [1, 3]$. There are no further contacts. Set

$$Q = P_0 \cup P_1 \cup P_2, \qquad S = (0, 0), \quad T = (2, 2).$$

Both S and T are midpoints of *open passages* between copies and consequently are interior points of Q.



Five-square base P




Figure 1: The five-square base P and the 22-sided room Q. The bold exterior outline is reflective. The dashed segments are open seams, not partitions. The two marked points S and T lie in the interior of Q.

For reproducibility, the counterclockwise sequence of all 22 vertices of Q is

$$\begin{gathered}(-1, -1), (0, -1), (0, -2), (1, -2), (1, -3), (2, -3), \\ (2, -1), (1, -1), (1, 1), (3, 1), (3, 0), (5, 0), \\ (5, 1), (4, 1), (4, 2), (3, 2), (3, 3), (1, 3), \\ (1, 2), (0, 2), (0, 1), (-1, 1).\end{gathered} \tag{4}$$

Every successive edge is horizontal or vertical and nonzero, with successive edges meeting at right angles. The union description (or a direct check of the vertex cycle) shows that the boundary does not self-intersect or self-touch. There are 13 convex vertices and nine reflex vertices, and the area is 15.

The *folding map* $f : Q \to P$ is defined copywise by

$$f|_{P_0}(x, y) = (x, y), \qquad f|_{P_1}(x, y) = (x, -y), \qquad f|_{P_2}(x, y) = (4 - x, -y). \tag{5}$$

These formulas agree on the seams, so f is well defined and continuous. In particular, $f(S) = a$ and $f(T) = b$.

3 The lattice obstruction on the staircase

The next lemma concerns *ordinary* billiard motion in P, reflecting only at smooth boundary points and never touching a vertex. Paths have finitely many nonzero straight legs. At each intermediate smooth-wall contact the incoming and outgoing legs satisfy the specular reflection rule.

Lemma 1 (Parity obstruction)**.** *There is no finite corner-avoiding billiard path in P from $a = (0, 0)$ to $b = (2, -2)$.*

Proof. First inspect the integer points in P. There are no interior integer points. Its only integer boundary points that are not vertices are a and b. More explicitly, its seven convex corners are

$$(-1, -1),\ (-1, 0),\ (0, -2),\ (1, -3),\ (1, 0),\ (2, -3),\ (2, -1),$$

and its three reflex corners are

$$(0, -1),\ (1, -2),\ (1, -1).$$

Suppose an ordinary billiard path from a to b exists. Unfold its finitely many reflections, reflecting the staircase instead of bending the path. The result is a single straight segment starting at $a = (0, 0)$ and ending at an image B of b. Reflection across any wall $x = c$ or $y = d$, where $c, d \in \mathbb{Z}$, preserves both coordinate parities. Since b has even coordinates, $B \in (2\mathbb{Z})^2$. Further, $B \neq (0, 0)$ because unfolding preserves the positive length of the path.

Write $B = 2^r z$, where $r \geq 1$ and $z \in \mathbb{Z}^2$ has at least one odd coordinate; choose r maximal. Then z is a lattice point strictly between 0 and B on the unfolded segment. Fold z back to P along the proposed itinerary. All inverse unfolding maps are compositions of reflections in integer-coordinate lines and therefore preserve integrality and coordinate parities. The image of z is an integer point of P with an odd coordinate. It is neither a nor b, hence must be a vertex of P. This would make the original path hit a vertex, a contradiction. □

Remark 1. The lemma is valid for arbitrary finite path length, direction, and number of reflections. The obstruction arises from the parity of an *unfolded endpoint*, not from a bounded search over directions.

4 What happens at the seams?

A billiard trajectory in Q need not fold into an ordinary billiard trajectory in P at every instant. The exceptions are precisely the smooth boundary points of Q lying at the ends of open seams. It is important to account for them rather than treating them as forbidden corners of Q.

Away from the seams, each branch of f is an isometry. A reflection at a smooth exterior wall of Q thus becomes an ordinary reflection at a smooth wall of P, unless that point is a seam endpoint. A straight passage across the *interior* of either seam also becomes an ordinary reflection in P: the adjacent folding maps agree at the seam and reverse the normal component of velocity.

The four seam endpoints are

$$E_1 = (-1, 0), \qquad E_2 = (1, 0), \qquad E_3 = (2, 1), \qquad E_4 = (2, 3). \tag{6}$$

Each E_j is smooth on the external boundary of Q, yet folds to a convex right-angle corner of P:

smooth point of Q	vertex of P	exterior mirror in Q
$(-1, 0)$	$(-1, 0)$	vertical
$(1, 0)$	$(1, 0)$	vertical
$(2, 1)$	$(2, -1)$	horizontal
$(2, 3)$	$(2, -3)$	horizontal

Lemma 2 (Seam-endpoint reversal)**.** *If a corner-avoiding billiard trajectory in Q reflects at a seam endpoint E_j, its folded trajectory in P instantaneously* reverses its entire velocity *at the corresponding convex corner and retraces its incoming route.*

Proof. For example, consider $E_2 = (1, 0)$. A ray approaching from P_1 with physical velocity $(u, -v)$ $(u, v > 0)$ folds via $(x, y) \mapsto (x, -y)$ to incoming velocity (u, v) in P. At E_2, the actual exterior mirror is vertical, so the physical outgoing velocity becomes $(-u, -v)$ and the ray travels into P_0. Folding this outgoing segment by the identity gives $(-u, -v)$, the negative of the incoming folded velocity. Reversing the time orientation gives the same conclusion for arrivals from P_0.

At E_1 the same two reflections occur in the opposite horizontal direction. At E_3 and E_4, interchange the horizontal and vertical coordinates: the physical exterior reflection negates the vertical component, while switching copies across the vertical seam negates the horizontal component. In every case, the combined effect is $(p, q) \mapsto (-p, -q)$. □




Figure 2: The exceptional event at the horizontal seam endpoint E_2. A normal mirror reflection in Q switches copies and becomes an exact retracing at a convex vertex of P. The diagrams show local geometry, not a possible complete connecting path.

Every original corner of P gives one corner incidence in each of P_0, P_1, P_2. Of these 30 incidences, two meet at each of the four smooth endpoints E_j (eight incidences total). All remaining 22 incidences are actual vertices of Q. In particular, *no reflex corner of P is the image of a smooth point of Q*. It follows that a valid Q-trajectory can fold only to ordinary billiard motion in P, occasionally interrupted by the corner reversals of Lemma 2.

5 Why reversals do not create a connecting path

Lemma 3 (Reversals do not help)**.** *There is no finite path from a to b in P obtained by ordinary billiard reflections at smooth walls together with the additional rule that a path may reverse its velocity and retrace itself at a convex corner. Reflex corners remain forbidden.*

Proof. Suppose such a path exists. Stop it at its first arrival at b, and retain only the subpath beginning at its *last* visit to a before that arrival. The resulting path γ starts at a, ends at b, and contains no interior occurrence of either marked point. There are finitely many such visits because the path has finitely many nonzero straight legs. This truncation takes place in P: a visit to a counts regardless of which preimage of a the original path visits in Q.

Assume γ has a corner reversal, and choose the first one. Up to that event, motion consists entirely of ordinary billiard reflections. Ordinary billiard motion is deterministic and reversible. Once the first reversal takes place, the path must retrace the entire preceding route in reverse order, ultimately reaching a. It cannot reach b during this retracing, since the corresponding point would already have occurred before the reversal, contradicting the first-arrival choice of b. Thus the path revisits a before reaching b, contradicting the last-visit choice of a.

Hence γ has no reversals and is an ordinary corner-avoiding billiard path from a to b, contrary to Lemma 1. □

Proof of the darkness assertion in Theorem 1. The explicit construction and boundary cycle establish that Q is a strictly simple orthogonal polygon with 22 maximal sides, no holes, and $S, T \in \operatorname{int}(Q)$.

Suppose a finite billiard trajectory joins S to T in Q. It cannot be axis-parallel: reflecting in a horizontal or vertical wall preserves the property that one velocity component vanishes, whereas S and T differ in both coordinates. Consequently every straight leg meets either seam transversely, if at all. Since the trajectory has finitely many legs and each leg crosses each seam at most once, its folding by f splits into finitely many straight segments and events.

By the seam analysis, the image is a finite path from $a = f(S)$ to $b = f(T)$ following the rules of Lemma 3. That lemma forbids such a path. Therefore no finite corner-avoiding billiard trajectory joins S and T. □

6 Every other interior pair is illuminated

Here and throughout, a pair means two *distinct* interior points. Self-return trajectories and boundary sources are not part of the classification.

6.1 The four-copy unfolding

Form a translation surface M from four copies of Q, labelled by $\varepsilon = (\varepsilon_1, \varepsilon_2) \in \{1, -1\}^2$. In the copy labelled ε, use developing coordinates $(\varepsilon_1 x, \varepsilon_2 y)$, up to integer translations. Across a vertical exterior wall, glue to the copy with the first sign reversed; across a horizontal exterior wall, reverse the second sign. Internal unit-cell edges are ordinary translation identifications on the same sheet. The resulting surface consists of 60 unit squares. Let Σ be the set of all lifts of actual polygon vertices. Include convex vertices in Σ even though their lifts have total angle 2π. Smooth boundary lattice points and interior lattice points are not in Σ. A straight geodesic avoiding Σ projects to a legal billiard trajectory.

The map τ that replaces ε by $-\varepsilon$ is an isometry with derivative $-I$, and projects to the same point of Q. It has no fixed point outside Σ: an interior point has four distinct lifts, while a smooth wall point identifies only one of the two sign changes. Only at polygon vertices can the two simultaneous sign changes be identified.

A *regular cylinder* in a rational direction is a maximal open annulus of parallel closed geodesics avoiding Σ. In coordinates it is $(\mathbb{R}/L\mathbb{Z}) \times (0, h)$; its closure is the image of a closed strip. The involution τ cannot preserve a regular cylinder. Indeed, on such a cylinder it would have the form $(u, v) \mapsto (c - u, h - v)$, which has two fixed points at $v = h/2$, contrary to the preceding observation.

Lemma 4 (Two-cylinder visibility)**.** *Suppose a rational direction decomposes M into two regular cylinders, exchanged by τ, whose closures cover M. If an interior point A has a lift on a regular leaf in that direction, then A illuminates every other interior point B of Q.*

Proof. Let $\widehat{A}$ lie in the open cylinder C. Choose a lift $\widehat{B}$; replace it by $\tau\widehat{B}$ if necessary so that it lies in $\overline{C}$. Since $A \neq B$, their lifts are distinct. Represent them in a closed covering strip $\mathbb{R} \times [0, h]$ as (u_A, v_A) and (u_B, v_B), where $0 < v_A < h$ and $0 \leq v_B \leq h$. For $0 < t < 1$, the interpolated height $(1-t)v_A + tv_B$ lies strictly between 0 and h. Thus the entire open segment between the representatives lies inside the open strip. Its image avoids Σ in its interior; the endpoint also avoids Σ because B is interior to Q. Projection to Q gives a finite legal billiard path. Compactness of the segment and its avoidance of the finite marked set ensure that only finitely many wall crossings occur. □




Figure 3: The visibility step, shown schematically in a cylinder's covering strip. Dashed lines indicate repeated periods, not mirrors. No assertion is made that the connecting segment follows the cylinder's periodic direction.

6.2 Two finite cylinder certificates

For this room, the directions $(p, q) = (4, 7)$ and $(7, 4)$ satisfy the hypothesis of Lemma 4. Here is an explicit finite prescription for checking this input; it requires no search over rooms or possible billiard paths.

Let $\mathcal{C}$ be the 15 cell lower-left corners determined by (1)–(3). Order the 60 squares of M lexicographically by cell, then by sheet in the order $(+,+), (-,+), (+,-), (-,-)$, numbering from 0. Because Q is the union of its cells with no internal mirrors, a shared unit edge is an open passage between the adjacent cells. The right-neighbour permutation r sends square $((a, b), (e_1, e_2))$ to $((a + e_1, b), (e_1, e_2))$ when that cell is in $\mathcal{C}$, and otherwise to $((a, b), (-e_1, e_2))$. Similarly the up-neighbour permutation u moves to $((a, b + e_2), (e_1, e_2))$ when possible, and otherwise to $((a, b), (e_1, -e_2))$.

In each square divide its bottom edge into q open intervals $(j/q, (j + 1)/q)$, $0 \le j < q$. Following direction (p, q) from one bottom crossing to the next induces the permutation

$$(i, j) \longmapsto \left(u\left(r^{\lfloor (j+p)/q \rfloor}(i)\right),\ (j + p) \bmod q\right).$$

The permutation preserves the within-interval offset $t - j/q$, so each cycle closes every leaf in its open interval. A cycle therefore describes an open annular strip of regular leaves. These strips cover every leaf whose transverse coordinate $qx - py$ is nonintegral. All marked points have integral developing coordinates, so none lies inside such a strip. The complete cycle lists are included in `certificate.json`, and the standalone standard-library Python program `verify_cylinders.py` regenerates them from the cell formulas. The results are

(p, q)	slots	cycle lengths	number of cycles
$(4, 7)$	420	$210, 210$	2
$(7, 4)$	240	$120, 120$	2

Every regular cylinder contains one of these strips: the complement of the integral transverse levels is dense. Thus there are at most two regular cylinders. There are at least two, since τ cannot preserve a cylinder. Consequently there are exactly two, exchanged by τ; their closures cover M, again by density. This proves the required cylinder input from the finite permutations.

A compact printed certificate. Write T for the slot permutation above. Since $\gcd(p, q) = 1$, the residue j visits every value modulo q and first returns to 0 after exactly q steps. Hence the first-return permutation $\Phi_{p,q}(i) = \pi_1(T^q(i, 0))$ acts on just the 60 squares. Each cycle of length n for

$\Phi_{p,q}$ corresponds to a cycle of length nq for T, and every cycle of T occurs this way. The successive exponents of r are $(0,1,0,1,0,1,1)$ for $(4,7)$ and $(1,2,2,2)$ for $(7,4)$. Thus, with composition read from right to left,

$$\Phi_{4,7} = (ur)^2u(ur)u(ur)u, \qquad \Phi_{7,4} = (ur^2)^3(ur).$$

Using the square numbering specified above, their complete cycle decompositions are listed below. Each parenthesis denotes one closed cycle; line breaks within a parenthesis only save space.

(p,q)	cycles of $\Phi_{p,q}$
$(4,7)$	$(0, 43, 47, 54, 52, 48, 41, 4, 13, 36, 59, 34, 38, 23, 11,$ $15, 32, 57, 19, 26, 6, 28, 24, 9, 21, 17, 30, 2, 50, 45),$ $(1, 27, 31, 29, 10, 5, 46, 49, 18, 22, 7, 44, 40, 42, 53,$ $3, 25, 16, 51, 55, 58, 37, 33, 35, 20, 56, 39, 14, 12, 8)$
$(7,4)$	$(0, 1, 47, 10, 13, 39, 52, 40, 32, 51, 59, 58, 38, 53, 19,$ $25, 21, 33, 41, 18, 6, 7, 24, 20, 50, 46, 11, 31, 30, 12),$ $(2, 29, 36, 3, 15, 14, 28, 8, 9, 44, 57, 35, 26, 5, 17,$ $16, 34, 22, 23, 27, 4, 42, 45, 56, 48, 49, 43, 55, 54, 37)$

The two displayed cycles for each direction cover all 60 squares and give the cycle lengths in the table. The included checker also regenerates these reduced permutations and compares them with `certificate.json`.

6.3 Eliminating all other sources

Fix one of these directions (p,q). On sheet (e_1,e_2), the transverse value at a lift of $A=(x,y)$ is $e_1qx - e_2py$ modulo integers. Thus the four values are

$$\pm(qx-py), \qquad \pm(qx+py).$$

If either $qx-py$ or $qx+py$ is nonintegral, a lift of A lies on a regular leaf, since every leaf through a marked point has integral transverse value. Lemma 4 then shows that A illuminates every other interior point. Therefore each endpoint of a dark pair must satisfy

$$7x-4y,\ 7x+4y,\ 4x-7y,\ 4x+7y\ \in \mathbb{Z}.$$

Adding and subtracting gives $14x, 8x, 8y, 14y \in \mathbb{Z}$, hence $2x, 2y \in \mathbb{Z}$. Write $x=a/2$, $y=b/2$ with $a,b\in\mathbb{Z}$. Integrality of $(7a+4b)/2$ forces a even; integrality of $(4a+7b)/2$ forces b even. Thus $(x,y)\in\mathbb{Z}^2$. Conversely every lattice point satisfies the four conditions.

The cell construction has only two interior lattice points, $S=(0,0)$ and $T=(2,2)$. This can be checked directly from the 15 squares: (x,y) is an interior lattice point exactly when all four cells with lower-left corners in $\{x-1,x\}\times\{y-1,y\}$ occur. The included program checks this as well. Every other interior point illuminates every other distinct interior point. The preceding darkness proof excludes a connection between S and T. This completes the proof of Theorem 1.

7 Scope and reproduction

The classification covers distinct real interior endpoints in the specified polygon. It does not classify boundary endpoints or self-return trajectories, and it gives no lower bound for other polygons. The darkness argument is analytic; the illumination argument combines the strip lemma and the integrality calculation with the two finite permutation certificates. Run

```
python3 verify_cylinders.py --check
```

from the manuscript directory to regenerate and compare both certificates. The program requires only Python 3 and the included JSON file. The illustrated companion `explainer.html` shows the strip construction and the exceptional-source conditions.

Acknowledgements and AI assistance. The authors used the AI systems ChatGPT (OpenAI) and Claude (Anthropic) extensively while exploring candidate geometries, developing explanations, and refining the proof. The authors take responsibility for the mathematical content of any eventual submitted version.

References

[1] G. W. Tokarsky, "Polygonal Rooms Not Illuminable from Every Point," *The American Mathematical Monthly* **102** (1995), no. 10, 867–879. https://doi.org/10.1080/00029890.1995.12004679.

[2] A. Wolecki, "Illumination in Rational Billiards," *arXiv:1905.09358v3* (2024), Theorem 1.3 and Figure 1. https://arxiv.org/abs/1905.09358v3.