

# Status of Shortspine Thornyhead (*Sebastolobus alascanus*) along the US West coast in 2023

by

Joshua A. Zahner<sup>1</sup>

Madison Heller-Shipley<sup>1</sup>

Sabrina Beyer<sup>1</sup>

Adam L. Hayes<sup>1</sup>

Pierre-Yves Hernvann<sup>2</sup>

Andrea N. Odell<sup>3</sup>

Haley Oleynik<sup>4</sup>

Jane Y. Sullivan<sup>5</sup>

Matthieu Veron<sup>6</sup>

<sup>1</sup>School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat Street,  
Seattle, Washington 98195

<sup>2</sup>Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and  
Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard  
East, Seattle, Washington 98112

<sup>3</sup>University of California Davis, One Shields Avenue, Davis, California 95616

<sup>4</sup>Institute for the Oceans and Fisheries, University of British Columbia, 2202 Main Mall,  
Vancouver, British Columbia Canada V6T 1Z4

<sup>5</sup>Alaska Fisheries Science Center, U.S. Department of Commerce, National Oceanic and  
Atmospheric Administration, National Marine Fisheries Service, 17109 Point Lena Loop  
Road, Juneau, Alaska 99801

<sup>6</sup>Alaska Fisheries Science Center, U.S. Department of Commerce, National Oceanic and  
Atmospheric Administration, National Marine Fisheries Service, 7600 Sand Point Way N.E.,  
Seattle, Washington 98115

© Pacific Fishery Management Council, 2023

Correct citation for this publication:

Zahner, J.A., M.A. Heller-Shipley, S. Beyer, A.L. Hayes, P-Y. Hernvann, A.N. Odell, H. Oleynik, J.Y. Sullivan, M. Veron. 2023. Status of Shortspine Thornyhead (*Sebastolobus alascanus*) along the US West coast in 2023. Pacific Fishery Management Council, Portland, Oregon. 109 p.

---

# Contents

<b>Executive summary</b>	<b>i</b>
Stock	i
Catches	i
Data and assessment	ii
Stock biomass and dynamics	iv
Recruitment	v
Exploitation status	vi
Ecosystem considerations	vii
Reference points	vii
Management performance	x
Unresolved problems and major uncertainties	xi
Decision table and projections	xi
Scientific uncertainty	xii
Research and data needs	xii
<b>1 Introduction</b>	<b>1</b>
1.1 Basic Information	1
1.2 Stock Structure	1
1.3 Life History	2
1.4 Ecosystem Considerations	2
1.5 Historical and Current Fishery Information	3
1.6 Summary of Management History and Performance	4
1.7 Foreign Fisheries	5
<b>2 Data</b>	<b>5</b>
2.1 Fishery-Dependent Data	6
2.1.1 Catch History	6
2.1.2 Discards and retention	7
2.1.3 Fishery Length Compositions	7
2.1.4 Age Compositions	8
2.2 Fishery-Independent Data	8
2.2.1 AFSC/NWFSC West Coast Triennial Shelf Survey	8
2.2.2 AFSC and NWFSC Slope Surveys	9
2.2.3 NWFSC West Coast Groundfish Bottom Trawl Survey	9
2.2.4 Survey Stratification	10
2.2.5 Design-based Indices of Abundance	10
2.2.6 Geostatistical Model-based Indices of Abundance	10
2.2.7 Length Composition Data	11
2.2.8 Frequency of Occurrence and Survey Information	11
2.3 Biological Data	11
2.3.1 Natural Mortality	11
2.3.2 Maturation and Fecundity	12
2.3.3 Length-Weight Relationship	14
2.3.4 Growth (Length-at-Age)	14

---

2.3.5	Ageing Precision and Bias . . . . .	15
2.4	Environmental and Ecosystem Data . . . . .	16
2.5	Changes in data from the 2013 assessment . . . . .	16
<b>3</b>	<b>Assessment Model</b>	<b>16</b>
3.1	Summary of Previous Assessments and Reviews . . . . .	16
3.1.1	History of Modeling Approaches . . . . .	16
3.1.2	Most Recent STAR Panel and SSC Recommendations . . . . .	16
3.1.3	Response to Groundfish Subcommittee Requests . . . . .	17
3.2	Model Structure and Assumptions . . . . .	17
3.2.1	Model Changes from the Last Assessment . . . . .	17
3.2.2	Modeling Platform and Structure . . . . .	17
3.2.3	Model Fleet Structure . . . . .	18
3.2.4	Model Parameters . . . . .	18
3.3	Base Model Results . . . . .	20
3.3.1	Parameter Estimates . . . . .	20
3.3.2	Fits to the Data . . . . .	21
3.3.3	Population Trajectory . . . . .	22
3.4	Model Diagnostics . . . . .	23
3.4.1	Convergence . . . . .	23
3.4.2	Sensitivity Analyses . . . . .	23
3.4.3	Retrospective Analysis . . . . .	25
3.4.4	Likelihood Profiles . . . . .	25
3.4.5	Unresolved Problems and Major Uncertainties . . . . .	26
<b>4</b>	<b>Management</b>	<b>27</b>
4.1	Reference Points . . . . .	27
4.2	Harvest Projections and Decision Tables . . . . .	27
4.3	Evaluation of Scientific Uncertainty . . . . .	27
4.4	Research and Data Needs . . . . .	28
<b>5</b>	<b>Acknowledgments</b>	<b>28</b>
<b>6</b>	<b>References</b>	<b>29</b>
<b>7</b>	<b>Tables</b>	<b>33</b>
7.1	Executive Summary . . . . .	33
<b>8</b>	<b>Figures</b>	<b>63</b>
8.1	Introduction and Data . . . . .	63
8.2	Base Model Results and Fits . . . . .	74
8.3	Likelihood Profiles, Retrospectives, and Sensitivity Analyses . . . . .	89

## Executive summary

### Stock

This assessment applies to shortspine thornyhead (*Sebastolobus alascanus*) off of the west coast of the United States from the U.S.-Canada border to the U.S.-Mexico border using data through 2023. Shortspine Thornyhead are modeled in this assessment as a single stock. They have been reported as deep as 1,524 m, and this assessment applies to their full depth range although survey and fishery data are only available down to 1,280 m.


### Catches

Landings of shortspine thornyhead rose to a peak of 4,815 mt in 1989, followed by a sharp decline during a period of trip limits and other management measures imposed in the 1990s. Since the institution of separate trip limits for shortspine and longspine thornyhead, the fishery has had more moderate removals of between 1,000 and 2,000 mt per year from 1995 through 1998. Landings fell below 1,000 mt per year from 1999 through 2006, then rose to 1,531 in 2009 and have declined since that time. Recreational fishery landings of thornyheads were negligible, so only commercial landings were included in the model.

Commercial landings are dominantly bottom trawl caught, and non-trawl landings include all other gear types, the majority of which is longline (Table i; Figure i). For this assessment, trawl landings were divided into northern (the waters off Washington and Oregon) and southern (the waters off California) fleets, and non-trawl landings represent one, coastwide, fleet. Discard rates (landings divided by total catch) for shortspine thornyhead have been estimated as high as 43% per year, but are more frequently below 20%. Discard rates in the trawl fleets, available from the West Coast Groundfish Observer Program (WCGOP), declined from 2003–2011 and have since increased from 2012-present, when the catch shares system began.

**Table i:** Recent landings by fleet, total landings summed across fleets, and the total mortality including discards.

Year	North Trawl	South Trawl	Non-Trawl	Total Landings	Total Dead
2013	547.98	294.83	166.40	1,009.21	1,085.62
2014	433.12	254.05	147.81	834.98	900.66
2015	503.14	244.29	131.30	878.73	945.40
2016	577.19	185.73	168.94	931.86	1,012.75
2017	606.86	158.30	223.82	988.97	1,085.60
2018	525.04	105.07	184.48	814.60	895.39
2019	402.95	127.94	143.48	674.37	736.82
2020	248.47	87.99	85.17	421.64	458.87
2021	226.00	73.39	78.74	378.13	411.62
2022	261.16	97.61	66.22	424.98	456.65



**Figure i:** Estimated landing history for shortspine thornyhead.

## Data and assessment

The most recent assessment for shortspine thornyhead was conducted in 2012 (Taylor and Stephens 2013). Stock status was determined to be above the management target and catches did not attain the full management limits, so reassessment of thornyheads has not been a higher priority. This assessment uses Stock Synthesis (Methot and Wetzel 2013) Version 3.30.21, used in many other recent west coast assessments.


Data were divided into three fishery fleets: North trawl (the waters off Washington and Oregon), South trawl (the waters off California), and coastwide Non-trawl, and three survey fleets: the AFSC/NWFSC West Coast Triennial Shelf Survey (Triennial Survey) from 1980–2003, which was divided into early (pre-1995) and late period (post-1995) to account for a change in depth-sampling, and the NWFSC West Coast Groundfish Bottom Trawl Survey (WCGBTS), 2004–2022 (Figure ii).

Most data used in the 2013 assessment were newly pulled and processed for this assessment, including length compositions from all fishing and survey fleets, indices of abundance derived from new geostatistical analyses, discard rates from both a 1980s observer study (Pikitch et al. 1988) and the current WCGOP, historical catch data from Washington, Oregon, and California, and all reported catches from 1981–2022. The only data taken from the previous assessment without reanalysis were discard rates from the Enhanced Data Collection Project (EDCP) study in the 1990s.

New maturity analyses of samples collected in the WCGOP in 2011, 2013, 2014, 2016 and 2018 were available for this assessment (M. Head, pers. comm.). The larger number and

better spatial coverage of these samples allowed the use of statistical modeling to better understand the spatial variation in the proportion of female spawning. This assessment also assumes a new fecundity relationship, in which fecundity increases with body size. New growth curves were estimated, using data from Butler (1995), which were similar to the curves assumed in the 2005 and 2013 assessments. In the previous assessment, a Beverton-Holt stock recruitment relationship was assumed and steepness ( $h$ ) was fixed at 0.60. This assessment fixed steepness at 0.72, as recommended by Thorson et al. (2019). Natural mortality ( $M$ ) was also slightly updated from the 2013 assessment to be fixed at 0.04.


This assessment estimated 180 parameters. The log of the unfished equilibrium recruitment,  $\ln(R_0)$ , controls the scale of the population and annual deviations around the stock-recruit curve (135 parameters) allow for more uncertainty in the population trajectory. In addition, 43 selectivity and retention parameters for the three fishery fleets and three surveys allowed for estimation of annual length compositions and discards rates. Two catchability parameters were analytically computed from the data, and one additional parameter, representing additional variability in the early Triennial survey, was directly estimated by the model.



**Figure ii:** Summary of data sources used in the base model.

## Stock biomass and dynamics

Unfished equilibrium spawning output ( $B_0$ ) is estimated to be 20.332 trillion eggs, with a 95% confidence interval of 16.338–24.327 trillion eggs. The  $B_0$  estimate here is not comparable to previous assessment as the integration of new fecundity and maturity assumptions have changed the output units from traditional biomass to spawned eggs. Spawning biomass is estimated to have remained stable until the early-1970s before beginning to decline near linearly through the present day. The estimated spawning output in 2023 is 8.372 trillion eggs (5.346–11.201 trillion eggs), which represents a stock status or “depletion” ( $B_{2023}/B_0$ ) of 41.4% (33%–49%; Table ii; Figure iii). The depletion in 2013 was estimated to be 43.6%, a large decrease from what was estimated in 2013 (~75%). The standard deviation of the log of spawning biomass in 2023 is 0.18, which is less than the 0.36 minimum assumed for use in  $p^*$  adjustments to overfishing limit (OFL) values.



**Figure iii:** Estimated spawning output trajectory for shortspine thornyhead.

**Table ii:** Spawning output (millions of eggs) and fraction unfished with associated 95% confidence intervals (CI) from the base model.

Year	Spawning Output	Spawning Output 95% CI	Fraction Unfished	Fraction Unfished 95% CI
2013	8,875	5,904–11,845	0.4	0.4–0.5
2014	8,767	5,807–11,727	0.4	0.4–0.5

**Table ii:** Spawning output (millions of eggs) and fraction unfished with associated 95% confidence intervals (CI) from the base model. (*continued*)


Year	Spawning Output	Spawning Output 95% CI	Fraction Unfished	Fraction Unfished 95% CI
2015	8,679	5,728–11,630	0.4	0.3–0.5
2016	8,593	5,650–11,536	0.4	0.3–0.5
2017	8,508	5,572–11,445	0.4	0.3–0.5
2018	8,423	5,492–11,355	0.4	0.3–0.5
2019	8,358	5,431–11,286	0.4	0.3–0.5
2020	8,311	5,386–11,236	0.4	0.3–0.5
2021	8,291	5,366–11,215	0.4	0.3–0.5
2022	8,280	5,355–11,205	0.4	0.3–0.5
2023	8,273	5,346–11,201	0.4	0.3–0.5

## Recruitment

This assessment assumed a Beverton-Holt stock recruitment relationship. Steepness (the fraction of expected equilibrium recruitment associated with 20% of equilibrium spawning biomass) was fixed at 0.72, slightly higher than what was assumed in previous assessments ( $h = 0.60$ ). The scale of the population is largely determined by the log of unfished recruitment ( $R_0$ ), which was estimated to be 9.354. This results in an unfished recruitment of 11,550,000 recruits (9,281,000–13,820,000). Recruitment deviations were estimated for the years 1901 through 2022, and ranged from -0.5 to 1.5 on the log scale. Estimated recruitments do not show high variability, and the uncertainty in each estimate is greater than the variability between estimates (Table iii; Figure iv).

**Table iii:** Estimated recent trend in recruitment and recruitment deviations and the 95% confidence intervals (CI) from the base model.

Year	Recruitment	95% CI	RecDevs	RecDev 95% CI
2013	6,024	2,469–14,698	-0.439	-1.352–0.474
2014	5,962	2,446–14,532	-0.447	-1.358–0.465
2015	5,954	2,438–14,542	-0.446	-1.360–0.468
2016	6,057	2,465–14,886	-0.427	-1.349–0.495
2017	5,836	2,385–14,279	-0.462	-1.379–0.454
2018	5,745	2,346–14,069	-0.476	-1.393–0.442
2019	8,863	3,557–22,086	-0.064	-1.003–0.874
2020	9,536	3,760–24,183	-0.013	-0.973–0.946
2021	10,335	3,984–26,811	0.044	-0.943–1.032
2022	10,118	3,924–26,090	0.000	-0.980–0.980
2023	10,117	3,924–26,086	0.000	-0.980–0.980



**Figure iv:** Estimated recruitment timeseries.

### Exploitation status


The summary harvest rate (total catch divided by age-1 and older biomass) closely follows the landings trajectory. The harvest rates are estimated to have never exceeded 5% and have remained below 2% in the past decade. Expressing exploitation rates in terms of spawning potential ratio (SPR) indicates that the exploitation consistently exceeded the  $SPR_{50\%}$  reference point from 1980-2018. However, the stock status is estimated to have never fallen below the  $B_{40\%}$  management target, though the uncertainty interval around the 2023 estimate does encapsulate the  $B_{40\%}$  target (Table iv; Figures v-viii).

**Table iv:** Estimated recent trend in the  $(1-SPR)/(1-SPR 50\%)$  where SPR is the spawning potential ratio the exploitation rate, and the 95% intervals.

Year	$(1-SPR)/(1-SPR 50\%)$	95% CI	Exploitation Rate	95% CI
2013	1.25	1.03–1.47	0.0124	0.0084–0.0165
2014	1.12	0.90–1.34	0.0103	0.0069–0.0137
2015	1.15	0.92–1.37	0.0109	0.0073–0.0145
2016	1.19	0.96–1.42	0.0117	0.0078–0.0155
2017	1.23	1.00–1.46	0.0125	0.0083–0.0167
2018	1.09	0.86–1.32	0.0103	0.0069–0.0138
2019	0.95	0.73–1.17	0.0085	0.0056–0.0114

**Table iv:** Estimated recent trend in the (1-SPR)/(1-SPR 50%) where SPR is the spawning potential ratio the exploitation rate, and the 95% intervals. (*continued*)

Year	(1-SPR)/(1-SPR 50%)	95% CI	Exploitation Rate	95% CI
2020	0.66	0.48–0.84	0.0053	0.0035–0.0071
2021	0.59	0.43–0.76	0.0047	0.0031–0.0063
2022	0.64	0.47–0.81	0.0052	0.0034–0.0070




**Figure v:** Estimated relative spawning output trajectory for shortspine thornyhead.

## Ecosystem considerations


This stock assessment does not explicitly incorporate trophic interactions, habitat factors or environmental factors into the assessment model. More predation, diet, and habitat work, and mechanistic linkages to environmental conditions would be needed to incorporate these elements into the stock assessment.

## Reference points


Reference points were calculated using the estimated catch distribution in the final year of the model (2023). In general, the population is on the boundary between “precautionary” ( $B/B_0 = 0.40$ ) and “healthy” ( $B/B_0 > 0.40$ ) status relative to the reference points (Figure



**Figure vi:** Summary F rate.



**Figure vii:** Estimated spawning potential ratio.



**Figure viii:** Phase diagram.


ix). Sustainable total yield (landings plus discards) was estimated at 1,060 mt when using an  $SPR_{50\%}$  reference harvest rate and ranged from 870–1,250 mt based on estimates of uncertainty (Table v). The spawning output equivalent to 40% of the unfished spawning output ( $B_{40\%}$ ) was 8.273 trillion eggs. The most recent total mortality (landings plus discards) have been lower than the estimated long-term yields calculated using an  $SPR_{50\%}$  reference point, but not as low as the lower bound of the 95% uncertainty interval. However, this is due to the fishery not fully attaining the full annual catch limit (ACL). The OFL and Acceptable Biological Catch (ABC) values over the past 6 years have been approximately 3100 mt and 2,500 mt, respectively.

**Table v:** Summary of reference points and management quantities, including estimates of the 95% intervals.

Variable of Interest	Estimate	95% CI
Unfished Spawning Output	20,332	16,338–24,327
Unfished Age 1+ Biomass (mt)	196,023	157,510–234,536
Unfished Recruitment (R0)	11,550	9,281–13,820
Spawning Output (2023)	8,273	5,346–11,201
Fraction Unfished (2023)	0.41	0.33–0.49
Reference Points Based SB40%		
Proxy Spawning Output SB40%	8,133	6,535–9,731
SPR Resulting in SB40%	0.458	0.458–0.458

**Table v:** Summary of reference points and management quantities, including estimates of the 95% intervals. (*continued*)

Variable of Interest	Estimate	95% CI
Exploitation Rate Resulting in SB40%	0.012	0.011–0.012
Yield with SPR Based On SB40% (mt)	1,060	869–1,251
Reference Points Based on SPR Proxy for MSY		
Proxy Spawning Output (SPR50)	9,071	7,289–10,854
SPR50	0.500	-
Exploitation Rate Corresponding to SPR50	0.010	0.010–0.011
Yield with SPR50 at SB SPR (mt)	1,013	831–1,195
Reference Points Based on Estimated MSY Values		
Spawning Output at MSY (SB MSY)	5,651	4,548–6,755
SPR MSY	0.348	0.345–0.351
Exploitation Rate Corresponding to SPR MSY	0.017	0.016–0.017
MSY (mt)	1,121	919–1,323



**Figure ix:** Estimated yield curve with reference points.

## Management performance

Catches for shortspine thornyhead have not fully attained the catch limits in recent years. ACLs have hovered around 2500 mt since 2013, while total mortality has never exceeded 1085

mt, and is often smaller than that (Table vi). The fishery for shortspine thornyhead may be limited more by the ACLs on sablefish, with which they co-occur, and by the challenging economics of deep sea fishing, than by the management measures currently in place.

**Table vi:** Recent trend in the overfishing limits (OFLs), the acceptable biological catches (ABCs), the annual catch limits (ACLs), the total landings, and total mortality (mt).

Year	OFL	ABC	ACL	Landings	Total Mortality
2013	2333	2230	1937	1,009.21	1,085.62
2014	2310	2208	1918	834.98	900.66
2015	3203	2668	2668	878.73	945.40
2016	3169	2640	2639	931.86	1,012.75
2017	3144	2619	2619	988.97	1,085.60
2018	3116	2596	2596	814.60	895.39
2019	3089	2573	2573	674.37	736.82
2020	3063	2551	2552	421.64	458.87
2021	3211	2183	2184	378.13	411.62
2022	3194	2130	2130	424.98	456.65

### Unresolved problems and major uncertainties

Major uncertainties in the model are centered around uncertainty in biological processes including growth, maturity, and mortality. The absence of reliable ageing methods for shortspine thornyhead, particularly, makes it difficult to estimate growth and natural mortality. Sensitivities demonstrated that changes to the growth curve have large effects on the estimated stock status. Likelihood profiles over natural mortality demonstrate the model to be quite sensitive to its assumed value. There is insufficient information in the data to estimate mortality directly, constraining us to use meta-analyses or other mortality estimators, which frequently make use of aging information that is unavailable, and again, highly uncertain for shortspine thornyhead. Due to imperfect seasonal and spatial coverage of histological data for shortspine thornyhead, there is significant uncertainty about the shape of the species' maturity curve, though the model appears to be largely insensitive to variations in maturity.

This model fails to fully capture the observed increase in abundance seen in the WCGBTS index time series in 2021 and 2022. The model also fails to fully capture the peak of the length compositions for the North trawl fleet, underestimating the number of mid-sized fish that the fleet takes. This underestimation appears to be consistent, particularly in the last 10 years, implying a possible recent change in selectivity.

### Decision table and projections

TBD

## **Scientific uncertainty**

The model estimated uncertainty around the 2024 spawning biomass was  $\sigma = 0.18$  and the uncertainty around the OFL was  $\sigma = 0.17$ . The category 2 default  $\sigma = 1.0$  is used to apply scientific uncertainty in the projections.

## **Research and data needs**

Research and data needs for future assessments include the following:

1. Research into aging methods and availability of reliable age data would be valuable for future stock assessments. Otoliths have been collected in good quantities from the WCGBTS, but there is currently no validated aging method for shortspine thornyhead.
2. Additional investigation into growth patterns would provide valuable information for future population projections. We acknowledge that additional work on aging shortspine thornyhead would be required to make such additional growth research possible.
3. More investigation into maturity of shortspine thornyhead is necessary to understand the patterns in maturity observed in WCGBTS samples.
4. Information on possible migration of shortspine thornyhead would be valuable for understanding stock dynamics. Analysis of trace elements and stable isotopes in shortspine thornyhead otoliths may provide valuable information on the extent of potential migrations. Possible connections between migration and maturity could likewise be explored.
5. A greater understanding of the connection between thornyheads and bottom type could be used to refine the indices of abundance. Thornyheads are very well sampled in trawlable habitat, but the extrapolation of density to a survey stratum could be improved by accounting for the proportion of different bottom types within a stratum and the relative density of thornyheads within each bottom type.
6. Additional investigation into spatial stock structure could be valuable for determining whether future assessments should develop a spatial assessment model, or if shortspine thornyhead should be assessed at distinct spatial scales in the fut

# 1 Introduction

## 1.1 Basic Information

This assessment reports the status of shortspine thornyhead (*Sebastolobus alascanus*) off the US West coast using data through 2022.

Shortspine Thornyhead (*Sebastolobus alascanus*) are found in the waters off the West Coast of the United States from northern Baja California to the Bering Sea at depths of 20 meters to over 1,500 meters. The majority of the spawning biomass occurs in the oxygen minimum zone between 600 and 1,400 meters. The distribution of the smallest shortspine thornyhead suggests that they tend to settle at around 100–400 meters and are believed to have ontogenetic migration down the slope, although large individuals are found across the depth range. Higher densities (kg/ha) of shortspine thornyhead occur in shallower areas (under 500 meters) off Oregon and Washington, whereas in California, they occur in deeper areas (**above x meters**; Figure 1).

Despite variation in density across the coast, shortspine thornyheads are present in almost all trawlable areas below 500 meters. They are caught in 91% of trawl survey hauls deeper than 500 m and **XX%** of commercial bottom trawl hauls deeper than 500m. Camera-tows show that thornyheads are spaced randomly across the sea floor, indicating a lack of schooling and territoriality (Wakefield 1990; Du Preez and Tunnicliffe 2011).

## 1.2 Stock Structure

**NOTE: This section was added.**

Genetic studies of stock structure show few genetic differences among shortspine thornyhead along the Pacific coast, and thus do not suggest separate stocks Stepien (1995). Stepien (1995) suggested that there may be a separate population of shortspine thornyhead in the isolated area around Cortes Bank off San Diego, California. Stepien (1995) also pointed out that juvenile dispersion might be limited in the area where the Alaska and California currents split, which occurs towards the northern boundary of the assessment area, near 48° N.

Stepien et al. (2000), using a more discerning genetic material (mtDNA), found evidence of a pattern of genetic divergence in shortspine thornyhead corresponding to geographic distance. However, this study, which included samples collected from southern California to Alaska, did not identify a clear difference between stocks even at the extremes of the range. No such pattern was seen in longspine thornyhead, which suggests that the shorter pelagic stage (~1 yr vs. ~2 yrs) of shortspine thornyhead may contribute to an increased genetic separation with distance.

Dorval et al. (2022) applied otolith microchemistry to immature fish to redefine population structure of shortspine thornyhead on the west coast. Their results indicate that the

population of immature shortspines belongs to two distinct groups distributed north and south of Cape Mendocino.

### 1.3 Life History

Shortspine Thornyheads along the West Coast spawn pelagic, gelatinous floating egg masses between December and May (Wakefield 1990; Erickson and Pikitch 1993; Pearson and Gunderson 2003). Cooper et al. (2005) and Pearson and Gunderson (2003) found no evidence for batch spawning in this species on the West Coast, but more recent histological examination of ovaries suggest that some shortspine thornyhead can be batch spawners with two to three batches developing simultaneously (Melissa Head, Northwest Fisheries Science Center (NWFSC), pers. comm.). Juveniles settle at around 1 year of age (22- 27 mm in length), likely in the range of 100-200 m (Vetter and Lynn 1997), and migrate down the slope with age and size, although large individuals are found across the depth range.

Shortspine Thornyhead are notoriously challenging to age, and a recent age validation study using  $^{14}\text{C}$  bomb radiocarbon was inconclusive (Kastelle et al. 2020). However, best available data suggests that the shortspine thornyhead life span may exceed 100 y (Butler 1995; Kline 1996). Estimates of natural mortality for shortspine thornyhead range from 0.013 (Pearson and Gunderson 2003) to 0.07 (Kline 1996). However, Pearson and Gunderson's estimate is based upon a regression model, using the gonadosomatic index as a proxy. Butler (1995) estimated M to be 0.05 based upon a maximum lifespan of 100 years for shortspine thornyhead. Butler (1995) also suggested that M may be lower for older, larger shortspine thornyhead residing in the oxygen minimum zone due to lack of predators. All estimates of M for thornyheads are highly uncertain.

Shortspine Thornyhead grow very slowly and may continue growing throughout their lives, reaching maximum lengths of over 70 cm. Females grow to larger sizes than males. Maturity in females has been estimated as occurring near 18 cm, with fish transitioning from immature to mature within a relatively narrow range of sizes between 15 and 20 cm Pearson and Gunderson (2003). However, more recent histological data collected in the WCGBTS and analyzed using current best practices suggests that functional maturation, which accounts for abortive maturation and skip spawning, occurs over a broader spectrum of sizes between 10 and 55 cm (length-at-50% maturity,  $L_{50} = 31.4$ ; personal communication, Melissa Head, NWFSC, pers. comm.).

### 1.4 Ecosystem Considerations

Shortspine Thornyheads have historically been caught alongside longspine thornyheads in a Dover sole, thornyhead, and trawl-caught sablefish complex (DTS). Other groundfishes that frequently co-occur in deep waters include a complex of slope rockfishes, Rex sole, longnose skate, roughtail skate, Pacific grenadier, giant grenadier, and Pacific flatnose. Non-groundfish species such as Pacific hagfish and a diverse complex of eelpouts also co-occur with shortspine thornyhead.

Shortspine Thornyheads typically occur in shallower water than the shallowest longspine

thornyheads, and migrate to deeper water as they age. The majority of spawning shortspine thornyheads occur between 600 and 1,400 meters, where longspine thornyheads are most abundant (Jacobson and Vetter 1996; Bradburn et al. 2011). When shortspine thornyheads have reached a depth where they overlap with longspine thornyheads, they are typically larger than the largest longspine thornyheads.

Species distribution models developed by Liu et al. (in press) suggest that expected environmental changes over the next decades will lead to a decline in shortspine and increase in longspine abundance. Shortspine Thornyheads are also projected to shift offshore, into deeper waters, potentially decreasing their availability in fisheries. To date, shortspine thornyheads have been observed by cameras below the 1280 meter limit of the current fishery and survey, but their distribution, abundance, and ecosystem interactions in these deep waters are relatively unknown. Thornyheads spawn gelatinous masses of eggs which float to the surface, which may represent a significant portion of the upward movement of organic carbon from the deep ocean (Wakefield 1990).

Shortspine Thornyhead diet composition, as derived from stomach content collection in the 1980s and 1990s, varied by year (Bizzarro et al. 2023). In some years their diet consisted primarily of invertebrate species including pandalid shrimp, pink shrimp, and Tanner crab, while in others their stomach content was dominated by finfish species such as Pacific cod and Pacific Hake. As prey themselves, shortspine thornyheads were only found in the stomachs of other species in two years, 1991 and 1992 as recorded in the CA Current Trophic Database (CCTD), where shortspine thornyhead occurred in sablefish, Pacific hake, and other shortspine thornyhead stomachs (Bizzarro et al. 2023).

## 1.5 Historical and Current Fishery Information

Thornyhead harvest has experienced fluctuations over time due to increased depth range of the fisheries, variable markets, and changes in fisheries management. In the early 1900's, landings were minimal because there were few markets for thornyheads and relatively little trawling at depths where the majority of thornyheads occur. Beginning in the 1930s, thornyhead landings increased as they were landed as incidental catch in the California sablefish fishery. The first significant market for thornyheads began in northern California in the early 1960s, when larger (30-35 cm) thornyhead were sold as "ocean catfish." By the early 1980s, the minimum marketable size decreased to 25 cm, and in the late 1980s a market for small thornyheads (~20 cm) developed due to the depletion of a related species (*Sebastolobus machrochir*) off the coast of Japan. The fishery moved into deeper waters with the demand for smaller thornyheads and began catching more longspine thornyheads. This is reflected in the changes in proportion of shortspine to total thornyheads through time, which decreased from around 90% in 1981 to 40% in 1994 Figure 2.

Landings of shortspine thornyheads off the coast of California peaked around 3,500 mt in 1989, and have exceeded those from further north in most years (Figure 3). In the northern area off of Oregon and Washington, the fishery grew in the early 1980s, with landings peaking in 1991 at around 2200 mt.

Non-trawl landings of shortspine thornyheads were relatively low prior to the mid-1990s, at which point non-trawl landings, dominantly longline, in California began to increase steadily from less than 5 mt in 1994 to 237 mt in 2011. The increase in non-trawl landings was driven by the development of live-fish markets for thornyheads and the fact that ex-vessel prices associated with the non-trawl landings are much higher than those for the trawl fishery.

**Nominal prices for line-caught shortspine thornyhead increased steadily from \$0.69/lb in 1993 to \$3.81/lb in 2008, and have remained near or above that level since.** Citation?

Trawl prices, on the other hand, changed from \$0.46/lb to \$0.72/lb in the same period, although, when Japanese demand was strong they were between \$0.80 and \$1.06/lb. In contrast, non-trawl landings of shortspine in Washington and Oregon have remained below the estimated peak of 54 mt in 1991.

The foreign fishery off of the West Coast is estimated to have caught approximately 7,400 mt of shortspine thornyhead during the 11 year period from 1966-1976 (Rogers 2003), which is on the order of the estimate of domestic catch (~8,600 mt) during that same period.

**Management measures** have contributed to a decline in coastwide landings from an estimated peak of 4,815 mt in 1989 to between 1,000 and 2,000 mt per year from 1995 through 1998. Landings fell below 1,000 mt per year from 1999 through 2006, then rose to 1,531 in 2009 and have declined since (Table 9).

In 2011, the west coast trawl fishery was rationalized, with the introduction of the Individual Fishing Quota (IFQ) Program. In order to provide more flexibility for fishers on the west coast, NOAA Fisheries implemented the West Coast Groundfish Trawl Fishery Catch Share Program, which allows for the division of catch allocated to the trawl fishery into shares controlled by individuals or cooperatives (West Coast Regional Office n.d.). All vessels that participate in the IFQ program are required to have 100% observer coverage at all times the vessels are at sea.(West Coast Regional Office n.d.)

## 1.6 Summary of Management History and Performance

Beginning in 1989, both thornyhead species were managed as part of a DTS. In 1991, the Pacific Fishery Management Council (PFMC) adopted separate ABC levels for thornyheads and catch limits were imposed on the thornyhead complex. harvest guideline (HG) were instituted in 1992 along with an increase in the minimum mesh size for bottom trawl fisheries. In 1995 separate landing limits were placed on shortspine and longspine thornyheads and trip limits became more restrictive. Trip limits (predominantly 2-month limits on cumulative vessel landings) have often been adjusted during the year since 1995 in order to not exceed the HG or optimum yield (OY). At first, the HG for shortspine thornyhead was set higher than the ABC (1,500 vs. 1,000 mt in 1995-1997) in order to allow a greater catch of longspine thornyhead, which was considered relatively undepleted. In 1999 the OY was set at less than 1,000 mt and remained close to that level through 2006. As a result of the 2005 shortspine

assessment, catch limits increased to about 2,000 mt per year and have remained near that level **to the present**.

Since early 2011, trawl harvest of each thornyhead species has been managed under the PFMC's catch share, or individual fishing quota (IFQ), program. Whereas the trip limits previously used to limit harvest restricted only the amount of fish each vessel could land, individual vessels fishing under the catch-share program are now held accountable for all of the quota-share species they catch.

**Landings of shortspine thornyhead have been below the catch limits since 1999. The estimated total catch, including discards, has likewise remained below the limit during this period.**

## 1.7 Foreign Fisheries

The Alaska Fisheries Science Center (AFSC) conducts assessments of thornyheads as a mixed stock complex, including shortspine and longspine thornyheads. Results of the 2022 Alaska Thornyhead complex assessment suggest that thornyheads are not being subjected to overfishing (Echave et al. 2022).

## 2 Data

Data comprise the foundational components of stock assessment models. The decision to include or exclude particular data sources in an assessment model depends on many factors. These factors often include, but are not limited to, the way in which data were collected (e.g., measurement method and consistency); the spatial and temporal coverage of the data; the quantity of data available per desired sampling unit; the representativeness of the data to inform the modeled processes of importance; timing of when the data were provided; limitations imposed by the Terms of Reference; and the presence of an avenue for the inclusion of the data in the assessment model. Attributes associated with a data source can change through time, as can the applicability of the data source when different modeling approaches are explored (e.g., stock structure or time-varying processes). Therefore, the specific data sources included or excluded from this assessment should not necessarily constrain the selection of data sources applicable to future stock assessments for shortspine thornyhead. Even if a data source is not directly used in the stock assessment they can provide valuable insights into biology, fishery behavior, or localized dynamics.

Data from a wide range of programs were available for possible inclusion in the current assessment model. Descriptions of each data source included in the model (Figure 4) and sources that were explored but not included in the base model are provided below. Data that were excluded from the base model were explicitly explored during the development of this stock assessment or have not changed since their past exploration in a previous shortspine thornyhead stock assessment. In some cases, the inclusion of excluded data sources were explored through sensitivity analyses.

## 2.1 Fishery-Dependent Data

### 2.1.1 Catch History

Pacific Fisheries Information Network (PacFIN) data from 1981-present was used to estimate landings in the North (Oregon and Washington) and South (California) by gear type (Trawl and Non-Trawl) (Figure 3). All landings reported for the shortspine thornyhead and nominal shortspine thornyhead categories were considered shortspine thornyhead, whereas landings categorized as unidentified thornyheads were split between longspine thornyhead and shortspine thornyhead by the ratio of identified longspine and shortspine landings for each year-state-gear combination. The values of this ratio for each state and gear-type from 1981-2023 are shown in Figure 2.

Catches prior to 1981 are based on historical reconstructions provided by the respective states and a reconstruction of foreign fleet catch. Oregon landings for 1892-1986 are provided by ODFW and outlined in Karnowski et al. (2014) shortspine thornyhead landings are not present in the PacFIN data for Oregon for the years 1981-1986 and the state reconstruction is used for this period instead. Washington landings for 1954-1980 are provided by WDFW. Landings prior to the beginning of this data are assumed to be zero. California landings are provided by CDFW and SWFSC, and consist of California commercial data for 1969-1980, and a catch reconstruction documented by Ralston et al. (2010) for 1934-1968. As in the two previous assessments, catch data from Rogers (2003) is used to account for catches by foreign fleets during the years 1966-1976. Foreign catch in the Monterey and Eureka International North Pacific Fishery Commission (INPFC) areas is attributed to the Southern Trawl fleet, while foreign catch in Columbia and Vancouver areas is attributed to the Northern Trawl fleet, as was the case in the 2013 assessment.

For historical catches prior to 1981, all shortspine thornyhead, nominal shortspine, and unidentified thornyhead landings in the state catch reconstructions are considered shortspine thornyhead. Neither California reconstructions prior to 1978, nor the Karnowski et al. (2014) reconstruction for Oregon, distinguish between shortspine and longspine thornyhead species. It is possible that assigning all thornyhead landings to shortspine overestimates shortspine landings, however, the overwhelming majority of thornyhead landings were shortspine until the late 1980s when vessels began to move into deeper waters and a distinct fishery targeting longspine thornyhead developed (Hamel 2005; Karnowski et al. 2014).

This treatment of unassigned thornyhead landings differs from the 2005 and 2013 assessments. The 2005 assessment did not have access to the historical reconstructions used here, and instead imputed shortspine landings as 30% of annual sablefish landings for the years 1901-1961. The 2013 assessment used the same imputed values as the 2005 assessment, but also conducted a sensitivity analysis in which all unassigned thornyheads in historical catch were considered shortspine thornyhead. Stock abundance estimates were found to be largely insensitive to which reconstructions were used (Taylor and Stephens 2013). The imputed historical values used for the 2005 and 2013 assessments will continue to be included as a sensitivity analysis here. Landings after 1961 remain very similar to the landings used in the 2013 assessment (Figure 3).

### **2.1.2 Discards and retention**

Predicted discards were based on estimated retention and selectivity for each fleet and are shown in Figure 5. Discards were informed by four data sources covering three different periods. Data sets included, 1) Pikitch et al. (1988) Discard and Mesh Studies, used to estimate both discard rates and length composition of the northern trawl fleet between 1985 and 1987 (J. R. Wallace, pers. comm.), 2) the EDCP covering 1995-1999, which only informed discard rates of the northern trawl fleet, 3) the WCGOP, which provided discard rates, length composition, and individual average weight for years between 2002 and 2021 for all fleets, and 4) the Groundfish Expanded Mortality Multi-Year (GEMM) data set, covering the same period and completing the WCGOP with catch-share participation information and estimates of discard survival rates.

While the estimates from the first two data sets were directly integrated into the model, fleet discard rates after 2011 were available separately for catch-share and non-catch-share programs. Final fleet-specific discard rates were thus computed as the average WCGOP discard rate weighted by the relative proportion of total landings belonging to the catch-share and non-catch-share, respectively. (Figure 6). Regardless of the type of data, all estimates derived from these data sets had associated uncertainty accounting for the variability observed within the sample of hauls and fishing trips of each fleet. WCGOP-derived discard rates are an exception as, after the catch share program was initiated in 2011, 100% of hauls from catch share fleets were observed., while non-catch share vessels were only partially covered (West Coast Regional Office n.d.).

The discard data sources were the same as those used in the 2013 assessment. The main improvements are the increased representativity of all 4 fleets (11 more years) and more accurate estimates of discard rates from EDCP that were not ready at the time of the previous assessment. Last, some errors in the previous assessment were corrected regarding the weight units considered for the average individual weight (WCGOP provides weight as pounds and not as kg).

### **2.1.3 Fishery Length Compositions**

Commercial fishery length-composition data were obtained from PacFIN for 1978-2023. Due to variations in sampling effort and because the number of fish sampled by port samplers is not proportional to the amount of landed catch in each trip, the observed length data were expanded using the following algorithm using the PacFIN.Utilities package in R:

1. Length data were acquired at the trip level by sex, year and state.
2. The raw numbers in each trip were scaled by a per-trip expansion factor calculated by dividing the total weight of trip landings by the total weight of the species sampled.
3. A per-year, per-state expansion factor was computed by dividing the total weight of state landings by the total weight of the species sampled for length in the state.
4. The per-trip expanded numbers were multiplied by the per-state expansion factor and summed to provide the coast-wide length-frequency distributions by year.

Only randomly collected samples were used. The sample sizes associated with the length compositions from the fishing fleets are shown in Table 12 (landings) and Table 13 (discards). Length samples from the Trawl North fleet in 1980, 1994, and 1995 showed a very different pattern than the surrounding years. The effective sample sizes for these years were substantially lower than other years ( $N_{eff} < 15$ ), so the observed differences are likely due to non-representative sampling.

Input sample sizes  $N_{input}$  for fishery length frequency distributions by year were calculated as a function of the number of trips and number of fish via the Stewart Method (Stewart, pers.com):

$$N_{input} = N_{trips} + 0.138N_{fish} \quad \text{when } \frac{N_{fish}}{N_{trips}} < 44$$

$$N_{input} = 7.06N_{trips} \quad \text{when } \frac{N_{fish}}{N_{trips}} \geq 44$$

The method is based on analysis of the input and model-derived effective sample sizes from west coast groundfish stock assessments. A piece-wise linear regression was used to estimate the increase in effective sample size per sample based on fish-per-sample and the maximum effective sample size for large numbers of individual fish.

All length data from commercial fisheries included in the model with sexes combined. This avoids the possibility of bias due to difficulty in sex determination of thornyheads.

#### 2.1.4 Age Compositions

No age composition data was used for this assessment because thornyheads have proven very difficult to age (P. MacDonald, pers. comm.). Even in directed studies such as those done by Kline (1996) and Butler (1995), there are large inter-reader differences, and a second reading by the same reader can produce a markedly different result. Kline (1996) reported only about 60% of the multiple reads were within 5 years of each other, and inter-reader differences were as large as 24 years for a sample of 50 otoliths. No production ageing of thornyheads is undertaken at this time for the west coast, although shortspine thornyhead otoliths are routinely collected in the NWFSC trawl survey.

### 2.2 Fishery-Independent Data

Four trawl surveys have been conducted on the U.S. west coast over the past four decades.

#### 2.2.1 AFSC/NWFSC West Coast Triennial Shelf Survey

The AFSC conducted a triennial groundfish trawl survey (the “triennial” survey) on the continental shelf from 1977 to 2001, although the 1977 survey had incomplete coverage and is not believed to be comparable to the later years. A final survey was conducted in 2004 by the NWFSC using the same survey design. In 1995, the timing of the survey shifted so that instead of occurring between mid-July and late September, it was conducted from early

June through mid-August. The years 1980–1992 had a maximum depth of 366 m, while from 1995 onward, the maximum depth was extended to 500 m. The shallow limit of the survey was 55 m in all years, but for purposes of computing indices, only tows deeper than 100 m were used as shortspine thornyhead are rarely seen at shallower depths. The triennial survey consists of 9 data points, from surveys operating every third year from 1980–2004.

For some species, the shift in timing between the 1992 and 1995 surveys would be expected to influence their catchability, availability, or distribution. However, thornyheads are believed to be sedentary enough that the change in timing would not be as influential. On the other hand, the increase in depth is expected to significantly increase the range of shortspine thornyhead habitat covered by the survey. In the 2013 assessment, the triennial survey was split into two-time series, separated by the 366 m depth contour, in order to preserve a time series of maximum length while eliminating the influence of the increased depth range. The first time series, “AFSC Triennial Shelf Survey 1,” consists of 9 data points spanning the range 1980–2004 and covering the depths 100–366 m. The second, “AFSC Triennial Shelf Survey 2,” consists of 4 data points spanning 1995–2004 and covering depths 366–500 m. This second time series is recognized as providing little information about stock status due to the limited number of points and depth range, but there was no compelling reason to exclude it from the assessment. However, in contrast to the 2013 assessment, this assessment will treat the triennial survey as a single time series for constructing the geostatistical model-based indices, and will use a different set of latitudinal and depth-based strata for survey length compositions.

### **2.2.2 AFSC and NWFSC Slope Surveys**

Starting in the late 1990s, two slope surveys were conducted on the west coast. The AFSC Slope Survey (AFSC Slope Survey) was conducted during the years 1997 and 1999–2001 using the research vessel Miller Freeman. The NWFSC Slope Survey (NWFSC Slope Survey) was conducted from 1998–2002, and was conducted cooperatively using commercial fishing vessels. The AFSC Slope Survey was a source of valuable information on the depth distribution and overlap of shortspine and longspine thornyheads in the 1980s, but these early years had a very limited latitudinal range and will not be included. This survey also had a different net and larger roller gear than the NWFSC Slope Survey.

### **2.2.3 NWFSC West Coast Groundfish Bottom Trawl Survey**

In 2003, the design of the NWFSC Slope Survey was modified, and the survey was expanded to cover the shelf and slope between 50 m and 1280 m. This combination shelf-slope survey, “NWFSC Combo Survey,” more recently known as the WCGBTS, has been conducted every year from 2003 to present with consistent design (note that the survey was not conducted in 2020 due to ongoing concerns about COVID-19). Data for the years 2003–2021 were available for this assessment. The WCGBTS represents the largest number of survey observations, the largest depth range, and the most consistent groundfish sampling program in the history of west coast fisheries. Continuing this time series in a consistent manner is vital for improving estimates of current stock status and detecting any future changes in size distribution or abundance of west coast groundfish.

#### **2.2.4 Survey Stratification**

Data from these four (nominally five for design-based indices) fishery-independent surveys were considered for use in this assessment (Figure 7) to estimate abundance. Two distinct survey abundance estimation methods were considered: design-based and geostatistical model-based indices. The 2013 assessment utilized delta-GLMMs following the methods of Thorson and Ward (2013), but these methods are no longer considered best practice within the field and were not considered in this assessment.

The five surveys were stratified based on depth and latitude, similar to how they were in 2013 (Table 14). The Triennial Survey was divided into two distinct survey time series, split on the year 1995. The early-Triennial time series (1981-1992) was further stratified into four strata: north and south of 42° N, and shallower and deeper than 200m. The late-Triennial time series (1995-2004) was also further stratified into four strata: north and south of 40° N, and shallower and deeper than 200m. The AFSC Slope Survey was split into two coast-wide strata: shallow and deeper than 550m. The NWFSC Slope Survey was divided into 6 strata, with breaks dividing southern, central, and northern strata at 40.5° N and 43° N, each of which was further divided with a break at 550 m. The WCGBTS was divided into 7 strata, with two southern strata below 34.5° N, one covering 183–550 m and the other covering 550–1280 m. Two central strata, between 34.5° N and 40.5° N, had the same depth ranges. The latitudinal divide around 34.5° N is associated with changes in sampling intensity. North of 40.5° N, three strata were used, covering the ranges 100–183 m, 183–550 m, and the other covering 550–1280 m. The depth breaks at 183 m and 550 m are also associated with changes in the sampling intensity of the survey and are recommended to be used. South of 40.5° N, there are very few shortspine thornyhead shallower than 183 m, so no shallow stratum was used in these latitudes. The 2013 stratification was reused for the design-based indices as there was not sufficient evidence to support modifying the existing strata.

#### **2.2.5 Design-based Indices of Abundance**

Design-based indices of abundance were derived for all surveys. Note that for these indices of abundance, the Triennial Survey was split into two independent time series, separated by the year 1995. The construction of design-based indices mirrors a weighted average approach. For each survey year, an average CPUE is calculated across all tows within a stratum and expanded by area to determine the total estimated biomass. These values are then summed across all strata within the survey to create a time series of design-based indices of abundance. Design-based indices were computed using the official nwfscSurvey R package.

#### **2.2.6 Geostatistical Model-based Indices of Abundance**

Model-based indices of abundance for all surveys were derived using geostatistical models Thorson et al. (2015) developed using the R package sdmTMB Anderson et al. (2022). This approach utilizes geostatistical GLMMs with spatially and spatiotemporally correlated random effects, which can account for variables that cause correlations in the data across space and time. For this reason, the Triennial Survey survey can be, and was, treated as a single time series rather than split into two timeseries based on the introduction of additional

sampling at greater depths. For the Triennial Survey and WCGBTS surveys, geostatistical models included spatial and spatiotemporal random effects and depth and depth squared as a scaled covariate. Geostatistical models for the NWFSC Slope Survey and AFSC Slope Survey were not run with depth.

Abundance indices were obtained for models using both gamma and log-normal error structures. There is limited agreement on how best to go about model selection for these types of geostatistical models, and both error structures were tested as sensitivity analyses alongside the simple design-based indices described above. The abundance indices derived from the gamma model were most similar to the design-based indices for the Triennial and WCGBT surveys and were thus used for the base model (indices derived from the log-normal model displayed a similar trend to the gamma model-based indices, and the design-based indices, but were consistently larger in scale).

### **2.2.7 Length Composition Data**

Length-composition data were available for each year of each survey. In each haul, there is a set number of random samples regardless of the amount of catch, decoupling the sample and catch size. Therefore, the length compositions were calculated using an expansion factor to account for differences in the amount of catch that samples represent. An expansion factor (calculated as weight of caught fish divided by weight of fish sampled) is calculated for each haul, multiplied by the number of fish in each size bin, and then summed across hauls. This algorithm is repeated for each spatial stratum. Length composition data were compiled into 34 length bins, ranging from 6 to 72 cm. Year-specific length frequency distributions generated for each survey are shown in Figure 8).

### **2.2.8 Frequency of Occurrence and Survey Information**

The frequency of occurrence of shortspine and longspine thornyheads in trawl surveys remains extremely high. 91% of the tows in the WCGBTS below 500 m have at least one shortspine thornyhead in the catch (and 96% for longspine thornyhead), similar to the 2013 assessment. The number of survey hauls and shortspine thornyheads sampled available for this assessment is described in Table 15.

## **2.3 Biological Data**

### **2.3.1 Natural Mortality**

**THIS WILL BE UPDATED**

Butler (1995) estimated the lifespan of shortspine thornyhead to exceed 100 years and suggested that  $M$  was likely less than 0.05.  $M$  may decrease with age as shortspine migrate ontogenetically down the slope to the oxygen minimum zone, which is largely devoid of predators for fish of their body size. The 2005 assessment fixed the natural mortality parameter at 0.05, while the 2013 assessment used a prior on natural mortality developed based on a maximum age of 100 years. The prior had a mean of 0.0505 and a standard

deviation on a log scale of 0.5361 (Hamel, pers. comm.). For the base case, natural mortality was fixed at the mean of this prior distribution. This assessment uses an updated prior on M by Owen S. Hamel (2022), where the median of M is:

$$\frac{5.40}{Age_{max}} \quad (1)$$

This assessment assumed the same maximum age of 100 as in the previous assessments, with an updated prior of 0.054 (median value) and log-space standard deviation = 0.31 (Owen S. Hamel 2022). The 2023 assessment will explore estimating M and fixing M at the prior mean.

### 2.3.2 Maturation and Fecundity

**2.3.2.1 Maturity** Pearson and Gunderson (2003) estimated length at 50% maturity to be 18.2 cm on the West coast, with most females maturing between 17 and 19 cm. This was represented in the 2005 and 2013 assessments by the logistic function,

$$M(L) = (1 + e^{-2.3(L-18.2)}) - 1 \quad (2)$$

where L is the length in cm.

The 2013 assessment considered new (at the time) maturity information from ovaries collected for maturity analysis on the 2011 and 2012 WCGBTS. Histological analysis of those samples (M. Head, pers. comm.) indicated puzzling patterns of spawning by female size and by latitude, with a higher fraction of fish spawning in the north than in the south and a higher fraction of spawning fish in the 20-30cm than in the 30-40cm range. However, due to the complexity of these observed patterns and the known ontogenetic migrations of shortspine, samples collected in 2011 and 2013 were not considered adequate for estimation of a new representative maturity curve for the entire shortspine thornyhead population in 2013. Nonetheless, such a maturity curve was considered in a sensitivity analysis. On the basis of the sensitivity analysis, the 2013 assessment suggested that the slow but steady rate of growth for TEST shortspine thornyheads, with growth still occurring at age 100, reduces the importance of assumptions about maturity because older individuals have significantly higher spawning output due to their much larger size, regardless of the fraction spawning.

New maturity analyses of samples collected on the WCGBTS in 2011, 2013, 2014, 2016 and 2018 were available for the 2023 assessment (M. Head, pers. comm.). The larger number (N=397) and better spatial coverage of these samples allowed the use of statistical modeling to better understand the spatial variation in the proportion of female spawning.

In the 2013 assessment, the exploration of maturity analyses from the WCGBTS samples highlighted maturity gradients along latitude and depth. To assess a potential relationship between fish location and the shape of the maturity curve, a general linear model (GLM)

was designed for estimating maturity curve parameters while integrating latitude and depth as covariates. This GLM consists of a logistic regression in which the functional maturity of samples, modeled with a Bernoulli distribution, is expressed as a linear combination of fish length, latitude, squared latitude, depth and squared depth of collection. Once fitted, the GLM was used to predict the response of the probability of being mature along the range of individual shortspine length considered in the model. For the 2023 assessment, this model prediction was made while setting the latitude and depth at the values of the center of gravity (using number of fish as weighing factor) of the population of shortspine thornyhead sampled during the WCGBTS to develop a single curve for the coastwide population assessment. Thus, this response of functional maturity to length was considered the mean maturity curve of the west coast shortspine thornyhead population. The parameters of the maturity curve L<sub>50</sub> and k were arithmetically derived from this response to fish length. The new maturity curve is expressed as follows:

$$M(L) = (1 + e^{-2.3(L-31.42)})^{-1} \quad (3)$$

Figure 11 shows the fit of the maturity curve of the model per class of depth and latitude.

A sensitivity analysis will assess the impact of this change in the maturity curve on the model estimates by considering the newly estimated parameters, the Pearson and Gunderson's relationship from the 2013, and one intermediate option (Figure 12).

**2.3.2.2 Fecundity** The previous assessments assumed spawning biomass was equivalent to spawning output. The 2023 assessment will explore using fecundity-at-length parameters reported in Cooper et al. (2005), where fecundity was modeled as a power function of length. Cooper et al. (2005) estimated the fecundity of 54 females collected from the West Coast and Alaska. The study found no difference in the length-fecundity relationship by region and pooled the samples. The fecundity-at-length parameters in that study were:

$$F = 0.0544L_{3.978} \quad (4)$$

where F is fecundity in the number of eggs per female and L is length in cm. Fecundity information from Cooper et al. (2005) suggests that fecundity increases at a faster rate with length than body weight with length. This means larger females are likely to have greater relative fecundity compared to small females (i.e., large females produce more eggs per kg of body weight). This assessment will explore modeling a fecundity-at-length relationship using the fecundity parameters from Cooper et al. (2005) and scaling the fecundity intercept by one million in SS3 to report fecundity in billions of eggs.

Uncertainty remains in the spawning strategy of shortspine thornyhead. Cooper et al. (2005) and Pearson and Gunderson (2003) found no evidence for batch spawning in this species (i.e., a determinate, total spawning strategy). However, updated histology information suggests a possibility of batch spawning in this species (Melissa Head, NWFSC, pers. comm.). Batch spawning could influence the fecundity-at-length relationship if not properly accounted for

and should be a focus of future research. The histology analysis also found evidence of parasites in the ovaries and atresia (degrading eggs), which could influence fecundity (Melissa Head, NWFSC, pers. comm.).

### 2.3.3 Length-Weight Relationship

Fisheries-independent length and weight specimen data are available from the AFSC Slope Survey (1997, 1999-2001; N=7,623) and the WGBTS (2003-2021, excluding 2020; N=20,142). The WGBTS data were used to estimate the length-weight relationship because it has the largest sample size and covers the greatest spatiotemporal resolution. The allometric function models weight (W) as an exponential function of length (L), where:

$$W = \alpha L^\beta \quad (5)$$

This function can be linearized by taking the natural logarithm of both sides. The predicted weight-at-length values were bias-corrected using a multiplier of 2 / 2. Length-weight was estimated for both sexes in R using the lm() function (R Core Team 2021).

The resulting allometric parameters for 2023 (females:  $\alpha = 6.49e-6$ ,  $\beta = 3.18$ ; males:  $\alpha = 6.71e-6$ ,  $\beta=3.17$ ) were similar to the 2013 assessment values, which estimated a single length-weight relationship for males and females combined using WGBTS data through 2012 (sexes combined:  $\alpha = 4.77e-6$ ,  $\beta=3.26$ ). The  $\beta$  value was higher in the 2013 assessment, indicating a slightly higher weight-at-length for longer fish. We found no temporal trend in the available data and were unable to account for these small differences in results. The available data suggest that length-weight is highly conserved in shortspine thornyhead; therefore, no sensitivity analysis was conducted for this set of parameters in the 2023 assessment.

### 2.3.4 Growth (Length-at-Age)

No validated ageing methodology currently exists for shortspine thornyhead; therefore, this species is not aged by the NWFSC or Southwest Fisheries Science Center (SWFSC) and length-at-age data are limited for this stock assessment. Two research age datasets exist for shortspine thornyhead in the West Coast region: (1) Kline (1996) includes 319 unsexed fish collected from Monterey Bay in central California in 1991, and (2) Butler (1995) includes 1,023 sexed fish collected in the waters off northern California and Oregon in 1978-1988 and 1990. The Kline specimens were aged by one age reader, and lengths were reported as total lengths, whereas the Butler specimens were aged independently by two separate age readers, and lengths are reported in fork length. The Butler data age data presented in this assessment are the mean ages between the two age readers.

The length-at-age curve developed in the 2005 stock assessment and used again in 2013 was based on a Schnute parameterization of the Von-Bertalanffy growth function fit to the Kline data. The resulting parameter estimates for this growth function were as follows: growth rate k was 0.018 for both males and females length at age-2 was 7 cm for both males and

females, and length at age-100 was 67.5 cm for males and 75 cm for females based on the assumption that the asymptotic length for males should be 90% of the asymptotic length for females (Hamel 2005). The data and associated analysis from 2005 were lost; however, the original Kline and Butler datasets were obtained for use in this assessment (Donna Kline, pers. comm., March 2023). Using these newly obtained data, we could not reproduce the parameters used in the 2005 assessment.

Because the Butler data were sex-specific, had a higher sample size, were aged by two readers instead of one, and were collected from a larger geographic area and over more years compared to the Kline data, we determined that Butler was the preferred dataset to estimate the length-at-age relationship for the 2023 stock assessment. We fit sex-specific Schnute growth functions to the Butler data:

$$\hat{L}_a = L_{a_1} + \frac{(L_{a_2} - L_{a_1})(1 - \exp(-k(a_2 - a_1)))}{(1 - \exp(-k(a_2 - a_1)))} \quad (6)$$

where:  $L_{a_1}$  and  $L_{a_2}$  are the lengths at reference ages  $a_1$  and  $a_2$  where  $a_1 = 1$ ;  $a_2 = 100$  and  $k$  is the growth rate. Growth curve estimation was conducted in R using the `optim()` function (R Core Team 2021). Errors were assumed to be lognormally distributed and predicted length-at-age was bias-corrected using a multiplier of  $\frac{\sigma_2}{2}$ .

Shortspine Thornyhead are slow-growing fish that appear to continue to grow throughout their lifespan (i.e., the growth curve does not asymptote). The new growth curves estimated using the Butler dataset exhibited similar trends to those assumed in the 2005 and 2013 assessments (Figure 9). The male curves were almost identical, with the 2005/2013 curve exhibiting slightly lower length-at-age at young ages and slightly higher length-at-age at older ages. The 2005/2013 female curve was defined by a higher growth rate, leading to the higher length-at-age in the intermediate age range.

Two alternative sensitivity analyses were developed for the 2023 assessment. During the exploratory data analysis phase, we found that specimens collected in the Kline study exhibited higher size-at-age when compared to the Butler specimens (Figure 10). It is unknown if these differences should be attributed to spatial differences in growth between central California and northern California/Oregon, bias among age readers, or discrepancies between the total and fork length measurements (Donna Kline, pers. comm., March 2023). In order to account for this alternative growth pattern, we increased the lengths at ages 2 and 100 by 25% in the upper sensitivity analysis (Figure 10). The lower sensitivity analysis was defined by decreasing the lengths at ages 2 and 100 by 10% from the base model.

### **2.3.5 Ageing Precision and Bias**

*REPEAT WITH CLARITY*

## **2.4 Environmental and Ecosystem Data**

No ecological or environmental information was used in this assessment.

## **2.5 Changes in data from the 2013 assessment**

Most of the data used in the previous assessment has been newly extracted and processed, including length compositions from each fishing fleet and survey, indices of abundance derived from new geostatistical models of survey data, discard rates from both the 1980s Pikitch study and the current WCGOP, and the time series of catch from 1900-2023.

New data or uses of data for this assessment include the geostatistical model-based indices of abundance for the four fisheries independent surveys, the histological maturity samples from the WCGBTS survey, and the historical state catch reconstructions. Previous assessments have treated the AFSC Triennial Shelf Survey as two separate indices of abundance separated by the 366m depth contour, but the transition to using geostatistical model-based indices have rendered this separation unnecessary by implicitly accounting for changes in depth sampling within the model. State-level historical reconstructions also replace previous analyses that imputed historical shortspine thornyhead catch as a fixed proportion of sablefish catch.

# **3 Assessment Model**

## **3.1 Summary of Previous Assessments and Reviews**

### **3.1.1 History of Modeling Approaches**

Shortspine Thornyhead was first assessed in 1990 by Jacobson (1990) and Jacobson (1991), and subsequently by Ianelli et al. (1994), Rogers et al. (1998), and Piner and Methot (2001). What would now be called a data-moderate assessment was conducted in 2005 Hamel (2005) using Stock Synthesis (SS2). More recently, shortspine thornyhead were assessed by Taylor and Stephens (2013) using SS3. The 2013 model retained many of the assumptions made by Hamel (2005) including a four fisheries fleet structure, sex-specific growth, and no fecundity relationship. A catch-only projection was conducted in 2019 (Taylor 2019).

### **3.1.2 Most Recent STAR Panel and SSC Recommendations**

1. More investigation into maturity of shortspine
2. Information on possible migration of shortspine thornyheads would be valuable for understanding stock dynamics.
3. A greater understanding of catchability of thornyheads would help define the scale of the populations..
4. Age data and additional research on ageing methods for thornyheads would be valuable.
5. A greater understanding of the connection between thornyheads and bottom type could be used to refine the indices of abundance.
6. A comprehensive catch reconstruction for shortspine and longspine thornyheads should be completed to estimate landings for each species prior to 1981 in each of the three

states.

7. Exploration of simpler assessment methods for thornyheads and evaluation of whether such methods would provide a more robust management strategy than the current approach.
8. More tows or visual surveys south of 34.5 deg. N. lat. including the large Cowcod Conservation Area. Because the southern Conception Area is a large potential habitat for thornyheads, more sampling effort would help refine the estimations of their abundance in this area.

Recommendations from the last assessment were taken into consideration in the development of the current assessment model. New histological data from the NWFSC Combo Survey in recent years enabled calculation of new maturity curves for the model. In addition, more comprehensive catch reconstructions from Washington, Oregon, and California were used for catches prior to 1980 in the model. Lastly, model structure for the current assessment was simplified by reducing the number of fishery and survey fleets.

### **3.1.3 Response to Groundfish Subcommittee Requests**

*TBD*

## **3.2 Model Structure and Assumptions**

### **3.2.1 Model Changes from the Last Assessment**

The most notable changes from the previous assessment, conducted in 2013, include significant modifications to the fleet and survey structure, and major changes to the maturity and fecundity relationships that underlie the model’s biological assumptions.

The 2013 assessment consisted of four fisheries fleets, and used information from four (nominally five) scientific surveys, while the new assessment uses a condensed structure consisting of just three fisheries fleets and only two (nominally three) surveys (see **section 3.2.2** for more details).

This assessment assumes a new fecundity relationship, in which fecundity increases with body size, as well as a new maturity relationship, in which fish mature at much larger sizes and thus older ages, than were assumed in the 2013 assessment. Further details on the fecundity and maturity relationships can be found in **section 2.3.2**. A sensitivity analysis was performed to determine the effect of different maturity assumptions on the final model output.

### **3.2.2 Modeling Platform and Structure**

This new assessment, including all exploratory models, profiles, and related analyses, was performed using Stock Synthesis Version 3.30.21 [Methot and Wetzel (2013); Methot et al. (2020)]. The majority of analyses were performed using multiple recent versions of R (R

Core Team 2021), and relied heavily on the ‘r4ss’ R package (Taylor et al. 2021) among others. The assessment model was developed and tested across multiple operating systems, including recent versions of Windows and macOS.

### 3.2.3 Model Fleet Structure

Commercial fisheries landings were divided into three distinct fisheries fleets: a northern trawl fleet (hereinafter referred to as North Trawl) operating off the coasts of Oregon and Washington, a southern trawl fleet (hereinafter referred to as South Trawl) operating off the coast of California, and a coastwide non-trawl fleet (hereinafter referred to as Non-trawl).

Data from two fisheries-independent scientific surveys were used in this model: the AFSC Slope Survey from 1980-2003, and the more recent WCGBTS from 2004-2022. The Triennial Survey was further divided into an early (pre-1995) and late period (post-1995) survey to account for the change in depth-sampling that occurred during the 1995 season. These two periods for the Triennial Survey were treated as separate surveys in the model. The contribution of each new data component to the changes in the assessment outcome was analyzed by data piecewise.

### 3.2.4 Model Parameters

There are 180 estimated parameters in this assessment. The log of unfished recruitment,  $\log(R_0)$ , controls the overall scale of the population, while annual deviations in recruitment about the assumed stock-recruit relationship (135 parameters) allow for additional uncertainty in the population trajectory. Selectivity and retention parameters (43 parameters) for three fisheries fleets and three scientific surveys allow for estimation of annual length compositions and discards rates. Two catchability parameters are analytically computed from the data, and one additional parameter, representing additional variability in the early Triennial Survey, is directly estimated by the model.

**3.2.4.1 Growth, Maturity, Fecundity, Mortality, and Recruitment** Growth, maturity, and fecundity parameters were fixed at values determined by external analyses (see **section 2.3** for more information). Due to a lack of aging data, growth could not be modeled internally by the assessment, though, like in the 2005 and 2013 models, there is no systematic misfit to the data suggesting that the externally derived growth curves were misspecified. Sensitivity analyses were performed to determine the overall effect of different assumptions regarding growth and maturity.

For this assessment, natural mortality was fixed at a value of 0.04. A likelihood profile exploring alternative natural mortality values was also conducted (Figure 35). This marks a change from the 2013 assessment, where natural mortality was fixed at 0.0505 and a standard deviation on a log scale of 0.5361 (Taylor and Stephens 2013).

As in the previous shortspine thornyhead assessment, a Beverton-Holt stock recruitment relationship was assumed. Unlike the 2013 assessment, where steepness was fixed at a value

of 0.60, this assessment fixed steepness at 0.72, as recommended by Thorson et al. (2019). A likelihood profile exploring alternative steepness parameters was conducted and the model results were found to be largely insensitive to the assumed value (Figure XX 38).

The overall scale of the population is estimated through the log of the initial recruitment parameter ( $R_0$ ). Recruitment deviations were additionally estimated for the years 1901-2022. Recruitment bias adjustments were phased in beginning in 1950, and were adjusted by a factor of **0.75 in the years 1982-2022** (Taylor and Methot 2013). The  $R$  parameter which controls the variability in recruitment deviations was fixed at 0.5 as in the previous assessment. Past assessments performed likelihood profiles over  $R$ , finding the model results to be relatively insensitive to its value, and thus further profiles over the parameter were not conducted here.

**3.2.4.2 Selectivity and Retention** Gear selectivity parameters used in this assessment were specified as a function of size with the additional assumption that age 0 fish were not selected, regardless of their size. Separate size-based selectivity curves were fit to each fishery fleet and survey.

The selectivity curves for all fisheries and surveys were allowed to be dome-shaped and modeled with double-normal selectivity. The double-normal selectivity curve parameterization has six parameters, including: (1) peak, the length at which individuals are first fully selected, (2) width of the selectivity plateau, (3) width of the ascending part of the curve, (4) width of the descending part of the curve, (5) starting selectivity, and (6) final selectivity. Parameters 5 and 6 were not estimated and fixed at 0.0. The 2013 model allowed for all selectivity parameters to be estimated, regardless of whether one or more were estimated to be on the parameter bound. This model fixed parameter 2 (the plateau width) to the value of -15 for the North Trawl and Non-trawl fleets to alleviate them hitting the lower parameter bound. Though exploratory models run with the plateau width on its lower bound still converged, fixing the parameter had negligible impact on the fits to the length composition data for those fleets. Sex-specific selectivity curves were fit to the WCGBTS and Triennial Survey length composition data.

Retention curves are defined as a logistic function of size. These are controlled by four parameters: (1) inflection, (2) slope, (3) asymptotic retention, and (4) male offset to inflection. Male offset to retention was fixed at 0 (i.e. no male offset was applied). The parameters for inflection and asymptotic retention were modeled as time-varying quantities via use of time blocks, as was done in the 2013 assessment. Therefore, both North Trawl and South Trawl fleets were broken into three periods: (1) 1901–2006, (2) 2007–2010, (3) 2011–2022. The first break was based on observation of a strong reduction in discard rates for both North and South Trawl in this year, while the later break was associated with the beginning of the IFQ program.

Alternative retention blocking schemes were investigated as part of a sensitivity analysis. Notably, a sequence of four-year blocks beginning in 2007 was investigated in order to better fit noticeably lower discards rates in the mid 2010s. Additionally, a short, 2-year, timeblock

for the years **2021** and **2022** was also attempted, as discard rates were noticeably higher in those years than in previous. [...]

**3.2.4.3 Catchability** Catchability coefficients ( $q$ ) were calculated for each of the two survey abundance time series. Unlike in the 2013 model, which estimated catchability for each survey, this model computes catchability analytically for each survey using the Stock Synthesis “floatQ” option (Methot et al. 2020).

This model depends on the assumptions that thornyheads are long-lived, slow-growing, and relatively sedentary groundfish. They are assumed to represent a single stock within the area considered for this assessment. If the assumptions about growth, natural mortality, or stock structure turn out to be far from the true life history and ecology of shortspine thornyheads, this assessment will be highly inaccurate.

### 3.3 Base Model Results

#### 3.3.1 Parameter Estimates

A complete set of parameter estimates are available in Table 17.

**3.3.1.1 Recruitment** The model estimated 135 annual recruitment deviations (1901–2034) as well as the log of unfished recruitment  $\text{LN}(R_0)$ . Unfished recruitment was estimated to be ~11,000,000 annual age-0 recruits ( $\text{LN}(R_0) = 9.34$ ) while annual log deviations were generally estimated between -0.5 and 0.5 (Figure 13 ; Table 18). Deviations in 2003 and 2007 were estimated to be substantially larger than other years. As in the 2013 assessment, uncertainty in the scale of annual deviations was substantially larger than the variation between the deviations. Recruitment bias adjustments were performed following the advice of Methot et al. (2011).

**3.3.1.2 Selectivity and Retention** Selectivity curves for all three fisheries fleets and the three scientific surveys were estimated as dome-shaped (Figure 14).

The early- and late-period Triennial Survey possessed the highest degree of dome-shape, with peak selectivity occurring at relatively small length (~26 cm, and ~22 cm respectively), before quickly declining. This shape is consistent with the design of the survey which focused its sampling on the relatively shallow shelf, where younger, smaller, shortspine thornyhead live before migrating to deeper waters as they age and mature. There was little difference in the estimated selectivity curves between male and female fish. Meanwhile, the WCGBTS was estimated to have a wide plateau (beginning at ~30cm) over which the species is fully selected for, including the lengths over which the species spends the bulk of its lifespan. This indicates that the WCGBTS is sampling a large proportion of the stock, and that annual length composition data from the survey is likely a good representation of the true distribution of lengths in the population.

The North Trawl fleet was estimated to have a dome-shaped curve with a small plateau around 28 cm in length, and a long tail that spanned nearly the entire range of observed lengths.

The South Trawl fleet was estimated to have a very large selectivity plateau (beginning at ~30cm and ranging from 30-55cm, with very steep ascending and descending limbs. Finally the Non-Trawl fleet was estimated to have a relatively small plateau beginning at a much higher length than any other fleet or survey (~45 cm). This can be explained by the fact that hook and line gear, the dominant gear type in the Non-Trawl fleet, selectively catches larger shortspine thornyhead.

Retention curves for all three fisheries were asymptotic in shape, with the two trawl fisheries asymptoting at a retention value of 1.0 and the non-trawl fishery asymptoting a value just below 1.0 (**Figure XX**), indicating that the Non-Trawl fishery still discards large fish in limited cases. Retention was modeled in distinct time blocks to allow for better fits to the discard rate data. The time-blocked fits to the North Trawl fleet show the fishery to have begun retaining smaller fish in more recent years than they have historically. A similar pattern is observed for the South Trawl fleet, but to a much smaller extent (**Figure XX**).

### 3.3.2 Fits to the Data

**3.3.2.1 Abundance Indices** The base model reasonably fit the available index data with the exception of the most recent two years of the WCGBTS. The fit to the Triennial Survey was relatively flat across the entire timeseries (1980-2004; Figure 15). An extra parameter was used to estimate additional variance beyond that estimated by the geostatistical model for this survey. The model fit to the WCGBTS indices appropriately captured the lack of trend in the early and middle portions of the timeseries, but struggled to accurately capture the recent increase in abundance displayed by the indices (Figure 16). The model fit for this survey fails to fall within the estimated confidence interval for the 2021 and 2022 indices. This could be, in part, due to the lack of index data from 2020 (surveys were not conducted due to the ongoing COVID-19 pandemic), which may have helped the model more accurately capture the increase.

**3.3.2.2 Fishery Discard Rates** The model reasonably fit the discard rates for all three fisheries fleets. The three timeblocks, carried over and extended where necessary from the 2013 model, allowed for the declining trend in discards for the North Trawl fleet to be adequately captured, however, there is a period from 2015-2018 where discard rates for the North Trawl fleet are exceptionally low, that the model fails to fully capture (Figure 17). Discard rates in the Southern Trawl fleet are well fit 2006-2014, before and after which the model systematically underestimates the observed discards rates (Figure 18). The discard rates for the coastwide Non-Trawl fleet are exceptionally well fit by the model, and there is no evidence that time blocking is necessary for this fleet (Figure 19).

**3.3.2.3 Fishery Length Compositions** The base model fit the fishery and discard length compositions reasonably well in aggregate (Figure 20), though there was significant annual variability in the quality fit, often due to difference in effective sample sizes. The Southern Trawl and Non-Trawl fleets were exceptionally well fit by the model, while the model fit to the length compositions from the Northern Trawl fleet underestimated the scale of the peak of the distribution. This type of misfit was similarly observed in the model

fits on an annual basis, with all years 2018-2022 displaying a similar underestimation of either the location or scale of the peak of the distribution (Figure 21). The exact causes of this under-estimation remain unknown at this time, but could be due to subtle changes in selectivity or availability. The effect of including time-varying selectivity was assessed in a sensitivity analysis (see **section 3.4.2.4** for details).

Trawl discards length compositions were well fit by the model in both the north and south regions, while the model struggled to adequately fit discard compositions from the Non-Trawl fleet (Figure 20). The Non-Trawl discards were of a larger size and were generally more dispersed than the discard compositions in the two trawl fleets, a feature the model did pick up on, but the model fit a wide plateau rather than narrow peak to these composition data. This is likely due to the wide variability in annual length compositions seen in this fleet, as well as the wide spatial coverage.

**3.3.2.4 Survey Length Compositions** Like the fishery derived length compositions, survey-derived length compositions were reasonably well fit in aggregate by the base model (Figure 20), though there was considerable annual variability in the quality of the model fit (Figure 22). The early-period Triennial Survey length composition data for both sexes were exceptionally well fit by the model. Length compositions from the late-period Triennial survey were slightly less well fit, with the model under-estimating the location of the peak for both sexes. For the WCGBTS length compositions, the male, female, and unsexed location of the compositional peaks were well estimated, though the overall scales were slightly underestimated. Pearson residuals did not demonstrate any obvious trends that would indicate systematic misfits to the data (Figure 23; Figure 24; Figure 25); as such, time varying selectivity was not included in the base model.

**3.3.2.5 Mean Body Weight** Mean body weight of discarded fish was well fit in the two trawl fleets, and no major trends were observed in either the data or the model estimates (Figure 26; Figure 27; Figure 28). Mean discard weight in the Non-Trawl fleet was observed to have increased in the last ten years, but this trend was not captured by the model. The model, instead, fit a declining trend in discard weight to the Non-Trawl fleet data (Figure 28). The reason for this disparity between the observed data and the model fit is unclear.

### 3.3.3 Population Trajectory

Unfished equilibrium spawning output ( $B_0$ ) is estimated to be 20,262 trillion eggs (16,291–24,232 trillion eggs). The  $B_0$  estimate is not directly comparable to estimates made in previous assessments, which assumed no fecundity relationship, and thus calculated  $B_0$  in terms of biomass rather than egg production. Spawning biomass is estimated to have remained stable until the mid-1960s and then declined in the 1970s to about 80% in the mid-1980s, followed by a slower decline under the lower catch levels in the 2000s (Table 10; Figure 29). While the spawning output of the stock has declined near linearly since 1975, total biomass has stabilized in recent decades around ~85,000 mt. The estimated spawning output in 2023 is 8.273 trillion eggs (5.346–11.201 trillion eggs), which represents a stock

status (depletion level) of 41% (33%-49%; Table 7, Figure 30). The depletion estimated for 2013 is 43.6%, which is significantly lower than the 74.2% estimated for 2013 in the previous assessment.

Twelve-year projections predict that the population is unlikely to experience a large increase in spawning output or spawning biomass in the near future, if the full ACL is taken each year (Figure 31; Figure 32).

## 3.4 Model Diagnostics

### 3.4.1 Convergence

The maximum likelihood parameter estimates found by AD Model Builder (ADMB) indicated a well-converged model. The base model had a small maximum gradient component (0.0002317) and a positive definite Hessian matrix, both of which are associated with converged models.

Runs with 100 alternative sets of starting parameter values found no models with a better likelihood (**Table JITTER RESULTS**). Of the 100 jittered model runs, 74 reconverged to the best estimates associated with the base model, while 76 reconverged to within 0.25% of the best estimate. This indicates that the data is relatively informative about the population dynamics of the stock, as the model fits to the data were virtually identical.

### 3.4.2 Sensitivity Analyses

**3.4.2.1 Sensitivity to growth parameters** Growth parameters are uncertain for shortspine thornyhead due to difficulties in determining age from otoliths and subsequent lack of length-at-age information for this species. The 2023 assessment used growth parameters based on length-at-age information from Butler (1995) of shortspine thornyhead collected in Oregon and Northern California from 1978-87, 1988, and 1990. The data is sex-specific and with two age readers (N=1,023). A sensitivity analysis was conducted to higher growth observed by Kline (1996) of shortspine thornyhead collected from central California in 1991. Fish were unsexed, with a smaller sample size (N=319) and only a single age reader. A sensitivity of length-at-age 25% higher and 10% lower was conducted to encompass the uncertainty in growth. Different assumptions about growth did not have much influence on relative spawning depletion. Depletion levels in the final year were slightly greater in a higher growth scenario and slightly lower in a lower growth scenario (Figure 48). Spawning output was more sensitive to assumptions about growth, with much higher spawning output when assuming higher growth (Figure 47). The high growth sensitivity had a slightly better overall fit to the data, including to the survey indices and length compositions (Table 16). Different assumptions about growth influence recruitment patterns and the timing of strong year classes. However, the influence of different growth assumptions is also sensitive to changes in other parameters, such as natural mortality.

**3.4.2.2 Sensitivity to maturity** The 2023 assessment uses updated maturity-at-length information from the WCGBTS and port-sampling (Melissa Head, pers. comms.),

which show a larger length at 50% maturity (31.4 cm) and shallower slope (-0.177) of a logistic regression compared to a smaller length at 50% maturity (18.1 cm) and steeper slope (-2.304) assumed in the 2013 assessment based on Pearson and Gunderson (2003). A sensitivity was conducted to maturity information from the two datasets and to an intermediate maturity-at-length logistic curve (mix\_curve, L50% = 24.8 cm and slope = -0.350). As in the 2013 assessment, the base model was not sensitive to different maturity assumptions (Figure 50; Figure 51). Differences in fits to the data were negligible. The slightly higher stock status in the final year with the Pearson and Gunderson (2003) maturity assumptions is likely due to females from strong cohorts in the 2000s being (or becoming) mature that are not yet mature with 2023 larger size-at-maturity assumptions. As stated in the 2013 assessment, the slow but steady rate of growth for shortspine thornyheads, with growth still occurring at age 100, reduces the importance of assumptions about maturity because older individuals will have significantly higher spawning output due to their much larger size, regardless of the fraction spawning.

**3.4.2.3 Sensitivity to Landings** The 2023 assessment uses state-level historical landings reconstructions for thornyhead. The two most recent assessments in 2005 and 2013 instead imputed historical catch by assuming shortspine thornyhead landings for the southern trawl fleet prior to 1962 equaled 30% of the sablefish landings. Further, new landings reconstructions have become available for Oregon and Washington, and data in PacFIN has undergone revisions since the 2013 assessment, leading to differences in landings between the 2023 landings data and the 2013 landings data. Two sensitivities were conducted for landings data. One sensitivity that was also conducted in the 2013 assessment replaces the historical landings reconstructions prior to 1962 with the imputed landings that were used in the 2005 and 2013 assessments (23.land.hist\_impute). A second sensitivity replaces all landings information prior to 2013 with the values that were used in the 2013 assessment (23.land.2013). There was little appreciable difference to base model fits across the two sensitivities (Figure 52; Figure 53).

**3.4.2.4 Sensitivity to retention and selectivity** The 2023 assessment substantially extends the period over both length data of retained and discarded catch, mean individual weight in discards and discard rates estimates are available. This data may reflect the dynamics of the thornyhead of the population but also structural, technical or behavioral changes in the fishing fleets. Due to the uncertainty around the impact of such changes on the retention and selectivity of the fleets, we assessed the sensitivity of the model to alternative combinations of time blocks for these parameters. The sensitivity to time blocks in retention was first performed and the most pertinent model (good fit to the data and parsimony) and the base model were then used to conduct the sensitivity to time blocks in selectivity. Time blocks explored were based on a combination of breaks that are summarized in **Table XX**.

Setting up additional time blocks for the retention parameters improved the fit of the model to the length distributions and the discard rates, hence reducing corresponding likelihood of the model. However, the impact on the other components of the model likelihood were very limited. None of the time blocks tested substantially reduced other components of the model likelihood or led to changes in the recruitment trends or the depletion status of the

population (**Figure XX**).

### 3.4.3 Retrospective Analysis

Retrospective analysis indicates that removing the most recent years of data has minimal impact on the estimates of spawning output (Figure 43) and stock status (Figure 44). This is consistent with the results of the likelihood profile over  $R_0$  (Figure 33) which showed that the data are relatively informative about stock scale. While the analysis does display some minimal evidence of a retrospective pattern, all estimates of spawning output in the retrospective analysis fell within the 95% uncertainty interval around the base model spawning output time series (Figure 46).

As in the previous assessment, there is little evidence that such retrospective patterns are the result of additional years of survey abundance data (Figure 45), and thus, it is most likely that removal of informative length composition data is the source of such pattern.

### 3.4.4 Likelihood Profiles

Likelihood profiles were conducted over the log of unfished recruitment ( $R_0$ ), the steepness of the stock recruit relationship ( $h$ ), the value of natural mortality ( $M$ ), and the growth curve.

A likelihood profile over  $\log(R_0)$  was performed to assess the influence of the various data sources on the unfished scale of the population. The profile shows most of the data sources to be in agreement regarding the best estimate of  $R_0$ , with the exception of the indices of abundance, which are best fit by larger values of  $R_0$  (Figure 33). Similarly, there is little inconsistency in likelihood contribution by fleet across the range of plausible  $R_0$  values (Figure 33). This indicates that the data, together, are relatively informative about the overall scale of the population. The highest spawning output and lowest depletion levels were associated with higher  $R_0$  values (Figure 34).

Likelihood values and model results were largely insensitive to changes in steepness (Figure 39). The change in negative log likelihood over the range of  $h = 0.5\text{--}1.0$  was less than 10 units with the largest contribution coming from recruitment and abundance indices. No other likelihood component had a change of greater than 1 unit. The lowest  $B_0$  and depletion values were associated with the most productive population, with  $h = 1.0$ , but there was no qualitative difference between any of these cases (total change in depletion values across the range of  $h$  tested was 0.02 units; Figure 40). The apparent lack of influence of  $h$  on population dynamics for shortspine thornyhead is likely the result of the relatively high estimated stock status across the entire time series, which makes estimation of  $h$  difficult.

A likelihood profile over natural mortality ( $M$ ) found the model results to be quite sensitive to the assumed value of  $M$ : with all values of  $M$  0.03–0.05 resulting in likelihoods within 1 unit of the base model (Figure 35). Meanwhile there was no support in the data for values of  $M$  above 0.06. Length composition data, particularly from the two trawl fleets, was the major contributor to the changes in likelihood observed over the range of tested values for  $M$ .

(Figure 35). All plausible values of M resulted in similar levels of depletion (Figure 36), but a wide range of levels for unfished spawning output (Figure 37)

Because of the high sensitivity of the model to assumptions about growth, a likelihood profile over growth coefficients, length at minimum age ( $a_{min}$ ) and length at maximum age ( $a_{max}$ ), for each sex, was conducted. For the profile, length at amin for females ranged from 10 cm to 15 cm and length at  $a_{max}$  ranged from 60 cm to 150 cm. For males, length at  $a_{min}$  ranged from 8cm to 12 cm and length at  $a_{max}$  ranged from 55 cm to 140 cm. As with the growth sensitivity analysis described above, likelihood was minimized for higher growth scenarios (Figure 41). Spawning biomass trajectories showed similar patterns to the growth sensitivity, with higher growth scenarios producing substantially higher spawning output (Figure 42). Based on both the fishery and survey length compositions, the largest fish, under a high growth model, are not necessarily being observed in the data. This, in combination with a low natural mortality, may be driving the higher spawning output patterns under high growth scenarios.

### 3.4.5 Unresolved Problems and Major Uncertainties

Few problems remain totally unresolved, though improvements to the model fit to the WCGBTS abundance indices and to the North Trawl fleet length compositions would be desirable. In addition, being able to freely estimate the width of the selectivity plateaus for many of the fleets and surveys would also improve the model.

The model fails, at this time, to fully capture the observed increase in abundance seen in the WCGBTS index time series, significantly underestimating the abundance in 2021 and 2022 (Figure 16). Better fits to WCGBTS length compositions in those years could possibly improve fits to the indices, but improvements to the length composition fits proved difficult without introducing time-blocked selectivity. As the WCGBTS is supposed to follow highly standardized survey methodologies, there seems to be minimal justification for introducing time-varying selectivity in the model at this time. The model also fails to fully capture the peak of the length compositions for the Northern Trawl fleet, underestimating the number of mid-sized fish that the fleet takes (Figure 20). This underestimation appears to be consistent, particularly in the last 10 years (Figure 21), implying a possible recent change in selectivity. While time-varying selectivity was not investigated here, if this trend persists, future assessments may wish to apply a selectivity time block to this fleet in order to better capture the peaks of the length compositions.

Major uncertainties in the model are centered around uncertainty in the biological parameters that govern growth, maturity, and mortality. Due to a lack of reliable aging methods, growth was estimated externally to data collected in the 1990s (see **section 2.3.4** for more information). Sensitivities conducted on the length-at-age-100 demonstrated that changes to the growth function could have large effects on the estimated stock status (Figure 48). Due to inconsistent histological data for shortspine thornyhead, there is some uncertainty about the shape of the species' maturity curve, though the model appears to be largely insensitive to variations in maturity (Figure 51). Finally, likelihood profiles over natural

mortality demonstrate the model to be quite sensitive to its assumed value. There is insufficient information in the data to estimate mortality directly, constraining us to using meta-analyses or other mortality estimators, which frequently make use of aging information that is unavailable or highly uncertain for shortspine thornyhead.

## 4 Management

### 4.1 Reference Points

Reference points were calculated using the estimated catch distribution in the final year of the model (2023). In general, the population is on the boundary between “precautionary” ( $B/B_0 = 0.40$ ) and “healthy” status relative to the reference points (Figure XX). Sustainable total yield (landings plus discards) was estimated at 1,060 mt when using an SPR50% reference harvest rate and ranged from 870–1,250 mt based on estimates of uncertainty (Table XX). The spawning output equivalent to 40% of the unfished spawning output ( $B_{40\%}$ ) was 8.273 trillion eggs. The most recent catches (landings plus discards) have been lower than the estimated long-term yields calculated using an SPR50% reference point, but not as low as the lower bound of the 95% uncertainty interval. However, this is due to the fishery not fully attaining the full ACL. The OFL and ABC values over the past 6 years have been approximately 3100 mt and 2,500 mt, respectively. Both of those values are higher than the OFL and ACL values predicted in short-term forecasts, which are around 900 mt and 777 mt respectively for 2025–2026 (Table XX). This is reflected in the timeseries of low harvest rates (Figure XX), high 1-SPR values (Figure XX), and the phase plot showing the history of being above the target biomass but also above the target fishing intensity reference points (Figure XX). The sharp decline in the OFL and ACL in coming years is the result of continued decline in the relative spawning output of the stock, which has placed it very near the “precautionary” zone for management.

### 4.2 Harvest Projections and Decision Tables

### 4.3 Evaluation of Scientific Uncertainty

Scientific uncertainty was evaluated via several likelihood profiles and a wide range of sensitivity analyses, not all of which are reported on here. Likelihood profiles were performed over unfished recruitment ( $R_0$ ), recruitment steepness ( $h$ ), and natural mortality ( $M$ ), as required by the Groundfish Terms of Reference. The profiles found the model results, particularly estimates of stock scale, to be relatively sensitive to changes to  $R_0$  and  $M$ , as would be expected, but largely insensitive to changes in steepness. An additional likelihood profile was also run over growth, and found the model results to be quite sensitive to the assumed values of the growth curve. Sensitivities were performed using alternative growth and maturity curves, alternative time blocks for selectivity and retention, and alternative historical landings timeseries. Model results (particularly estimates of stock status) were largely insensitive to all changes except for growth.

The model estimated uncertainty around the 2023 spawning biomass was  $\sigma = 0.18$  and the uncertainty around the OFL was  $\sigma = 0.16$ .

#### **4.4 Research and Data Needs**

Research and data needs for future assessments include the following:

1. Research into aging methods and availability of reliable age data would be valuable for future stock assessments. Otoliths have been collected in good quantities from the NWFSC survey, but there is currently no validated aging method for Shortspine thornyhead.
2. Additional investigation into growth patterns would provide valuable information for future population projections. We acknowledge that additional work on aging shortspine thornyhead would be required to make such additional growth research possible.
3. More investigation into maturity of Shortspine thornyhead is necessary to understand the patterns in maturity observed in WCGBTS samples.
4. Information on possible migration of Shortspine thornyheads would be valuable for understanding stock dynamics. Analysis of trace elements and stable isotopes in short-spine otoliths may provide valuable information on the extent of potential migrations. Possible connections between migration and maturity could likewise be explored.
5. A greater understanding of the connection between thornyheads and bottom type could be used to refine the indices of abundance. Thornyheads are very well sampled in trawlable habitat, but the extrapolation of density to a survey stratum could be improved by accounting for the proportion of different bottom types within a stratum and the relative density of thornyheads within each bottom type.
6. Additional investigation into spatial stock structure could be valuable for determining whether future assessments should develop a spatial assessment model, or if shortspine thornyhead should be assessed at distinct spatial scales in the future.

### **5 Acknowledgments**

Here are all the mad props!

## 6 References

- Anderson, S.C., Ward, E.J., English, P.A., and Barnett, L.A.K. 2022. sdmTMB: An r package for fast, flexible, and user-friendly generalized linear mixed effects models with spatial and spatiotemporal random fields. *bioRxiv*. Cold Spring Harbor Laboratory. doi:10.1101/2022.03.24.485545.
- Bizzarro, J., Dewitt, L., Wells, B., Curtis, A., Santora, J., and Field, J. 2023. California current trophic database (CCTD). Marine Data Archive; NOAA Southwest Fisheries Science Center: United States.
- Bradburn, M.J., Keller, A.A., and Horness, B.H. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.
- Butler, C.K., J. L. 1995. Age determination of shortspine thornyhead, *sebastolobus alascanus*, using otolith sections and  $^{210}\text{Pb}$ :  $^{226}\text{Ra}$  ratio. Admin. Rep. No. LJ-95-12. National Marine Fisheries Service, Southwest Fisheries Science Center, La Jolla, Calif.
- Cooper, D.W., Pearson, K.E., and Gunderson, D.R. 2005. Fecundity of shortspine thornyhead (*sebastolobus alascanus*) and longspine thornyhead (s. *Altivelis*) (scorpaenidae) from the northeastern pacific ocean, determined by stereological and gravimetric techniques\*. Available from <http://hdl.handle.net/1834/26245>.
- Dorval, E., Methot, R., Taylor, I., and Piner, K. 2022. Otolith chemistry indicates age and region of settlement of immature shortspine thornyhead *sebastolobus alascanus* in the eastern pacific ocean. *Mar. Ecol. Prog. Ser.* **693**: 157–175. doi:10.3354/meps14092.
- Du Preez, C., and Tunnicliffe, V. 2011. Shortspine thornyhead and rockfish (scorpaenidae) distribution in response to substratum, biogenic structures and trawling. *Mar. Ecol. Prog. Ser.* **425**: 217–231. doi:10.3354/meps09005.
- Echave, K., Siwicke, K.A., Sullivan, J., Ferris, B., and Hulson, P.F. 2022. Assessment of the thornyhead stock complex in the gulf of alaska.
- Erickson, D.L., and Pikitch, E.K. 1993. A histological description of shortspine thornyhead, *sebastolobus alascanus*, ovaries: Structures associated with the production of gelatinous egg masses. *Environmental Biology of Fishes* **36**(3): 273–282. doi:10.1007/BF00001723.
- Hamel, O.S. 2005. Status and future prospects for the shortspine thornyhead resource in waters off washington, oregon, and california as assessed in 2005. Northwest Fisheries Science Center, US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.

Ianelli, J.N., Lauth, R., and Jacobson, L.D. 1994. Status of the thornyhead (*sebastelobus* sp.) Resource in 1994. National Marine Fisheries Service, Alaska Fisheries Science Center, Seattle, WA; Southwest Fisheries Science Center, La Jolla, CA.

Jacobson, L.D. 1990. Thornyheads—stock assessment for 1990. National Marine Fisheries Service, Southwest Fisheries Science Center, La Jolla, CA.

Jacobson, L.D. 1991. Thornyheads—stock assessment for 1991. National Marine Fisheries Service, Southwest Fisheries Science Center, La Jolla, CA.

Jacobson, L.D., and Vetter, R.D. 1996. Bathymetric demography and niche separation of thornyhead rockfish: *Sebastolobus alascanus* and *sebastolobus altivelis*. **53**.

Karnowski, M., Gertseva, V.V., and Stephens, A. 2014. Historical Reconstruction of Oregon's Commercial Fisheries Landings. Oregon Department of Fish; Wildlife, Salem, OR.

Kastelle, C., Helser, T., TenBrink, T., Hutchinson, C., Goetz, B., Gburski, C., and Benson, I. 2020. Age validation of four rockfishes (genera *sebastes* and *sebastolobus*) with bomb-produced radiocarbon. *Mar. Freshwater Res.* **71**(10): 1355–1366. Available from <https://doi.org/10.1071/MF19280>.

Kline, D.E. 1996. Radiochemical age verification for two deep-sea rockfishes, *sebastolobus altivelis* and *s. alascanus*. San Jose State University.

Liu, O., Ward, S., and Anderson, S. in pressin press. Species redistribution creates unequal outcomes for multispecies fisheries under projected climate change, PREPRINT (version 1).

Methot, R.D., 1953-, Wetzel, C.R., Taylor, I.G., 1974-, and Doering, K. 2020. Stock synthesis user manual : Version 3.30.15. doi:10.25923/5wpn-qt71.

Methot, R.D., Taylor, I.G., and Chen, Y. 2011. Adjusting for bias due to variability of estimated recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(10): 1744–1760. doi:10.1139/f2011-092.

Methot, R.D., and Wetzel, C.R. 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* **142**: 86–99. doi:10.1016/j.fishres.2012.10.012.

Owen S. Hamel, J.M.C. 2022. Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research* **256**: 106477. doi:10.1016/j.fishres.2022.106477.

- Pearson, K.E., and Gunderson, D.R. 2003. Reproductive biology and ecology of shortspine thornyhead rockfish, *sebastolobus alascanus*, and longspine thornyhead rockfish, *s. Alutivelis*, from the northeastern pacific ocean. *Environmental Biology of Fishes* **67**(2): 117–136. doi:10.1023/A:1025623426858.
- Pikitch, E.K., Erickson, D.L., and Wallace, J.R. 1988. An evaluation of the effectiveness of trip limits as a management tool. Northwest; Alaska Fisheries Center, National Marine Fisheries Service NWAFC Processed Report. Available from <https://www.afsc.noaa.gov/Publications/ProcRpt/PR1988-27.pdf> [accessed 28 February 2017].
- Piner, K., and Methot, R. 2001. Stock status of shortspine thornyhead off the pacific west coast of the united states 2001. National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle, WA.
- R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available from <https://www.R-project.org/>.
- Ralston, S., Pearson, D.E., Field, J.C., and Key, M. 2010. Documentation of the California catch reconstruction project. US Department of Commerce, National Oceanic; Atmospheric Adminstration, National Marine.
- Rogers, B.R., Builder, T.L., Crone, P.R., Brodziak, J., Methot, R.D., and Conser, R.J. 1998. Status of the shortspine thornyhead (*sebastolobus alascanus*) resource in 1998. National Marine Fisheries Service, Northwest Fisheries Science Center, Newport, OR.; Alaska Fisheries Science Center, Seattle, WA.
- Rogers, J.B. 2003. Species allocation of *Sebastodes* and *sebastolobus* species caught by foreign countries off Washington, Oregon, and California, U.S.A. In 1965-1976. Unpublished document.
- Siebenaller, J.F. 1978. Genetic variability in deep-sea fishes of the genus *sebastolobus* (scorpaenidae). In *Marine Organisms: Genetics, Ecology, and Evolution*. Edited by B. Battaglia and J. Beardmore. Plenum Press, New York. pp. 95–122.
- Stepien, C.A. 1995. Population genetic divergence and geographic patterns from DNA sequences: Examples from marine and freshwater fishes. American Fisheries Society Symposium. pp. 263–287.
- Stepien, C.A., Dillon, A.K., and Patterson, A.K. 2000. Population genetics, phylogeography, and systematics of the thornyhead rockfishes (*sebastolobus*) along the deep continental slopes of the north pacific ocean. *Canadian Journal of Fisheries and Aquatic Sciences* **57**(8): 1701–1717. doi:10.1139/f00-095.
- Taylor, I.G. 2019. A 2019 catch-only projection from the 2013 stock assessment of shortspine thornyhead. National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle, WA.

- Taylor, I.G., Doering, K.L., Johnson, K.F., Wetzel, C.R., and Stewart, I.J. 2021. Beyond visualizing catch-at-age models: Lessons learned from the r4ss package about software to support stock assessments. *Fisheries Research* **239**: 105924. Available from <https://doi.org/10.1016/j.fishres.2021.105924>.
- Taylor, I.G., and Methot, R.D. 2013. Hiding or dead? A computationally efficient model of selective fisheries mortality. *Fisheries Research* **142**: 75–85. doi:<https://doi.org/10.1016/j.fishres.2012.08.021>.
- Taylor, I.G., and Stephens, A. 2013. Stock assessment of shortspine thornyhead in 2013. Portland: Pacific Fishery Management Council.
- Thorson, J.T., Dorn, M.W., and Hamel, O.S. 2019. Steepness for West Coast rockfishes: Results from a twelve-year experiment in iterative regional meta-analysis. *Fisheries Research*. doi:[10.1016/j.fishres.2018.03.014](https://doi.org/10.1016/j.fishres.2018.03.014).
- Thorson, J.T., Shelton, A.O., Ward, E.J., and Skaug, H.J. 2015. Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. *ICES Journal of Marine Science* **72**(5): 1297–1310. doi:[10.1093/icesjms/fsu243](https://doi.org/10.1093/icesjms/fsu243).
- Thorson, J.T., and Ward, E.J. 2013. Accounting for space–time interactions in index standardization models. *Fisheries Research* **147**: 426–433. doi:<https://doi.org/10.1016/j.fishres.2013.03.012>.
- Vetter, R.D., and Lynn, E.A. 1997. Bathymetric demography, enzyme activity patterns, and bioenergetics of deep-living scorpaenid fishes (genera *sebastes* and *sebastolobus*): Paradigms revisited. *Mar Ecol Prog Ser* **155**: 173–188. Available from <https://www.int-res.com/abstracts/meps/v155/p173-188/>.
- Wakefield, W.W., II. 1990. Patterns in the distribution of demersal fishes on the upper continental slope off central California with studies on the role of ontogenetic vertical migration in particle flux. PhD thesis, University of California, San Diego, United States – California. Available from <https://www.proquest.com/dissertations-theses/patterns-distribution-demersal-fishes-on-upper/docview/303821089/se-2?accountid=14784>.
- West Coast Regional Office. (n.d.). West coast groundfish trawl catch share program. NOAA. Available from <https://www.fisheries.noaa.gov/west-coast/sustainable-fisheries/west-coast-groundfish-trawl-catch-share-program>.

## 7 Tables

### 7.1 Executive Summary

**Table 1:** Recent landings by fleet, total landings summed across fleets, and the total mortality including discards.

Year	North Trawl	South Trawl	Non-Trawl	Total Landings	Total Dead
2013	547.98	294.83	166.40	1,009.21	1,085.62
2014	433.12	254.05	147.81	834.98	900.66
2015	503.14	244.29	131.30	878.73	945.40
2016	577.19	185.73	168.94	931.86	1,012.75
2017	606.86	158.30	223.82	988.97	1,085.60
2018	525.04	105.07	184.48	814.60	895.39
2019	402.95	127.94	143.48	674.37	736.82
2020	248.47	87.99	85.17	421.64	458.87
2021	226.00	73.39	78.74	378.13	411.62
2022	261.16	97.61	66.22	424.98	456.65

**Table 2:** Spawning output (millions of eggs) and fraction unfished with associated 95% confidence intervals (CI) from the base model.

Year	Spawning Output	Spawning Output 95% CI	Fraction Unfished	Fraction Unfished 95% CI
2013	8,875	5,904–11,845	0.4	0.4–0.5
2014	8,767	5,807–11,727	0.4	0.4–0.5
2015	8,679	5,728–11,630	0.4	0.3–0.5
2016	8,593	5,650–11,536	0.4	0.3–0.5
2017	8,508	5,572–11,445	0.4	0.3–0.5
2018	8,423	5,492–11,355	0.4	0.3–0.5
2019	8,358	5,431–11,286	0.4	0.3–0.5
2020	8,311	5,386–11,236	0.4	0.3–0.5
2021	8,291	5,366–11,215	0.4	0.3–0.5
2022	8,280	5,355–11,205	0.4	0.3–0.5
2023	8,273	5,346–11,201	0.4	0.3–0.5

**Table 3:** Estimated recent trend in recruitment and recruitment deviations and the 95% confidence intervals (CI) from the base model.

Year	Recruitment	95% CI	RecDevs	RecDev 95% CI
2013	6,024	2,469–14,698	-0.439	-1.352–0.474
2014	5,962	2,446–14,532	-0.447	-1.358–0.465
2015	5,954	2,438–14,542	-0.446	-1.360–0.468
2016	6,057	2,465–14,886	-0.427	-1.349–0.495
2017	5,836	2,385–14,279	-0.462	-1.379–0.454
2018	5,745	2,346–14,069	-0.476	-1.393–0.442
2019	8,863	3,557–22,086	-0.064	-1.003–0.874
2020	9,536	3,760–24,183	-0.013	-0.973–0.946
2021	10,335	3,984–26,811	0.044	-0.943–1.032
2022	10,118	3,924–26,090	0.000	-0.980–0.980
2023	10,117	3,924–26,086	0.000	-0.980–0.980

**Table 4:** Estimated recent trend in the  $(1-\text{SPR})/(1-\text{SPR } 50\%)$  where SPR is the spawning potential ratio the exploitation rate, and the 95% intervals.

Year	$(1-\text{SPR})/(1-\text{SPR } 50\%)$	95% CI	Exploitation Rate	95% CI
2013	1.25	1.03–1.47	0.0124	0.0084–0.0165
2014	1.12	0.90–1.34	0.0103	0.0069–0.0137
2015	1.15	0.92–1.37	0.0109	0.0073–0.0145
2016	1.19	0.96–1.42	0.0117	0.0078–0.0155
2017	1.23	1.00–1.46	0.0125	0.0083–0.0167
2018	1.09	0.86–1.32	0.0103	0.0069–0.0138
2019	0.95	0.73–1.17	0.0085	0.0056–0.0114
2020	0.66	0.48–0.84	0.0053	0.0035–0.0071
2021	0.59	0.43–0.76	0.0047	0.0031–0.0063
2022	0.64	0.47–0.81	0.0052	0.0034–0.0070

**Table 5:** Summary of reference points and management quantities, including estimates of the 95% intervals.

Variable of Interest	Estimate	95% CI
Unfished Spawning Output	20,332	16,338–24,327
Unfished Age 1+ Biomass (mt)	196,023	157,510–234,536
Unfished Recruitment (R0)	11,550	9,281–13,820
Spawning Output (2023)	8,273	5,346–11,201
Fraction Unfished (2023)	0.41	0.33–0.49
Reference Points Based SB40%		
Proxy Spawning Output SB40%	8,133	6,535–9,731
SPR Resulting in SB40%	0.458	0.458–0.458
Exploitation Rate Resulting in SB40%	0.012	0.011–0.012
Yield with SPR Based On SB40% (mt)	1,060	869–1,251
Reference Points Based on SPR Proxy for MSY		
Proxy Spawning Output (SPR50)	9,071	7,289–10,854
SPR50	0.500	-
Exploitation Rate Corresponding to SPR50	0.010	0.010–0.011
Yield with SPR50 at SB SPR (mt)	1,013	831–1,195
Reference Points Based on Estimated MSY Values		
Spawning Output at MSY (SB MSY)	5,651	4,548–6,755
SPR MSY	0.348	0.345–0.351
Exploitation Rate Corresponding to SPR MSY	0.017	0.016–0.017
MSY (mt)	1,121	919–1,323

**Table 6:** Recent trend in the overfishing limits (OFLs), the acceptable biological catches (ABCs), the annual catch limits (ACLs), the total landings, and total mortality (mt).

Year	OFL	ABC	ACL	Landings	Total Mortality
2013	2333	2230	1937	1,009.21	1,085.62
2014	2310	2208	1918	834.98	900.66
2015	3203	2668	2668	878.73	945.40
2016	3169	2640	2639	931.86	1,012.75
2017	3144	2619	2619	988.97	1,085.60
2018	3116	2596	2596	814.60	895.39
2019	3089	2573	2573	674.37	736.82
2020	3063	2551	2552	421.64	458.87
2021	3211	2183	2184	378.13	411.62
2022	3194	2130	2130	424.98	456.65

**Table 7:** Projections of potential OFLs (mt), ABCs (mt), estimated spawning output, and fraction unfished.

Year	Predicted OFL (mt)	ABC (mt)	Age 1 Biomass (mt)	Spawning Output	Fraction Unfished
2023	856.34	420.00	88,366.00	8,273.34	0.41
2024	872.85	420.00	88,980.60	8,275.64	0.41
2025	888.93	776.93	89,609.70	8,284.28	0.41
2026	899.90	778.42	89,881.20	8,273.27	0.41
2027	910.21	780.05	90,156.60	8,267.04	0.41
2028	919.89	780.99	90,434.90	8,265.16	0.41
2029	929.00	781.29	90,716.20	8,267.24	0.41
2030	937.62	781.04	91,000.70	8,272.85	0.41
2031	945.82	781.25	91,288.50	8,281.59	0.41
2032	953.66	780.10	91,578.90	8,292.98	0.41
2033	961.20	778.58	91,873.20	8,306.70	0.41
2034	968.48	777.69	92,171.70	8,322.42	0.41

**Table 8:** Summary of recent estimates and management quantities.

**Table 9:** Landings (mt) by fleet for all years, total landings (mt), and total mortality (mt) summed by year.

Year	North Trawl	South Trawl	Non-Trawl	Total Landings	Total Dead
1901	0.00	0.00	0.09	0.09	0.11
1902	0.00	0.00	0.11	0.11	0.14
1903	0.00	0.00	0.13	0.13	0.16
1904	0.00	0.00	0.15	0.15	0.19
1905	0.00	0.00	0.17	0.17	0.21
1906	0.00	0.00	0.19	0.19	0.24
1907	0.00	0.00	0.21	0.21	0.27
1908	0.00	0.00	0.23	0.23	0.29
1909	0.00	0.00	0.26	0.26	0.32
1910	0.00	0.00	0.28	0.28	0.34
1911	0.00	0.00	0.30	0.30	0.37
1912	0.00	0.00	0.32	0.32	0.39
1913	0.00	0.00	0.34	0.34	0.42
1914	0.00	0.00	0.36	0.36	0.44
1915	0.00	0.00	0.38	0.38	0.47
1916	0.00	0.00	0.40	0.40	0.49
1917	0.00	0.00	0.42	0.42	0.52
1918	0.00	0.00	0.44	0.44	0.54
1919	0.00	0.00	0.46	0.46	0.57
1920	0.00	0.00	0.48	0.48	0.60
1921	0.00	0.00	0.50	0.50	0.62
1922	0.00	0.00	0.52	0.52	0.65
1923	0.00	0.00	0.54	0.54	0.67
1924	0.00	0.00	0.56	0.56	0.70
1925	0.00	0.00	0.58	0.58	0.72
1926	0.00	0.00	0.60	0.60	0.75
1927	0.00	0.00	0.63	0.63	0.78
1928	0.00	0.00	1.05	1.05	1.29
1929	0.00	0.00	1.66	1.66	2.05
1930	0.00	0.00	1.39	1.39	1.72
1931	0.00	0.00	1.13	1.13	1.40
1932	0.00	0.00	0.42	0.42	0.52
1933	0.00	0.00	0.62	0.62	0.77
1934	0.00	4.57	0.71	5.28	5.62
1935	0.00	6.33	0.67	6.99	7.40
1936	0.00	2.70	1.45	4.15	4.60
1937	0.01	5.42	1.44	6.87	7.42
1938	0.00	5.62	1.34	6.96	7.49
1939	0.01	5.81	0.42	6.24	6.56

**Table 9:** Landings (mt) by fleet for all years, total landings (mt), and total mortality (mt) summed by year. (*continued*)

Year	North Trawl	South Trawl	Non-Trawl	Total Landings	Total Dead
1940	0.19	0.95	1.60	2.74	3.17
1941	0.29	1.96	2.65	4.90	5.64
1942	0.69	1.03	3.77	5.49	6.51
1943	3.06	1.43	10.17	14.66	17.50
1944	5.34	0.21	1.82	7.38	8.48
1945	5.34	0.99	0.68	7.01	7.87
1946	4.07	0.61	0.99	5.68	6.44
1947	4.75	0.04	0.72	5.51	6.27
1948	13.38	0.02	1.02	14.42	16.34
1949	13.52	0.02	0.34	13.88	15.67
1950	6.93	0.01	0.82	7.75	8.83
1951	11.16	0.00	0.59	11.76	13.32
1952	13.85	0.00	0.28	14.13	15.97
1953	2.63	2.96	0.24	5.83	6.35
1954	112.45	0.00	0.38	112.83	127.54
1955	62.93	4.99	0.28	68.20	76.69
1956	133.04	6.82	0.48	140.34	158.20
1957	63.88	0.00	0.49	64.37	72.91
1958	27.80	9.51	0.14	37.45	41.55
1959	28.81	32.26	0.24	61.31	66.53
1960	31.09	149.61	0.10	180.80	191.23
1961	28.55	56.76	0.38	85.68	91.92
1962	22.47	113.69	0.45	136.60	144.43
1963	7.77	223.17	0.27	231.21	241.60
1964	25.19	173.49	0.78	199.45	210.08
1965	31.75	307.58	0.13	339.46	356.18
1966	623.08	542.10	0.13	1,165.31	1,265.97
1967	375.82	867.40	0.34	1,243.57	1,325.03
1968	207.45	1,834.26	0.26	2,041.97	2,139.58
1969	215.73	430.43	0.95	647.10	689.99
1970	179.79	599.25	0.26	779.30	823.42
1971	347.53	607.82	0.08	955.43	1,018.56
1972	390.43	1,380.65	0.11	1,771.19	1,866.34
1973	704.17	2,321.09	0.44	3,025.70	3,188.41
1974	219.17	1,146.08	1.32	1,366.58	1,431.72
1975	399.30	1,872.54	0.53	2,272.37	2,381.61
1976	103.83	1,824.32	0.51	1,928.66	2,002.99
1977	112.56	1,472.13	9.21	1,593.90	1,658.75
1978	176.33	1,013.53	2.96	1,192.82	1,247.39
1979	65.96	1,715.69	6.56	1,788.21	1,855.78

**Table 9:** Landings (mt) by fleet for all years, total landings (mt), and total mortality (mt) summed by year. (*continued*)

Year	North Trawl	South Trawl	Non-Trawl	Total Landings	Total Dead
1980	179.97	1,204.88	2.25	1,387.10	1,449.25
1981	132.48	1,608.49	2.74	1,743.71	1,815.68
1982	293.65	1,654.79	2.71	1,951.15	2,044.80
1983	480.91	1,416.64	2.76	1,900.30	2,009.74
1984	1,039.29	1,841.97	1.18	2,882.44	3,078.65
1985	956.35	2,397.97	6.00	3,360.32	3,575.65
1986	613.11	2,331.52	7.98	2,952.62	3,129.25
1987	638.84	1,322.44	22.19	1,983.47	2,131.22
1988	877.45	1,965.85	24.92	2,868.22	3,084.11
1989	1,655.55	3,123.78	35.74	4,815.07	5,208.77
1990	2,092.81	1,911.73	31.36	4,035.90	4,451.92
1991	2,416.99	995.94	53.54	3,466.47	3,907.34
1992	1,782.91	1,455.11	61.25	3,299.27	3,677.11
1993	2,010.20	1,571.84	26.59	3,608.63	4,034.78
1994	2,083.16	1,182.92	21.00	3,287.08	3,716.96
1995	953.79	929.27	57.01	1,940.08	2,176.77
1996	810.14	699.59	100.09	1,609.82	1,824.69
1997	689.59	654.17	62.74	1,406.50	1,592.89
1998	582.09	593.35	62.18	1,237.62	1,404.29
1999	373.90	309.07	64.39	747.36	858.57
2000	340.72	421.91	66.59	829.22	944.10
2001	276.61	197.49	57.98	532.08	618.30
2002	288.51	364.76	114.25	767.53	883.27
2003	346.05	302.40	166.56	815.01	953.33
2004	328.55	286.59	139.49	754.62	882.55
2005	292.59	214.11	149.69	656.39	774.51
2006	334.06	210.53	159.17	703.77	833.84
2007	626.03	222.56	158.80	1,007.39	1,102.25
2008	972.95	259.94	196.97	1,429.86	1,563.90
2009	1,022.69	308.38	200.62	1,531.70	1,673.25
2010	834.86	284.22	228.36	1,347.44	1,481.92
2011	483.47	232.99	260.52	976.98	1,069.97
2012	455.93	263.59	192.07	911.59	987.50
2013	547.98	294.83	166.40	1,009.21	1,085.62
2014	433.12	254.05	147.81	834.98	900.66
2015	503.14	244.29	131.30	878.73	945.40
2016	577.19	185.73	168.94	931.86	1,012.75
2017	606.86	158.30	223.82	988.97	1,085.60
2018	525.04	105.07	184.48	814.60	895.39
2019	402.95	127.94	143.48	674.37	736.82

**Table 9:** Landings (mt) by fleet for all years, total landings (mt), and total mortality (mt) summed by year. (*continued*)

Year	North	South	Non-Trawl	Total	Total Dead
	Trawl	Trawl		Landings	
2020	248.47	87.99	85.17	421.64	458.87
2021	226.00	73.39	78.74	378.13	411.62
2022	261.16	97.61	66.22	424.98	456.65

**Table 10:** Time series of population estimates from the base model.

Year	Total Biomass (mt)	Spawning Output	Total Biomass (mt)	Fraction Unfished	Age 0 Recruits	Total Mortality	(1-SPR)/(1-50%)	Exploitation Rate
1901	196,035	20,332	196,023	100.0	11,286	0.11	0.00	0.00
1902	196,033	20,332	196,020	100.0	11,260	0.14	0.00	0.00
1903	196,029	20,332	196,016	100.0	11,227	0.16	0.00	0.00
1904	196,024	20,332	196,011	100.0	11,187	0.19	0.00	0.00
1905	196,017	20,332	196,005	100.0	11,147	0.21	0.00	0.00
1906	196,009	20,332	195,996	100.0	11,113	0.24	0.00	0.00
1907	195,998	20,332	195,986	100.0	11,081	0.27	0.00	0.00
1908	195,985	20,332	195,973	100.0	11,040	0.29	0.00	0.00
1909	195,970	20,332	195,958	100.0	11,001	0.32	0.00	0.00
1910	195,951	20,332	195,939	100.0	10,955	0.34	0.00	0.00
1911	195,930	20,332	195,917	100.0	10,919	0.37	0.00	0.00
1912	195,904	20,332	195,892	100.0	10,909	0.39	0.00	0.00
1913	195,876	20,332	195,864	100.0	10,909	0.42	0.00	0.00
1914	195,843	20,331	195,831	100.0	10,862	0.44	0.00	0.00
1915	195,806	20,331	195,794	100.0	10,810	0.47	0.00	0.00
1916	195,765	20,330	195,753	100.0	10,784	0.49	0.00	0.00
1917	195,719	20,330	195,707	100.0	10,764	0.52	0.00	0.00
1918	195,669	20,329	195,657	100.0	10,745	0.54	0.00	0.00
1919	195,613	20,328	195,601	100.0	10,718	0.57	0.00	0.00
1920	195,552	20,327	195,541	100.0	10,700	0.60	0.00	0.00
1921	195,486	20,325	195,475	100.0	10,679	0.62	0.00	0.00
1922	195,415	20,324	195,403	100.0	10,667	0.65	0.00	0.00
1923	195,338	20,322	195,326	99.9	10,665	0.67	0.00	0.00
1924	195,256	20,320	195,244	99.9	10,660	0.70	0.00	0.00
1925	195,168	20,317	195,156	99.9	10,661	0.72	0.00	0.00
1926	195,074	20,314	195,062	99.9	10,668	0.75	0.00	0.00
1927	194,975	20,311	194,963	99.9	10,681	0.78	0.00	0.00
1928	194,871	20,308	194,859	99.9	10,702	1.29	0.00	0.00

**Table 10:** Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass (mt)	Frac-tion Un-fished	Age 0 Recruits	Total Mortal-ity	(1- SPR)/(1- 50%)	Ex-ploita-tion Rate
1929	194,761	20,304	194,749	99.9	10,732	2.05	0.00	0.00
1930	194,645	20,299	194,633	99.8	10,770	1.72	0.00	0.00
1931	194,525	20,294	194,513	99.8	10,819	1.40	0.00	0.00
1932	194,400	20,288	194,388	99.8	10,878	0.52	0.00	0.00
1933	194,273	20,282	194,261	99.8	10,949	0.77	0.00	0.00
1934	194,142	20,276	194,130	99.7	11,031	5.62	0.00	0.00
1935	194,003	20,268	193,990	99.7	11,124	7.40	0.01	0.00
1936	193,859	20,260	193,847	99.6	11,228	4.60	0.00	0.00
1937	193,717	20,252	193,705	99.6	11,340	7.42	0.01	0.00
1938	193,571	20,242	193,559	99.6	11,458	7.49	0.01	0.00
1939	193,425	20,232	193,413	99.5	11,578	6.56	0.00	0.00
1940	193,281	20,222	193,268	99.5	11,693	3.17	0.00	0.00
1941	193,142	20,212	193,129	99.4	11,796	5.64	0.00	0.00
1942	193,003	20,200	192,990	99.4	11,878	6.51	0.00	0.00
1943	192,866	20,188	192,853	99.3	11,928	17.50	0.01	0.00
1944	192,722	20,175	192,709	99.2	11,936	8.48	0.01	0.00
1945	192,591	20,162	192,578	99.2	11,892	7.87	0.01	0.00
1946	192,466	20,149	192,453	99.1	11,789	6.44	0.00	0.00
1947	192,348	20,135	192,335	99.0	11,624	6.27	0.00	0.00
1948	192,234	20,122	192,221	99.0	11,400	16.34	0.01	0.00
1949	192,113	20,106	192,101	98.9	11,124	15.67	0.01	0.00
1950	191,997	20,091	191,985	98.8	10,805	8.83	0.01	0.00
1951	191,889	20,077	191,878	98.7	10,427	13.32	0.01	0.00
1952	191,779	20,061	191,768	98.7	10,036	15.97	0.01	0.00
1953	191,664	20,046	191,654	98.6	9,643	6.35	0.00	0.00
1954	191,558	20,031	191,548	98.5	9,259	127.54	0.11	0.00
1955	191,319	20,006	191,309	98.4	8,891	76.69	0.07	0.00
1956	191,127	19,986	191,118	98.3	8,548	158.20	0.14	0.00
1957	190,840	19,958	190,831	98.2	8,235	72.91	0.06	0.00
1958	190,634	19,938	190,625	98.1	7,957	41.55	0.04	0.00
1959	190,449	19,920	190,441	98.0	7,719	66.53	0.06	0.00
1960	190,225	19,901	190,217	97.9	7,524	191.23	0.15	0.00
1961	189,853	19,870	189,845	97.7	7,380	91.92	0.07	0.00
1962	189,570	19,849	189,562	97.6	7,293	144.43	0.11	0.00
1963	189,213	19,823	189,205	97.5	7,270	241.60	0.18	0.00
1964	188,734	19,788	188,726	97.3	7,320	210.08	0.16	0.00
1965	188,269	19,756	188,260	97.2	7,448	356.18	0.26	0.00
1966	187,627	19,710	187,619	96.9	7,658	1265.97	0.82	0.01
1967	185,993	19,584	185,984	96.3	7,945	1325.03	0.83	0.01

**Table 10:** Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass (mt)	Fraction Unfished	Age 0 Recruits	Total Mortality	(1-SPR)/(1-50%)	Exploitation Rate
1968	184,271	19,449	184,262	95.7	8,302	2139.58	1.13	0.01
1969	181,659	19,237	181,650	94.6	8,713	689.99	0.51	0.00
1970	180,573	19,154	180,563	94.2	9,172	823.42	0.58	0.00
1971	179,330	19,057	179,319	93.7	9,664	1018.56	0.71	0.01
1972	177,866	18,941	177,855	93.2	10,172	1866.34	1.08	0.01
1973	175,487	18,745	175,476	92.2	10,640	3188.41	1.46	0.02
1974	171,686	18,425	171,674	90.6	10,968	1431.72	0.93	0.01
1975	169,754	18,259	169,742	89.8	11,071	2381.61	1.28	0.01
1976	166,807	18,003	166,795	88.5	10,910	2002.99	1.16	0.01
1977	164,266	17,776	164,254	87.4	10,589	1658.75	1.05	0.01
1978	162,097	17,578	162,086	86.5	10,266	1247.39	0.88	0.01
1979	160,376	17,415	160,365	85.7	10,081	1855.78	1.14	0.01
1980	158,012	17,193	158,001	84.6	10,100	1449.25	1.00	0.01
1981	156,095	17,005	156,084	83.6	10,290	1815.68	1.16	0.01
1982	153,801	16,781	153,790	82.5	10,486	2044.80	1.26	0.01
1983	151,280	16,534	151,268	81.3	10,566	2009.74	1.28	0.01
1984	148,816	16,288	148,805	80.1	10,513	3078.65	1.58	0.02
1985	145,227	15,941	145,215	78.4	10,418	3575.65	1.67	0.02
1986	141,121	15,543	141,110	76.4	10,340	3129.25	1.61	0.02
1987	137,513	15,183	137,502	74.7	10,353	2131.22	1.41	0.02
1988	135,000	14,914	134,989	73.4	10,553	3084.11	1.64	0.02
1989	131,494	14,556	131,482	71.6	10,798	5208.77	1.87	0.04
1990	125,735	14,000	125,723	68.9	10,945	4451.92	1.85	0.04
1991	120,810	13,513	120,799	66.5	10,529	3907.34	1.84	0.03
1992	116,490	13,077	116,479	64.3	10,018	3677.11	1.82	0.03
1993	112,431	12,657	112,420	62.3	9,681	4034.78	1.87	0.04
1994	108,009	12,205	107,999	60.0	8,899	3716.96	1.86	0.03
1995	103,948	11,782	103,938	57.9	8,778	2176.77	1.64	0.02
1996	101,552	11,498	101,544	56.5	6,632	1824.69	1.55	0.02
1997	99,563	11,248	99,555	55.3	7,224	1592.89	1.48	0.02
1998	97,847	11,022	97,839	54.2	7,653	1404.29	1.40	0.01
1999	96,366	10,817	96,356	53.2	9,260	858.57	1.07	0.01
2000	95,510	10,664	95,497	52.4	11,621	944.10	1.14	0.01
2001	94,612	10,507	94,597	51.7	13,999	618.30	0.86	0.01
2002	94,120	10,383	94,108	51.1	11,515	883.27	1.09	0.01
2003	93,426	10,240	93,391	50.4	32,115	953.33	1.15	0.01
2004	92,710	10,097	92,697	49.7	11,038	882.55	1.10	0.01
2005	92,210	9,964	92,196	49.0	12,291	774.51	1.01	0.01
2006	91,827	9,845	91,813	48.4	12,616	833.84	1.06	0.01

**Table 10:** Time series of population estimates from the base model. (*continued*)

Year	Total Biomass (mt)	Spawning Output	Total Biomass (mt)	Fraction Unfished	Age 0 Recruits	Total Mortality	(1-SPR)/(1-50%)	Exploitation Rate
2007	91,421	9,726	91,407	47.8	12,325	1102.25	1.26	0.01
2008	90,816	9,591	90,777	47.2	34,678	1563.90	1.50	0.02
2009	89,785	9,424	89,774	46.3	10,066	1673.25	1.54	0.02
2010	88,786	9,251	88,776	45.5	8,768	1481.92	1.47	0.02
2011	87,968	9,098	87,960	44.7	8,002	1069.97	1.25	0.01
2012	87,599	8,980	87,592	44.2	6,687	987.50	1.19	0.01
2013	87,359	8,875	87,353	43.6	6,024	1085.62	1.25	0.01
2014	87,051	8,767	87,044	43.1	5,962	900.66	1.12	0.01
2015	86,968	8,679	86,961	42.7	5,954	945.40	1.15	0.01
2016	86,867	8,593	86,861	42.3	6,057	1012.75	1.19	0.01
2017	86,722	8,508	86,715	41.8	5,836	1085.60	1.23	0.01
2018	86,519	8,423	86,513	41.4	5,745	895.40	1.09	0.01
2019	86,530	8,358	86,520	41.1	8,863	736.82	0.95	0.01
2020	86,721	8,311	86,711	40.9	9,536	458.87	0.66	0.01
2021	87,231	8,291	87,220	40.8	10,335	411.62	0.59	0.00
2022	87,816	8,280	87,805	40.7	10,118	456.65	0.64	0.01
2023	88,377	8,273	88,366	40.7	10,117	420.00	0.58	0.00
2024	88,992	8,276	88,981	40.7	10,117	420.00	0.58	0.00
2025	89,621	8,284	89,610	40.7	10,120	776.93	0.91	0.01
2026	89,892	8,273	89,881	40.7	10,117	778.42	0.90	0.01
2027	90,168	8,267	90,157	40.7	10,115	780.05	0.90	0.01
2028	90,446	8,265	90,435	40.6	10,115	780.99	0.89	0.01
2029	90,727	8,267	90,716	40.7	10,115	781.29	0.89	0.01
2030	91,012	8,273	91,001	40.7	10,117	781.04	0.88	0.01
2031	91,300	8,282	91,288	40.7	10,119	781.25	0.88	0.01
2032	91,590	8,293	91,579	40.8	10,122	780.10	0.87	0.01
2033	91,884	8,307	91,873	40.9	10,125	778.57	0.86	0.01
2034	92,183	8,322	92,172	40.9	10,129	777.69	0.86	0.01

**Table 11:** Likelihood components by source.

Source	Likelihood Component
TOTAL	536.6430000
Catch	0.0000000
Equil catch	0.0000000
Survey	-48.5980000
Discard	406.1260000
Mean body wt	-79.6714000
Length comp	270.3960000
Recruitment	-13.1937000
InitEQ Regime	0.0000000
Forecast Recruitment	0.0125207
Parm priors	1.5716500
Parm devs	0.0000000
F Ballpark	0.0000000
F Ballpark(info only) 1999 estF tgtF	0.0148413
Crash Pen	0.0000000

**Table 12:** Sample sizes of length composition samples for shortspine thornyhead landings.

Year	North Trawl	South Trawl	Non-trawl	NA	NA	NA
1981	30	1	737	42	NA	NA
1982	150	5	723	57	NA	NA
1983	NA	NA	1230	91	NA	NA
1984	NA	NA	2755	118	NA	NA
1985	NA	NA	3176	136	3	3
1986	NA	NA	978	42	9	9
1987	NA	NA	343	26	54	15
1988	NA	NA	140	15	8	7
1989	NA	NA	741	38	18	12
1990	390	33	517	32	24	14
1991	1059	47	532	41	NA	NA
1992	1227	52	448	36	75	4
1993	281	12	993	56	3	3
1994	40	1	1367	65	46	12
1995	24	2	2248	103	36	6
1996	497	15	2078	94	26	1
1997	2322	49	1720	92	36	2
1998	757	28	1130	57	130	7
1999	819	27	821	48	1852	73
2000	660	23	1027	63	447	20
2001	1632	45	1413	73	132	8
2002	2183	50	2320	108	1036	42
2003	2431	67	1909	80	834	29
2004	1509	53	1073	61	133	10
2005	1649	50	1393	75	620	34
2006	1573	53	3109	156	596	32
2007	2452	76	1893	98	393	17
2008	2660	90	2212	106	2037	80
2009	2854	85	2137	87	1298	60
2010	3130	91	1720	88	1346	75
2011	2381	67	2950	106	3276	163
2012	2262	67	2423	102	2419	137
2013	2519	84	2505	102	2010	115
2014	2036	73	1490	77	3283	149
2015	1915	65	1615	69	2971	152
2016	1495	59	2362	82	3087	147
2017	2158	75	1571	48	2504	139
2018	1828	78	1222	29	1333	91
2019	1539	68	740	29	1702	134
2020	951	47	897	26	696	64

**Table 12:** Sample sizes of length composition samples for shortspine thornyhead landings.  
*(continued)*

Year	North Trawl	South Trawl	Non-trawl	NA	NA	NA
2021	1505	80	294	15	1067	108
2022	1103	57	273	13	863	75

**Table 13:** Sample sizes of length composition samples for discards.

Year	North Trawl	South Trawl	Non-trawl
2005	NA	NA	4
2006	148	56	102
2007	249	64	145
2008	354	79	102
2009	485	102	71
2010	271	43	98
2011	282	74	168
2012	378	126	224
2013	366	155	55
2014	311	126	120
2015	204	154	148
2016	216	108	189
2017	182	39	153
2018	211	47	148
2019	175	79	77
2020	163	99	43
2021	210	97	104

**Table 14:** Survey stratification information for each of the fishery independent surveys

Survey	Strata Definitions
Survey	Strata Definitions
AFSC Slope	2 strata: <a0><a0><a0><a0><a0>32.0-49.0 degrees N: 150-500 m, 500-1280 m
NWFSC Combo<a0>	7 strata <a0><a0><a0><a0><a0>32.0-34.5 degrees N: 183-550 m, 550-1280 m
(West Coast Groundfish Bottom Trawl Survey)	<a0><a0><a0><a0><a0>34.5-40.5 degrees N: 183-550 m, 550-1280 m <a0><a0><a0><a0><a0>40.5-49.0 degrees N: 100-183 m, 183-550 m, 550-1280 m The depth breaks at 183 m and 550 m are associated with changes in sampling intensity of the survey and are recommended to be used.
NWFSC Slope	6 strata <a0><a0><a0><a0><a0>32.0-40.5 degrees N: 55-500m, 550-1280m <a0><a0><a0><a0><a0>40.5-43.0 degrees N: 55-550m, 550-1280m <a0><a0><a0><a0><a0>43.0-49.0 degrees N: 55-550m, 550-1280m
AFSC Triennial 1<a0>	1 stratum: <=366 m
AFSC Triennial 2<a0>	1 stratum: 366-500 m

**Table 15:** Survey samples and hauls for each of the fishery independent surveys for available years spanning from 1989 to 2022.

Year	survey	samples	hauls
1989	Early	1770	51
	Triennial		
1992	Early	1143	23
	Triennial		
1995	Late	9984	128
	Triennial		
1997	AFSC Slope	7454	171
1998	WCGBTS	8946	270
1998	Late	9871	147
	Triennial		
1999	AFSC Slope	6752	188
1999	WCGBTS	10061	302
2000	AFSC Slope	7017	196
2000	WCGBTS	8057	295
2001	AFSC Slope	6072	196
2001	WCGBTS	8091	297
2001	Late	10147	190
	Triennial		
2002	WCGBTS	11835	374
2003	nwfsc combo	7693	293
2004	nwfsc combo	6694	214
2004	Late	8508	137
	Triennial		
2005	nwfsc combo	8047	315
2006	nwfsc combo	6198	332
2007	nwfsc combo	5499	367
2008	nwfsc combo	4697	362
2009	nwfsc combo	4195	340
2010	nwfsc combo	3859	360
2011	nwfsc combo	4697	347
2012	nwfsc combo	4678	349
2013	nwfsc combo	3119	247
2014	nwfsc combo	4617	346
2015	nwfsc combo	4511	332
2016	nwfsc combo	4604	355
2017	nwfsc combo	4730	363
2018	nwfsc combo	4996	368
2019	nwfsc combo	2401	175
2021	nwfsc combo	4690	345
2022	nwfsc combo	4202	312

**Table 16:** Comparison of likelihoods for all sensitivity analyses

Source	Base Model	Low Growth	High Growth	2013 Maturity	Inde- term. Maturity	Imputed Landings	2013 Landings
Total L	536.643	729.873	695.432	718.427	718.481	536.308	535.813
Survey L	-48.598	-43.945	-52.567	-47.060	-47.016	-48.690	-49.003
Length Comp L	270.396	443.197	435.191	440.203	440.192	270.292	270.614
Discards L	406.126	417.364	405.127	414.636	414.648	406.031	406.048
Mean Body Wt L	-79.671	-78.922	-79.368	-78.982	-78.983	-79.679	-79.684
Recruitment L	-13.194	-9.747	-15.400	-12.339	-12.328	-13.234	-13.754
Prior L	1.572	1.894	1.865	1.874	1.874	1.576	1.580
R0	11550.400	15443.300	7230.490	11159.700	11167.100	11666.700	11879.200
B0	8273.340	7205.460	10956.200	8442.120	8069.820	8167.330	8257.250
Depletion	0.407	0.418	0.370	0.402	0.391	0.398	0.395
Relative SPR	0.584	1.000	0.981	1.000	0.995	0.584	0.577

**Table 17:** All parameter estimates for the proposed base model

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
NatM_break_1_Fem_GP_1	0.0400000	-3	NA	0.040000	0.040000
NatM_break_2_Fem_GP_1	0.0400000	-3	NA	NA	NA
L_at_Amin_Fem_GP_1	11.3832000	-2	NA	NA	NA
L_at_Amax_Fem_GP_1	73.6079000	-2	NA	NA	NA
VonBert_K_Fem_GP_1	0.0098986	-3	NA	NA	NA
CV_young_Fem_GP_1	0.1090340	-3	NA	NA	NA
CV_old_Fem_GP_1	0.1090340	-3	NA	NA	NA
Wtlen_1_Fem_GP_1	0.0000065	-3	NA	NA	NA
Wtlen_2_Fem_GP_1	3.1797500	-3	NA	NA	NA
Mat50%_Fem_GP_1	31.4247000	-3	NA	NA	NA
Mat_slope_Fem_GP_1	-0.1772910	-3	NA	NA	NA
Eggs_scalar_Fem_GP_1	0.0000001	-3	NA	NA	NA
Eggs_exp_len_Fem_GP_1	3.9780000	-3	NA	NA	NA
NatM_break_1_Mal_GP_1	0.0400000	-3	NA	NA	NA
NatM_break_2_Mal_GP_1	0.0400000	-3	NA	NA	NA
L_at_Amin_Mal_GP_1	9.1733000	-3	NA	NA	NA
L_at_Amax_Mal_GP_1	66.0728000	-2	NA	NA	NA
VonBert_K_Mal_GP_1	0.0167854	-3	NA	NA	NA
CV_young_Mal_GP_1	0.1090340	-3	NA	NA	NA
CV_old_Mal_GP_1	0.1090340	-3	NA	NA	NA
Wtlen_1_Mal_GP_1	0.0000067	-3	NA	NA	NA
Wtlen_2_Mal_GP_1	3.1732500	-3	NA	NA	NA
CohortGrowDev	1.0000000	-1	NA	NA	NA
FracFemale_GP_1	0.5000000	-99	NA	NA	NA
SR_LN(R0)	9.3544800	4	0.0001044	NA	NA
SR_BH_stEEP	0.7200000	-2	NA	NA	NA
SR_sigmaR	0.5000000	-4	NA	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
SR_regime	0.0000000	-4	NA	NA	NA
SR_autocorr	0.0000000	-1	NA	NA	NA
Early_InitAge_1	-0.0212086	6	-0.0000012	NA	NA
Early_RecrDev_1901	-0.0231360	6	0.0000000	NA	NA
Early_RecrDev_1902	-0.0254220	6	0.0000011	NA	NA
Early_RecrDev_1903	-0.0283702	6	-0.0000005	NA	NA
Early_RecrDev_1904	-0.0319698	6	0.0000020	NA	NA
Early_RecrDev_1905	-0.0355489	6	0.0000036	NA	NA
Early_RecrDev_1906	-0.0385733	6	-0.0000019	NA	NA
Early_RecrDev_1907	-0.0414951	6	0.0000025	NA	NA
Early_RecrDev_1908	-0.0451950	6	-0.0000019	NA	NA
Early_RecrDev_1909	-0.0487619	6	0.0000017	NA	NA
Early_RecrDev_1910	-0.0529210	6	-0.0000073	NA	NA
Early_RecrDev_1911	-0.0562301	6	0.0000016	NA	NA
Early_RecrDev_1912	-0.0571398	6	0.0000020	NA	NA
Early_RecrDev_1913	-0.0571633	6	0.0000058	NA	NA
Early_RecrDev_1914	-0.0614764	6	0.0000008	NA	NA
Early_RecrDev_1915	-0.0662355	6	-0.0000012	NA	NA
Early_RecrDev_1916	-0.0686894	6	-0.0000014	NA	NA
Early_RecrDev_1917	-0.0705061	6	0.0000064	NA	NA
Early_RecrDev_1918	-0.0722232	6	0.0000061	NA	NA
Early_RecrDev_1919	-0.0747498	6	-0.0000012	NA	NA
Early_RecrDev_1920	-0.0764656	6	-0.0000017	NA	NA
Early_RecrDev_1921	-0.0784239	6	0.0000041	NA	NA
Early_RecrDev_1922	-0.0795038	6	0.0000050	NA	NA
Early_RecrDev_1923	-0.0797302	6	0.0000051	NA	NA
Early_RecrDev_1924	-0.0801474	6	0.0000042	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Early_RecrDev_1925	-0.0800694	6	0.0000076	NA	NA
Early_RecrDev_1926	-0.0794285	6	0.0000037	NA	NA
Early_RecrDev_1927	-0.0781531	6	0.0000000	NA	NA
Early_RecrDev_1928	-0.0761695	6	-0.0000007	NA	NA
Early_RecrDev_1929	-0.0734039	6	0.0000010	NA	NA
Early_RecrDev_1930	-0.0697849	6	0.0000091	NA	NA
Early_RecrDev_1931	-0.0652490	6	-0.0000003	NA	NA
Early_RecrDev_1932	-0.0597442	6	0.0000033	NA	NA
Early_RecrDev_1933	-0.0532400	6	0.0000054	NA	NA
Early_RecrDev_1934	-0.0457363	6	0.0000005	NA	NA
Early_RecrDev_1935	-0.0372755	6	0.0000061	NA	NA
Early_RecrDev_1936	-0.0279592	6	0.0000062	NA	NA
Early_RecrDev_1937	-0.0179638	6	0.0000021	NA	NA
Early_RecrDev_1938	-0.0075590	6	0.0000060	NA	NA
Early_RecrDev_1939	0.0028729	6	0.0000057	NA	NA
Early_RecrDev_1940	0.0128254	6	0.0000044	NA	NA
Early_RecrDev_1941	0.0216624	6	0.0000029	NA	NA
Early_RecrDev_1942	0.0286306	6	0.0000073	NA	NA
Early_RecrDev_1943	0.0328934	6	0.0000042	NA	NA
Early_RecrDev_1944	0.0335935	6	0.0000061	NA	NA
Early_RecrDev_1945	0.0299395	6	0.0000045	NA	NA
Early_RecrDev_1946	0.0213064	6	0.0000046	NA	NA
Early_RecrDev_1947	0.0073268	6	0.0000034	NA	NA
Early_RecrDev_1948	-0.0120490	6	0.0000022	NA	NA
Early_RecrDev_1949	-0.0365423	6	0.0000079	NA	NA
Early_RecrDev_1950	-0.0655857	6	0.0000017	NA	NA
Early_RecrDev_1951	-0.0981321	6	0.0000054	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Early_RecrDev_1952	-0.1333580	6	0.0000045	NA	NA
Early_RecrDev_1953	-0.1702910	6	0.0000035	NA	NA
Early_RecrDev_1954	-0.2079570	6	0.0000025	NA	NA
Early_RecrDev_1955	-0.2453950	6	0.0000053	NA	NA
Early_RecrDev_1956	-0.2817010	6	0.0000040	NA	NA
Early_RecrDev_1957	-0.3159610	6	0.0000021	NA	NA
Early_RecrDev_1958	-0.3472870	6	0.0000048	NA	NA
Early_RecrDev_1959	-0.3747180	6	0.0000048	NA	NA
Early_RecrDev_1960	-0.3971880	6	0.0000033	NA	NA
Early_RecrDev_1961	-0.4134930	6	0.0000026	NA	NA
Early_RecrDev_1962	-0.4223490	6	0.0000025	NA	NA
Early_RecrDev_1963	-0.4223980	6	0.0000027	NA	NA
Early_RecrDev_1964	-0.4124790	6	0.0000030	NA	NA
Early_RecrDev_1965	-0.3919790	6	0.0000025	NA	NA
Early_RecrDev_1966	-0.3610010	6	0.0000057	NA	NA
Early_RecrDev_1967	-0.3206150	6	0.0000034	NA	NA
Early_RecrDev_1968	-0.2730580	6	0.0000047	NA	NA
Early_RecrDev_1969	-0.2207630	6	0.0000034	NA	NA
Early_RecrDev_1970	-0.1660450	6	0.0000042	NA	NA
Early_RecrDev_1971	-0.1102690	6	0.0000048	NA	NA
Early_RecrDev_1972	-0.0555105	6	-0.0000016	NA	NA
Early_RecrDev_1973	-0.0064973	6	0.0000078	NA	NA
Early_RecrDev_1974	0.0285943	6	0.0000078	NA	NA
Early_RecrDev_1975	0.0417962	6	0.0000059	NA	NA
Early_RecrDev_1976	0.0316066	6	0.0000050	NA	NA
Early_RecrDev_1977	0.0060750	6	0.0000071	NA	NA
Early_RecrDev_1978	-0.0207783	6	-0.0000014	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Early_RecrDev_1979	-0.0349278	6	0.0000093	NA	NA
Early_RecrDev_1980	-0.0286589	6	0.0000057	NA	NA
Early_RecrDev_1981	-0.0059002	6	0.0000048	NA	NA
Early_RecrDev_1982	0.0173986	6	0.0000057	NA	NA
Early_RecrDev_1983	0.0267644	6	0.0000044	NA	NA
Early_RecrDev_1984	0.0234703	6	0.0000057	NA	NA
Early_RecrDev_1985	0.0169923	6	0.0000046	NA	NA
Early_RecrDev_1986	0.0125559	6	0.0000031	NA	NA
Early_RecrDev_1987	0.0167085	6	0.0000042	NA	NA
Early_RecrDev_1988	0.0381407	6	0.0000038	NA	NA
Early_RecrDev_1989	0.0642914	6	0.0000015	NA	NA
Early_RecrDev_1990	0.0829343	6	0.0000003	NA	NA
Early_RecrDev_1991	0.0490698	6	0.0000016	NA	NA
Early_RecrDev_1992	0.0039841	6	0.0000012	NA	NA
Early_RecrDev_1993	-0.0254987	6	-0.0000009	NA	NA
Early_RecrDev_1994	-0.1043340	6	-0.0000016	NA	NA
Early_RecrDev_1995	-0.1125850	6	-0.0000029	NA	NA
Main_RecrDev_1996	-0.3889790	6	0.0000025	NA	NA
Main_RecrDev_1997	-0.2999770	6	-0.0000012	NA	NA
Main_RecrDev_1998	-0.2388930	6	-0.0000013	NA	NA
Main_RecrDev_1999	-0.0452484	6	-0.0000046	NA	NA
Main_RecrDev_2000	0.1842930	6	-0.0000038	NA	NA
Main_RecrDev_2001	0.3730210	6	-0.0000105	NA	NA
Main_RecrDev_2002	0.1797160	6	-0.0000084	NA	NA
Main_RecrDev_2003	1.2078500	6	-0.0000102	NA	NA
Main_RecrDev_2004	0.1423860	6	-0.0000039	NA	NA
Main_RecrDev_2005	0.2522910	6	-0.0000020	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Main_RecrDev_2006	0.2805240	6	-0.0000005	NA	NA
Main_RecrDev_2007	0.2594420	6	0.0000046	NA	NA
Main_RecrDev_2008	1.2964800	6	-0.0000063	NA	NA
Main_RecrDev_2009	0.0628162	6	0.0000040	NA	NA
Main_RecrDev_2010	-0.0717626	6	0.0000043	NA	NA
Main_RecrDev_2011	-0.1599470	6	0.0000048	NA	NA
Main_RecrDev_2012	-0.3369110	6	0.0000075	NA	NA
Main_RecrDev_2013	-0.4389960	6	0.0000070	NA	NA
Main_RecrDev_2014	-0.4468810	6	0.0000072	NA	NA
Main_RecrDev_2015	-0.4461790	6	0.0000113	NA	NA
Main_RecrDev_2016	-0.4270100	6	0.0000011	NA	NA
Main_RecrDev_2017	-0.4622620	6	0.0000074	NA	NA
Main_RecrDev_2018	-0.4757690	6	0.0000082	NA	NA
Late_RecrDev_2019	-0.0641552	5	0.0000019	NA	NA
Late_RecrDev_2020	-0.0132446	5	-0.0000027	NA	NA
Late_RecrDev_2021	0.0443739	5	-0.0000015	NA	NA
Late_RecrDev_2022	0.0000000	5	0.0000000	NA	NA
ForeRecr_2023	0.0000000	5	0.0000000	NA	NA
ForeRecr_2024	0.0000000	5	0.0000000	NA	NA
ForeRecr_2025	0.0000000	5	0.0000000	NA	NA
ForeRecr_2026	0.0000000	5	0.0000000	NA	NA
ForeRecr_2027	0.0000000	5	0.0000000	NA	NA
ForeRecr_2028	0.0000000	5	0.0000000	NA	NA
ForeRecr_2029	0.0000000	5	0.0000000	NA	NA
ForeRecr_2030	0.0000000	5	0.0000000	NA	NA
ForeRecr_2031	0.0000000	5	0.0000000	NA	NA
ForeRecr_2032	0.0000000	5	0.0000000	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
ForeRecr_2033	0.0000000	5	0.0000000	NA	NA
ForeRecr_2034	0.0000000	5	0.0000000	NA	NA
LnQ_base_Triennial1(4)	-0.4725640	-1	NA	NA	NA
Q_extraSD_Triennial1(4)	0.0387429	4	0.0000001	NA	NA
LnQ_base_NWFSCcombo(6)	0.1823020	-1	NA	NA	NA
Size_DblN_peak_Trawl_N(1)	27.8920000	1	0.0001238	NA	NA
Size_DblN_top_logit_Trawl_N(1)	-15.0000000	-3	NA	NA	NA
Size_DblN_ascend_se_Trawl_N(1)	4.6011700	3	-0.0001243	NA	NA
Size_DblN_descend_se_Trawl_N(1)	6.9088400	4	0.0001059	NA	NA
Size_DblN_start_logit_Trawl_N(1)	-	-99	NA	NA	NA
	999.0000000				
∞ Size_DblN_end_logit_Trawl_N(1)	-	-99	NA	NA	NA
	999.0000000				
Retain_L_infl_Trawl_N(1)	29.0502000	3	-0.0002307	NA	NA
Retain_L_width_Trawl_N(1)	2.7569300	3	-0.0000938	NA	NA
Retain_L_asymptote_logit_Trawl_N(1)	9.9999800	3	0.0000000	NA	NA
Retain_L_maleoffset_Trawl_N(1)	0.0000000	-4	NA	NA	NA
Size_DblN_peak_Trawl_S(2)	29.4637000	1	0.0000586	NA	NA
Size_DblN_top_logit_Trawl_S(2)	-0.4208990	3	0.0000671	NA	NA
Size_DblN_ascend_se_Trawl_S(2)	3.9103400	3	-0.0000537	NA	NA
Size_DblN_descend_se_Trawl_S(2)	4.7143800	4	0.0000235	NA	NA
Size_DblN_start_logit_Trawl_S(2)	-	-99	NA	NA	NA
	999.0000000				
Size_DblN_end_logit_Trawl_S(2)	-	-99	NA	NA	NA
	999.0000000				
Retain_L_infl_Trawl_S(2)	25.9045000	3	-0.0000512	NA	NA
Retain_L_width_Trawl_S(2)	2.7373100	3	-0.0000433	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Retain_L_asymptote_logit_Trawl_S(2)	10.0000000	-3	NA	NA	NA
Retain_L_maleoffset_Trawl_S(2)	0.0000000	-4	NA	NA	NA
Size_DblN_peak_Non-trawl(3)	45.3712000	2	-0.0000012	NA	NA
Size_DblN_top_logit_Non-trawl(3)	-3.3853700	3	-0.0000026	NA	NA
Size_DblN_ascend_se_Non-trawl(3)	5.1284900	3	-0.0000006	NA	NA
Size_DblN_descend_se_Non-trawl(3)	4.7080400	4	-0.0000042	NA	NA
Size_DblN_start_logit_Non-trawl(3)	-	-99	NA	NA	NA
	999.0000000				
Size_DblN_end_logit_Non-trawl(3)	-	-99	NA	NA	NA
	999.0000000				
Retain_L_infl_Non-trawl(3)	25.2560000	3	0.0000003	NA	NA
Retain_L_width_Non-trawl(3)	2.7244000	3	-0.0000002	NA	NA
Retain_L_asymptote_logit_Non-trawl(3)	1.4966000	3	0.0000018	NA	NA
Retain_L_maleoffset_Non-trawl(3)	0.0000000	-4	NA	NA	NA
Size_DblN_peak_Triennial1(4)	26.7034000	2	0.0000003	NA	NA
Size_DblN_top_logit_Triennial1(4)	-7.0000000	-3	NA	NA	NA
Size_DblN_ascend_se_Triennial1(4)	4.1400700	3	0.0000018	NA	NA
Size_DblN_descend_se_Triennial1(4)	3.1583700	4	-0.0000002	NA	NA
Size_DblN_start_logit_Triennial1(4)	-	-99	NA	NA	NA
	999.0000000				
Size_DblN_end_logit_Triennial1(4)	-	-99	NA	NA	NA
	999.0000000				
SzSel_Male_Peak_Triennial1(4)	-1.9920600	3	-0.0000010	NA	NA
SzSel_Male_Ascend_Triennial1(4)	-0.0989113	3	0.0000043	NA	NA
SzSel_Male_Descend_Triennial1(4)	0.0000000	-3	NA	NA	NA
SzSel_Male_Final_Triennial1(4)	0.0000000	-3	NA	NA	NA
SzSel_Male_Scale_Triennial1(4)	1.0000000	-4	NA	NA	NA

**Table 17:** All parameter estimates for the proposed base model (*continued*)

Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Size_DblN_peak_Triennial2(5)	23.1266000	2	0.0000001	NA	NA
Size_DblN_top_logit_Triennial2(5)	-7.0000000	-3	NA	NA	NA
Size_DblN_ascend_se_Triennial2(5)	4.3565000	3	0.0000018	NA	NA
Size_DblN_descend_se_Triennial2(5)	4.1375700	4	0.0000007	NA	NA
Size_DblN_start_logit_Triennial2(5)	-	-99	NA	NA	NA
	999.0000000				
Size_DblN_end_logit_Triennial2(5)	-	-99	NA	NA	NA
	999.0000000				
SzSel_Male_Peak_Triennial2(5)	-1.7395500	3	-0.0000005	NA	NA
SzSel_Male_Ascend_Triennial2(5)	-0.2840780	3	0.0000007	NA	NA
SzSel_Male_Descend_Triennial2(5)	0.0000000	-3	NA	NA	NA
SzSel_Male_Final_Triennial2(5)	0.0000000	-3	NA	NA	NA
SzSel_Male_Scale_Triennial2(5)	1.0000000	-4	NA	NA	NA
Size_DblN_peak_NWFSCcombo(6)	29.9823000	2	0.0000138	NA	NA
Size_DblN_top_logit_NWFSCcombo(6)	-0.8537680	3	0.0000046	NA	NA
Size_DblN_ascend_se_NWFSCcombo(6)	4.9374300	3	-0.0000211	NA	NA
Size_DblN_descend_se_NWFSCcombo(6)	4.5447800	4	0.0000023	NA	NA
Size_DblN_start_logit_NWFSCcombo(6)	-	-99	NA	NA	NA
	999.0000000				
Size_DblN_end_logit_NWFSCcombo(6)	-	-99	NA	NA	NA
	999.0000000				
SzSel_Male_Peak_NWFSCcombo(6)	-4.1440800	3	0.0000040	NA	NA
SzSel_Male_Ascend_NWFSCcombo(6)	-0.5434570	3	-0.0000282	NA	NA
SzSel_Male_Descend_NWFSCcombo(6)	0.0000000	-3	NA	NA	NA
SzSel_Male_Final_NWFSCcombo(6)	0.0000000	-3	NA	NA	NA
SzSel_Male_Scale_NWFSCcombo(6)	1.0000000	-4	NA	NA	NA
Retain_L_infl_Trawl_N(1)_BLK1delta_2007	-5.7314400	4	-0.0000607	-7.379180	-4.083690

**Table 17:** All parameter estimates for the proposed base model (*continued*)


Source	Value	Phase	Gradient	Lower 95% CI	Upper 95% CI
Retain_L_infl_Trawl_N(1)_BLK1delta_2011	-2.4994900	4	-0.0000675	-3.712360	-1.286620
Retain_L_asymptote_logit_Trawl_N(1)_BLK1delta_2007	0.0002155	4	0.0000000	-0.391742	0.392173
Retain_L_asymptote_logit_Trawl_N(1)_BLK1delta_2011	0.0001030	4	-0.0000002	-0.391877	0.392083
Retain_L_infl_Trawl_S(2)_BLK2delta_2007	-5.2047800	4	-0.0000120	-7.293730	-3.115830
Retain_L_infl_Trawl_S(2)_BLK2delta_2011	-3.4190500	4	-0.0000137	-4.345530	-2.492580
Retain_L_asymptote_logit_Trawl_S(2)_BLK2delta_2007	0.0017796	4	0.0000004	-0.389867	0.393426
Retain_L_asymptote_logit_Trawl_S(2)_BLK2delta_2011	0.0015285	4	0.0000000	-0.390168	0.393225

**Table 18:** Estimated recent trend in recruitment and recruitment deviations and the 95% confidence intervals (CI) from the base model.


Year	Recruitment	Interval	Recruit- ment.Devia- tions	Interval.1
2013	6,024	2,469–14,698	-0.439	-1.352–0.474
2014	5,962	2,446–14,532	-0.447	-1.358–0.465
2015	5,954	2,438–14,542	-0.446	-1.360–0.468
2016	6,057	2,465–14,886	-0.427	-1.349–0.495
2017	5,836	2,385–14,279	-0.462	-1.379–0.454
2018	5,745	2,346–14,069	-0.476	-1.393–0.442
2019	8,863	3,557–22,086	-0.064	-1.003–0.874
2020	9,536	3,760–24,183	-0.013	-0.973–0.946
2021	10,335	3,984–26,811	0.044	-0.943–1.032
2022	10,118	3,924–26,090	0.000	-0.980–0.980
2023	10,117	3,924–26,086	0.000	-0.980–0.980

## **8 Figures**


### **8.1 Introduction and Data**




**Figure 1:** Biomass of shortspine thornyhead found in the NWFSC West Coast Groundfish Bottom Trawl Survey annual survey (2003-2022) coastwide.




**Figure 2:** Unidentified thornyhead catches (mt) and the proportion identified as shortspines, calculated as the ratio of shortspine thornyhead catches to combined longspine and shortspine catches.




**Figure 3:** Estimated landing history for shortspine thornyhead.




**Figure 4:** Summary of data sources used in the base model.




**Figure 5:** Predicted discards based estimated retention and selectivity for each fleet.




**Figure 6:** Discard rates from the WCGOP data set with catch share and non-catch share considerations from the GEMM dataset.




**Figure 7:** Summary of survey data sources used in the base model.




**Figure 8:** Survey length composition data.




**Figure 9:** Growth curve comparison. .



**Figure 10:** Growth curve sensitivities.




**Figure 11:** Fit of the maturity curves per size and depth classes. Classes are designed for visual check of the model predictions only since the model assumes continuous and not categorical response to these variables.




**Figure 12:** Maturity curves considered in the present assessment (Head (2023)) and alternative versions considered in the sensitivity analysis.


## 8.2 Base Model Results and Fits




**Figure 13:** Annual recruitment deviations with 95% intervals.




**Figure 14:** Selectivity at length for multiple fleets.




**Figure 15:** Fit to index data for Triennial1. Lines indicate 95% uncertainty interval around index values based on the model assumption of lognormal error. Thicker lines (if present) indicate input uncertainty before addition of estimated additional uncertainty parameter.




**Figure 16:** Fit to index data for NWFSCcombo. Lines indicate 95% uncertainty interval around index values based on the model assumption of lognormal error. Thicker lines (if present) indicate input uncertainty before addition of estimated additional uncertainty parameter.




**Figure 17:** Discard fraction for the North Trawl fleet.




**Figure 18:** Discard fraction for the South Trawl fleet.




**Figure 19:** Discard fraction for the South Trawl fleet.




**Figure 20:** Length comps, aggregated across time by fleet. Labels ‘retained’ and ‘discard’ indicate discarded or retained sampled for each fleet. Panels without this designation represent the whole catch.




**Figure 21:** The most recent ten years of length comps and fit for North Trawl retained catch. ‘N adj.’ is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




**Figure 22:** Length comps, whole catch, for the WCGBTS. ‘N adj.’ is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Ianelli tuning method.




**Figure 23:** Pearson residuals, whole catch, for the early Triennial survey (max=1.04). Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




**Figure 24:** Pearson residuals, whole catch, for the late Triennial survey (max=1.34). Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




**Figure 25:** Pearson residuals, whole catch, for the WCGBTS (max=1.78). Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




**Figure 26:** Mean weight in discard for the North Trawl fleet.




**Figure 27:** Mean weight in discard for the South Trawl fleet.




**Figure 28:** Mean weight in discard for the Non-trawl fleet.




**Figure 29:** Spawning output with ~95% asymptotic intervals.



**Figure 30:** Relative spawning output:  $B/B_0$  with ~95% asymptotic intervals.




**Figure 31:** Forecasted spawning output with ~95% asymptotic intervals.




**Figure 32:** Forecasted spawning output:  $B/B_0$  with  $\sim 95\%$  asymptotic intervals.


### 8.3 Likelihood Profiles, Retrospectives, and Sensitivity Analyses




**Figure 33:** Piner panel plot showing the impact of changing  $R_0$  on the overall (top), length composition (middle), and survy (bottom) likelihoods.




**Figure 34:** High to low values of  $R_0$  and impact on spawning output.




**Figure 35:** Piner panel plot showing the impact of changing natural mortality on the overall (top), length composition (middle), and survey (bottom) likelihoods.




**Figure 36:** High to low values of M and impact on relative spawning output.




**Figure 37:** High to low values of  $M$  and impact on spawning output.




**Figure 38:** Likelihood profile for steepness showing the impact of the steepness value on the change in log likelihood (top-left), the fraction of unfished biomass (top-right), initial spawning biomass (bottom-left) and the terminal year of spawning biomass (bottom-right).




**Figure 39:** Piner panel plot showing the impact of changing  $h$  on the overall (top), length composition (middle), and survy (bottom) likeihoods.




**Figure 40:** High to low values of  $h$  and impact on relative spawning output.




**Figure 41:** Piner panel plot showing the impact of low to high  $a_{\{max\}}$  assumptions on the overall (top), length composition (middle), and survy (bottom) likehoods.




**Figure 42:** Low to high values of  $a_{\max}$  and impact on spawning output.




**Figure 43:** Retrospectives showing impact on spawning output.




**Figure 44:** Retrospectives showing impact on stock status.




**Figure 45:** Retrospectives showing impact on fits to WCGBTS index.




**Figure 46:** Retrospectives showing impact on spawning output.




**Figure 47:** Spawning output comparisons of the base model and high growth and low growth assumptions.




**Figure 48:** Relative spawning output comparisons of the base model and high growth and low growth assumptions.




**Figure 49:** Comparison of fits to combo survey data between the base model and high growth and low growth sensitivities.




**Figure 50:** Spawning output comparisons of the base model and maturity sensitivities.



**Figure 51:** Relative spawning output comparisons of the base model and maturity sensitivities.



**Figure 52:** Spawning output comparisons of the base model and landing sensitivities.



**Figure 53:** Relative spawning output comparisons of the base model and landing sensitivities.