


Using Multiple Sources to Solve Problems

Sometimes information from more than one source of data is needed to answer a question. A table, graph, or text may contain only part of the information you need.

Suppose Greg plans to drive from Los Angeles to San Francisco, but he needs to rent a car. Greg knows that gasoline costs about \$3.70 per gallon. What would be the cost of driving the various rental cars based on distance, gas mileage, and gas prices?

Both the chart and graph below contain information that can be used to answer the question. The mileage chart shows the distance between the two cities. The bar graph shows the number of miles per gallon each type of rental car gets.

	Atl	Bos	Chi	Dal	LA	NO	SF
Atl		1,104	720	781	2,217	467	2,595
Bos	1,104		985	1,770	2,984	1,525	3,098
Chi	720	985		968	2,015	925	2,130
Dal	781	1,770	968		1,435	520	1,814
LA	2,217	2,984	2,015	1,435		1,895	380
NO	467	1,525	925	520	1,895		2,275
SF	2,595	3,098	2,130	1,814	380	2,275	



Example

Hide

How much will it cost Greg to drive from Los Angeles to San Francisco if he rents a subcompact car and the price of gasoline is \$3.70 per gallon?

Step 1

Gather the information.

The total distance is 380 miles. A subcompact car gets 28 mpg.

Step 2

Find the number of gallons of gas it will take to get to San Francisco.

$$380 \div 28 \approx 13.6 \text{ gallons}$$

Step 3

Find the total cost of the gasoline.

$$13.6 \times \$3.70 = \$50.32$$

It will cost **\$50.32** in gas for Greg to drive a subcompact from Los Angeles to San Francisco.

Rate Your Confidence

High

Medium

Low

Continue >>

Category

Lesson 2: Interpreting Graphs

Related Questions:

[Practice questions taken related to this topic](#)