

Sex emerges at scale

Philip J Gerrish^{a,b,c,1}

This manuscript was compiled on October 2, 2026

Sex is costly yet ubiquitous. This evolutionary paradox remains largely unresolved despite decades of intense scrutiny. Theoretical endeavors to resolve the paradox have looked to non-additive contingencies, like epistasis, as primary drivers of sex evolution. Models proliferated accordingly: one for each contingency. Contingency dependance and model proliferation are symptoms of a complex system, which can be governed by completely different rule sets at different scales. We find that re-scaling brings clarity to the problem of sex, and the collective potential that survives re-scaling is additive fitness. We find that the additive advantage of sex is non-negative singular and heavy tailed, easily overcoming large canonical costs of sex, without requiring epistasis.

Evolution of sex | Condensed matter physics | Macro-evolution | Renormalization | Additive fitness

The absence of sex (hereafter *asex*) has several well-studied evolutionary costs that fall naturally into two categories: *within*- and *across*-lineage selective interference. For clarity, we introduce the following notation: the genetic background (or wildtype allele) is denoted $\bullet$ if the lineage is growing and $\circ$ if it has constant size; $+$ denotes advantageous allele, and $-$ denotes deleterious allele. Genotypes have two loci and will be represented by two of these symbols.

Within a growing lineage of genotype $(\bullet, \bullet)$, the efficacy of natural selection is reduced, with the following consequences: 1) decreased fixation probabilities $\mathcal{P}$ of newly-arising beneficial mutants: $\mathcal{P}(\bullet, +) < \mathcal{P}(\circ, +)$, 2) increased fixation probabilities of newly-arising deleterious mutations $\mathcal{P}(\bullet, -) > \mathcal{P}(\circ, -)$, and 3) increased rate $\mathcal{R}$ of irreversible accumulation of deleterious mutations $\mathcal{R}[(\bullet, \bullet) \mapsto (\bullet, -) \mapsto (-, -)] > \mathcal{R}[(\circ, \circ) \mapsto (\circ, -) \mapsto (-, -)]$. Consequences 1 and 2 constitute Hill-Robertson Interference (HRI) after the authors of its first quantitative treatment (1–5); Consequence 3 has been dubbed “lineage contamination” and studied quantitatively in (6) and (7–11).

Across growing lineages, the advantageous traits causing one lineage to grow can be lost if a second lineage appears (not a sublineage of the first) with superior traits. This can happen when: 1) a beneficial mutation arises on a deleterious genetic background, creating competing genotypes $\{(\circ, \circ), (-, +)\}$ or $\{(\bullet, \bullet), (-, +)\}$, 2) one lineage carries a beneficial mutation in one locus and a second lineage carries a beneficial mutation in the other locus, creating competing genotypes: $\{(+, \circ), (\circ, +)\}$ or $\{(+, \bullet), (\bullet, +)\}$. Scenario 1 was envisioned by Fisher, and scenario 2 by Muller; together, they are referred to as Fisher-Muller Interference (FMI). Scenario 2 by itself is often referred to as “clonal interference” (12–17).

Both within- and across-lineage interference are examples of “selection interference” (18) – inefficiencies that arises in the absence of sex. Efficiency in evolution that is lost to selection interference can be regained by breaking the genetic linkage causing it. And linkage breaking is a consequence of sex, defined here as any form of genetic exchange.

The costs of *asex*, while many, are nevertheless thought to be incremental and subtle when compared to the large and obvious costs of sex. Sexually-produced offspring carry only half of the mother’s genes (genome dilution). Sex requires finding a mate in order to reproduce, which makes prospective parents vulnerable to predation and disease. And when only females bear offspring, males are numerically superfluous and impose a canonical two-fold demographic cost (19–23). The cost asymmetry between *asex* and sex constitutes a fitness deficit that clearly favors *asex*.

Significance

On a fundamental level, the evolution of sex is like the familiar game of chutes and ladders. The players are sex-enabling genes and they advance on the board by jumping from one genome to the next; one square to the next. Sometimes the genome they land on benefits from the sex they enable and they get a free ride to higher frequencies; they climb a ladder. And sometimes the genome they land on is harmed by the sex they enable and they are taken to lower frequencies; they slide down a chute. The present study finds that extremely tall ladders can be encountered with non-negligible probability while extremely long chutes are astronomically rare. This asymmetry results in a universal tendency for evolution to favor sex – a prediction that is spectacularly corroborated by nature. Previous studies used a measure of height (total fitness) that hid the really tall ladders, thereby obscuring the asymmetry.

Author affiliations: ^aDepartment of Mathematics, Universidad Autónoma del Estado de Hidalgo, México; ^bDepartment of Biology, University of New Mexico, USA; ^cTheoretical Division, Los Alamos National Labs, USA

The author declares no competing interests.

¹To whom correspondence should be addressed. E-mail: pgerrishunm.edu

The evolutionary advantage of sex is quantified naturally by a dynamical measure proposed by Mustonen and Lässig (24, 25) called “fitness flux”, which can accommodate both fitness landscapes (static topologies) and fitness seascapes (dynamic topologies). We define *sex potential* as the potential fitness flux created by an interference trajectory, or “path” (HRI or FMI). Here we depart from precedent: we concern ourselves not with contingent individual paths but with the collection, or “ensemble”, of all possible paths and the usual probability space induced. Sex potential is therefore a random variable; it is distributed as the *path-ensemble* value of potential fitness flux favoring sex. This quantity reduces to the change in log-odds of a sex^+ modifier frequency.

We consider the selective sorting of n haploid genomes, each of which has two loci, x and y . Here, we depart from precedent by defining genomes as random vectors of genic fitnesses, (X_i, Y_i) , $i = 1, 2, \dots, n$, whose elements need not be independent or identically distributed across loci. (Adding more loci does not change things qualitatively.) We note that this agnostic setting implicitly absorbs all forms of selective interference. Initial genome frequencies can be fixed or random. We derive the average selective advantage of a sex-enabling gene that attaches to a randomly-chosen genome at random time $T \in [0, \infty)$ and subsequently dissociates from that genome with rate $r \in [0, \infty)$.

Mean fitness $L(t)$ may be obtained by partitioning the Fisher-Ewens fundamental theorem of natural selection (FTNS) (26–30) in integrated form as follows:

$$L(t) = T(t) - V(t),$$

where $T(t) = \int_0^t \sigma_X^2(u) + \sigma_Y^2(u) du$ and $V(t) = -2 \int_0^t \sigma_{XY}(u) du$ represent cumulative kinetic- and potential-fitness flux, respectively (24, 25, 31). Under free genetic exchange, $V(t) = 0$ and mean fitness evolves as $L^*(t) = T(t)$. The cumulative fitness flux to be gained from sex at time t is therefore $L^*(t) - L(t) = V(t)$.

Mathematically, *sex potential* is defined as: $V^* := \lim_{t \rightarrow \infty} V(t)$. It reflects total selection for sex-enabling genes that are highly promiscuous, in that they jump from genome to genome through the very recombination they enable. Such genes do not track autosomal (non-sex-related) genes but instead evolve independently in a kind of genetic symbiosis. The fate of sex-enabling genes in this context is therefore determined primarily by collective fitness imbalance among autosomal genes.

Not all sex-enabling genes are highly promiscuous, however, with many remaining *linked* for a number of generations to the shuffled autosomal genomes they helped generate. During this time, their frequencies track the autosomal genomes to which they are linked. The result is a process of serial hitchhiking that determines their fate. To probe the full range of promiscuity, we expand our definition of *sex potential* to V_r^* , where r is the dissociation rate of sex-enabling genes. Extremes $r = \infty$ and $r = 0$ correspond to fully promiscuous (no linkage) and fully linked (no promiscuity).

Sex Emergence Theorems:

Under trivial nondegeneracy requirements detailed in the SI, the three fundamental theorems governing the emergence of sex at scale are:

1 (sign): $\mathbb{E}[V_r^*] \geq 0$ ($+\infty$) for $r \geq 0$ (extended expectation)

2 (surges): $\Pr(V_r^* > v) \propto \begin{cases} v^{-1} & r > 0 \\ v^{-1/2} & r = 0 \end{cases}$

3 (reach): $\Pr(T_0 > t) \propto t^{-1}$ and $\Pr(X_0 > x) \propto x^{-2}$

All three results arise near $U = \Delta_X + \Delta_Y = 0$: two genomes can be nearly tied in total fitness while carrying opposing effects at the two loci. Such paths are sorted slowly, so their positive contribution to sex potential accumulates. Complete linkage gives the heaviest surges; any positive dissociation rate ultimately changes the tail to v^{-1} , with a crossover of order r^{-2} .

The final theorem concerns the duration and spatial reach of individual episodes. For $r > 0$, dissociation cuts them off at $1/r$ and $\sqrt{D/r}$, respectively. At $r = 0$ the mean duration and the second moment of spatial reach diverge,

although mean reach remains finite. These laws do not specify the spacing between separate surge onsets.

The true novelty of the present study lies primarily in the rigorous proof of this theorem (Methods), because it clearly shows why the central claim is non-parametric and does not fall short of universality. There are also two heuristic arguments, however, that cannot get us all the way to universality but that can help to understand why the claim should be true most of the time. The first is a verbal line of reasoning from Otto and Barton (18, 21, 32, 33): they suggest that selectively well-matched alleles ($\sigma_{XY} > 0$) should be rapidly fixed or eliminated, while selectively mismatched alleles ($\sigma_{XY} < 0$) should tend to have longer sojourn times, thereby contributing more to statistical associations. The second heuristic argument is mathematical. Letting random variables ΔX and ΔY denote genic fitness differences at two loci, we have $V^* \propto -\Delta X \Delta Y / |\Delta X + \Delta Y|$. V^* blows up as the denominator approaches zero, which happens as $\Delta Y \rightarrow -\Delta X$, or as $\Delta X \Delta Y \rightarrow -\Delta X^2 \leq 0$. This creates an obvious bias towards positive values of V^* .

Remarkably, sex can initially be strongly deleterious on average ($\langle \sigma_{XY}(0) \rangle \gg 0$) and yet sex potential remains non-negative in expectation ($\langle V_r^* \rangle \geq 0$). This observation underscores the inadequacy of local measures – such as linkage disequilibrium – for predicting the evolutionary fate of sex at scale.

Taken together, these theorems portray a general pattern governing the evolution of sex at scale: weak but non-negative selection for sex persists in expectation (Thm 1); occasionally, strong surges of selection for sex occur (Thm 2); the strongest of these surges favors linked sex-enforcing genes ($1/r$ dependence in Thm 2). Notably, selection surges most likely derive not from large differences at two loci, but from the collective effect of smaller fitness differences across more loci (SI).

The currency of our analyses is additive fitness (28, 34–36). This fact adds to the novelty of the present study by its orthogonality to the non-additive effects, such as epistasis and dominance, often considered essential for sex to be successful on micro-evolutionary scales. Furthermore, additive fitness emerges as the component that survives renormalization and drives the evolution of sex at scale.

To illustrate, we let w_{ij} denote fitness of haploid genome (i/j), and we set: $w_{11} = w_{22} = 1$ and $w_{12} = w_{21} = -1$. We consider two extreme scenarios: A) If the population initially consists solely of genotypes (1/1) and (2/2), sex would be detrimental with coefficient of selection -1. B) On the other hand, if the population initially consists solely of genotypes (1/2) and (2/1), sex would be advantageous with coefficient of selection +1. In both cases, there is strong selection on sex genes on micro-evolutionary scales, and yet additive fitness is zero, indicating no net selection at scale. If A were more likely than B, sex would be generally detrimental at scale, and it would indicate that additive fitness does not tell the whole story, i.e., that non-additive fitness is still relevant at scale. The large-scale systemic biases that this would require, however, lack *a priori* rationale (37) and are not supported by the data (38–40). In the language of renormalization, non-additive effects fall naturally into the “irrelevant” bin at scale.

We note that, in the context of organisms that already reproduce sexually, a second interpretation of *ensemble* is possible. Genes or linkage regions in these organisms can evolve independently, and each can experience its own HRI path; a single evolving population of such organisms can thus be considered an ensemble. Sex potential in this context can represent total additive fitness to be gained from increased within-locus recombination in a single population. In this light, it is a measure of selection pressure to maintain sex and potentially reduce gene size (41).

Similar reasoning can be found in the original Hill and Robertson paper (1). The authors present it cautiously, however, because they appear to have doubts about their key assumption which, on the surface, does seem to strain credulity. Namely, they assume that the different HRI paths across loci must have statistical properties that are somehow similar enough to be considered part of the same ensemble. They even invoke rampant gene duplication as a potential scenario that would satisfy their assumption of groupable HRI paths. Had they known that comparable statistical properties are not required, as our results show, they might have been less cautious.

In light of our findings, the observed ubiquity of sex in nature may be interpreted as yet another challenge to assumptions of equilibrium in evolution (42–44). Previous theory that assumes micro-evolutionary equilibrium and a specific kind of fitness mutation finds a “general reduction principle” predicting *ever*-decreasing rates of sex and recombination (45–47). We find that eliminating the equilibrium requirement at scale reverses the prediction to *never*-decreasing rates of sex and recombination.

When presented with a new theory purporting to explain the evolution and persistence of sex, often the first question that comes to mind is: does it account for and overcome the canonical two-fold cost of sex [REFS]? For many reasons the question itself is problematic and is often dismissed [REFS]. Nevertheless, our answer is clear: the very heavy tail of sex potential means that the two-fold cost (or 3- or 4-fold cost for that matter) is easily overcome repeatedly on evolutionary timescales.

Our theory does not replace existing explanations of sex; it reveals their common large-scale structure. Microevolutionary mechanisms may differ, but the power laws and divergent correlation lengths derived here expose a shared critical organization at scale. A useful analogy is melting: individual molecular trajectories alone do not explain the phase transition. Evolutionary stasis is the frozen state; in the critical regime identified by our power laws, sex^- clusters fragment and sex^+ clusters emerge with scale-free sizes and lifetimes. This boundary regime is therefore especially favourable to the evolution of sex—and perhaps to life itself.

Sex has long seemed exceptional because its causes were sought one mechanism at a time. At scale, those mechanisms resolve into a common law: selection for sex is non-negative on average, punctuated by heavy-tailed surges, and divergent correlation lengths across space and time. The resulting criticality is not assumed but derived, recasting the ubiquity of sex as a universal large-scale pattern rather than the sum of special cases.

Methods

Heuristic proofs for the sex emergence theorems. Let $U = \Delta_X + \Delta_Y$ be the total-fitness difference and $W = \Delta_X$ the difference at one locus. Assume a smooth, nonzero path density near $U = 0$ with $W \neq 0$, and the relevant finite moments; for the spatial result, assume local diffusion. Exact conditions and proofs are given in the SI.

1 (sign). Sex potential is a positive persistence kernel times $W^2 - UW$. Near a total-fitness tie this numerator approaches $W^2 > 0$, whereas its negative part is bounded by $U^2/4$. The kernel diverges as $1/|U|$ at fixed positive dissociation rate and as $1/U^2$ under complete linkage. The positive mean therefore diverges; the negative mean remains finite. Thus the extended expectation of sex potential is $+\infty$.

2 (surges). At fixed nonzero W , a potential above v requires a total-fitness gap of order v^{-1} when dissociation is positive, or $v^{-1/2}$ under complete linkage. Because path probability near a tie is proportional to the width of that gap, the upper tails have the same respective exponents. For small positive dissociation, the crossover occurs at a characteristic potential of order r^{-2} .

3 (persistence and reach). Selection resolves a fitness gap $S = |U|$ at a rate proportional to S ; modifier dissociation adds rate r . Persistence time is therefore inversely proportional to their combined rate, and diffusive reach is proportional to the square root of that time. Since the probability of a gap below ε is proportional to ε , complete linkage yields a t^{-1} tail for duration and an x^{-2} tail for spatial reach. Positive dissociation truncates both. These are episode extents, not intervals between surge onsets.

Supplementary materials

Materials and Methods

Supplementary Text

1. WG Hill, A Robertson, The effect of linkage on limits to artificial selection. *Genet. Res.* **8**, 269–294 (1966).
2. AOB Whitlock, KM Peck, RBR Azevedo, CL Burch, An Evolving Genetic Architecture Interacts with Hill-Robertson Interference to Determine the Benefit of Sex. *Genetics* **203**, 923–936 (2016).
3. IM Rouzine, Long-range Hill-Robertson effect in adapting populations with recombination and standing variation (bioRxiv preprint) (2022).

4. JM Comeron, A Williford, RM Kliman, The Hill-Robertson effect: evolutionary consequences of weak selection and linkage in finite populations. *Hered. (Edinb.)* **100**, 19–31 (2008).

5. D Roze, NH Barton, The Hill-Robertson effect and the evolution of recombination. *Genetics* **173**, 1793–1811 (2006).

6. T Johnson, NH Barton, The effect of deleterious alleles on adaptation in asexual populations. *Genetics* **162**, 395–411 (2002).

7. S Pénilson, PD Sniegowski, A Colato, PJ Gerrish, Lineage dynamics and mutation–selection balance in non-adapting asexual populations. *J. Stat. Mech.* **2013**, P01013 (2013).

8. S Pénilson, T Singh, P Sniegowski, P Gerrish, Dynamics and Fate of Beneficial Mutations Under Lineage Contamination by Linked Deleterious Mutations. *Genetics* **205**, 1305–1318 (2017).

9. A Colato, JF Fontanari, Soluble model for the accumulation of mutations in asexual populations. *Phys. Rev. Lett.* **87**, 238102 (2001).

10. JF Fontanari, A Colato, RS Howard, Mutation accumulation in growing asexual lineages. *Phys. Rev. Lett.* **91**, 218101 (2003).

11. PJ Gerrish, A Colato, PD Sniegowski, Genomic mutation rates that neutralize adaptive evolution and natural selection. *J. R. Soc. Interface* **10**, 20130329 (2013).

12. R Miralles, PJ Gerrish, A Moya, SF Elena, Clonal interference and the evolution of RNA viruses. *Science* **285**, 1745–1747 (1999).

13. JAGM de Visser, CW Zeyl, PJ Gerrish, JL Blanchard, RE Lenski, Diminishing returns from mutation supply rate in asexual populations. *Science* **283**, 404–406 (1999).

14. DE Rozen, JAGM de Visser, PJ Gerrish, Fitness effects of fixed beneficial mutations in microbial populations. *Curr. Biol.* **12**, 1040–1045 (2002).

15. PJ Gerrish, A Colato, AS Perelson, PD Sniegowski, Complete genetic linkage can subvert natural selection. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 6266–6271 (2007).

16. PJ Gerrish, RE Lenski, The fate of competing beneficial mutations in an asexual population. *Genetica* **102-103**, 127–144 (1998).

17. P Gerrish, The rhythm of microbial adaptation. *Nature* **413**, 299–302 (2001).

18. SP Otto, Selective Interference and the Evolution of Sex. *J. Hered.* **112**, 9–18 (2021).

19. J Lehtonen, MD Jennions, H Kokko, The many costs of sex. *Trends Ecol. Evol.* **27**, 172–178 (2012).

20. D Roze, Disentangling the benefits of sex. *PLoS Biol.* **10**, e1001321 (2012).

21. SP Otto, The evolutionary enigma of sex. *Am. Nat.* **174 Suppl 1**, S1–S14 (2009).

22. SP Otto, T Lenormand, Resolving the paradox of sex and recombination. *Nat. Rev. Genet.* **3**, 252–261 (2002).

23. NH Barton, B Charlesworth, Why sex and recombination? *Science* **281**, 1986–1990 (1998).

24. V Mustonen, M Lässig, From fitness landscapes to seascapes: non-equilibrium dynamics of selection and adaptation. *Trends Genet.* **25**, 111–119 (2009).

25. V Mustonen, M Lässig, Fitness flux and ubiquity of adaptive evolution. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 4248–4253 (2010).

26. RA Fisher, *The genetical theory of natural selection*. (Oxford University Press, Oxford), p. 302 (1930).

27. GR Price, Fisher's 'fundamental theorem' made clear. *Ann. Hum. Genet.* **36**, 129–140 (1972).

28. WJ Ewens, Mean fitness increases when fitnesses are additive. *Nature* **221**, 1076 (1969).

29. WJ Ewens, An interpretation and proof of the Fundamental Theorem of Natural Selection. *Theor. Popul. Biol.* **36**, 167–180 (1989).

30. WJ Ewens, The fundamental theorem of natural selection: the end of a story. *Evolution* **78**, 803–808 (2024).

31. V Mustonen, M Lässig, Molecular evolution under fitness fluctuations. *Phys. Rev. Lett.* **100**, 108101 (2008).

32. SP Otto, NH Barton, The evolution of recombination: removing the limits to natural selection. *Genetics* **147**, 879–906 (1997).

33. NH Barton, Genetic linkage and natural selection. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **365**, 2559–2569 (2010).

34. M Geeta Arun, A Angus-Henry, DJ Obbard, JD Hadfield, Estimating the additive genetic variance for relative fitness from changes in allele frequency. *Genetics* **232**, iyaf232 (2026).

35. AWF Edwards, The fundamental theorem of natural selection. *Theor. Popul. Biol.* **61**, 335–337 (2002).

36. FA Kondrashov, AS Kondrashov, Multidimensional epistasis and the disadvantage of sex. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 12089–12092 (2001).

37. PC Phillips, Epistasis—the essential role of gene interactions in the structure and evolution of genetic systems. *Nat. Rev. Genet.* **9**, 855–867 (2008).

38. DM Weinreich, NF Delaney, MA Depristo, DL Hartl, Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114 (2006).

39. J Franke, A Klözer, JAGM de Visser, J Krug, Evolutionary accessibility of mutational pathways. *PLoS Comput. Biol.* **7**, e1002134 (2011).

40. J Otwinowski, I Nemenman, Genotype to phenotype mapping and the fitness landscape of the *E. coli* lac promoter. *PLoS One* **8**, e61570 (2013).

41. Lenski, A distinction between the origin and maintenance of sex. *J. Evol. Biol.* **12**, 1034–1035 (1999).

42. S Song, P Chen, X Shen, J Zhang, Adaptive tracking with antagonistic pleiotropy results in seemingly neutral molecular evolution. *Nat. Ecol. Evol.* **9**, 2358–2373 (2025).

43. L Perfeito, L Fernandes, C Mota, I Gordo, Adaptive mutations in bacteria: high rate and small effects. *Science* **317**, 813–815 (2007).

44. T Held, D Klemmer, M Lässig, Survival of the simplest in microbial evolution. *Nat. Commun.* **10**, 2472 (2019).

45. MW Feldman, U Liberman, An evolutionary reduction principle for genetic modifiers. *Proc. Natl. Acad. Sci.* **83**, 4824–4827 (1986).

46. L Altenberg, U Liberman, MW Feldman, Unified reduction principle for the evolution of mutation, migration, and recombination. *Proc. Natl. Acad. Sci.* **114**, E2392–E2400 (2017).

47. U Liberman, MW Feldman, A general reduction principle for genetic modifiers of recombination. *Theor. Popul. Biol.* **30**, 341–371 (1986).

571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634




Fig. 1. Enter Caption