The Seven Wonders of the World Exercises

Updated 11 February 2025 10:27

Contents

1	Physics, quantities, units	5
	1.0	5
	1.1	5
	1.2	5
	1.3	6
	1.4	6
	1.5	7
	1.6	7
	Example solutions	10
2	Time and space	13
	2.0	13
	2.1	13
	2.2	14
	2.3	15
	2.4	15
	2.5	16
	2.6	16
	2.7	16
	Example solutions	18
	URLs for chapter 2	21
3	Main physical quantities	22
	3.0	22
	3.1	22
	3.2	22
	3.3	23
	3.4	23
	3.5	23

	3.6	23
	3.7	24
	Example solutions	25
	URLs for chapter 3	27
4	37.1	20
4	Volume contents, fluxes, supplies	28
	4.0	28
	4.1	28
	4.2	29
	4.3	31
	4.4	31
	4.5	31
	4.6	32
	4.7	33
	4.8	33
	4.9	34
	4.10	34
	4.11	34
	4.12	35
	Example solutions	37
	URLs for chapter 4	45
	CRES for chapter 4	10
5	Physical laws	46
	5.0	46
	5.1	46
	5.2	47
	5.3	47
	5.4	48
	5.5	49
	5.6	49
	Example solutions	50
	URLs for chapter 5	54
	OKLS for Chapter 3	9
6	The Seven Wonders of the world	55
	6.0	55
	Example solutions	56
7	Consequention & helenoop of mostless	
7	Conservation & balances of matter	57
	7.0	57 58
	Hyampia calutione	>

8	Conservation of electric charge	59	
	8.0	59	
	Example solutions	60	
9	Conservation of magnetic flux	61	
	9.0	61	
	Example solutions	62	
10	Balance of momentum	63	
	10.0	63	
	Example solutions	64	
11	Balance of energy	65	
	11.0	65	
	Example solutions	66	
12	Balance of angular momentum	67	
	12.0	67	
	Example solutions	68	
13	Remarks on momentum and energy	69	
	13.0	69	
	Example solutions	70	
14	Balance of entropy	71	
	14.0	71	
	Example solutions	72	
15	Constitutive relations	73	
	15.0	73	
	Example solutions	74	
Bibliography			

Physics, quantities, units 1

For some of the following exercises you can refer to tables 1.1 and 1.2 on page 9 (reproduced from the textbook).

1.0

(Do the exercises in the main text.)

1.1

Preferably together with a colleague:

If you have some large-language-model service (such as ChatGPT), ask it which physical laws are universally valid in Newtonian Mechanics and in General Relativity and in Thermodynamics and in Chemistry and in Electromagnetics.

Discuss the answer you get, based on what you have learned so far. (Note: if the answer mention a 'balance of boost momentum', that's actually correct.)

Argue with the LLM and see where the discussion goes.

1.2

Take *time* and *velocity* as primitive quantities.

- **1.** Try to define *distance* as a derived quantity
- **2.** Try to define *acceleration* as a derived quantity.

Which of the following quantities are scalars, and which are vectors?

- Time
- Distance
- Position
- Energy
- Velocity
- Speed
- Momentum
- Entropy
- Angular momentum
- Force
- Temperature
- Magnetic flux
- Electric charge
- Electric current
- Heat
- Power
- Volume
- Pressure

1.4

Find the correct units for the following quantities:

- Volumic energy or energy density, defined as energy divided by volume
- *Energy flux*, defined as energy divided by time.
- *Power*, defined as energy divided by time.
- *Heating*, defined as energy divided by time.
- *Magnetic flux*, which we take as a primitive quantity.
- *Electric potential difference*, defined as magnetic flux divided by time.
- Force, defined as momentum divided by time.
- *Momentum flux*, defined as momentum divided by time.
- *Momentum supply*, defined as momentum divided by time.
- Pressure, defined as force divided by area.
- *Amount of substance (or of matter)*, which we take as primitive.
- Molar mass, defined as mass divided by amount of substance.
- Specific momentum, defined as momentum divided by mass.

- Volumic charge or charge density, defined as charge divided by volume.
- *Entropy*, which we take as primitive, has dimension of energy divided by temperature.
- Matter density, defined as amount of substance divided by volume.
- *Matter flux*, defined as amount of substance divided by time.

With a friend or colleague:

- 1. Try to explain to your friend the difference between a *primitive quantity* and a *derived quantity*; then let your friend criticize unclear or incorrect points in your explanation, and comment on the good points. Then invert your roles: your friend tries to explain to you, and you criticize and comment.
- **2.** Similarly as the previous exercise, but explaining the difference between a *scalar quantity* and a *vector quantity*.
- **3.** If you have some large-language-model service (such as ChatGPT or DeepSeek), ask it to explain the difference between primitive and derived quantity, and between scalar and vector quantity. Find out weak or unsure points in its answer, given what you've learned so far.

1.6

Find which of the following mathematical expressions and equalities are dimensionally incorrect, and explain why they are incorrect:

- 11J + 4 kg
- $ightharpoonup \tan\left(\frac{a}{b}\right)$, where a has dimension length and b has dimension time
- ▶ 299 792 458 m/s
- $ho \exp\left(\frac{71 \,\mathrm{s}}{3 \,\mathrm{s}}\right)$
- ► cos(3.14) m
- \rightarrow m-v, where m has dimension of mass and v of velocity

- $ightharpoonup 10 \, N \, s 2 \, kg \, m/s = 8 \, J \, s/m$
- ▶ exp(-8 J)
- ▶ (9 m, 0.1 rad, -0.5 rad)
- > 8 J/s = 12 N m 4 N m
- ⊳ e⁻⁸ J
- $\frac{15 \text{ J}}{5 \text{ kg/s}^2} = 3 \text{ m}^2$
- $\sqrt{25} \, K = 5$
- ▶ $(e^7)^s$
- $\rightarrow \tan\left(\frac{10\,\mathrm{m}}{5\,\mathrm{m}}\right)$
- ▶ √300 K
- $ightharpoonup \sin(t/s)$, where t has dimension of time
- $\Rightarrow \frac{3}{8}$
- sin(10 s)

Quantity	SI Dimension	Unit
Time	time	second s
Length	length	metre m
Temperature	temperature	kelvin K
Matter Electric charge Magnetic flux	amount of substance electric charge magnetic flux	mole mol coulomb C weber Wb
Energy	energy, mass	joule J, kilogram kg
Momentum	force · time, mass · length/time, energy · time/length	$N \cdot s$, $kg \cdot m/s$, $J \cdot s/m$
Angular momentum	force · length · time, mass · length ² /time, energy · time	$N \cdot m \cdot s$, $kg \cdot m^2/s$, $J \cdot s$
Entropy	energy/temperature	J/K

Table 1.1 Dimensions and units of the main physical quantities used in these notes. Their fluxes have the dimensions divided by time, and therefore units divided by seconds. Quantities in **boldface** are vectors, the others are scalars.

Quantity	Volume content [unit]	Flux [unit]
matter	N [mol]	J [mol/s]
electric charge	Q [C]	I [C/s or A]
magnetic flux	${\mathcal B}$ [Wb]	\mathcal{E} [Wb/s or V]
energy	E [J]	Φ [J/s or W]
momentum	P [Ns]	F [N]
angular momentum	L [Nms]	τ [N m]
entropy	S [J/K]	П [J/(Кs)]

Table 1.2 Units for volume contents and fluxes of the main seven quantities.

Example solutions

② 1.2

- **1.** "Distance is the product of a time lapse and a particular velocity". See section 2.3 about *Radar distance* in our lecture notes.
- **2.** "Acceleration is the ratio between a change in the product of a time lapse and a particular velocity, and the time taken by that change".

♀ 1.3

These quantities are scalars:

- Time
- Distance
- Energy
- Speed
- Entropy
- Temperature
- Magnetic flux
- Electric charge
- Electric current
- Heat
- Power
- Volume

These quantities are vectors:

- Position
- Velocity
- Momentum
- Angular momentum
- Force

For *pressure*, it depends on the context. In some applications it is considered a scalar, but in other applications it is considered a vector – or actually a generalized kind of vector, called *tensor*, which can be represented by a matrix.

Ω 1.4

• Volumic energy: J/m³

1. Physics, quantities, units

• Energy flux: J/s

• Power: J/s

• Heating: J/s

• Magnetic flux: Wb

• Electric potential difference: Wb/s

• Force: N

• Momentum flux: N

• Momentum supply: N

• Pressure: N/m²

• Amount of substance: mol

• Molar mass: kg/mol

• *Specific momentum*: $N \cdot s/kg \equiv m/s$

• *Volumic charge*: C/m³

• Entropy: J/K

• *Matter density*: mol/m³

• *Matter flux*: mol/s

Ω 1.6

 \triangleright 11 J + 4 kg

Incorrect: cannot sum quantities of different dimension

▶ $\tan\left(\frac{a}{b}\right)$, where a dimension length and b has dimension time **Incorrect**: trigonometric function must have a dimensionless argument, but a/b has dimension length/time

▶ 299 792 458 m/s

$$ightharpoonup \exp\left(\frac{71 \, \mathrm{s}}{3 \, \mathrm{s}}\right)$$

▶ cos(3.14) m

▶ m - v, where m has dimension of mass and v of velocity **Incorrect**: cannot subtract quantities of different dimension

> 10 N s - 2 kg m/s = 8 J s/m

▶ exp(-8 J)

Incorrect: exponential function must have a dimensionless argument, but this argument has dimension energy

- \triangleright (9 m, 0.1 rad, -0.5 rad)
- ► 8 J/s = 12 N m 4 N mIncorrect: $\text{J/s} \neq \text{N m}$ (correct is J = N m)
- ▶ e⁻⁸ J

$$\frac{15 \text{ J}}{5 \text{ kg/s}^2} = 3 \text{ m}^2$$

$$\sqrt{25} \, K = 5$$

Incorrect: both sides of an equation must have the same dimension; here the left side has dimension length $^{1/2}$, right side is dimensionless

▶ (e⁷)^s
 Incorrect: cannot raise to a dimensional power

$$ightharpoonup tan \left(\frac{10 \, \text{m}}{5 \, \text{m}} \right)$$

- ▶ √300 K
- $ightharpoonup \sin(t/s)$, where t has dimension of time
- $\Rightarrow \frac{3}{s}$
- sin(10 s)

Incorrect: trigonometric function must have a dimensionless argument

Time and space 2

Make sure you're familiar with the 'dot-notation' explained in § 2.8 of our text.

2.0

(Do the exercises in the main text.)

2.1

Preferably together with a colleague:

The *Veritasium*¹ channel has many informative and entertaining videos on diverse scientific topics. Most of these videos are accurate and pedagogically very useful. But a couple of them contain some inaccuracies or partially faulty reasoning.

One example of partially inaccurate video is *Why no one has measured* the speed of light². It contains many correct and insightful statements and explanations, but also some faulty reasoning.

Watch the video and

- 1. Identify and ponder about some explanations that reflect what you learned so far. (For instance, do you recognize *radar distance* between t=3:10 and t=3:20?)
- 2. Consider the discussion between t=4:57³ and t=5:14, and the statement "and get a response 20 minutes later". What kind of time is this statement referring to? is it proper time? if so, whose proper time? or is it coordinate time?
- **3.** Consider the same snip and the statement "we imagine our signal takes 10 minutes to get there". Draw a spacetime diagram (similar to fig. 2.1 in our main text) illustrating this statement. In the diagram,

place the proper times on the worldline of the Earth station and on Mark's worldline; and mark the points where the signal is sent and where it is received.

How can we imagine that it takes 10 minutes to get there? Which proper time are we speaking about?

- **4.** Consider again the snip and the statement "it's possible that our message took all 20 minutes to get there". Draw a spacetime diagram illustrating this statement. What's the difference from the previous spacetime diagram? Are the two spacetime diagrams actually different?
- 5. Now consider the discussion between t=9:47⁴ and t=10:16, and the statement "one of the clocks will be ahead of the other". When we say *ahead*, to which kind of time are we referring? is it proper time? if so, whose proper time? Does it make sense to say that one clock is "ahead" of the other?
- **6.** Draw one or two spacetime diagrams illustrating the discussion in the snip above. Can we make sense of the discussion using the diagrams?
- 7. Find parts in which the reasoning offered in the video is inconsistent. For instance, find discussions where Derek says "right now": does "right now" make sense in those discussions?

1. 2.2

Preferably together with a colleague:

A particular coordinate system (t, x, y, z) with spatial Cartesian coordinates is defined as follows:

- The time coordinate *t* is your proper time.
- The origin of the coordinates is your navel
- The *x*-axis points in front of you, the *y*-axis to your left, the *z*-axis upwards (through the top of your head).
- The unit coordinate is 1 m, measured as usual.

Answer the following questions:

1. What are your position $\mathbf{r}(t)$ and velocity $\mathbf{v}(t)$ in this coordinate system while you sleep? (Let's say that by "your position" we mean the position of your navel.)

- **2.** What are your position $\mathbf{r}(t)$ and velocity $\mathbf{v}(t)$ while you run or bike or drive to school?
- **3.** What is your acceleration $\boldsymbol{a}(t)$ in different situations?
- **4.** Determine the *z* coordinate of the floor in this coordinate system, when you are standing still.
- **5.** Determine the spatial coordinates of the tip of the index finger of your right hand, when it is extended horizontally outwards.

- **1.** You're told that the position $\mathbf{r}(t)$ of an object is constant in time t. How much is the velocity $\mathbf{v}(t)$?
- **2.** If the velocity $\mathbf{v}(t_0)$ is zero at a time t_0 , must also the acceleration $\mathbf{a}(t_0)$ be zero at time t_0 ?
- **3.** Is it possible for a coordinate velocity $v_x(t_1)$ to be positive at a time t_1 , and the acceleration $a_x(t_1)$ negative at the same time? If not, explain why not. If yes, show by constructing a concrete example and explain what this situation means physically.

3. 2.4

We have a coordinate system (t, x) with one spatial dimension only. A small object S has position $x_S(t)$ which changes with the coordinate time t. The time dependence of the position is given by

$$x_S(t) = at + b$$
 with $a = -3 \,\mathrm{m/s}$, $b = 7 \,\mathrm{m}$.

- 1. Verify that the equation above is dimensionally consistent.
- **2.** What is the spatial coordinate of S at times t = 0 s, t = -10 s, and t = 5 s?
- **3.** What is the spatial coordinate of S at time t = 10?
- **4.** Calculate the time dependence of the coordinate velocity of S.
- **5.** What is the cooordinate velocity of S at time t = 5 s?
- **6.** What is the *speed* of S at time t = 5 s?
- 7. Calculate the time dependence of the coordinate acceleration of S.

We have a coordinate system (t, x) with one spatial dimension only. A small object S has position $x_S(t)$ given by

$$x_{\rm S}(t) = L \sin(\omega t) + b$$
 with $L = 2 \,\mathrm{m}$, $\omega = \frac{\pi}{3} \,\mathrm{s}^{-1}$, $b = 7 \,\mathrm{m}$.

- 1. Verify that the equation above is dimensionally consistent.
- **2.** Calculate the expressions for velocity $\dot{x}_{\rm S}(t)$ and acceleration $\ddot{x}_{\rm S}(t)$.
- **3.** Find a time t_0 in which the velocity is $0 \,\text{m/s}$ and the acceleration is approximately $-2.2 \,\text{m/s}^2$.
- **4.** Find a time t_1 in which the velocity is approximately -2.1 m/s and the acceleration is 0 m/s².
- **5.** Plot $x_S(t)$ and $\dot{x}_S(t)$ as functions of time for $t \in [-4, 4]$ s.

2.6

We have a coordinate system (t, x, y, z), where the three spatial coordinates have each dimension length. A small object S has position $\mathbf{r}_{S}(t)$ given by

$${\pmb r}_{\rm S}(t) = \begin{bmatrix} at+b\\ L\sin(\omega t)+b\\ 0 \end{bmatrix}$$
 with $L=2\,{\rm m}$, $\omega=\frac{\pi}{3}\,{\rm s}^{-1}$, $a=-3\,{\rm m/s}$, $b=7\,{\rm m/s}$.

- 1. Verify that the equation above is dimensionally consistent.
- **2.** Calculate the expressions for velocity $\dot{r}_{\rm S}(t)$ and acceleration $\ddot{r}_{\rm S}(t)$.
- **3.** Plot the three components of the velocity as functions of time for $t \in [-4, 4]$ s.

2.7

We have a coordinate system (t, z) with one spatial dimension only. The coordinate velocity $v_z(t)$ of a small pulse of light travelling in a particular material is given by

$$v_z(t) = c \exp(-t/\tau)$$
 with $c = 299792458 \,\mathrm{m/s}$, $\tau = 0.08 \,\mathrm{s}$,

and the pulse is located at $z = -2 \,\mathrm{m}$ at $t = 1 \,\mathrm{s}$.

- **1.** Find the expression for the position z(t) of the pulse as a function of coordinate time.
- **2.** Find the location of the pulse at time $t = 1.01 \, \text{s}$.

Example solutions

Ω 2.3

- **1.** The derivative of a constant is zero, so the velocity is $\mathbf{v}(t) = 0 \,\text{m/s}$. We must not forget the correct units!
- **2.** No, we can have zero velocity and non-zero acceleration at a given time. See exercise 2.5 as an example.
- **3.** No, we can have positive velocity and negative acceleration at a given time. See exercise 2.5 as an example. It means that, at that time, the movement is in the positive-*x* direction (positive *x*-velocity), and the *x*-velocity is decreasing that is, it will be positive but smaller a very short time later.

Ω 2.4

- 1. It is, provided that t has dimension time and x has dimension length. In this case, since a has dimension length/time, then a t has dimension length, which is added to b which also has dimension length; the left and right side have then both dimension length.
- **2.** $x_S(0 s) = 7 m$, $x_S(-10 s) = 37 m$, $x_S(5 s) = -8 m$.
- **3.** The question doesn't make sense, because "t = 10" is dimensionless; it should have dimension length instead.
- **4.** Denoting with \dot{x}_S the coordinate velocity of S, then $\dot{x}_S(t) = a$, which is constant in time.
- **5.** $\dot{x}_{\rm S}(t) = -3 \, {\rm m/s}$ at any time.
- **6.** The speed is $|\dot{x}_{\rm S}(t)| = 3$ m/s at any time.
- 7. Denoting with \ddot{x}_S the coordinate acceleration of S, then $\ddot{x}_S(t) = 0 \text{ m/s}^2$, which is zero at all times.

Ω 2.5

- **1.** The expression is dimensionally correct, provided *t* has dimension time and *x* has dimension length. The argument of the sine function is dimensionless, and the two terms on the right have dimension length.
- 2. From the rules for the derivative,

$$\dot{x}_{\rm S}(t) = \omega L \cos(\omega t)$$
, $\ddot{x}_{\rm S}(t) = -\omega^2 L \sin(\omega t)$.

3. The time t_0 must satisfy the system of equations

$$\omega L \cos(\omega t) = 0 \text{ m/s}$$
 $-\omega^2 L \sin(\omega t) \approx -2.2 \text{ m/s}^2$.

The cosine is zero when its argument is $\pi/2$, $3\pi/2$, and so on. Let's try taking $\omega t_0 = \pi/2$, which means $t_0 = \pi/(2\omega)$. We find indeed

$$\dot{x}_{\rm S}(t_0) = \omega L \cos(\omega t_0) = 0 \,\mathrm{m/s}$$
 $\ddot{x}_{\rm S}(t_0) = -\omega^2 L \sin(\omega t_0) \approx -2.19 \,\mathrm{m/s^2}$.

4. The time t_1 must satisfy the system of equations

$$\omega L \cos(\omega t) = -2.1 \,\mathrm{m/s}$$
 $-\omega^2 L \sin(\omega t) \approx 0 \,\mathrm{m/s^2}$.

The sine is zero when its argument is 0, π , and so on. Let's try taking $\omega t_1 = 0$, which means $t_1 = 0$ s. We find

$$\omega L \cos(\omega t_1) \approx 2.09 \,\mathrm{m/s}$$
 $-\omega^2 L \sin(\omega t_1) = 0 \,\mathrm{m/s}^2$,

which is not what we want. Trying next $\omega t_1 = \pi$, which means $t_1 = \pi/\omega$, leads to the desired result.

5. We can plot x_S and \dot{x}_S in two separate graphs:



Or we could plot them on the same graph – but only if we indicate separately the vertical axis for x_S and the one for \dot{x}_S (for instance one on the left and one on the right), because these quantities have different dimensions.

♀ 2.6

- **1.** No, the expression is not dimensionally correct, because the z component of r_S is "0", which is a dimensionless number, whereas z has dimension length. The z component should be "0 m".
- 2. See exercises 2.4 and 2.5

♀ 2.7

1. Let's call the specific position $z_0 := -2 \,\mathrm{m}$ at $t_0 1 \,\mathrm{s}$. The expression for z(t) is found integrating $v_z(t)$ between t_0 and t, and adding the position at t_0 :

$$z(t) = z_0 + \int_{t_0}^t c \exp(-t/\tau) dt$$

$$= z_0 - c\tau \exp(-t/\tau) \Big|_{t_0}^t$$

$$= z_0 - c\tau \Big[\exp(-t/\tau) - \exp(-t_0/\tau) \Big]$$
with $c = 299792458 \,\text{m/s}$, $\tau = 0.08 \,\text{s}$, $z_0 = -2 \,\text{m}$, $t_0 = 1 \,\text{s}$.

2. Substituting in the expression above, $z(1.01 \, \text{s}) = 8.50 \, \text{m}$.

URLs for chapter 2

- 1. https://www.youtube.com/c/veritasium/videos
- 2. https://www.youtube.com/watch?v=pTn6Ewhb27k
- 3. https://youtu.be/pTn6Ewhb27k?t=297
- 4. https://youtu.be/pTn6Ewhb27k?t=588

Main physical quantities 3



(Do the exercises in the main text.)

5. 3.1

We have seen that six of the seven main physical quantities have two important properties:

- We can speak about their *content*, *flux*, and *supply*.
- They are *extensive*: for instance if a 3D region consists of two non-overlapping 3D subregions, then the content in the region is the sum of the content in the subregions. Similarly for flux and surfaces (2D regions).
- **1.** Can you think of some physical quantity that has the property of extensivity, but for which it doesn't make sense to speak of "content" or of "flux" or of "supply"?
- **2.** Vice versa, can you think of some physical quantity for which we can speak of content, flux, supply, but which doesn't have the property of extensivity?

3.2

Do a little research, and find out whether there are any physics disciplines in which *mass* is usually measured in units of *energy*.

Do a little research, and find out whether there are any physics disciplines in which *mass* is usually measured in units of *energy*.

3.4

With a colleague or a large language model:

Take turns to explain to each other what is the difference between *matter* and *mass*. Try to find weak points in each other's explanations.

3.5

1. Imagine that someone tells you this:

An important difference between *matter* and *electric charge* is that electric charge can be both positive and negative, whereas matter can only be positive.

How would you reply? Can you give counterexamples to this statement?

2. The same person tells you:

In nature we observe both positive and negative electric charge equally easily. But we mostly observe 'positive' matter, and very rarely 'negative' matter (antimatter).

Do some research and find out whether this statement is true.

3.6

Someone tells you:

Mass and energy are two different things. I can experimentally prove it to you: Take a battery for example, and weigh it to measure its mass. Now use the battery for some device. As you use it, the battery loses energy; in fact eventually it can't power the device anymore. But if you weigh it again, you'll find that it has the same mass as before. Therefore mass and energy must be two different things.

How would you reply to this person?

Let's say you use a coordinate system (x, y, z). In a given 3D region of space you measure a net amount (content) of momentum $\mathbf{P} = [2, -3, 0] \,\mathrm{N}\,\mathrm{s}$. Which of the following statements are true? which false? Explain why.

- **1.** There must be some non-zero electric charge in the 3D region.
- **2.** The net amount of momentum has zero *z*-component.
- **3.** There must be some matter (or antimatter) in the 3D region.
- **4.** The 3D region must be enough small.
- **5.** Some kind of motion, with respect to your coordinate system, must be occurring in the 3D region.
- **6.** Any scientist measuring the net momentum in the 3D region would agree that its value is [2, -3, 0] N s.
- 7. Whatever it is that contributes to the net momentum, it must be uniformly spread out through the whole 3D region.

Example solutions

\circ 3.1

[Before reading this answer, keep in mind that the word volume has two different meanings: sometimes we use it in the sense of "3D region of space"; sometimes in the sense of the size of a 3D region of space (measured in cubic metres for instance).]

- 1. One example is the *volume* of a 3D region. It is extensive, because the volume of two non-overlapping 3D regions together is the sum of their volumes. But it doesn't make much sense to speak of the "flux" of volume through a surface. Similarly for *area*.
- **2.** The writer of this solution doesn't know any example of such a physical quantity. It is possible to define mathematical objects that behave this strange way, but no physical quantity seems to be represented by such objects.

② 3.3

One example is particle physics¹, where the rest mass of subatomic particles is measured in *electronvolts*, denoted 'eV', which is a unit for energy equal to $1.602\,176\,634\times10^{-19}$ J. Other examples are special relativity and general relativity.

\circ 3.7

- **1.** *False.* There can be momentum in a region even if the net electric charge is zero.
- **2.** *True. z* is the third coordinate, and the third momentum component is 0 N s.
- **3.** *False.* Electromagnetic fields have momentum, so in the region there could be only an electromagnetic field, such as a beam of light, but no matter.
- **4.** *False.* We can speak of the net amount of momentum in arbitrarily large regions.
- **5.** *True.* Momentum is associated with the motion of matter or of electromagnetic fields.

- **6.** *False.* The amount of momentum depends on the coordinate system we choose; so scientists that use coordinates different from yours will measure a different net momentum it could even be completely zero.
- 7. *False*. The matter or electromagnetic field that possess the momentum might be concentrated in one or several small regions within the 3D region, for example.

URLs for chapter 3

1. https://cms.cern/content/glossary#E

Volume contents, fluxes, supplies



4.0

(Do the exercises in the main text.)

4.1

In the following figures, region A is indicated in blue and a solid contour, and region B is indicated in red and a dashed contour (they are 2D simplified representations of 3D regions). For each figure determine, if possible, the net volume content of the corresponding quantities in the region comprising A and B. If not possible, explain why.

1. Energy-mass:



2. Matter:



3. Electric charge:



4. Energy-mass:



4.2

In the following figures, region A is indicated in blue and a solid contour, and region B is indicated in red and a dashed contour (they are 2D simplified representations of 3D regions). Use a coordinate system (x, y, z) where x is horizontal to the right, y is vertical upwards on the page, and z comes out of the page towards you. For each figure:

- draw the vectors representing the momentum contents of A and B;
- determine, if possible, the net volume content of momentum in the region comprising A and B; if not possible, explain why.







 $P_{\rm A} = [38, 5.5, 0] \, \rm Ns$



4.



 $P_{\rm A} = [-7, 4, 0] \, {\rm Ns}$

5.





В

In the side figure, consider volume A, which contains the ball, and volume B, occupied by the light beam. Imagine there's no air (although in reality if there were no air we wouldn't be able to see the light beam).

In a given coordinate system, the momentum content of volume A is $\mathbf{P}_{A} = [1,0,0] \,\mathrm{N}\,\mathrm{s}$, and that of volume B is $\mathbf{P}_{B} = [2,-2,1] \times 10^{-16} \,\mathrm{N}\,\mathrm{s}$.

Does it really make sense to speak of the momentum in volume B (remember there's no air there)? If it does make sense, how much is the net momentum content of volumes A and B considered together?



4.4

Take a look at the energy flux through a surface represented in the side figure. Which of the representations below is completely equivalent to the one on the side? Which does represent a different flux instead?





4.5

An ordinary battery is attached to wires and is powering some device. This means that there is a flow of electromagnetic energy from the battery to the device. Imagine a control surface around the battery, as illustrated in the side figure. Can you guess across which parts of the control surface is the flux of energy? Is it across the whole surface? or only across the parts where the wire is? Do a little research to find out.



A cuboid region is delimited by a closed control surface that can be divided into six parts; call them 'up', 'down', 'front', 'back', 'left', 'right'. They are all have an inward crossing direction, except for 'back' which has an outward crossing direction.

Here are the fluxes of various quantities through the six surfaces in the given crossing directions. Calculate the total net **in**fluxes.

1.
$$J_u = 23.9 \text{ mol/s}$$
 $J_d = -8.1 \text{ mol/s}$ $J_f = 0.9 \text{ mol/s}$ $J_b = -30.5 \text{ mol/s}$ $J_l = 2.3 \text{ mol/s}$ $J_r = 37.6 \text{ mol/s}$


2.
$$\Phi_{\rm u} = -24.6 \, {\rm J/s}$$
 $\Phi_{\rm d} = 2.4 \, {\rm J/s}$ $\Phi_{\rm f} = 1.3 \, {\rm J/s}$ $\Phi_{\rm b} = 10.8 \, {\rm J/s}$ $\Phi_{\rm l} = 15.4 \, {\rm J/s}$ $\Phi_{\rm r} = -2.1 \, {\rm J/s}$

3.
$$I_u = -9.7 \,\text{C/s}$$
 $I_d = 27.4 \,\text{C/s}$ $I_f = -6.3 \,\text{C/s}$ $I_b = 16.4 \,\text{C/s}$ $I_l = -25.1 \,\text{C/s}$ $I_r = -20.0 \,\text{C/s}$

4.
$$\Pi_u = 31.7 \text{ J/(K s)}$$
 $\Pi_d = 17.9 \text{ J/(K s)}$ $\Pi_f = 7.2 \text{ J/(K s)}$ $\Pi_b = -20.4 \text{ J/(K s)}$ $\Pi_l = -16.5 \text{ J/(K s)}$ $\Pi_r = -4.8 \text{ J/(K s)}$

5.
$$\mathbf{F}_{u} = [25.2, -42.7, 4.1] \, N$$
 $\mathbf{F}_{d} = [-46.3, -5.9, -33.3] \, N$ $\mathbf{F}_{f} = [10.2, -16.2, -36.7] \, N$ $\mathbf{F}_{b} = [-9.8, 19.5, -60.0] \, N$ $\mathbf{F}_{l} = [29.5, 18.3, 58.4] \, N$ $\mathbf{F}_{r} = [16.0, 1.2, -3.2] \, N$

Which of the following graphical representations of fluxes do make sense? Which don't? Explain why.



4.8

- **1.** A control volume contains [2, -3, 1] N s of momentum. Momentum is a quantity related to movement. Does this mean that the control volume is moving?
- **2.** Suppose that through a control surface, in a given crossing direction, there is a non-zero flux of momentum. Does this mean that the control surface must be in motion?
- **3.** If through a control surface there's a non-zero flux of some quantity, does it mean that the fluxes of all other quantities must be zero?
- **4.** Can there be a flux of temperature through a control surface?

Which of the following graphical representations of fluxes do make sense? Which don't? Explain why.



4.10

The Möbius band¹ is a surface that, in a certain sense, has only *one* side. It is quite easy to make with a strip of paper and some tape.

Do you manage to choose a definite crossing direction on this surface? Do you think it could be chosen as a control surface?



4.11

For each figure, calculate and draw, or describe precisely, the net total flux through the composite control surface. (The different parts are different

tiated by different colours and thicknesses.) Don't forget to specify the crossing direction when you speak about a flux.

1. 4.9J/s 2.2J/s



2.









4.12

1. Through a particular closed control surface there's a continuous influx of electric charge I(t) with the following time dependence:

$$I(t) = A \sin(\omega t)$$
 with $A = 30 \text{ C/s}$, $\omega = 376.991 \, 184 \, \text{s}^{-1}$.

How much is the time-integrated influx of charge between times $t_0 = 10 \,\mathrm{s}$ and $t_1 = 40 \,\mathrm{s}$?

2. Imagine a closed control surface around the planet Mercury. The energy $\Phi(t)$ influx through this control surface is approximately given by

$$\Phi(t) = W + u \cos(\sigma t)$$

with $W = 2 \times 10^{17} \,\text{J/s}$, $u = 8 \times 10^{16} \,\text{J/s}$, $\sigma = 8 \times 10^{-7} \,\text{s}^{-1}$.

Mercury's year lasts around 88 Earth-days. How much is the time-integrated influx of energy over a year on Mercury?

3. A football has a mass-energy m = 0.4 kg. The football has a continuous supply of momentum $\boldsymbol{G}(t)$ from the Earth's gravitational field. This supply is constant in time and given by

$$\boldsymbol{G}(t) = m\boldsymbol{g} \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$$

where $\mathbf{g} = 9.8 \,\mathrm{N/kg}$, and we are using a coordinate system (t, x, y, z) where z is vertical with respect to the ground and points upwards.

How much is the time-integrated influx of momentum in the football between times $t_0 = 0$ s and $t_1 = 3600$ s?

Ω 4.1

- **1.** Net energy-mass content is $E_A + E_B = 22.1 \, \text{J}$.
- **2.** Unknown: A and B are overlapping, so we cannot simply add their contents. The result depends on the net matter content in the overlap region, which is unknown.
- **3.** Net electric-charge content is $Q_A + Q_B = 2 C$.
- **4.** A and B are overlapping, so we cannot simply add their mass-energy contents. But in this case A is a subregion of B, so the full region occupied by both is B itself. The net mass-energy content in this region is therefore $m_{\rm B}=0.47$ kg. We can also deduce that the net mass-energy content in the region between A and B is $m_{\rm B}-m_{\rm A}=0.35$ kg.

② 4.2

- **1.** Net momentum content is $\mathbf{P}_A + \mathbf{P}_B = [0, 0, 0] N s$.
- **2.** Unknown: A and B are overlapping, so we cannot simply add their momentum contents. The result depends on the net momentum content in the overlap region, which is unknown.
- 3. Net momentum content is $\mathbf{P}_A + \mathbf{P}_B = [-3, -3, 0] \,\mathrm{N} \,\mathrm{s}$.
- **4.** A and B are overlapping, so we cannot simply add their momentum contents. But in this case B is a subregion of A, so the full region occupied by both is A itself. The net momentum content in this region is therefore $\mathbf{P}_{A} = [-7, 4, 0] \, \mathrm{N} \, \mathrm{s}$. We can also deduce that the net momentum content in the region between B and A is $\mathbf{P}_{A} \mathbf{P}_{B} = [-9, 4, 0] \, \mathrm{N} \, \mathrm{s}$.
- **5.** Net momentum content is $P_A + P_B = [6.4, 0.8, 0] \,\mathrm{N}\,\mathrm{s}$. The region consisting of A and B together is a spatially disconnected region, but we can speak of volume content even for such kind of regions.

\bigcirc 4.3

Yes, it does make sense. First of all, we can *always* speak of the momentum content in *any* 3D region; it isn't important whether there's matter or electromagnetic fields or not in that region. If it's completely empty, its momentum content is simply zero. Second, in volume B in the figure

there's an electromagnetic field, which does have momentum just like matter.

The net momentum content in the volumes A and B considered together is $\mathbf{P}_A + \mathbf{P}_B \approx [1, -2 \times 10^{-16}, 1 \times 10^{-16}] \,\mathrm{N}\,\mathrm{s}.$

② 4.4

1. Same

Same crossing direction, same flux magnitude. Wiggly arrows only indicate the crossing direction, it doesn't matter how many there are.

4.

Different •

This flux has same magnitude but opposite direction.

2.
Different

This flux has a different magnitude.

5.

Same I Same flux, by the principle of symmetry of flux.

3. Same 👍

Same crossing direction, same flux magnitude. Crossing direction is indicated verbally.

6. Same **⊯**

Same flux, by the principle of symmetry of flux.



The flux of electromagnetic energy occurs – that is, it is non-zero – across *every part of the control surface*, not just across the parts close to the wire.

In general the flow of electromagnetic energy occurs especially around wires, which act as sort of guides; but it extends to all space.

Two good videos for learning more about this are *Veritasium's The big misconception about electricity*² and *How electricity actually works*³. More detailed examples, calculations, and figures can be found for example in Jackson 1996 (see especially fig. 4), Davis & Kaplan 2011, Harbola 2010, Boyer 2019.



♀ 4.6

To calculate the efflux across the whole surface we use the principle of extensivity. The fluxes across the 'back' surface get a minus sign, because that surface had an inward crossing direction.

- 1. $J_{\text{tot}} = J_{\text{u}} + J_{\text{d}} + J_{\text{f}} J_{\text{b}} + J_{\text{l}} + J_{\text{r}} = 87.1 \,\text{mol/s}$
- **2.** $\Phi_{tot} = \Phi_u + \Phi_d + \Phi_f \Phi_b + \Phi_l + \Phi_r = -18.4 \text{ J/s}$
- 3. $I_{\text{tot}} = I_{\text{u}} + I_{\text{d}} + I_{\text{f}} I_{\text{b}} + I_{\text{l}} + I_{\text{r}} = -50.1 \,\text{C/s}$
- **4.** $\Pi_{\text{tot}} = \Pi_{\text{u}} + \Pi_{\text{d}} + \Pi_{\text{f}} \Pi_{\text{b}} + \Pi_{\text{l}} + \Pi_{\text{r}} = -50.1 \,\text{C/s}$
- **5.** $\mathbf{F}_{tot} = \mathbf{F}_{u} + \mathbf{F}_{d} + \mathbf{F}_{f} \mathbf{F}_{b} + \mathbf{F}_{l} + \mathbf{F}_{r} = [44.4, -64.8, 49.3] \text{ C/s}$

Q 4.7

1. Makes sense

No •• Crossing direction is unclear.

2.

4. 5. No •••

Crossing direction Crossing direction is unclear. is missing.

7.
No ••
'N' is the unit of a

vector flux, but there's no vector or vector components here. 8.

Alternative way of indicating the crossing direction, but perfectly clear.

Makes sense

3. No •••

'J' is not the unit of a flux.

6. Makes sense

The arrows for the crossing direction are very tilted, but crossing direction is still clear.

9. Makes sense **1**

♀ 4.8

1. No. If a control volume contains momentum, then something like matter or electromagnetic field is moving within the volume; but the volume itself may be static. This is because a control volume is just an

imaginary delimitation of a region of space, not a physical object; we decide how it moves. In some situations we may decide that it should move together with the matter or electromagnetic field it contains, but in other situations we may keep it static.

- **2.** No, for analogous reasons as in the answer above.
- **3.** No: *in principle* the fluxes of the seven quantities are independent and can all be non-zero; or some can be zero and others non-zero.
- **4.** No: we can speak of the temperature at a point at a given time, but we cannot speak of volume content or flux or supply of temperature. Temperature is not an extensive quantity.

♀ 4.9

1. Makes sense

This vector flux is expressed in components, and the coordinates are shown: we can reconstruct the vector if needed.

2. Makes sense

3. No !•

There should be only one vector indicating the flux. Unclear if this flux should have magnitude 4 N.

4. No !•

'J/s' is the unit of a **scalar** flux, but there's a vector here. Either the unit is wrong or the vector is there by mistake. 5. Makes sense 🏚

This is a total influx.

6.

No !

The wiggly arrows that indicate the crossing direction are not mutually consistent.

7.

Unclear if one of the two arrows indicates a crossing direction. 8. Makes sense

Total influx again. The flux is expressed in components, and the coordinates are shown.

9. No !•


The magnitude of a vector cannot be negative.

♀ 4.10

You have noticed that if you try to choose consistently a crossing direction on the whole band, you end up where you started but with an *opposite* crossing direction. For this reason the Möbius band cannot be used as a control surface: it cannot be given an overall crossing direction. The lack of crossing direction is related to another peculiarity: it is impossible to extend this surface in such a way as to enclose a three-dimensional region of space.

If we clip a part of the Möbius band, that part can be used as a control surface. Or if we clip the band so that it gets an extra border, then it becomes the same as a twisted rectangle, and so it can be used as a control surface.

♀ 4.11



Ω 4.12

1. The time-integrated flux of charge is found with a time integral. Note that the result has a dimension of electric charge: this is the net amount of charge that flowed through the control surface, in the given crossing direction, between the two times:

$$\int_{t_0}^{t_1} I(t) dt = \int_{t_0}^{t_1} A \sin(\omega t) dt = -\frac{A}{\omega} \cos(\omega t) \Big|_{t_0}^{t_1}$$
$$\approx -796 \,\mathrm{C} \cdot (1 - 1) = 0 \,\mathrm{C}.$$

So even if the flux of charge most of the time is non-zero, reaching a maximum of $\pm 30 \,\text{C/s}$, the *net* amount of charge crossing the surface in the given direction during the $30 \,\text{s}$ is zero. This is because the flux is sometimes positive, sometimes negative.

2. 88 Earth-days are approximately equal to $88 \cdot 24 \cdot 60 \cdot 60 \, \text{s} \approx 7600\,000 \, \text{s}$. So we time-integrate between $t_0 = 0 \, \text{s}$ and $t_1 = 7600\,000 \, \text{s}$:

$$\int_{t_0}^{t_1} \Phi(t) dt = \int_{t_0}^{t_1} \left[W + u \cos(\sigma t) \right] dt = Wt \Big|_{t_0}^{t_1} + \frac{u}{\sigma} \sin(\sigma t) \Big|_{t_0}^{t_1}$$

$$\approx 1.5 \times 10^{24} \,\text{J} + 1 \times 10^{23} \,\text{J} \cdot (-0.2 - 0)$$

$$\approx 1.5 \times 10^{24} \,\text{J} .$$

3. Recall that time-integrating a vector simply means time-integrating each component. In this case the supply is constant, so integration is easy:

$$\int_{t_0}^{t_1} \mathbf{G}(t) dt = \int_{t_0}^{t_1} m \mathbf{g} \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix} dt$$

$$= m \mathbf{g} \begin{bmatrix} \int_{t_0}^{t_1} 0 dt \\ \int_{t_0}^{t_1} 0 dt \\ -\int_{t_0}^{t_1} 1 dt \end{bmatrix} = m \mathbf{g} \begin{bmatrix} 0 s \\ 0 s \\ -t \end{bmatrix} \Big|_{t_0}^{t_1} = m \mathbf{g} \begin{pmatrix} \begin{bmatrix} 0 s \\ 0 s \\ -t_1 \end{bmatrix} - \begin{bmatrix} 0 s \\ 0 s \\ -t_0 \end{bmatrix} \right)$$

$$\approx 3.9 \, \text{N} \cdot \begin{bmatrix} 0 \\ 0 \\ -3600 \end{bmatrix} s \approx \begin{bmatrix} 0 \\ 0 \\ -1.4 \times 10^4 \end{bmatrix} \text{Ns}.$$

This net amount of momentum is vertical, pointing downward. But note that we don't know whether the football contains this much momentum

after $3600\,\mathrm{s}$, because we don't know how much the flux of momentum into the football was. It's possible that the flux of momentum completely balanced this supply of momentum.

URLs for chapter 4

- 1. https://mathworld.wolfram.com/MoebiusStrip.html
- 2. https://www.youtube.com/watch?v=bHIhgxav9LY
- 3. https://www.youtube.com/watch?v=oI_X2cMHNe0

Physical laws 5



(Do the exercises in the main text.)

5.1

For each question:

- Identify and describe the closed control surface and control volume used.
- Identify which among volume content, flux, supply, are given, and how much they are; identify which are unknown.
- Find the requested quantity, if possible, by using a balance or conservation law as specified in the question, showing all mathematical steps. If not possible, explain why.
- 1. At a time of 0 s, a tank contains 10 mol of water, plus other substances. The tank is sealed. In a time lapse of 20 s an amount of 3 mol of water is produced in the tank by chemical reaction. Assume that the amount of water satisfies a balance law. How much water is in the tank at the end of this time lapse?
- 2. In 30 min a battery has emitted 4300 J of electromagnetic energy. At the end of this time there is no energy in the battery (with respect to a given zero of energy). Assume that energy satisfies a conservation law in this example. How much energy was in the battery at the beginning of the 30 min?
- 3. A small block of thorium- 231^1 contains 141.0 mol of neutrons. The thorium undergoes beta decay². After 9.2×10^4 s, the number of neutrons in the block is 140.5 mol. No neutrons were emitted. Assume that

- neutrons satisfy a balance law. How much was the time-integrated supply of neutron during that time lapse? (Be careful about the signs.)
- **4.** A bowling ball, at a given time and in a given coordinate system, has momentum [11.6, 0, 0.6] N s. Three seconds later the ball is completely at rest, with momentum [0, 0, 0] N s. How much was the time-integrated influx of momentum during the three seconds?
- 5. A small laser beam, of given width and length, contains a net angular momentum [0, 2.1e 36, 0] N m s, in a given coordinate system. The same region of space, 0.1 s later, does not contain any electromagnetic field anymore, and therefore contains zero angular momentum. Assume that angular momentum satisfies a conservation law. How much was the time-integrated efflux of angular momentum through the control surface that contained the beam?

5.2

Find the requested quantity, if possible, by using any combination of:

- given data,
- balance or conservation law (as specified),
- principle of extensivity,
- principle of symmetry of flux,
- calculus.

If not possible, explain why.

1. Water is produced in a tank at a rate of 2 mol/s

5.3

With a colleague or a large language model:

- 1. Explain to a colleague the difference between a balance law and a conservation law. Let your colleague criticize unclear or incorrect points in your explanation, and comment on the good points. Then exchange roles.
- **2.** If you have a large-language-model service, ask it what's the definition of a balance law, and then what's the difference between a balance law and a conservation law. Compare its answer with what you learned so far. Argue with it and see where the discussion goes.

Keep in mind that there exist many slightly different definitions of 'balance law' and 'conservation law'. Large language models are trained on texts containing many different definitions – including erroneous ones!

5.4

Feel free to discuss this with a colleague:

Imagine you're supervising a research team. Your team is exploring and developing the new, ground-breaking kind of material *super-energium*, studying its characteristics about energy and temperature.

Super-energium is supposed to be somewhat similar to the better known *energium*, which is therefore used as a comparison in your team's investigations. The energy flux Φ of a block of *energium* satisfies the differential conservation law

$$\frac{\mathrm{d}E(t)}{\mathrm{d}t} = \Phi(t) \tag{5.1}$$

where E is the energy of the block. For this material, the energy is connected with the temperature T of the block by the constitutive relation

$$U(t) = aT(t)^4 (5.2)$$

where *a* is a constant.

Your team performs experiments on blocks of *super-energium*, and finds that the equations above don't match the experimental results. The equations that describe *super-energium* must therefore be different. The engineers in your team propose and test different modifications of the physical laws, and finally find two different modifications that work:

(A) One group of engineers show that if we keep the original constitutive relation (5.2), but modify the conservation law (5.1) as follows:


$$\left[\frac{\mathrm{d}E(t)}{\mathrm{d}t}\right]^2 = \Phi(t) \tag{5.3}$$

then we can perfectly describe and predict all experimental results.

(B) Another group of engineers show that if we keep the original conservation law (5.1), but modify the constitutive relation (5.2) as follows:

$$U(t) = \frac{16}{7}a^2T(t)^7 (5.4)$$

then we can also perfectly describe and predict all experimental results.



You must decide whether to continue experiments and development using the modifications proposed by engineers (A), or those proposed by engineers (B) (you can't pursue both). Which do you choose, and why?

5.5

Consider a small control volume completely occupied by a particular kind of gas, which we shall call *ideal gas*. Focus also on a small part of the closed control surface associated with this control volume.

Two constitutive relations are known to be valid for such a control volume:

• The ideal-gas law

$$F = \frac{A}{V}RNT$$

says that the magnitude of the surface force (momentum influx) on the small area is proportional to the amount of gas N in the control volume, the temperature T of the gas, the constant R, the surface's area A; and inversely proportional to the volume V.

• The energy equation for an ideal gas

$$E = U_0 + CNT$$

says that the total energy-mass in the control volume containing the gas is proportional to the amount of gas N there, the gas's temperature T, and two constants C and U_0 .

Find a constitutive relation that directly connects the volume content of energy-mass and the magnitude of the magnitude of flux of momentum, without involving any auxiliary quantities; constants and geometric quantities such as areas and volumes are allowed.

5.6

Ω 5.2

1. We consider an imaginary closed control surface corresponding to the inner surface of the tank.

The problem gives:

- Initial time $t_0 = 0$ s, final time $t_1 = 20$ s.
- Volume content of water $N(t_0) = 10$ mol at time t_0 .
- Time-integrated supply of water in control volume $\int_{t_0}^{t_1} J(t) dt =$ 3 mol.
- Time-integrated influx of water is 0 mol ("the tank is sealed").

Unknown:

• Volume content of water $N(t_1)$ at time t_1 .

If water obeys a balance law, the quantities above are related by

$$N(t_1) = N(t_0) + \int_{t_0}^{t_1} J(t) dt + \int_{t_0}^{t_1} A(t) dt$$

of which three are known. We can find the unknown:

$$N(t_1) = N(t_0) + \int_{t_0}^{t_1} J(t) dt + \int_{t_0}^{t_1} A(t) dt$$

= 10 mol + 0 mol + 3 mol = 13 mol.

2. We consider an imaginary closed control surface perfectly wrapping the battery.

The problem gives:

- Initial time t_0 , final time $t_1 = t_0 + 1800 \, \mathrm{s}$. Time-integrated **in**flux of energy $\int_{t_0}^{t_1} \Phi(t) \, \mathrm{d}t = -4300 \, \mathrm{J}$; minus sign because the text specifies the efflux.
- Volume content of energy $E(t_1) = 0$ mol at time t_1 .

Unknown:

• Volume content of energy $E(t_0)$ at time t_0 .

If energy obeys a conservation law, the quantities above are related by

$$E(t_1) = E(t_0) + \int_{t_0}^{t_1} \Phi(t) dt$$

of which two are known. We can find the unknown by simple algebra:

$$E(t_0) = E(t_1) - \int_{t_0}^{t_1} \Phi(t) dt$$

= 0J + 4300J = 4300J.

3. We consider an imaginary closed control surface perfectly wrapping the block of material.

The problem gives:

- Initial time t_0 , final time $t_1 = t_0 + 9.2 \times 10^4$ s.
- Volume content of neutrons $N(t_0) = 141.0$ mol at time t_0 .
- Volume content of neutrons $N(t_1) = 140.5$ mol at time t_1 .
- Time-integrated influx of neutrons $\int_{t_0}^{t_1} J(t) dt = 0 \mod (\text{"no neutrons were emitted"}).$

Unknown:

• Time-integrated supply of neutrons $\int_{t_0}^{t_1} A(t) dt$.

If the amount of neutrons obeys a balance law, the quantities above are related by

$$N(t_1) = N(t_0) + \int_{t_0}^{t_1} J(t) dt + \int_{t_0}^{t_1} A(t) dt$$

of which three are known. We can find the unknown by algebra:

$$\int_{t_0}^{t_1} A(t) dt = N(t_1) - yN(t_0) - \int_{t_0}^{t_1} J(t) dt$$

= 141.0 mol - 140.5 mol - 0 mol = -0.5 mol.

The negative sign for the supply makes sense, because the number of neutrons has decreased.

4. We consider an imaginary closed control surface perfectly wrapping the bowling ball (note that this control volume and surface are not static).

The problem gives:

• Initial time t_0 , final time $t_1 = t_0 + 3$ s.

- Volume content of momentum $\mathbf{P}(t_0) = [11.6, 0, 0.6] \,\mathrm{N}\,\mathrm{s}$ at time t_0 .
- Volume content of momentum $P(t_1) = [0, 0, 0] \text{ N s at time } t_1$.

Unknown:

• Time-integrated influx of momentum $\int_{t_0}^{t_1} \mathbf{F}(t) dt$.

Momentum obeys a balance law, so the quantities above are related by

$$\mathbf{P}(t_1) = \mathbf{P}(t_0) + \int_{t_0}^{t_1} \mathbf{F}(t) dt + \int_{t_0}^{t_1} \mathbf{G}(t) dt.$$

Only two of these four quantities are given in the problem, so we can't find the influx of momentum.

One could, intelligently, think of maybe using the constitutive relation for the supply of momentum: $\mathbf{G} = mg[0,0,-1]$. But unfortunately the mass m of the bowling ball is not given in the problem.

5. We consider an imaginary and static closed control surface enclosing the region where the laser beam (the electromagnetic field) was initially.

The problem gives:

- Initial time t_0 , final time $t_1 = t_0 + 0.1$ s.
- Angular-momentum content $\mathbf{L}(t_0) = [0, 2.1e 36, 0] \,\mathrm{N}\,\mathrm{m}\,\mathrm{s}$ at time t_0 .
- Angular-momentum content $L(t_1) = [0, 0, 0] \text{ N m s at time } t_1$.

Unknown:

• Time-integrated **ef**flux of angular momentum $-\int_{t_0}^{t_1} \boldsymbol{\tau}(t) \, \mathrm{d}t$; the minus sign is because $\boldsymbol{\tau}$ denotes the influx.

Assuming angular momentum to obey a conservation law, the quantities above are related by

$$\mathbf{L}(t_1) = \mathbf{L}(t_0) + \int_{t_0}^{t_1} \mathbf{\tau}(t) \, \mathrm{d}t$$

of which two are known. We can find the efflux by algebra:

$$-\int_{t_0}^{t_1} \boldsymbol{\tau}(t) dt = \boldsymbol{L}(t_0) - yL(t_1)$$

$$= [0, 2.1e - 36, 0] \text{ N m s} - [0, 0, 0] \text{ N m s} = [0, 2.1e - 36, 0] \text{ N m s}.$$

It makes sense that the efflux is positive, because all angular momentum left the control volume.

Ω 5.5

We want to find a relation that gets rid of the auxiliary quantity T, the temperature. This can be done with several equivalent algebraic manipulations; here's one. Solve the ideal-gas law for the temperature:

$$T = \frac{V}{A} \frac{F}{RN}$$

then substitute this expression into the energy equation:

$$E = U_0 + CN\left(\frac{V}{A}\frac{F}{RN}\right) \qquad \Longrightarrow \qquad \boxed{E = U_0 + \frac{V}{A}\frac{C}{R}F} \ .$$

This is the desired constitutive relation. It says that the energy content E is directly proportional to the magnitude of the momentum influx F, with some proportionality constants that also involve the size of the control volume and the area of the control surface.

URLs for chapter 5

- 1. https://pubchem.ncbi.nlm.nih.gov/compound/Thorium-231
- 2. http://hyperphysics.phy-astr.gsu.edu/hbase/Nuclear/beta.html

The Seven Wonders of the world 6



Conservation & balances of matter





Conservation of electric charge 8





Conservation of magnetic flux 9



Balance of momentum 10



Balance of energy 11



Balance of angular momentum 12



Remarks on momentum and energy 13



Balance of entropy 14



Constitutive relations 15



Bibliography

Believe nothing, O monks, merely because you have been told it, or because it is traditional, or because you yourselves have imagined it. Do not believe what your teacher tells you merely out of respect for the teacher.

(Attributed to Gautama Buddha)

("de X" is listed under D, "van X" under V, and so on, regardless of national conventions.)

- Boyer, T. H. (2019): *Illustrations of Maxwell's term and the four conservation laws of electromagnetism*. Am. J. Phys. **87**⁹, 729–738. DOI:10.1119/1.5115339.
- Davis, B. S., Kaplan, L. (2011): *Poynting vector flow in a circular circuit*. Am. J. Phys. **79**¹¹, 1155–1162. DOI:10.1119/1.3630927.
- Harbola, M. K. (2010): Energy flow from a battery to other circuit elements: role of surface charges. Am. J. Phys. **78**¹¹, 1203–1206. DOI:10.1119/1.3456567.
- Jackson, J. D. (1996): *Surface charges on circuit wires and resistors play three roles*. Am. J. Phys. **64**⁷, 855–870. DOI:10.1119/1.18112.