

# **The Seven Wonders of the World**

## ***Exercises***

Updated 29 January 2025 22:20

# Contents

<b>1</b>	<b>Physics, quantities, units</b>	<b>5</b>
1.0	.....	5
1.1	.....	5
1.2	.....	5
1.3	.....	6
1.4	.....	6
1.5	.....	7
1.6	.....	7
	Example solutions	10
<b>2</b>	<b>Time and space</b>	<b>13</b>
2.0	.....	13
2.1	.....	13
2.2	.....	14
2.3	.....	15
2.4	.....	15
2.5	.....	16
2.6	.....	16
2.7	.....	16
	Example solutions	18
	URLs for chapter 2	21
<b>3</b>	<b>Main physical quantities</b>	<b>22</b>
3.0	.....	22
3.1	.....	22
3.2	.....	22
3.3	.....	23
	Example solutions	24
	URLs for chapter 3	26

---

<b>4</b>	<b>Volume contents, fluxes, supplies</b>	<b>27</b>
4.0	.....	27
4.1	.....	27
	Example solutions	29
<b>5</b>	<b>Physical laws</b>	<b>30</b>
5.0	.....	30
	Example solutions	31
<b>6</b>	<b>The Seven Wonders of the world</b>	<b>32</b>
6.0	.....	32
	Example solutions	33
<b>7</b>	<b>Conservation &amp; balances of matter</b>	<b>34</b>
7.0	.....	34
	Example solutions	35
<b>8</b>	<b>Conservation of electric charge</b>	<b>36</b>
8.0	.....	36
	Example solutions	37
<b>9</b>	<b>Conservation of magnetic flux</b>	<b>38</b>
9.0	.....	38
	Example solutions	39
<b>10</b>	<b>Balance of momentum</b>	<b>40</b>
10.0	.....	40
	Example solutions	41
<b>11</b>	<b>Balance of energy</b>	<b>42</b>
11.0	.....	42
	Example solutions	43
<b>12</b>	<b>Balance of angular momentum</b>	<b>44</b>
12.0	.....	44
	Example solutions	45
<b>13</b>	<b>Remarks on momentum and energy</b>	<b>46</b>
13.0	.....	46
	Example solutions	47

---

<b>14 Balance of entropy</b>	<b>48</b>
14.0 . . . . .	48
Example solutions . . . . .	49
<b>15 Constitutive relations</b>	<b>50</b>
15.0 . . . . .	50
Example solutions . . . . .	51

# Physics, quantities, units 1

For some of the following exercises you can refer to tables [1.1](#) and [1.2](#) on page 9 (reproduced from the textbook).

## 1.0

(Do the **exercises** in the main text.)

## 1.1

*Preferably together with a friend or colleague:*

If you have some large-language-model service (such as ChatGPT), ask it which physical laws are universally valid in Newtonian Mechanics and in General Relativity and in Thermodynamics and in Chemistry and in Electromagnetics.

Discuss the answer you get, based on what you have learned so far. (Note: if the answer mention a ‘balance of boost momentum’, that’s actually correct.)

Argue with the LLM and see where the discussion goes.

## 1.2

Take *time* and *velocity* as primitive quantities.

1. Try to define *distance* as a derived quantity
2. Try to define *acceleration* as a derived quantity.

### 1.3

Which of the following quantities are *scalars*, and which are *vectors*?

- Time
- Distance
- Position
- Energy
- Velocity
- Speed
- Momentum
- Entropy
- Angular momentum
- Force
- Temperature
- Magnetic flux
- Electric charge
- Electric current
- Heat
- Power
- Volume
- Pressure

### 1.4

Find the correct units for the following quantities:

- *Volumic energy* or *energy density*, defined as energy divided by volume
- *Energy flux*, defined as energy divided by time.
- *Power*, defined as energy divided by time.
- *Heating*, defined as energy divided by time.
- *Magnetic flux*, which we take as a primitive quantity.
- *Electric potential difference*, defined as magnetic flux divided by time.
- *Force*, defined as momentum divided by time.
- *Momentum flux*, defined as momentum divided by time.
- *Momentum supply*, defined as momentum divided by time.
- *Pressure*, defined as force divided by area.
- *Amount of substance (or of matter)*, which we take as primitive.
- *Molar mass*, defined as mass divided by amount of substance.
- *Specific momentum*, defined as momentum divided by mass.

- *Volumic charge* or *charge density*, defined as charge divided by volume.
- *Entropy*, which we take as primitive, has dimension of energy divided by temperature.
- *Matter density*, defined as amount of substance divided by volume.
- *Matter flux*, defined as amount of substance divided by time.

## 1.5

*With a friend or colleague:*

1. Try to explain to your friend the difference between a *primitive quantity* and a *derived quantity*; then let your friend criticize unclear or incorrect points in your explanation, and comment on the good points. Then invert your roles: your friend tries to explain to you, and you criticize and comment.
2. Similarly as the previous exercise, but explaining the difference between a *scalar quantity* and a *vector quantity*.
3. If you have some large-language-model service (such as ChatGPT), ask it to explain the difference between primitive and derived quantity, and between scalar and vector quantity. Find out weak or unsure points in its answer, given what you've learned so far.

## 1.6

Find which of the following mathematical expressions and equalities are dimensionally incorrect, and explain why they are incorrect:

- ▷  $11 \text{ J} + 4 \text{ kg}$
- ▷  $\tan\left(\frac{a}{b}\right)$ , where  $a$  has dimension length and  $b$  has dimension time
- ▷  $299\,792\,458 \text{ m/s}$
- ▷  $\exp\left(\frac{71 \text{ s}}{3 \text{ s}}\right)$
- ▷  $\cos(3.14) \text{ m}$
- ▷  $m - v$ , where  $m$  has dimension of mass and  $v$  of velocity

- ▷  $10 \text{ N s} - 2 \text{ kg m/s} = 8 \text{ J s/m}$
- ▷  $\exp(-8 \text{ J})$
- ▷  $(9 \text{ m}, 0.1 \text{ rad}, -0.5 \text{ rad})$
- ▷  $8 \text{ J/s} = 12 \text{ N m} - 4 \text{ N m}$
- ▷  $e^{-8} \text{ J}$
- ▷  $\frac{15 \text{ J}}{5 \text{ kg/s}^2} = 3 \text{ m}^2$
- ▷  $\sqrt{25} \text{ K} = 5$
- ▷  $(e^7)^s$
- ▷  $\tan\left(\frac{10 \text{ m}}{5 \text{ m}}\right)$
- ▷  $\sqrt{300 \text{ K}}$
- ▷  $\sin(t/\text{s})$ , where  $t$  has dimension of time
- ▷  $\frac{3}{\text{s}}$
- ▷  $\sin(10 \text{ s})$



Quantity	SI Dimension	Unit
Time	time	<i>second</i> s
Length	length	<i>metre</i> m
Temperature	temperature	<i>kelvin</i> K
Matter	amount of substance	<i>mole</i> mol
Electric charge	electric charge	<i>coulomb</i> C
Magnetic flux	magnetic flux	<i>weber</i> Wb
Energy	energy, mass	<i>joule</i> J, <i>kilogram</i> kg
<b>Momentum</b>	force · time, mass · length/time, energy · time/length	N · s, kg · m/s, J · s/m
<b>Angular momentum</b>	force · length · time, mass · length <sup>2</sup> /time, energy · time	N · m · s, kg · m <sup>2</sup> /s, J · s
Entropy	energy/temperature	J/K

Table 1.1 Dimensions and units of the main physical quantities used in these notes. Their fluxes have the dimensions divided by time, and therefore units divided by seconds. Quantities in **boldface** are vectors, the others are scalars.

Quantity	Volume content [unit]	Flux [unit]
matter	$N$ [mol]	$J$ [mol/s]
electric charge	$Q$ [C]	$\mathcal{I}$ [C/s or A]
magnetic flux	$\mathcal{B}$ [Wb]	$\mathcal{E}$ [Wb/s or V]
energy	$E$ [J]	$\Phi$ [J/s or W]
momentum	$\mathbf{P}$ [N s]	$\mathbf{F}$ [N]
angular momentum	$\mathbf{L}$ [N m s]	$\boldsymbol{\tau}$ [N m]
entropy	$S$ [J/K]	$\Pi$ [J/(K s)]

Table 1.2 Units for volume contents and fluxes of the main seven quantities.

## Example solutions

### 💡 1.2

1. “Distance is the product of a time lapse and a particular velocity”. See section 2.3 about *Radar distance* in our lecture notes.
2. “Acceleration is the ratio between a change in the product of a time lapse and a particular velocity, and the time taken by that change”.

### 💡 1.3

These quantities are scalars:

- Time
- Distance
- Energy
- Speed
- Entropy
- Temperature
- Magnetic flux
- Electric charge
- Electric current
- Heat
- Power
- Volume

These quantities are vectors:

- Position
- Velocity
- Momentum
- Angular momentum
- Force

For *pressure*, it depends on the context. In some applications it is considered a scalar, but in other applications it is considered a vector – or actually a generalized kind of vector, called *tensor*, which can be represented by a matrix.

### 💡 1.4

- *Volumic energy*:  $\text{J/m}^3$

- *Energy flux*: J/s
- *Power*: J/s
- *Heating*: J/s
- *Magnetic flux*: Wb
- *Electric potential difference*: Wb/s
- *Force*: N
- *Momentum flux*: N
- *Momentum supply*: N
- *Pressure*: N/m<sup>2</sup>
- *Amount of substance*: mol
- *Molar mass*: kg/mol
- *Specific momentum*: N · s/kg  $\equiv$  m/s
- *Volumic charge*: C/m<sup>3</sup>
- *Entropy*: J/K
- *Matter density*: mol/m<sup>3</sup>
- *Matter flux*: mol/s

## 💡 1.6

- ▷  $11 \text{ J} + 4 \text{ kg}$   
**Incorrect:** cannot sum quantities of different dimension
- ▷  $\tan\left(\frac{a}{b}\right)$ , where  $a$  dimension length and  $b$  has dimension time  
**Incorrect:** trigonometric function must have a dimensionless argument, but  $a/b$  has dimension length/time
- ▷  $299\,792\,458 \text{ m/s}$
- ▷  $\exp\left(\frac{71 \text{ s}}{3 \text{ s}}\right)$
- ▷  $\cos(3.14) \text{ m}$
- ▷  $m - v$ , where  $m$  has dimension of mass and  $v$  of velocity  
**Incorrect:** cannot subtract quantities of different dimension
- ▷  $10 \text{ N s} - 2 \text{ kg m/s} = 8 \text{ J s/m}$
- ▷  $\exp(-8 \text{ J})$   
**Incorrect:** exponential function must have a dimensionless argument, but this argument has dimension energy

▷  $(9 \text{ m}, 0.1 \text{ rad}, -0.5 \text{ rad})$

▷  $8 \text{ J/s} = 12 \text{ N m} - 4 \text{ N m}$

**Incorrect:**  $\text{J/s} \neq \text{N m}$  (correct is  $\text{J} = \text{N m}$ )

▷  $e^{-8} \text{ J}$

▷  $\frac{15 \text{ J}}{5 \text{ kg/s}^2} = 3 \text{ m}^2$

▷  $\sqrt{25} \text{ K} = 5$

**Incorrect:** both sides of an equation must have the same dimension; here the left side has dimension  $\text{length}^{1/2}$ , right side is dimensionless

▷  $(e^7)^s$

**Incorrect:** cannot raise to a dimensional power

▷  $\tan\left(\frac{10 \text{ m}}{5 \text{ m}}\right)$

▷  $\sqrt{300 \text{ K}}$

▷  $\sin(t/\text{s})$ , where  $t$  has dimension of time

▷  $\frac{3}{\text{s}}$

▷  $\sin(10 \text{ s})$

**Incorrect:** trigonometric function must have a dimensionless argument

# Time and space 2

Make sure you're familiar with the 'dot-notation' explained in § 2.8 of our text.

## 2.0

(Do the **exercises** in the main text.)

## 2.1

*Preferably together with a friend or colleague:*

The [Veritasium](#)<sup>1</sup> channel has many informative and entertaining videos on diverse scientific topics. Most of these videos are accurate and pedagogically very useful. But a couple of them contain some inaccuracies or partially faulty reasoning.

One example of partially inaccurate video is [Why no one has measured the speed of light](#)<sup>2</sup>. It contains many correct and insightful statements and explanations, but also some faulty reasoning.

Watch the video and

1. Identify and ponder about some explanations that reflect what you learned so far. (For instance, do you recognize *radar distance* between  $t=3:10$  and  $t=3:20$ ?)
2. Consider the discussion between  $t=4:57$ <sup>3</sup> and  $t=5:14$ , and the statement “and get a response 20 minutes later”. What kind of time is this statement referring to? is it proper time? if so, whose proper time? or is it coordinate time?
3. Consider the same snip and the statement “we imagine our signal takes 10 minutes to get there”. Draw a spacetime diagram (similar to fig. 2.1 in our main text) illustrating this statement. In the diagram,

place the proper times on the worldline of the Earth station and on Mark's worldline; and mark the points where the signal is sent and where it is received.

How can we imagine that it takes 10 minutes to get there? Which proper time are we speaking about?

4. Consider again the snip and the statement "it's possible that our message took all 20 minutes to get there". Draw a spacetime diagram illustrating this statement. What's the difference from the previous spacetime diagram? Are the two spacetime diagrams actually different?
5. Now consider the discussion between  $t=9:47^4$  and  $t=10:16$ , and the statement "one of the clocks will be ahead of the other". When we say *ahead*, to which kind of time are we referring? is it proper time? if so, whose proper time? Does it make sense to say that one clock is "ahead" of the other?
6. Draw one or two spacetime diagrams illustrating the discussion in the snip above. Can we make sense of the discussion using the diagrams?
7. Find parts in which the reasoning offered in the video is inconsistent. For instance, find discussions where Derek says "right now": does "right now" make sense in those discussions?

## 2.2

*Preferably together with a friend or colleague:*

A particular coordinate system  $(t, x, y, z)$  with spatial Cartesian coordinates is defined as follows:

- The time coordinate  $t$  is your proper time.
- The origin of the coordinates is your navel
- The  $x$ -axis points in front of you, the  $y$ -axis to your left, the  $z$ -axis upwards (through the top of your head).
- The unit coordinate is 1 m, measured as usual.

Answer the following questions:

1. What are your position  $\mathbf{r}(t)$  and velocity  $\mathbf{v}(t)$  in this coordinate system while you sleep? (Let's say that by "your position" we mean the position of your navel.)

2. What are your position  $\mathbf{r}(t)$  and velocity  $\mathbf{v}(t)$  while you run or bike or drive to school?
3. What is your acceleration  $\mathbf{a}(t)$  in different situations?
4. Determine the  $z$  coordinate of the floor in this coordinate system, when you are standing still.
5. Determine the spatial coordinates of the tip of the index finger of your right hand, when it is extended horizontally outwards.

### 2.3

1. You're told that the position  $\mathbf{r}(t)$  of an object is constant in time  $t$ . How much is the velocity  $\mathbf{v}(t)$ ?
2. If the velocity  $\mathbf{v}(t_0)$  is zero at a time  $t_0$ , must also the acceleration  $\mathbf{a}(t_0)$  be zero at time  $t_0$ ?
3. Is it possible for a coordinate velocity  $v_x(t_1)$  to be positive at a time  $t_1$ , and the acceleration  $a_x(t_1)$  negative at the same time? If not, explain why not. If yes, show by constructing a concrete example and explain what this situation means physically.

### 2.4

We have a coordinate system  $(t, x)$  with one spatial dimension only. A small object S has position  $x_S(t)$  which changes with the coordinate time  $t$ . The time dependence of the position is given by

$$x_S(t) = at + b \quad \text{with} \quad a = -3 \text{ m/s}, \quad b = 7 \text{ m}.$$

1. Verify that the equation above is dimensionally consistent.
2. What is the spatial coordinate of S at times  $t = 0 \text{ s}$ ,  $t = -10 \text{ s}$ , and  $t = 5 \text{ s}$ ?
3. What is the spatial coordinate of S at time  $t = 10$ ?
4. Calculate the time dependence of the coordinate velocity of S.
5. What is the coordinate velocity of S at time  $t = 5 \text{ s}$ ?
6. What is the *speed* of S at time  $t = 5 \text{ s}$ ?
7. Calculate the time dependence of the coordinate acceleration of S.

## 2.5

We have a coordinate system  $(t, x)$  with one spatial dimension only. A small object S has position  $x_S(t)$  given by

$$x_S(t) = L \sin(\omega t) + b \quad \text{with} \quad L = 2 \text{ m}, \quad \omega = \frac{\pi}{3} \text{ s}^{-1}, \quad b = 7 \text{ m}.$$

1. Verify that the equation above is dimensionally consistent.
2. Calculate the expressions for velocity  $\dot{x}_S(t)$  and acceleration  $\ddot{x}_S(t)$ .
3. Find a time  $t_0$  in which the velocity is 0 m/s and the acceleration is approximately  $-2.2 \text{ m/s}^2$ .
4. Find a time  $t_1$  in which the velocity is approximately  $-2.1 \text{ m/s}$  and the acceleration is  $0 \text{ m/s}^2$ .
5. Plot  $x_S(t)$  and  $\dot{x}_S(t)$  as functions of time for  $t \in [-4, 4] \text{ s}$ .

## 2.6

We have a coordinate system  $(t, x, y, z)$ , where the three spatial coordinates have each dimension length. A small object S has position  $\mathbf{r}_S(t)$  given by

$$\mathbf{r}_S(t) = \begin{bmatrix} at + b \\ L \sin(\omega t) + b \\ 0 \end{bmatrix} \quad \text{with} \quad L = 2 \text{ m}, \quad \omega = \frac{\pi}{3} \text{ s}^{-1}, \quad a = -3 \text{ m/s}, \quad b = 7 \text{ m/s}.$$

1. Verify that the equation above is dimensionally consistent.
2. Calculate the expressions for velocity  $\dot{\mathbf{r}}_S(t)$  and acceleration  $\ddot{\mathbf{r}}_S(t)$ .
3. Plot the three components of the velocity as functions of time for  $t \in [-4, 4] \text{ s}$ .

## 2.7

We have a coordinate system  $(t, z)$  with one spatial dimension only. The coordinate velocity  $v_z(t)$  of a small pulse of light travelling in a particular material is given by

$$v_z(t) = c \exp(-t/\tau) \quad \text{with} \quad c = 299\,792\,458 \text{ m/s}, \quad \tau = 0.08 \text{ s},$$

and the pulse is located at  $z = -2 \text{ m}$  at  $t = 1 \text{ s}$ .



1. Find the expression for the position  $z(t)$  of the pulse as a function of coordinate time.
2. Find the location of the pulse at time  $t = 1.01$  s.

## Example solutions

### 💡 2.3

1. The derivative of a constant is zero, so the velocity is  $\boldsymbol{v}(t) = 0 \text{ m/s}$ . We must not forget the correct units!
2. No, we can have zero velocity and non-zero acceleration at a given time. See exercise 2.5 as an example.
3. No, we can have positive velocity and negative acceleration at a given time. See exercise 2.5 as an example. It means that, at that time, the movement is in the positive- $x$  direction (positive  $x$ -velocity), and the  $x$ -velocity is decreasing – that is, it will be positive but smaller a very short time later.

### 💡 2.4

1. It is, provided that  $t$  has dimension time and  $x$  has dimension length. In this case, since  $a$  has dimension length/time, then  $a t$  has dimension length, which is added to  $b$  which also has dimension length; the left and right side have then both dimension length.
2.  $x_S(0 \text{ s}) = 7 \text{ m}$  ,  $x_S(-10 \text{ s}) = 37 \text{ m}$  ,  $x_S(5 \text{ s}) = -8 \text{ m}$  .
3. The question doesn't make sense, because " $t = 10$ " is dimensionless; it should have dimension length instead.
4. Denoting with  $\dot{x}_S$  the coordinate velocity of  $S$ , then  $\dot{x}_S(t) = a$ , which is constant in time.
5.  $\dot{x}_S(t) = -3 \text{ m/s}$  at any time.
6. The speed is  $|\dot{x}_S(t)| = 3 \text{ m/s}$  at any time.
7. Denoting with  $\ddot{x}_S$  the coordinate acceleration of  $S$ , then  $\ddot{x}_S(t) = 0 \text{ m/s}^2$ , which is zero at all times.

### 💡 2.5

1. The expression is dimensionally correct, provided  $t$  has dimension time and  $x$  has dimension length. The argument of the sine function is dimensionless, and the two terms on the right have dimension length.
2. From the rules for the derivative,

$$\dot{x}_S(t) = \omega L \cos(\omega t) , \quad \ddot{x}_S(t) = -\omega^2 L \sin(\omega t) .$$

3. The time  $t_0$  must satisfy the system of equations

$$\omega L \cos(\omega t) = 0 \text{ m/s} \quad - \omega^2 L \sin(\omega t) \approx -2.2 \text{ m/s}^2 .$$

The cosine is zero when its argument is  $\pi/2, 3\pi/2$ , and so on. Let's try taking  $\omega t_0 = \pi/2$ , which means  $t_0 = \pi/(2\omega)$ . We find indeed

$$\dot{x}_S(t_0) = \omega L \cos(\omega t_0) = 0 \text{ m/s} \quad \ddot{x}_S(t_0) = -\omega^2 L \sin(\omega t_0) \approx -2.19 \text{ m/s}^2 .$$

4. The time  $t_1$  must satisfy the system of equations


$$\omega L \cos(\omega t) = -2.1 \text{ m/s} \quad - \omega^2 L \sin(\omega t) \approx 0 \text{ m/s}^2 .$$

The sine is zero when its argument is  $0, \pi$ , and so on. Let's try taking  $\omega t_1 = 0$ , which means  $t_1 = 0 \text{ s}$ . We find

$$\omega L \cos(\omega t_1) \approx 2.09 \text{ m/s} \quad - \omega^2 L \sin(\omega t_1) = 0 \text{ m/s}^2 ,$$

which is not what we want. Trying next  $\omega t_1 = \pi$ , which means  $t_1 = \pi/\omega$ , leads to the desired result.

5. We can plot  $x_S$  and  $\dot{x}_S$  in two separate graphs:



Or we could plot them on the same graph – but only if we indicate separately the vertical axis for  $x_S$  and the one for  $\dot{x}_S$  (for instance one on the left and one on the right), because these quantities have different dimensions.

## 💡 2.6

1. No, the expression is not dimensionally correct, because the  $z$  component of  $\mathbf{r}_S$  is "0", which is a dimensionless number, whereas  $z$  has dimension length. The  $z$  component should be "0 m".
2. See exercises 2.4 and 2.5 😊

## 💡 2.7

1. Let's call the specific position  $z_0 := -2 \text{ m}$  at  $t_0 = 1 \text{ s}$ . The expression for  $z(t)$  is found integrating  $v_z(t)$  between  $t_0$  and  $t$ , and adding the position at  $t_0$ :

$$z(t) = z_0 + \int_{t_0}^t c \exp(-t/\tau) dt$$

$$= z_0 - c\tau \exp(-t/\tau) \Big|_{t_0}^t$$

$$= z_0 - c\tau [\exp(-t/\tau) - \exp(-t_0/\tau)]$$

$$\text{with } c = 299\,792\,458 \text{ m/s}, \tau = 0.08 \text{ s}, z_0 = -2 \text{ m}, t_0 = 1 \text{ s}.$$

2. Substituting in the expression above,  $z(1.01 \text{ s}) = 8.50 \text{ m}$ .

## URLs for chapter 2

1. <https://www.youtube.com/c/veritasium/videos>
2. <https://www.youtube.com/watch?v=pTn6Ewhb27k>
3. <https://youtu.be/pTn6Ewhb27k?t=297>
4. <https://youtu.be/pTn6Ewhb27k?t=588>

# Main physical quantities 3

## 3.0

(Do the **exercises** in the main text.)

## 3.1

We have seen that six of the seven main physical quantities have two important properties:

- We can speak about their *content*, *flux*, and *supply*.
  - They are *extensive*: for instance if a 3D region consists of two non-overlapping 3D subregions, then the content in the region is the sum of the content in the subregions. Similarly for flux and surfaces (2D regions).
1. Can you think of some physical quantity that has the property of extensivity, but for which it doesn't make sense to speak of "content" or of "flux" or of "supply"?
  2. Vice versa, can you think of some physical quantity for which we can speak of content, flux, supply, but which doesn't have the property of extensivity?

## 3.2

Do a little research, and find out whether there are any physics disciplines in which *mass* is usually measured in units of *energy*.

### 3.3

Let's say you use a coordinate system  $(x, y, z)$ . In a given 3D region of space you measure a net amount (content) of momentum  $\mathbf{P} = [2, -3, 0]$  N s. Which of the following statements are true? which false? Explain why.

1. There must be some non-zero electric charge in the 3D region.
2. The net amount of momentum has zero  $z$ -component.
3. There must be some matter (or antimatter) in the 3D region.
4. The 3D region must be enough small.
5. Some kind of motion, with respect to your coordinate system, must be occurring in the 3D region.
6. Any scientist measuring the net momentum in the 3D region would agree that its value is  $[2, -3, 0]$  N s.
7. Whatever it is that contributes to the net momentum, it must be uniformly spread out through the whole 3D region.

## Example solutions

### 💡 3.1

*[Before reading this answer, keep in mind that the word volume has two different meanings: sometimes we use it in the sense of “3D region of space”; sometimes in the sense of the size of a 3D region of space (measured in cubic metres for instance).]*

1. One example is the *volume* of a 3D region. It is extensive, because the volume of two non-overlapping 3D regions together is the sum of their volumes. But it doesn't make much sense to speak of the “flux” of volume through a surface. Similarly for *area*.
2. The writer of this solution doesn't know any example of such a physical quantity. It is possible to define mathematical objects that behave this strange way, but no physical quantity seems to be represented by such objects.

### 💡 3.2

One example is [particle physics](#)<sup>1</sup>, where the rest mass of subatomic particles is measured in *electronvolts*, denoted ‘eV’, which is a unit for energy equal to  $1.602\,176\,634 \times 10^{-19}$  J. Other examples are special relativity and general relativity.

### 💡 3.3

1. *False*. There can be momentum in a region even if the net electric charge is zero.
2. *True*.  $z$  is the third coordinate, and the third momentum component is 0 N s.
3. *False*. Electromagnetic fields have momentum, so in the region there could be only an electromagnetic field, such as a beam of light, but no matter.
4. *False*. We can speak of the net amount of momentum in arbitrarily large regions.
5. *True*. Momentum is associated with the motion of matter or of electromagnetic fields.



6. *False.* The amount of momentum depends on the coordinate system we choose; so scientists that use coordinates different from yours will measure a different net momentum – it could even be completely zero.
7. *False.* The matter or electromagnetic field that possess the momentum might be concentrated in one or several small regions within the 3D region, for example.

## URLs for chapter 3

1. <https://cms.cern/content/glossary#E>

# Volume contents, fluxes, supplies 4


## 4.0

(Do the **exercises** in the main text.)


## 4.1

In the following figures, **region A** is indicated in blue and a solid contour, and **region B** is indicated in red and a dashed contour (they are 2D simplified representations of 3D regions). Determine, if possible, the net volume content of the corresponding quantities in the region comprising A and B. If not possible, explain why.


1. Energy:



2. Matter:



3. Electric charge:




## Example solutions

### 4.1

1. Net content is  $E_A + E_B = 22.1 \text{ J}$ .
2. Unknown: A and B are overlapping, so the result depends on the net matter content in the overlap region.
3. Net content is  $Q_A + Q_B = 2 \text{ C}$ .


# Physical laws 5



(Do the **exercises** in the main text.)

## Example solutions

# The Seven Wonders of the world 6



6.0

(Do the **exercises** in the main text.)



## **Example solutions**

# Conservation & balances of matter 7

## 7.0

(Do the **exercises** in the main text.)

## Example solutions

# Conservation of electric charge 8

## 8.0

(Do the **exercises** in the main text.)

## Example solutions

# Conservation of magnetic flux 9

## 9.0

(Do the **exercises** in the main text.)

## Example solutions

# Balance of momentum 10

## 10.0

(Do the **exercises** in the main text.)



## Example solutions

# Balance of energy 11



## 11.0

(Do the **exercises** in the main text.)

## Example solutions

# Balance of angular momentum 12

## 12.0

(Do the **exercises** in the main text.)

## Example solutions

# Remarks on momentum and energy 13

## 13.0

(Do the **exercises** in the main text.)

## Example solutions

# Balance of entropy 14



## 14.0

(Do the **exercises** in the main text.)



## Example solutions

# Constitutive relations 15

## 15.0

(Do the **exercises** in the main text.)

## Example solutions