

# Dimensional analysis in general relativity and differential geometry

P.G.L. Porta Mana 

Kavli Institute, Trondheim <[pgl@portamana.org](mailto:pgl@portamana.org)>

13 December 2019; updated 2 May 2020

This note illustrates how to perform dimensional analysis in general relativity and differential geometry, and tries to revive Dorgelo & Schouten's notion of *absolute dimension* of a tensorial quantity. The absolute dimension is independent of the dimensions of the coordinate functions. The dimensional analysis of several important tensors and tensor operations is summarized. In particular it is shown that the components of a tensor need not have all the same dimension, and that the Riemann (once contravariant and twice covariant) and Ricci (fully covariant) tensors are dimensionless. The relation between dimension and operational meaning for the metric and stress-energy-momentum tensors is also discussed.

*For Emma*

## 1 Introduction

From the point of view of dimensional analysis, do all components of a tensor need to have the same dimension? What is the dimension of the metric and the curvature? And what is the dimension of the constant in the Einstein equations?

There seem to be insecurity and incorrect notions, among some students and even some researchers in relativity, regarding the dimensions of tensors and of tensor components, the effect of tensor operators on dimensions, and the dimension of constants in differential-geometric and field equations. I've met, for example, with the statement that the components of a tensor should all have the same dimension; and with calculations of the dimensions of curvature tensors assuming that coordinates have dimensions of length. That statement is wrong; and that assumption is unnecessary.

Several factors probably cause or contribute to such difficulties. Modern texts in Lorentzian and general relativity commonly use geometrized units. They say that, for finding the dimension of some constant in a tensorial equation, it's sufficient to compare the dimensions of the

tensors in the equation. But this is not so immediate, because some tensors don't have universally agreed dimensions – prime example the metric tensor. Older texts often use coordinates with dimension of length and time, and base their dimensional analysis on that specific choice. They even multiply coordinates with dimension of time, as well as some tensorial components, by powers of  $c$ . They thus give the impression that coordinates ought to always be lengths, and that all components of a tensor ought to have the same dimension<sup>1</sup>.

The main purpose of this Note is to illustrate an extremely simple and intuitive way of reasoning with which one can quickly and consistently settle any dimensional-analysis questions in general relativity and differential geometry; for example the dimension of the Riemann tensor, or the dimensional results of contraction, of covariant derivative, of raising an index. This way of reasoning is presented and illustrated in §§ 5–6. It's just an application of the usual rules of dimensional analysis: if two quantities are summed, then they must have the same dimension; the dimension of a product is the product of the dimensions; and so on. This way of reasoning relies on the coordinate-free, *intrinsic* view of tensors and other differential-geometrical objects, a brief reminder of which is given in § 3, with references.

Another important purpose of this Note is to revive the forgotten notion of *absolute dimension* of a tensor, and the distinction between it and the dimensions of that tensor's *components* in some coordinate system. This notion is explained in § 5. The absolute dimension, introduced by Schouten and Dorgelo<sup>2</sup> and used in Truesdell & Toupin<sup>3</sup>, is invariant with respect to the dimensions of the coordinates; it's an intrinsic property of a tensor. The dimensions of the components depend instead on the dimensions of the coordinate functions – which can be completely arbitrary, as discussed in § 4. The absolute dimension should thus be the primary focus in dimensional analysis in general relativity and in differential geometry.

The rest of this Note is mainly an application of the intuitive way of reasoning mentioned above and of the notion of absolute dimension. We'll reveal and rectify a couple of misconceptions; for example, *the components of a tensor need not have the same dimensions*. The results for the

<sup>1</sup> e.g. Tolman 1949 § 37 eq. (37.8); Landau & Lifshitz 1996 § 32 eq. (32.15); Adler et al. 1975 § 10.1 eq. (10.15). <sup>2</sup> Dorgelo & Schouten 1946; Schouten 1989 ch. VI.

<sup>3</sup> Truesdell & Toupin 1960 Appendix II.

main tensor operations and operators are summarized in § 6. The results for the curvature tensors and other objects depending on a connection are presented in § 8. It is shown, in particular, that the absolute dimension of the (contravariant and thrice covariant) Riemann and (fully covariant) Ricci tensors is 1, that is, they are dimensionless. Some geometric objects or operators may be unfamiliar to researchers working in general relativity only; their discussion may simply be skipped by these readers.

The absolute dimension of the metric tensor is discussed in § 9. The literature on general relativity presents two standard choices for it. The absolute dimension of the stress-energy-momentum tensor is discussed in § 10. The dimension of the constant in the Einstein equation is derived in § 11, and the two standard results from the literature are recovered.

The final § 12 gives a summary of the main results, with some additional comments.

This Note assumes some familiarity with basic tensor calculus and related notions, for example of co- and contravariance, symmetry and antisymmetry, contraction. Some passages assume familiarity with the exterior calculus of differential forms. Above all, familiarity with the intrinsic presentation of differential geometry is assumed; but the general ideas should be understandable even without such familiarity. The notation used here is briefly presented in § 2.

Finally, quoting Truesdell & Toupin<sup>4</sup>, ‘dimensional analysis remains a controversial and somewhat obscure subject. We do not attempt a complete presentation here.’

## 2 Notation

For the notation in dimensional analysis I use ISO conventions:  $\dim(\mathbf{A})$  is the dimension of the quantity  $\mathbf{A}$ , and among the base quantities are mass  $M$ , length  $L$ , time  $T$ , temperature  $\Theta$ , electric current  $I$ . Note that I don’t discuss *units* – it doesn’t matter here whether the unit for length is the metre or the centimetre, for example.

Since the type of a tensor – that is, the number and ordering of its covariant and contravariant ‘slots’<sup>5</sup> – is important in dimensional analysis, I often denote it explicitly with a notation like  $\mathbf{A}^{**}$  to indicate that  $\mathbf{A}$  is covariant in its first slot and contravariant in its second and

---

<sup>4</sup> Truesdell & Toupin 1960 Appendix § 7 footnote 4. <sup>5</sup> Misner et al. 1973 § 3.2.

third slots. Its components would thus be  $(A_i^{jk})$ . For brevity I'll call this a 'co-contra-contravariant' tensor, with an obvious naming generalization for other types. In spirit this notation is similar to the abstract-index one of Penrose & Rindler<sup>6</sup>; but I avoid using letters for the indices, so that the notational difference between a tensor and its components, which is very important for our discussion, doesn't disappear.

I denote the contraction of the  $\alpha$ th and  $\beta$ th slots (which must have opposite variant types) of a tensor  $\mathbf{A}$  by  $\text{tr}_{\alpha\beta} \mathbf{A}$ , and the transposition (swapping) of the  $\alpha$ th and  $\beta$ th slots with  $\mathbf{A}^{\top\alpha\beta}$ . In index notation these operations are

$$\begin{array}{c} \alpha\text{th slot} \\ A \dots \overset{i}{\dots} \dots j \dots \\ \beta\text{th slot} \end{array} \quad \text{and} \quad \begin{array}{c} \alpha\text{th slot} \\ A \dots \overset{i}{\dots} j \dots \\ \beta\text{th slot} \end{array} \mapsto \begin{array}{c} \beta\text{th slot} \\ A \dots j \dots \overset{i}{\dots} \dots \\ \alpha\text{th slot} \end{array}.$$

### 3 Intrinsic view of differential-geometric objects: brief reminder

From the intrinsic point of view, a tensor is defined by its geometric properties. For example, a vector field  $v \equiv v(\cdot)$  is an object that operates on functions defined on the (spacetime) manifold, yielding new functions, with the properties  $v(af + bg) = av(f) + bv(g)$  and  $v(fg) = v(f)g + fv(g)$  for all functions  $f, g$  and reals  $a, b$ . A covector field (1-form)  $\omega$  is an object that operates on vector fields, yielding functions ('duality'), with the property  $\omega(fu + gv) = f\omega(u) + g\omega(v)$  for all vector fields  $u, v$  and functions  $f, g$ . The sum of vector or covector fields, and their products by functions – let's call this 'linearity' – are defined in an obvious way. Tensors are constructed from these objects.

A system of coordinates  $(x^i)$  is just a set of linearly independent functions. This set gives rise to a set of vectors fields  $\left(\frac{\partial}{\partial x^i}\right)$  and to a set of covector fields  $(dx^i)$  by the obvious requirements that  $\frac{\partial}{\partial x^i}(x^j) = \delta_i^j$  and  $dx^i\left(\frac{\partial}{\partial x^j}\right) = \delta_j^i$ . These two sets can be used as bases to express all other vectors and covectors as linear combinations. A vector field  $v$  can thus be written as

$$v \equiv \sum_i v^i \frac{\partial}{\partial x^i} \equiv v^i \frac{\partial}{\partial x^i}, \quad (1)$$

where the functions  $v^i := v(x^i)$  are its components with respect to the basis  $\left(\frac{\partial}{\partial x^i}\right)$ . Analogously for a covector field.

<sup>6</sup> Penrose & Rindler 2003 § 2.2.

For the presentation of the intrinsic view I recommend the excellent texts by Choquet-Bruhat et al. (1996), Boothby (2003), Abraham et al. (1988), Bossavit (1991), Burke (1987; 1980 ch. 2), and more on the general-relativity side Misner et al. (1973 ch. 9),ourgoulhon (2012 ch. 2), Penrose & Rindler (2003 ch. 4).

## 4 Coordinates

From a physical point of view, a coordinate is just a function that associates a value of a physical quantity with every event in a region (the domain of the coordinate chart) of spacetime. Together with the other coordinates, such function allows us to uniquely identify every event within that region. Any physical quantity will do: the distance from something, the time elapsed since something, an angle, an energy density, the strength of a magnetic flux, a temperature, and so on. A coordinate can thus have any dimension: length  $L$ , time  $T$ , angle  $1$ , energy density  $E := ML^{-1}T^{-2}$ , magnetic flux  $\Phi := ML^2T^{-2}I^{-1}$ , temperature  $\Theta$ , and so on.

The functional relation between two sets of coordinates must of course be dimensionally consistent. For example, if  $\dim(x^0) = T$  and  $\dim(x^1) = L$ , and we introduce a coordinate  $\xi(x^0, x^1)$  with dimension 1, additive in the previous two, then we must have  $\xi = ax^0 + bx^1$  with  $\dim(a) = T^{-1}$  and  $\dim(b) = L^{-1}$ .

## 5 Tensors: absolute dimension and components' dimensions

Consider a system of coordinates  $(x^i)$  with dimensions  $(X_i)$ , and the ensuing sets of covector fields (1-forms)  $dx^i$  and of vector fields  $\left(\frac{\partial}{\partial x^i}\right)$ , bases for the cotangent and tangent spaces. Their tensor products are bases for the tangent spaces of higher tensor types.

The differential  $dx^i$  traditionally has the same dimension as  $x^i$ :  $\dim(dx^i) = X_i$ , and the vector  $\frac{\partial}{\partial x^i}$  traditionally has the inverse dimension:  $\dim \frac{\partial}{\partial x^i} = X_i^{-1}$ .

For our discussion let's take a concrete example: a contra-covariant tensor field  $\mathbf{A} \equiv \mathbf{A}^\bullet_\bullet$ . The discussion generalizes to tensors of other types in an obvious way.

The tensor  $\mathbf{A}$  can be expanded in terms of the basis vectors and covectors, as mentioned in § 3:

$$\mathbf{A} = A^i_j \frac{\partial}{\partial x^i} \otimes dx^j \equiv A^0_0 \frac{\partial}{\partial x^0} \otimes dx^0 + A^0_1 \frac{\partial}{\partial x^0} \otimes dx^1 + \dots \quad (2)$$

Each function

$$A^i_j := \mathbf{A} \left( dx^i, \frac{\partial}{\partial x^j} \right) \quad (3)$$

is a *component* of the tensor in this coordinate system.

To make dimensional sense, all terms in the sum (2) must have the same dimension. This is possible only if the generic component  $A^i_j$  has dimension

$$\dim(A^i_j) = A X_i X_j^{-1}, \quad (4)$$

where  $A$  is *common* to all components. In fact, the  $X_i X_j^{-1}$  term cancels the  $X_i^{-1} X_j$  term coming from  $\frac{\partial}{\partial x^i} \otimes dx^j$  in the sum (2), and each summand therefore has dimension  $A$ .

The generalization of the formula above to tensors of other types is obvious:

$$\dim(A^{ij\dots}_{kl\dots}) = A X_i X_j \dots X_k^{-1} X_l^{-1} \dots,$$

where the ordering of the indices doesn't matter. (5)

For example, if we're using coordinates with dimensions

$$\dim(x^0) = \Theta, \quad \dim(x^1) = L, \quad \dim(x^2) = L, \quad \dim(x^3) = ML^{-1}T^{-2}, \quad (6)$$

then the components of  $\mathbf{A}$  have dimensions

$$\left( \dim(A^i_j) \right) = A \times \begin{pmatrix} 1 & L^{-1}\Theta & L^{-1}\Theta & M^{-1}L^2T^2\Theta \\ L\Theta^{-1} & 1 & 1 & M^{-1}L^2T^2 \\ L\Theta^{-1} & 1 & 1 & M^{-1}L^2T^2 \\ ML^{-1}T^{-2}\Theta^{-1} & ML^{-2}T^{-2} & ML^{-2}T^{-2} & 1 \end{pmatrix}. \quad (7)$$

Clearly most components have different dimensions. But this doesn't matter. What matters is that the sum (2) be dimensionally consistent. (Fokker<sup>7</sup>, for example, uses a metric tensor with components having different dimensions.)

---

<sup>7</sup> Fokker 1965 § VII.1 p. 88.

1 The dimension  $A$ , which is also the final dimension of the sum (2), is  
2 called the *absolute dimension*<sup>8</sup> of the tensor  $\mathbf{A}$ , and we write

$$3 \dim(\mathbf{A}) = A. \quad (8)$$

6 This is the intrinsic dimension of the tensor, independent of any coordin-  
7 ate system. It reflects the physical or operational<sup>9</sup> meaning of the tensor.  
8 We'll see an example of what this mean in § 9 with the metric tensor.

10 Different coordinate systems lead to different dimensions of the  
11 *components* of a tensor  $\mathbf{A}$ , but the absolute dimension of the tensor  
12 remains the same. Formula (5) for the dimensions of the components is  
13 consistent under changes of coordinates. For example, in new coordinates  
14  $(x^{i'})$  with dimensions  $(X_{i'})$ , the new components of  $\mathbf{A}$  are

$$16 A^{i'}_{j'} = A^i_j \frac{\partial x^{i'}}{\partial x^i} \frac{\partial x^j}{\partial x^{j'}} \quad (9)$$

19 and a quick check shows that  $\dim(A^{i'}_{j'}) = A X_{i'} X_{j'}^{-1}$ , consistent with the  
20 general formula (5).

21 In the following I'll drop the adjective 'absolute' when it's clear from  
22 the context.

## 25 6 Tensor operations

27 By the reasoning of the previous section, which simply applies standard  
28 dimensional considerations to the basis expansion (2), it's easy to find  
29 the resulting absolute dimension of various operations and operators on  
30 tensors and tensor fields.

31 Here is a summary of the dimensional rules for the main differential-  
32 geometric operations and operators, except for the covariant derivative,  
33 the metric, and related tensors, discussed more in depth in §§ 8–9  
34 below. Some of these rules are really definition or conventions, as briefly  
35 discussed in their description. The others can be proved; I only give  
36 a proof for one of them, leaving the other proofs as an exercise. For  
37 reference, in brackets I give the section of Choquet-Bruhat et al. (1996)  
38 where these operations are defined.

---

41 <sup>8</sup> Dorgelo & Schouten 1946; Schouten 1989 ch. VI.  
42 Sygne 1960a § A.2; Truesdell & Toupin 1960 §§ A.3–4.

<sup>9</sup> Bridgman 1958; see also

- The *tensor product* [III.B.5] multiplies dimensions:

$$\dim(\mathbf{A} \otimes \mathbf{B}) = \dim(\mathbf{A}) \dim(\mathbf{B}). \quad (10)$$

This is actually a definition or convention. In fact we tacitly used this rule already in § 5 for the tensors  $\frac{\partial}{\partial x^i} \otimes dx^j$  in the coordinate expansion (2). It is a natural definition, because for tensors of order 0 – that is, functions – the tensor product is just ordinary multiplication, and the dimension of a product is the product of the dimensions. And it is a definition that doesn't lead to inconsistencies.

- The *contraction* [III.B.5] or trace of the  $\alpha$ th and  $\beta$ th slots of a tensor has the same dimension as the tensor:

$$\dim(\text{tr}_{\alpha\beta} \mathbf{A}) = \dim(\mathbf{A}). \quad (11)$$

Note that the formula above only holds *without* raising or lowering indices; see § 9 for those operations.

This operation can be traced back to the duality of vectors and covectors mentioned in § 3: a covector field  $\omega$  operates linearly on a vector field  $v$  to yield a function  $f = \omega(v)$ . Also in this case we have that  $\dim(f) = \dim(\omega) \dim(v)$  by definition or convention, and the rule (11) follows from this convention. Also in this case this convention seems very natural, owing to the linearity properties of the trace, and doesn't lead to inconsistencies.

- The *transposition* (called 'building an isomer' by Schouten<sup>10</sup>) of the  $\alpha$ th and  $\beta$ th slots of a tensor has the same dimension as the tensor:

$$\dim(\mathbf{A}^{\tau_{\alpha\beta}}) = \dim(\mathbf{A}). \quad (12)$$

- The *Lie bracket* [III.B.3] of two vectors has the product of their dimensions:

$$\dim([u, v]) = \dim(u) \dim(v). \quad (13)$$

In fact, in coordinates ( $x^i$ ) the bracket can be expressed as

$$[u, v] = \left( u^j \frac{\partial v^i}{\partial x^j} - v^j \frac{\partial u^i}{\partial x^j} \right) \frac{\partial}{\partial x^i}, \quad (14)$$

and equating the dimensions of the left and right side, considering that

$$\dim(u^i) = \dim(u) X_i, \quad \dim(v^i) = \dim(v) X_i, \quad (15)$$

we find again that all  $X$  terms cancel out leaving the result (13).

<sup>10</sup> Schouten 1954 § I.3 p. 13; 1989 § II.4 p. 20.



- The *Lie derivative* [III.C.2] of a tensor with respect to a vector field has the product of the dimensions of the tensor and of the vector:

$$\dim(L_v \mathbf{A}) = \dim(v) \dim(\mathbf{A}). \quad (16)$$

Regarding operations and operators on differential forms:

- The *exterior product* [IV.A.1] of two differential forms multiplies their dimensions:

$$\dim(\omega \wedge \tau) = \dim(\omega) \dim(\tau). \quad (17)$$

- The *interior product* [IV.A.4] of a vector and a form multiplies their dimensions:

$$\dim(i_v \omega) = \dim(v) \dim(\omega). \quad (18)$$

- The *exterior derivative* [IV.A.2] of a form has the same dimension of the form:

$$\dim(d\omega) = \dim(\omega). \quad (19)$$

This can be proven using the identity  $d i_v + i_v d = L_v$  or similar identities<sup>11</sup> together with eqs (16) and (18).

- The *integral* [IV.B.1] of a form over a submanifold has the same dimension as the form:

$$\dim(\int_c \omega) = \dim(\omega). \quad (20)$$

## 7 Curves and integral curves

Consider a curve into spacetime,  $C: s \mapsto P(s)$ , with the parameter  $s$  having dimension  $\dim(s) = S$ .

---

<sup>11</sup> Curtis & Miller 1985 ch. 9 p. 180 Theorem 9.78; Abraham et al. 1988 § 6.4 Theorem 6.4.8.

If we consider the events of the spacetime manifold as dimensionless quantities, then the dimension of the tangent or velocity vector  $\dot{C}$  to the curve is

$$\dim(\dot{C}) = S^{-1}, \quad (21)$$

owing to the definition<sup>12</sup>

$$\dot{C} := \frac{\partial x^i[C(s)]}{\partial s} \frac{\partial}{\partial x^i}. \quad (22)$$

This has a quirky but interesting consequence. Given a vector field  $v$  we say that  $C$  is an integral curve for it if

$$v = \dot{C} \quad (23)$$

(or more precisely  $v_{C(s)} = \dot{C}_{C(s)}$  in usual differential-geometric notation<sup>13</sup>) at all events  $C(s)$  in the image of the curve. From the point of view of dimensional analysis this definition can only be valid if  $v$  has dimension  $S^{-1}$ . If  $v$  and  $s^{-1}$  have different dimensions – a case which can happen for physical reasons – the condition (22) must be modified into  $v = k\dot{C}$ , where  $k$  is a possibly dimensionful constant. This is equivalent to considering an affine and dimensional reparameterization of  $C$ .

## 8 Connection, covariant derivative, curvature tensors

Consider an arbitrary connection<sup>14</sup> with covariant derivative  $\nabla$ . For the moment we don't assume the presence of any metric structure.

The covariant derivative of the product  $fv$  of a function and a vector satisfies<sup>15</sup>

$$\nabla(fv) = df \otimes v + f\nabla v. \quad (24)$$

The first summand, from formulae (19) and (10), has dimension  $\dim(f)\dim(v)$ ; for dimensional consistency this must also be the dimension of the second summand. Thus

$$\dim(\nabla v) = \dim(v). \quad (25)$$

It follows that the *directional* covariant derivative  $\nabla_u$  has dimension

$$\dim(\nabla_u v) = \dim(u)\dim(v), \quad (26)$$

---

<sup>12</sup> Choquet-Bruhat et al. 1996 § III.B.1; Boothby 2003 § IV.(1.9). <sup>13</sup> Choquet-Bruhat et al. 1996 § III.B.1. <sup>14</sup> Choquet-Bruhat et al. 1996 § V.B. <sup>15</sup> Choquet-Bruhat et al. 1996 § V.B.1.

and by its derivation properties<sup>16</sup> we see that formula (25) extends from vectors to tensors of arbitrary type.

In the coordinate system  $(x^i)$ , the action of the covariant derivative is carried by the *connection coefficients* or Christoffel symbols  $(\Gamma^i_{jk})$  defined by

$$\nabla \frac{\partial}{\partial x^k} = \Gamma^i_{jk} dx^j \otimes \frac{\partial}{\partial x^i}. \quad (27)$$

From this equation and the previous ones it follows that these coefficients have dimensions

$$\dim(\Gamma^i_{jk}) = X_i X_j^{-1} X_k^{-1}. \quad (28)$$

The *torsion*  $\tau^{\bullet..}$ , *Riemann curvature*  $R^{\bullet...}$ , and *Ricci curvature*  $Ric..$  tensors are defined by<sup>17</sup>

$$\tau(u, v) := \nabla_u v - \nabla_v u - [u, v], \quad (29)$$

$$R(u, v; w) := \nabla_u \nabla_v w - \nabla_v \nabla_u w - \nabla_{[u, v]} w, \quad (30)$$

$$Ric.. := \text{tr}_{13} R^{\bullet...}. \quad (31)$$

From these definitions and the results of § 6 we find the dimensional requirements

$$\dim(\tau^{\bullet..}) \dim(u) \dim(v) = \dim(u) \dim(v), \quad (32)$$

$$\dim(R^{\bullet...}) \dim(u) \dim(v) \dim(w) = \dim(u) \dim(v) \dim(w), \quad (33)$$

$$\dim(Ric..) = \dim(R^{\bullet...}), \quad (34)$$

which imply that *torsion*, *Riemann curvature*, and *Ricci curvature* are dimensionless:

$$\dim(\tau^{\bullet..}) = \dim(R^{\bullet...}) = \dim(Ric..) = 1. \quad (35)$$

The exact contra- and covariant type used above for these tensors is very important in these equations. If we raise any of their indices using a metric, their dimensions will generally change.

Misner et al. (1973 pp. 35, 407) say that ‘curvature’, the Riemann or Einstein tensors (for which see § 9 below), has dimension  $L^{-2}$ , a statement seemingly at variance with the dimensionless results (35). But I believe that they refer to the *components* of those tensors, in specific coordinates

<sup>16</sup> Choquet-Bruhat et al. 1996 § V.B.1 p. 303.

<sup>17</sup> Choquet-Bruhat et al. 1996 § V.B.1.

of dimension  $L$ , and using geometrized units. In such specific coordinates every component  $R^i_{jkl}$  does indeed have dimension  $L^{-2}$ , according to the general formula (5), if and only if the *absolute* dimension of  $\mathbf{R}$  is unity,  $\dim(\mathbf{R}) = 1$ . So I believe that there's no real contradiction with Mister et al.'s statement and the results (35). This possible misunderstanding shows that it's important to distinguish between absolute dimensions, which don't depend on any specific coordinate choice, and component dimensions, which do.

The formulae above are also valid if a metric is defined and the connection is compatible with it. The connection coefficients in this case are defined in terms of the metric tensor, but using the results of § 9 it's easy to see that eqs (25), (26), (28), (35) still hold.

## 9 Metric and related tensors and operations

Let's now consider a metric tensor  $\mathbf{g}_{..}$ . What is its absolute dimension  $\dim(\mathbf{g})$ ? There seem to be two choices in the literature; both can be derived from the operational meaning of the metric.

Consider a (timelike) worldline  $s \mapsto C(s)$ ,  $s \in [a, b]$ , between events  $C(a)$  and  $C(b)$ . The metric tells us the *proper time*  $\Delta t$  elapsed for an observer having that worldline, according to the formula

$$\Delta t = \int_a^b ds \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|}. \quad (36)$$

From the results of § 6 this formula implies that  $T \equiv \dim(\Delta t) = \sqrt{\dim(\mathbf{g}_{..})}$ , (independently of the dimension of  $s$ ) and therefore

$$\dim(\mathbf{g}_{..}) = T^2. \quad (37)$$

Many authors<sup>18</sup>, however, prefer to include a dimensional factor  $1/c$  in front of the integral (36):

$$\Delta t = \frac{1}{c} \int_a^b ds \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|}, \quad (38)$$

thus obtaining

$$\dim(\mathbf{g}_{..}) = L^2. \quad (39)$$

---

<sup>18</sup> e.g. Fock 1964 § V.62 eq. (62.02); Curtis & Miller 1985 ch. 11 eq. (11.21); Rindler 1986 § 5.3 eq. (5.6); Hartle 2003 ch. 6 eq. (6.24).

1 The choice (39) seems also supported by the traditional expression  
2 for the ‘line element  $ds^2$ ’ as it appears in many works (for an exception,  
3 with dimension  $T^2$ , see Kilmister<sup>19</sup>),  
4

$$5 \quad ds^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2, \quad (40)$$

6 possibly with opposite signature. If the coordinates  $(t, x, y, z)$  have the  
7 dimensions suggested by their symbols, this formula has dimension  $L^2$ ,  
8 so that if we interpret ‘ $ds^2$ ’ as  $\mathbf{g}$  we find  $\dim(\mathbf{g}) = L^2$ . The line-element  
9 expression above often has an ambiguous differential-geometric meaning,  
10 however, because it may represent *the metric applied to some unspecified*  
11 *vector*, that is,  $\mathbf{g}(v, v)$ , where  $v$  is left unspecified<sup>20</sup>. In this case we have  
12  
13

$$14 \quad L^2 = \dim(\mathbf{g}) \dim(v)^2 \quad (41)$$

15 and the dimension of  $\mathbf{g}$  is ambiguous.  
16

17 The standard choices for  $\dim \mathbf{g}$  are thus  $T^2$  or  $L^2$ . My favourite choice  
18 is the first, (37), for reasons discussed by Synge and Bressan<sup>21</sup>. Synge  
19 gives a vivid summary:<sup>22</sup>  
20

21 We are now launched on the task of giving physical meaning to the Riemannian geometry [...]. It is indeed a Riemannian *chronometry* rather than  
22 *geometry*, and the word *geometry*, with its dangerous suggestion that we  
23 should go about measuring *lengths* with *yardsticks*, might well be abandoned  
24 altogether in the present connection  
25

26 In fact, to measure the proper time  $\Delta t$  defined above we only need to  
27 ensure that a clock has the worldline  $C$ , and then take the difference  
28 between the clock’s final and initial times. On the other hand, consider  
29 the case when the curve  $C$  is *spacelike*. Its proper length is still defined by  
30 the integral (36) (apart from a dimensional constant). Its measurement,  
31 however, is more involved than the timelike case. It requires dividing  
32 the curve into very short pieces, and having specially-chosen observers  
33 (orthogonal to the pieces) measure each piece. But the measurement  
34 of each piece actually relies on the measurement of *proper time*: each  
35 observer uses ‘radar distance’<sup>23</sup>, sending a lightlike signal which bounces  
36 back at the end of the piece and measuring the time it takes to come back.  
37 Even if rigid rods are used, their calibration still relies on a measurement  
38  
39

40 <sup>19</sup> Kilmister 1973 ch. II p. 25. <sup>20</sup> cf. Misner et al. 1973 Box 3.2 D p. 77. <sup>21</sup> Synge  
41 1960b §§ III.2–4; Bressan 1978 §§ 15, 18. <sup>22</sup> Synge 1960b § III.3 pp. 108–109. <sup>23</sup> Landau  
42 & Lifshitz 1996 § 84.

of time – this is also reflected in the current definition of the standard metre<sup>24</sup>.

The metric  $\mathbf{g}$  can be considered as an operator mapping vectors to covectors, which we can compactly write as  $\boldsymbol{\omega} = \mathbf{g}\mathbf{v}$  (instead of the cumbersome  $\boldsymbol{\omega} = \text{tr}_{23}(\mathbf{g} \otimes \mathbf{v})$ ). The *inverse metric tensor*  $\mathbf{g}^{-1\bullet\bullet}$  is then defined by the formula

$$\mathbf{g}^{-1}\mathbf{g} = \text{id}^{\bullet\bullet}, \quad \mathbf{g}\mathbf{g}^{-1} = \text{id}_{\bullet\bullet}, \quad (42)$$

so that, obviously,

$$\dim(\mathbf{g}^{-1}) = \dim(\mathbf{g})^{-1}. \quad (43)$$

The *metric volume element*<sup>25</sup> in spacetime is a 4-form  $\gamma$ , equivalent to a completely antisymmetric tensor  $\gamma_{\bullet\bullet\bullet\bullet}$ , such that  $\gamma(e_0, e_1, e_2, e_3) = 1$  for every set of positively-oriented orthonormal vector fields ( $e_k$ ), that is, such that  $\mathbf{g}(e_k, e_l) = \pm\delta_{kl}$  (remember that the orientation is not determined by the metric). It has only one non-zero component, given by the square root of the determinant of the (positively ordered) components ( $g_{ij}$ ) of the metric:

$$\gamma = \sqrt{|\det(g_{ij})|} \, dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3. \quad (44)$$

From this expression and the results of § 6 it can be shown that, in spacetime,

$$\dim(\gamma) = \dim(\mathbf{g})^2 \equiv \begin{cases} \mathbb{T}^4 \\ \mathbb{L}^4 \end{cases} \quad \text{if } \dim(\mathbf{g}) := \begin{cases} \mathbb{T}^2 \\ \mathbb{L}^2 \end{cases}. \quad (45)$$

(This is also the dimension of the *density*  $|\gamma|$ , which, as opposed to the volume element, has the property that  $|\gamma|(e_0, e_1, e_2, e_3) = 1$  for all sets of orthonormal vector fields, not only positively-oriented ones.)

The operation of *raising or lowering an index* of a tensor represents a contraction of the tensor product of that tensor with the metric or the metric inverse, for example  $\mathbf{A}_{\bullet\bullet} \equiv \text{tr}_{13} \mathbf{A}^{\bullet\bullet} \otimes \mathbf{g}_{\bullet\bullet}$  and similarly for tensors of other types. Therefore

$$\dim(\mathbf{A}_{\bullet\bullet\bullet\bullet}) = \dim(\mathbf{A}_{\bullet\bullet\bullet\bullet}) \dim(\mathbf{g}), \quad \dim(\mathbf{A}_{\bullet\bullet\bullet\bullet}) = \dim(\mathbf{A}_{\bullet\bullet\bullet\bullet}) \dim(\mathbf{g})^{-1}. \quad (46)$$

The formulae for the covariant derivative (25), connection coefficients (28), and curvature tensors (35) remain valid for a connection

<sup>24</sup> BIPM 1983 p. 98; Giacomo 1984 p. 25.

<sup>25</sup> Abraham et al. 1988 § 6.2.

compatible with the metric. In this case the connection coefficients can be obtained from the metric by the formulae<sup>26</sup>

$$\Gamma^i_{jk} = \frac{1}{2} \left( \frac{\partial}{\partial x^k} g_{jl} + \frac{\partial}{\partial x^j} g_{kl} - \frac{\partial}{\partial x^l} g_{jk} \right) g^{li}, \quad (47)$$

and it's easily verified that the dimensions (28) of these coefficients still hold. Also the result for the curvature tensors (35) still holds, since their expressions in terms of the connection coefficients is the same with or without a metric.

The scalar curvature  $\rho$  and the Einstein tensor  $\mathbf{G}_\bullet$

$$\rho := \text{tr } \mathbf{Ric}_\bullet \equiv \text{tr}_{23}(\mathbf{Ric} \otimes \mathbf{g}^{-1}), \quad \mathbf{G}_\bullet := \mathbf{Ric}_\bullet - \frac{1}{2} \rho \text{ id}_\bullet \quad (48)$$

have therefore dimension

$$\dim(\rho) = \dim(\mathbf{G}_\bullet) = \dim(\mathbf{g})^{-1} \equiv \begin{cases} T^{-2} \\ L^{-2} \end{cases} \quad \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases} \quad (49)$$

## 10 Stress-energy-momentum tensor

To find the dimension of the stress-energy-momentum  $\mathbf{T}$ , or '4-stress' for short, let's start with the analysis of the (3-)stress  $\sigma$  in Newtonian mechanics. The stress  $\sigma$  is the projection of the 4-stress  $\mathbf{T}$  onto a spacelike tangent plane with respect to some observer<sup>27</sup>. If we assume that such spatial projection preserves the absolute dimension, then the 4-stress and the stress have the same absolute dimension.

In Newtonian mechanics the stress  $\sigma$  is an object that, integrated over the boundary of a body, gives the total surface force acting on the body (such integration requires a flat connection). This means that it must be represented by a 'force-valued' 2-form. Force, in turn, can be interpreted as an object that, integrated over a (spacelike) trajectory, gives an energy – the work done by the force along the trajectory. It's therefore a 1-form. Putting these two requirements together, the stress turns out to be a covector-valued 2-form, equivalent to a tensor  $\sigma_{\bullet\bullet}$  antisymmetric in its last two indices. Integrated over a surface, and then over a trajectory, it

<sup>26</sup> Choquet-Bruhat et al. 1996 § V.B.2. <sup>27</sup>ourgoulhon 2012 § 3.4.1; Smarr & York 1978; York 1979; Smarr et al. 1980; Wilson & Mathews 2007 § 1.3; the projection doesn't need to be orthogonal: Marsden & Hughes 1994 § 2.4; Hehl & Obukhov 2003 § B.1.4.

yields an energy. From § 6, integration of a form does not change the dimension of the form. Therefore

$$\dim(\sigma_{\bullet\bullet}) = E \equiv ML^2T^{-2}. \quad (50)$$

But usually the stress is represented by a co-contravariant tensor  $\sigma_{\bullet}^{\bullet}$ . This is obtained by contracting the last two slots of  $\sigma_{\bullet\bullet}$  with the inverse of the volume element of the 3-metric – this is the duality<sup>28</sup> between  $k$ -vectors and  $(n - k)$ -covectors induced by the metric (and an orientation choice), where  $n$  is the geometric dimension of the manifold. If we assume the Newtonian 3-metric to have dimension  $L^2$ , it can be shown similarly to § 9 that its volume element has dimension  $L^3$ , and the inverse volume element has dimension  $L^{-3}$ . Thus we obtain

$$\dim(\sigma_{\bullet}^{\bullet}) = EL^{-3} \equiv ML^{-1}T^{-2}, \quad (51)$$

an energy density (or ‘volumic energy’ according to ISO<sup>29</sup>).

Since the stress  $\sigma_{\bullet}^{\bullet}$  is the projection of  $T_{\bullet}^{\bullet}$  and the projection preserves the absolute dimension, we finally find that  $T_{\bullet}^{\bullet}$  also has the dimension of an energy density:

$$\dim(T_{\bullet}^{\bullet}) = EL^{-3} \equiv ML^{-1}T^{-2}. \quad (52)$$

Note that other co- or contravariant versions of the 4-stress will have different absolute dimension, because they’re obtained by lowering or raising indices. For example,  $\dim(T_{\bullet\bullet}) = \dim(T_{\bullet}^{\bullet}) \dim(g) = ML^{-1}$  if  $\dim(g) := T^2$ .

Let me add a passing remark. Even though in most texts the 4-stress is represented by a tensor of order 2, as above, its most fitting geometrical nature is still shrouded by mystery from a kinematic and dynamical point of view. There are indications that it could be more properly represented by a covector-valued 3-form (equivalent to a tensor  $T_{\bullet\bullet\bullet}$  antisymmetric in the last three slots), or by a 3-vector-valued 3-form (equivalent to a tensor  $T^{\bullet\bullet\bullet}$  antisymmetric in the first three and last three slots), for reasons connected with integration, similar to those mentioned above for the stress  $\sigma_{\bullet\bullet}$ . See for example the discussion about ‘ $T$ ’ by Misner et al. (1973 ch. 15), the works by Segev (2002; 1986; 1999; 2000a,b), the discussion by Burke (1987 § 41).

<sup>28</sup> Bossavit 1991 § 4.1.2. <sup>29</sup> ISO 2009 item A.6.2.



## 11 Einstein equation and Einstein's constant

We finally arrive at the Einstein equation,

$$\mathbf{G} = \kappa \mathbf{T} \quad (53)$$

where  $\kappa$  (sometimes seen with a minus<sup>30</sup> depending on the signature of the metric or on how the orientation of the stress is chosen) is Einstein's constant. For the dimension of  $\kappa$  we thus find

$$\dim(\kappa) = \dim(\mathbf{G}_{\bullet}^{\bullet}) \dim(\mathbf{T}_{\bullet}^{\bullet})^{-1} \equiv \begin{cases} M^{-1}L & \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases} \\ M^{-1}L^{-1}T^2 & \end{cases} \quad (54)$$

This constant can be obtained from the dimensions of Newton's gravitational constant  $\dim(G) = M^{-1}L^3T^{-2}$  (this is not the Einstein tensor  $\mathbf{G}$ !) and of the speed of light  $\dim(c) = LT^{-1}$  only in the following ways, with an  $8\pi$  factor coming from the Newtonian limit:

$$\kappa = \begin{cases} 8\pi G/c^2 & \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases} \\ 8\pi G/c^4 & \end{cases} \quad (55)$$

The second choice is by far the most common, consistently with the most common choice of  $\dim(\mathbf{g}) = L^2$  discussed before. The first choice appears for example in Fock<sup>31</sup> and Adler et al.<sup>32</sup>.

## 12 Summary and conclusions

We have seen that dimensional analysis in general relativity and differential geometry can be seamlessly done, following its usual standard rules, if we adopt a coordinate-free or coordinate-invariant view, typical of more modern texts. In this view, each tensor has an 'absolute' dimension (Schouten & Dorgelo's terminology) that doesn't depend on the dimensions of the coordinates. It's the intrinsic dimension of the tensor. It would therefore be more profitable to focus on the absolute dimension of a tensor rather than on the dimensions of its components. The dimension of a specific component is easily found by formula (5): it's the product of the absolute dimension, times the dimension of the

---

<sup>30</sup> e.g. Tolman 1949 § 78 eq. (78.3); Fock 1964 § 52 eq. (52.06); Rindler 2006 § 14.2 eq. (14.8). <sup>31</sup> Fock 1964 § 55 eqs (55.15) and (52.06). <sup>32</sup> Adler et al. 1975 § 10.5 eq. (10.98).

$i$ th coordinate function for each contravariant index  $i$ , times the inverse of the dimension of the  $j$ th coordinate function for each covariant index  $j$ .

This generalized dimensional analysis rests mainly on two conventions: the tensor product and the action of covectors on vectors behave analogously to the usual multiplication for the purposes of dimensional analysis.

We found or re-derived some essential results for general relativity, in particular that the Riemann  $R^{\bullet}...$  and Ricci  $Ric_{\bullet}$  tensors are dimensionless, and that the Einstein tensor  $G_{\bullet}^{\bullet}$  has the inverse dimension of the metric. Maybe these can be of importance for some current research involving scales and conformal factors<sup>33</sup>.

Since the dimensions of the components are usually different from the absolute dimension and depend on the coordinates, I recommend to avoid statements such as ‘the tensor  $A_i^{jk}$  has dimension  $X$ ’, because it leaves unclear whether ‘ $A_i^{jk}$ ’ is meant to represent the tensor in general (as in Penrose & Rindler’s notation), or to represent its set of components, or to represent just a specific component.

Finally, the above analysis, specific to general relativity, can be made along the same lines also for Newtonian mechanics.

## Thanks

... to prof. Ingemar Bengtsson for valuable comments on a previous draft. To the staff of the NTNU library for their always prompt support. To Mari, Miri, Emma for continuous encouragement and affection, and to Buster Keaton and Saitama for filling life with awe and inspiration. To the developers and maintainers of L<sup>A</sup>T<sub>E</sub>X, Emacs, AUC<sub>T</sub>E<sub>X</sub>, Open Science Framework, R, Python, Inkscape, Sci-Hub for making a free and impartial scientific exchange possible. ☒

This work is financially supported by the Kavli Foundation and the Centre of Excellence scheme of the Research Council of Norway (Roudi group).

<sup>33</sup> e.g. Röhr & Ugglä 2005; Cadoni & Tuveri 2019.

# Bibliography

- Abraham, R., Marsden, J. E., Ratiu, T. (1988): *Manifolds, Tensor Analysis, and Applications*, 2nd ed. (Springer, New York). First publ. 1983.
- Adler, R., Bazin, M., Schiffer, M. (1975): *Introduction to General Relativity*, 2nd ed. (McGraw-Hill, Tokyo). First publ. 1965.
- BIPM (Bureau international des poids et mesures) (1983): *17<sup>e</sup> conférence générale des poids et mesures (CGPM)*. (BIPM, Sèvres). <https://www.bipm.org/en/worldwide-metrology/cgpm/resolutions.html#comptes-rendus>.
- Boothby, W. M. (2003): *An Introduction to Differentiable Manifolds and Riemannian Geometry*, rev. 2nd ed. (Academic Press, Orlando, USA). First publ. 1975.
- Bossavit, A. (1991): *Differential Geometry: for the student of numerical methods in Electromagnetism*. [https://www.researchgate.net/publication/200018385\\_Differential\\_Geometry\\_for\\_the\\_student\\_of\\_numerical\\_methods\\_in\\_Electromagnetism](https://www.researchgate.net/publication/200018385_Differential_Geometry_for_the_student_of_numerical_methods_in_Electromagnetism).
- Bressan, A. (1978): *Relativistic Theories of Materials*. (Springer, Berlin).
- Bridgman, P. W. (1958): *The Logic of Modern Physics*, 8th pr. (Macmillan, New York). First publ. 1927.
- Burke, W. L. (1980): *Spacetime, Geometry, Cosmology*. (University Science Books, Mill Valley, USA).
- (1987): *Applied Differential Geometry*, repr. (Cambridge University Press, Cambridge). First publ. 1985.
- Cadoni, M., Tuveri, M. (2019): *Galactic dynamics and long-range quantum gravity*. *Phys. Rev. D* **100**<sup>2</sup>, 024029.
- Choquet-Bruhat, Y., DeWitt-Morette, C., Dillard-Bleick, M. (1996): *Analysis, Manifolds and Physics. Part I: Basics*, rev. ed. (Elsevier, Amsterdam). First publ. 1977.
- Curtis, W. D., Miller, F. R. (1985): *Differential Manifolds and Theoretical Physics*. (Academic Press, Orlando, USA).
- Dorgelo, H. B., Schouten, J. A. (1946): *On unities and dimensions*. [I, II, III]. *Verh. Kon. Akad. Wetensch. Amsterdam* **49**<sup>2, 3, 4</sup>, 123–131, 282–291, 393–403.
- Flügge, S., ed. (1960): *Handbuch der Physik: Band III/1: Prinzipien der klassischen Mechanik und Feldtheorie [Encyclopedia of Physics: Vol. III/1: Principles of Classical Mechanics and Field Theory]*. (Springer, Berlin).
- Fock [Fok], V. A. (1964): *The Theory of Space, Time and Gravitation*, 2nd rev. ed. (Pergamon, Oxford). Transl. by N. Kemmer. First publ. in Russian 1955.
- Fokker, A. D. (1965): *Time and Space, Weight and Inertia: A chronogeometrical introduction to Einstein's theory*. (Pergamon, Oxford). Transl. by D. Bijl, translation ed. by D. Field. First publ. in Dutch 1960.
- Giacomo, P. (1984): *News from the bipm*. *Metrologia* **20**<sup>1</sup>, 25–30.
- Gourgoulhon, É. (2012): *3+1 Formalism in General Relativity: Bases of Numerical Relativity*. (Springer, Heidelberg). First publ. 2007 as arXiv:gr-qc/0703035.
- Hartle, J. B. (2003): *Gravity: An Introduction to Einstein's General Relativity*. (Addison-Wesley, San Francisco).
- Hehl, F. W., Obukhov, Y. N. (2003): *Foundations of Classical Electrodynamics: Charge, Flux, and Metric*. (Birkhäuser, Boston).
- ISO (International Organization for Standardization) (2009): *ISO 80000-1:2009: Quantities and units 1: General*. International Organization for Standardization.
- Kilmister, C. W. (1973): *General Theory of Relativity*. (Pergamon, Oxford).

- Landau, L. D., Lifshitz [Lifšic], E. M. (1996): *The Classical Theory of Fields*, corr. repr. of the 4th English ed. (Butterworth-Heinemann, Oxford). Transl. from the 1987 seventh Russian edition by Morton Hamermesh. First publ. 1939.
- Marsden, J. E., Hughes, T. J. R. (1994): *Mathematical Foundations of Elasticity*, unabridged corr. republ. (Dover, New York). <http://resolver.caltech.edu/CaltechB00K:1983.002>. First publ. 1983.
- Misner, C. W., Thorne, K. S., Wheeler, J. A. (1973): *Gravitation*, repr. (W. H. Freeman and Company, New York). [https://archive.org/details/Gravitation\\_201803](https://archive.org/details/Gravitation_201803).
- Penrose, R., Rindler, W. (2003): *Spinors and Space-Time. Vol. 1: Two-spinor calculus and relativistic fields*, corr. repr. (Cambridge University Press, Cambridge). First publ. 1984.
- Rindler, W. (1986): *Essential Relativity: Special, General, and Cosmological*, rev. 2nd ed. (Springer, New York). First publ. 1969.
- (2006): *Relativity: Special, General, and Cosmological*, 2nd ed. (Oxford University Press, Oxford). First publ. 2001.
- Röhr, N., Uggla, C. (2005): *Conformal regularization of Einstein's field equations*. *Class. Quantum Grav.* **22**<sup>17</sup>, 3775–3787.
- Schouten, J. A. (1954): *Ricci-Calculus: An Introduction to Tensor Analysis and Its Geometrical Applications*, 2nd ed. (Springer, Berlin). First publ. in German 1924.
- (1989): *Tensor Analysis for Physicists*, corr. 2nd ed. (Dover, New York). First publ. 1951.
- Segev, R. (1986): *Forces and the existence of stresses in invariant continuum mechanics*. *J. Math. Phys.* **27**<sup>1</sup>, 163–170. <http://www.bgu.ac.il/~rsegev/Papers/JMP865.pdf>.
- (2000a): *The geometry of Cauchy's fluxes*. *Arch. Rational Mech. Anal.* **154**<sup>3</sup>, 183–198. <http://www.bgu.ac.il/~rsegev/Papers/TheGeometryofCauchyFluxes.pdf>.
- (2000b): *Notes on stresses for manifolds*. *Rend. Sem. Mat. Univ. Pol. Torino* **58**<sup>2</sup>, 199–206.
- (2001): *A correction of an inconsistency in my paper "Cauchy's theorem on manifolds"*. *J. Elasticity* **63**<sup>1</sup>, 55–59. Ssee Segev, Rodnay (1999).
- (2002): *Metric-independent analysis of the stress-energy tensor*. *J. Math. Phys.* **43**<sup>6</sup>, 3220–3231. <http://www.bgu.ac.il/~rsegev/Papers/MetricIndependent.pdf>.
- Segev, R., Rodnay, G. (1999): *Cauchy's theorem on manifolds*. *J. Elasticity* **56**<sup>2</sup>, 129–144. <http://www.bgu.ac.il/~rsegev/Papers/CauchyTheoremOnManifolds.pdf>. See also erratum Segev (2001).
- Smarr, L., Taubes, C., Wilson, J. R. (1980): *General relativistic hydrodynamics: the comoving, Eulerian, and velocity potential formalisms*. In: Tipler (1980), ch. 11, 157–183.
- Smarr, L., York Jr., J. W. (1978): *Kinematical conditions in the construction of spacetime*. *Phys. Rev. D* **17**<sup>10</sup>, 2529–2551.
- Smarr, L. L., ed. (1979): *Sources of Gravitational Radiation*. (Cambridge University Press, Cambridge).
- Synge, J. L. (1960a): *Classical Dynamics*. In: Flügge (1960), I–VII, 1–225, 859–902.
- (1960b): *Relativity: The General Theory*. (North-Holland, Amsterdam).
- Tipler, F. J., ed. (1980): *Essays in General Relativity: A Festschrift for Abraham Taub*. (Academic Press, New York).
- Tolman, R. C. (1949): *Relativity, Thermodynamics and Cosmology*. (Oxford University Press). First publ. 1934.
- Truesdell III, C. A., Toupin, R. A. (1960): *The Classical Field Theories*. In: Flügge (1960), I–VII, 226–902. With an appendix on invariants by Jerald LaVerne Ericksen.
- Wilson, J. R., Mathews, G. J. (2007): *Relativistic Numerical Hydrodynamics*, repr. (Cambridge University Press, Cambridge). First publ. 2003.
- York Jr., J. W. (1979): *Kinematics and dynamics of general relativity*. In: Smarr (1979), 83–126.