

Dimensional analysis on differential manifolds

[draft]

P.G.L. Porta Mana

[<pgl@portamana.org>](mailto:pgl@portamana.org)

13 December 2019; updated 10 January 2020

Some notes on dimensional analysis on differential manifolds, with an eye on general relativity and the Einstein equation.

Note: Dear Reader & Peer, this manuscript is being peer-reviewed by you. Thank you.

1 ***

‘Dimensional analysis remains a controversial and somewhat obscure subject. We do not attempt a complete presentation here.’¹

There seem to be insecurity and wrong ideas among some students and even researchers in relativity, regarding the dimensions of tensors, of tensor components, and of dimensional constants in equations. I’ve met, for example, with the statement that the components of a tensor should all have the same dimension; and with calculations of the dimensions of curvature tensors starting from coordinates with dimensions of length. That statement is wrong, and that procedure is unnecessary.

Several factors probably cause or contribute to such difficulties. Modern texts in Lorentzian and general relativity commonly use geometrized units. They say that to find the dimension of some constant in a tensorial equations it’s sufficient to compare the dimensions of the tensors in the equation. But this is not so immediate, because some tensors don’t have universally agreed dimensions – prime example the 4-metric tensor. Older texts often use coordinates with dimension of length. They even multiply a timelike coordinate or some tensorial components by c , thus giving the impression that coordinates should always be lengths and that the components of a tensor would all have the same dimension.²

In this note I want to clarify some misconceptions about dimensional analysis in differential manifolds, and to illustrate a simple way of reasoning to solve dimensional-analysis doubts and problems. This way

¹ Truesdell et al. 1960 Appendix § 7 footnote 4. eq. (32.15).

² e.g. Landau et al. 1996 § 32

of reasoning relies on the coordinate-free, *intrinsic* view of tensors and other differential-geometrical objects.

Let's start from more general facts about dimensional analysis on differential manifolds.

For dimensional analysis I use ISO conventions and notation. I sometimes use notation such as $\mathbf{T} \cdot$ to indicate that the tensor \mathbf{T} is covariant in its first slot and contravariant in its second; I call this a 'co-contra-variant tensor'.

[check³]

2 Intrinsic view of differential-geometric objects: brief reminder

From the intrinsic point of view, a tensor is defined by its geometric properties. For example, a vector field $v \equiv v(\cdot)$ is an object that operates on functions defined on the manifold, yielding new functions, with the properties $v(af + bg) = av(f) + bv(g)$ and $v(fg) = v(f)g + fv(g)$ for all functions f, g and reals a, b . A covector field (1-form) ω is an object that operates on vector fields, yielding functions ('duality'), with the property $\omega(fu + gv) = f\omega(u) + g\omega(v)$ for all vector fields u, v and functions f, g . The sum of vector or covector fields, and their products by functions – let's call this 'linearity' – are defined in an obvious way. Tensors are constructed from these objects.

A system of coordinates (x^i) is just a set of linearly independent functions. This set gives rise to a set of vectors fields $\left(\frac{\partial}{\partial x^i}\right)$ and to a set of covector fields (dx^i) by the obvious requirements that $\frac{\partial}{\partial x^i}(x^j) = \delta_i^j$ and $dx^i\left(\frac{\partial}{\partial x^j}\right) = \delta_j^i$. These two sets can be used as bases to express all other vectors and covectors as linear combinations. A vector field v can thus be written as

$$v \equiv \sum_i v^i \frac{\partial}{\partial x^i} \equiv v^i \frac{\partial}{\partial x^i}, \quad (1)$$

where the *functions* $v^i := v(x^i)$ are its components with respect to the basis $\left(\frac{\partial}{\partial x^i}\right)$. Analogously for a covector field.

For the full presentation of the intrinsic view I recommend the excellent texts by Choquet-Bruhat et al. (1996), Boothby (2003), Marsden et al.⁴, Bossavit (1991), and more on the general-relativity side Burke

³ Aldersley 1977.

⁴ Marsden et al. 2007.

(1980 ch. 2), Misner et al. (1973 ch. 9), Gourgoulhon (2012 ch. 2), Penrose et al. (2003 ch. 4).

 check⁵

3 Coordinates

From a physical point of view, a coordinate is just a function that associates a value of a physical quantity with every event in a region (the domain of the coordinate chart) of spacetime. Together with the other coordinates, such function allows us to uniquely identify every event within that region. Any physical quantity will do: the distance from something, the time elapsed since something, an angle, an energy density, the strength of a magnetic flux, a temperature, and so on. A coordinate can thus have any dimensions: length L , time T , angle 1 , energy density $E := L^{-1}MT^{-2}$, magnetic flux $\Phi := L^2MT^{-2}I^{-1}$, temperature Θ , and so on.

The dimensions of the coordinates don't matter, as we'll now see.

4 Tensors

Consider a system of coordinates (x^i) with dimensions (X_i) , and the ensuing sets of covector fields (1-form) dx^i and of vector fields $(\frac{\partial}{\partial x^i})$, bases for the cotangent and tangent spaces. Their tensor products are bases for the tangent spaces of higher tensor types.

The differential dx^i traditionally has the same dimension as x^i : $\dim(dx^i) = X_i$, and the operator (a vector) $\frac{\partial}{\partial x^i}$ traditionally has the inverse dimension: $\dim \frac{\partial}{\partial x^i} = X_i^{-1}$. We'll see later that these conventions are self-consistent.

For our discussion let's take a concrete example: a contra-co-tensor field $\mathbf{A} \equiv \mathbf{A}^\bullet$. The discussion generalizes to tensors of other types in an obvious way.

The tensor \mathbf{A} can be expanded in terms of the basis vectors and covectors, as mentioned in § 2:

$$\mathbf{A} = A^i_j \frac{\partial}{\partial x^i} \otimes dx^j \equiv A^0_0 \frac{\partial}{\partial x^0} \otimes dx^0 + A^0_1 \frac{\partial}{\partial x^0} \otimes dx^1 + \dots \quad (2)$$

Each function

$$A^i_j := \mathbf{A}\left(dx^i, \frac{\partial}{\partial x^j}\right) \quad (3)$$

⁵ Fischer et al. 1972.

is a component of the tensor in this coordinate system.

To make dimensional sense, all terms in the sum (2) must have the same dimension. This is possible only if the generic component A^i_j has dimension

$$\dim(A^i_j) = A X_i X_j^{-1}, \quad (4)$$

where A is common to all components. Suppose for example that we're using coordinates with dimensions

$$\dim(x^0) = \Theta, \quad \dim(x^1) = L, \quad \dim(x^2) = L, \quad \dim(x^3) = L^{-1}MT^{-2}; \quad (5)$$

then the components of \mathbf{A} have dimensions

$$(\dim(A^i_j)) = A \times \begin{pmatrix} 1 & L^{-1}\Theta & L^{-1}\Theta & LM^{-1}T^2\Theta \\ L\Theta^{-1} & 1 & 1 & L^2M^{-1}T^2 \\ L\Theta^{-1} & 1 & 1 & L^2M^{-1}T^2 \\ L^{-1}MT^{-2}\Theta^{-1} & L^{-2}MT^{-2} & L^{-2}MT^{-2} & 1 \end{pmatrix}. \quad (6)$$

The dimension A , which is also the dimension of the sum (2), is called the *absolute dimension*⁶ of the tensor \mathbf{A} , and we write

$$\dim(\mathbf{A}) = A. \quad (7)$$

This is the intrinsic dimension of the tensor, independent of any coordinate system. It reflects the physical or operational⁷ meaning of the tensor. We'll see an example of what this mean in § 8.

Different coordinate systems lead to different dimensions of the *components* of \mathbf{A} , but its absolute dimension remains the same. Formula (4) for the dimensions of the components is consistent under changes of coordinates. For example, in new coordinates (x'^k) with dimensions (X'_k) , the new components of \mathbf{A} are

$$A'^k_l = A^i_j \frac{\partial x'^k}{\partial x^i} \frac{\partial x^j}{\partial x'^l} \quad (8)$$

and a quick check shows that $\dim(A'^k_l) = A X'_k X'^{-1}_l$, consistent with the general formula (4).

In the following I'll drop the adjective 'absolute' when it's clear from the context.

⁶ Dorgelo et al. 1946; Schouten 1989 ch. VI. § A.2; Truesdell et al. 1960 §§ A.3–4.

⁷ Bridgman 1958; see also Synge 1960a

5 Tensor operations

By the reasoning of the previous section, which simply applies standard dimensional considerations to the basis expansion (2), it's easy to find out the resultant absolute dimension of various operations and operators on tensors and tensor fields.

The tensor product of \mathbf{A}^{\cdot} and \mathbf{B}_{\cdot} , for example, can be written as the sum

$$\mathbf{A} \otimes \mathbf{B} = A^i_j B_{kl}^m \frac{\partial}{\partial x^i} \otimes dx^j \otimes dx^k \otimes dx^l \otimes \frac{\partial}{\partial x^m} \quad (9)$$

from which it follows that

$$\dim(A^i_j B_{kl}^m) = A B X_i X_j^{-1} X_k^{-1} X_l^{-1} X_m \quad (10)$$

with $A = \dim(\mathbf{A})$ and $B = \dim(\mathbf{B})$. The absolute dimension of $\mathbf{A} \otimes \mathbf{B}$ is therefore $AB \equiv \dim(\mathbf{A}) \dim(\mathbf{B})$.

Here is then a summary of the dimensional results of the main differential-geometric operations and operators, except for the covariant derivative and related tensors, discussed more in depth in § 7 below. In brackets I give the section of Choquet-Bruhat et al. (1996) where they are defined.

- *Tensor multiplication* [III.B.5] multiplies dimensions:

$$\dim(\mathbf{A} \otimes \mathbf{B}) = \dim(\mathbf{A}) \dim(\mathbf{B}). \quad (11)$$

- The *contraction* [III.B.5] of the i th and j th slots (one covariant and one contravariant) of a tensor has the same dimension as the tensor:

$$\dim(\text{tr}_{ij} \mathbf{A}) = \dim(\mathbf{A}). \quad (12)$$

Note that this only holds *without* raising or lowering indices.

- The *transposition* (swapping) of the i th and j th slots of a tensor has the same dimension as the tensor:

$$\dim(\mathbf{A}^{\top ij}) = \dim(\mathbf{A}). \quad (13)$$

- The *Lie bracket* [III.B.3] of two vectors has the product of their dimensions:

$$\dim([\mathbf{u}, \mathbf{v}]) = \dim(\mathbf{u}) \dim(\mathbf{v}). \quad (14)$$

- The *Lie derivative* [III.C.2] of a tensor with respect to a vector field has the product of the dimensions of the tensor and of the vector:

$$\dim(L_v \mathbf{A}) = \dim(v) \dim(\mathbf{A}). \quad (15)$$

Regarding operations and operators on differential forms:

- The *exterior product* [IV.A.1] of two differential forms multiplies their dimensions:

$$\dim(\omega \wedge \tau) = \dim(\omega) \dim(\tau). \quad (16)$$

- The *interior product* [IV.A.4] of a vector and a form multiplies their dimensions:

$$\dim(i_v \omega) = \dim(v) \dim(\omega). \quad (17)$$

- The *exterior derivative* [IV.A.2] of a form has the same dimension of the form:

$$\dim(d\omega) = \dim(\omega). \quad (18)$$

This can be proven using the identity $d i_v + i_v d = L_v$ or similar identities⁸ together with eqs (15) and (17).

- The *integral* [IV.B.1] of a form over a submanifold has the same dimension as the form:

$$\dim(\int_c \omega) = \dim(\omega). \quad (19)$$

The resultant absolute dimensions of other operators, for example the determinant⁹, can be obtained by similar reasoning.

6 Curves and integral curves

Consider a curve into spacetime, $c: s \mapsto P(s)$, with the parameter s having dimension $\dim(s) = S$.

⁸ Curtis et al. 1985 ch. 9 p. 180 Theorem 9.78; Marsden et al. 2007 § 7.4 Theorem 7.4.8.

⁹ Marsden et al. 2007 § 7.2.

If we consider the manifold as a dimensionless quantity (see §*** for what I mean by this), then the dimension of the tangent or velocity vector \dot{C} to the curve is

$$\dim(\dot{C}) = S^{-1}, \quad (20)$$

owing to the definition¹⁰

$$\dot{C} := \frac{\partial x^i[C(s)]}{\partial s} \frac{\partial}{\partial x^i}. \quad (21)$$

This has a quirky but interesting consequence. Given a vector field v we say that C is an integral curve for it if

$$v = \dot{C} \quad (22)$$

(or more precisely $v_{C(s)} = \dot{C}_{C(s)}$ in usual differential-geometric notation¹¹) at all events $C(s)$ in the image of the curve. From the point of view of dimensional analysis this definition can only be valid if v has dimension S^{-1} . If v and s^{-1} have different dimensions – a case which can happen for physical reasons – the condition (21) must be modified into $v = k\dot{C}$, where k is a possibly dimensionful constant. This is equivalent to considering an affine and dimensional reparameterization of C .

7 Connection, covariant derivative, curvature tensors

Consider an arbitrary connection¹² with covariant derivative ∇ . For the moment we don't assume the presence of any metric structure.

The covariant derivative of the product fv of a function and a vector satisfies¹³

$$\nabla(fv) = df \otimes v + f\nabla v. \quad (23)$$

The first summand, from formulae (18) and (11), has dimension $\dim(f)\dim(v)$; for dimensional consistency this must also be the dimension of the second summand. Thus

$$\dim(\nabla v) = \dim(v). \quad (24)$$

It follows that the *directional* covariant derivative has dimension

$$\dim(\nabla_u v) = \dim(u)\dim(v), \quad (25)$$

¹⁰ Choquet-Bruhat et al. 1996 § III.B.1; Boothby 2003 § IV.(1.9). ¹¹ Choquet-Bruhat et al. 1996 § III.B.1. ¹² Choquet-Bruhat et al. 1996 § V.B. ¹³ Choquet-Bruhat et al. 1996 § V.B.1.

and by its derivation properties¹⁴ we see that formula (24) extends from vectors to tensors of arbitrary type.

In the coordinate system (x^i) , the action of the covariant derivative is carried by the *connection coefficients* or Christoffel symbols (Γ^i_{jk}) defined by

$$\nabla \frac{\partial}{\partial x^k} = \Gamma^i_{jk} dx^j \otimes \frac{\partial}{\partial x^i}. \quad (26)$$

From this equation and the previous ones it follows that these coefficients have dimensions

$$\dim(\Gamma^i_{jk}) = X_i X_j^{-1} X_k^{-1}. \quad (27)$$

The *torsion* $\tau^*_{..}$, *Riemann curvature* $R^*_{...}$, and *Ricci curvature* $Ric_{..}$ tensors are defined by¹⁵

$$\tau(u, v) := \nabla_u v - \nabla_v u - [u, v], \quad (28)$$

$$R(u, v; w) := \nabla_u \nabla_v w - \nabla_v \nabla_u w - \nabla_{[u, v]} w, \quad (29)$$

$$Ric_{..} := \text{tr}_{13} R^*_{...}. \quad (30)$$

From these definitions and the results of § 5 we find the dimensional requirements

$$\dim(\tau^*_{..}) \dim(u) \dim(v) = \dim(u) \dim(v), \quad (31)$$

$$\dim(R^*_{...}) \dim(u) \dim(v) \dim(w) = \dim(u) \dim(v) \dim(w), \quad (32)$$

$$\dim(Ric_{..}) = \dim(R^*_{...}), \quad (33)$$

which imply that *torsion*, *Riemann curvature*, and *Ricci curvature* are dimensionless:

$$\dim(\tau^*_{..}) = \dim(R^*_{...}) = \dim(Ric_{..}) = 1. \quad (34)$$

The exact contra- and co-variant type used above for these tensors is very important in these equations. If we raise any of their indices using a metric, their dimensions will generally change.

Misner et al. (1973 pp. 35, 407) say that ‘curvature’, the Riemann or Einstein tensors (for which see § 8 below), has dimension L^{-2} , a statement seemingly at variance with the dimensionless results (34). But I believe that they refer to the *components* of those tensors, in specific coordinates of dimension L , and using geometrized units. In such specific coordinates

¹⁴ Choquet-Bruhat et al. 1996 § V.B.1 p. 303.

¹⁵ Choquet-Bruhat et al. 1996 § V.B.1.

every component R^i_{jkl} does indeed have dimension L^{-2} , according to the general formula (4), if and only if the *absolute* dimension of \mathbf{R} is unity, $\dim(\mathbf{R}) = 1$. So I believe that there's no real contradiction with that statement and the results (34). This possible misunderstanding shows that it's important to distinguish between absolute dimensions, which don't depend on any specific coordinate choice, and component dimensions, which do.

The formulae above are also valid if a metric is defined and the connection is compatible with it. The connection coefficients in this case are defined in terms of the metric tensor, but using the results of §*** it's easy to see that eqs (24), (25), (27), (34) still hold.

8 Metric tensor

Let's now consider a metric tensor $\mathbf{g}_{..}$. What is its absolute dimension $\dim(\mathbf{g})$? There seem to be two or three choices in the literature; all three can be derived from an operational meaning of the metric.

Consider a (timelike) worldline $s \mapsto C(s)$, $s \in [a, b]$, between events $C(a)$ and $C(b)$. The metric tells us the *proper time* elapsed for an observer having that worldline, according to the formula

$$\Delta t = \int_a^b ds \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|}. \quad (35a)$$

From the results of §5 this formula implies that $T \equiv \dim(\Delta t) = \sqrt{\dim(\mathbf{g}_{..})}$, (independently of the dimension of s) and therefore

$$\dim(\mathbf{g}_{..}) = T^2. \quad (36a)$$

Many authors¹⁶ prefer to include a dimensional factor $1/c$ in front of the integral (35a):

$$\Delta t = \frac{1}{c} \int_a^b ds \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|}, \quad (35b)$$

thus obtaining

$$\dim(\mathbf{g}_{..}) = L^2. \quad (36b)$$

¹⁶ e.g. Fock 1964 § V.62, eq. (62.02); Curtis et al. 1985 ch. 11 eq. (11.21); Rindler 1986 § 5.3 eq. (5.6); Hartle 2003 ch. 6 eq. (6.24).

The choice (36b) seems also supported by the traditional expression for the ‘line element ds^2 ’ as it appears in many works,

$$ds^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2, \quad (37)$$

possibly with opposite signature¹⁷. If the coordinates (t, x, y, z) have the dimensions suggested by their symbols, this formula has dimension L^2 , so that if we interpret ‘ ds^2 ’ as \mathbf{g} we find $\dim(\mathbf{g}) = L^2$. The line-element expression above often has an ambiguous differential-geometric meaning, however, because it may represent *the metric applied to some unspecified vector*, that is, $\mathbf{g}(v, v)$, where v is left unspecified¹⁸. In this case we have

$$L^2 = \dim(\mathbf{g}) \dim(v)^2 \quad (38)$$

and the dimension of \mathbf{g} is ambiguous – probably dimensionless if v has dimension of length.

The standard choices for $\dim \mathbf{g}$ are thus T^2 , L^2 , or possibly 1. My favourite choice is the first, (36a), for reasons discussed by Synge and Bressan¹⁹. Synge gives a vivid summary:²⁰

We are now launched on the task of giving physical meaning to the Riemannian geometry [...]. It is indeed a Riemannian *chronometry* rather than *geometry*, and the word *geometry*, with its dangerous suggestion that we should go about measuring *lengths* with *yardsticks*, might well be abandoned altogether in the present connection

In fact, to measure the proper time Δt defined above we only need to ensure that a clock has the worldline C , and then take the difference between its final and initial times. On the other hand, consider the case when the curve C is *spacelike*. Its proper length is still defined by the integral (35a) (apart from a dimensional constant). Its measurement, however, is more involved than the timelike case. It requires dividing the curve into very short pieces, and having specially-chosen observers (they must be orthogonal to the piece) measure each piece. But the measurement of each piece actually relies on the measurement of *proper time*: each observer uses ‘radar distance’²¹, explained in appendix***. Even if rigid rods are used, their calibration still relies on a measurement

¹⁷ for an exception, with dimension T^2 , see Kilmister 1973 ch. II p. 25.

et al. 1973 Box 3.2 D, p. 77.

1906b § III.3 pp. 108–109.

¹⁹ Synge 1960b §§ III.2–4; Bressan 1978 §§ 15, 18.

²¹ Landau et al. 1996 § 84.

¹⁸ cf. Misner

²⁰ Synge

of time – this is also reflected in the current definition of the standard metre²².

The metric \mathbf{g} can be considered as an operator mapping vectors to covectors, which we can compactly write as $\omega = \mathbf{g}v$, rather than $\omega = \text{tr}_{23}(\mathbf{g} \otimes v)$. The *inverse metric tensor* \mathbf{g}^{-1} is then defined by the formula

$$\mathbf{g}^{-1}\mathbf{g} = \text{id}^{\cdot}, \quad \mathbf{g}\mathbf{g}^{-1} = \text{id}^{\cdot}, \quad (39)$$

and obviously

$$\dim(\mathbf{g}^{-1}) = \dim(\mathbf{g})^{-1}. \quad (40)$$

The operation of *raising or lowering an index* of a tensor represents a contraction of the tensor product of that tensor with the metric or the metric inverse, for example $\mathbf{A}_{\cdot\cdot} \equiv \text{tr}_{13} \mathbf{A}^{\cdot} \otimes \mathbf{g}_{\cdot\cdot}$ and similarly for tensors of other types. Therefore

$$\dim(\mathbf{A}_{\dots}) = \dim(\mathbf{A}_{\dots}^{\cdot}) \dim(\mathbf{g}), \quad \dim(\mathbf{A}_{\dots}^{\cdot}) = \dim(\mathbf{A}_{\dots}) \dim(\mathbf{g})^{-1}. \quad (41)$$

The formulae for the covariant derivative (24), connection coefficients (27), and curvature tensors (34) remain valid for a connection compatible with the metric. In this case the connection coefficients can be obtained from the metric by the formulae²³

$$\Gamma_{jk}^i = \frac{1}{2} \left(\frac{\partial}{\partial x^k} g_{jl} + \frac{\partial}{\partial x^j} g_{kl} - \frac{\partial}{\partial x^l} g_{jk} \right) g^{li}, \quad (42)$$

and it's easily verified that formula (27) still holds, and the other two owing to the same expression of the curvature tensors from the connection coefficients.

The *scalar curvature* ρ and the *Einstein tensor* $\mathbf{G}_{\cdot}^{\cdot}$,

$$\rho = \text{tr } \mathbf{Ric}_{\cdot}^{\cdot} \equiv \text{tr}_{23}(\mathbf{Ric} \otimes \mathbf{g}^{-1}), \quad \mathbf{G}_{\cdot}^{\cdot} := \mathbf{Ric}_{\cdot}^{\cdot} - \frac{1}{2}\rho \text{id}^{\cdot}. \quad (43)$$

have therefore dimension

$$\dim(\rho) = \dim(\mathbf{G}_{\cdot}^{\cdot}) = \dim(\mathbf{g})^{-1} \equiv \begin{cases} \text{T}^{-2} \\ \text{L}^{-2} \\ 1 \end{cases} \quad \text{if } \dim(\mathbf{g}) := \begin{cases} \text{T}^2 \\ \text{L}^2 \\ 1 \end{cases}. \quad (44)$$

²² BIPM 1983 p. 98; Giacomo 1984 p. 25.

²³ Choquet-Bruhat et al. 1996 § V.B.2.

9 Stress-energy-momentum tensor

What are the absolute dimensions of the co-contra-variant stress-energy-momentum tensor \mathbf{T} ? We must look for an operational meaning here too. I'll try to sketch an informal argument that reflects my point of view. The argument can be made more rigorous but that would take too long to do here.

The dynamics equation $\nabla \cdot \mathbf{T} = 0$ holds in general-relativistic (thermo)mechanics, and also in Newtonian (thermo)mechanics when no body forces and no body heating are present. In Newtonian mechanics it's the formal combination of the balances of momentum density and energy density – which incidentally have the same dimensions $[\text{ML}^{-1}\text{T}^{-3}]$, energy/(volume \times time).

The divergence of the stress-energy-momentum gives us a 4-force density, just like the 3-divergence of the stress gives us a force density. Please check Misner & al (1973), chap. 14, for a very interesting discussion of these matters, and also Eckart (1940) and Burke (1980, 1987).

Further, the 4-force is an object that, integrated over a path, gives us an energy density (cf Milne 1951 chap. IV, and Burke again). The integral of a force in Newtonian mechanics is the work done by the force. In general-relativistic mechanics, the timelike component of the 4-force additionally gives us the increase in energy owing to heating (Eckart 1940).

So $\nabla \cdot \mathbf{T} \equiv T_i{}^j{}_{;j} dx^i$ has the dimensions of energy density, $[\text{ML}^{-1}\text{T}^{-2}]$. The *co-contra-variant* stress-energy-momentum \mathbf{T} has therefore the same dimensions. But the *co-co-variant* tensor, obtained by contraction with the metric, $\mathbf{T}_{..} \equiv \mathbf{T} \cdot \mathbf{g}$, has dimensions of energy density times squared time: $[\text{ML}^{-1}]$, a mass over length.

10 Einstein equation

24, 25

In fact the metric tells us the *proper time* Δt elapsed along any (timelike) worldline $s \mapsto C(s)$, $s \in [a, b]$, between events $C(a)$ and $C(b)$, by the formula

²⁴ Rindler 1986 § 3.5 p. 65.

²⁵ Fock 1964 § 2, p. 10.

We can use the metric to measure the "length" of (timelike or spacelike) paths in spacetime. The "length" of a path $c(s)$ with $s \in [a, b]$ is

$$\int_a^b ds \sqrt{|g_{ij}[c(s)] \dot{c}^i(s) \dot{c}^j(s)|}.$$

We see that this "length" has dimensions $[Z^{1/2}]$ and not unexpectedly it doesn't depend on the dimensions of the curve parameter s .

If the path is timelike, this "length" can be measured by a clock having that path as worldline – it's its proper time. Thus, for me $[Z^{1/2}] = [T]$, a time, and therefore the absolute dimensions of the metric tensor are time squared:

$$\dim(\mathbf{g}) = [T^2].$$

I believe that these dimensions also make sense for spacelike paths: in this case we would have to measure the "length" by dividing it in very small pieces and using radar coordinates on each piece. So we're measuring the "length" by checking clocks, to see how long it takes for the light to bounce back: time $[T]$, again.

By our usual argument it's possible to see that the Riemann curvature tensor $\mathbf{R}^{\bullet} \dots$, the Ricci tensor $\mathbf{R}_{\bullet\bullet}$, and the Einstein tensor $\mathbf{G}_{\bullet\bullet}$ are adimensional – $[1]$ – and the scalar curvature has dimensions $[T^{-2}]$. Note that the Riemann and Ricci tensors (with the contra/co-variant type specified above) do not require a metric for their definition, but an affine connection. They are adimensional no matter what dimensions we give the metric. By construction the (fully co-variant) Einstein tensor is always adimensional, too.

An important operation done with the metric:

- "lowering an index" of a tensor multiplies its dimensions by $[T^2]$, and "rising an index" multiplies them by $[T^{-2}]$ (if you agree with my discussion above).

^{26, 27, 28, 29, 30}

Einstein's constant κ therefore relates a dimensionless quantity and a mass over length:

$$\mathbf{G}_{\bullet\bullet} = \kappa \mathbf{T}_{\bullet\bullet}.$$

Its dimension must be $[M^{-1} L]$, and it's easily seen that these are the dimensions of G/c^2 . So I'm one of those people (like Fock 1964 p. 199)

²⁶ Fock 1964 § V.62, eq. (62.02). ²⁷ Curtis et al. 1985 ch. 11 eq. (11.21). ²⁸ Cook 2004 eqs (1) or (9). ²⁹ Hartle 2003 ch. 6 eq. (6.24). ³⁰ Kilmister 1973 ch. II p. 25.

who define

$$\kappa = 8\pi G/c^2.$$

³¹

** References **

- Burke (1980): **Spacetime, Geometry, Cosmology** (University Science Books) - Burke (1987): **Applied Differential Geometry** (Cambridge) - [Eckart (1940)]: **The thermodynamics of irreversible processes. III. Relativistic theory of the simple fluid**, Phys. Rev. 58, 919. - Fock (1964): **The Theory of Space, Time and Gravitation** (Pergamon) - Misner, Thorne, Wheeler (1973): **Gravitation** (Freeman) - Schouten (1989): **Tensor Analysis for Physicists** (Dover, 2nd ed.)

I first present my favourite choice, which is very close to the points of view of Synge and Bressan³², and then the other, probably the most common. I also discuss a possible misinterpretation of traditional expressions for the ‘length element ds^2 ’, with consequences for the dimension of the metric tensor.

Quoting Synge³³: ‘We are now launched on the task of giving physical meaning to the Riemannian geometry [. . .]. It is indeed a Riemannian *chronometry* rather than *geometry*, and the word *geometry*, with its dangerous suggestion that we should go about measuring *lengths* with *yardsticks*, might well be abandoned altogether in the present connection’.

³¹ Whitney 1968a,b.
1960b § III.3 pp. 108–109.

³² Synge 1960b §§ III.2–4; Bressan 1978 §§ 15, 18.

³³ Synge

Bibliography

- (‘de X ’ is listed under D, ‘van X ’ under V, and so on, regardless of national conventions.)
- Aldersley, S. J. (1977): *Dimensional analysis in relativistic gravitational theories*. Phys. Rev. D **15**², 370–376.
- BIPM (Bureau international des poids et mesures), ed. (1983): *17^e conférence générale des poids et mesures (CGPM)*. (BIPM, Sèvres). <https://www.bipm.org/en/worldwide-metrology/cgpm/resolutions.html#comptes-rendus>.
- Boothby, W. M. (2003): *An Introduction to Differentiable Manifolds and Riemannian Geometry*, rev. 2nd ed. (Academic Press, Orlando, USA). First publ. 1975.
- Bossavit, A. (1991): *Differential Geometry: for the student of numerical methods in Electromagnetism*. https://www.researchgate.net/publication/200018385_Differential_Geometry_for_the_student_of_numerical_methods_in_Electromagnetism.
- Bressan, A. (1978): *Relativistic Theories of Materials*. (Springer, Berlin).
- Bridgman, P. W. (1958): *The Logic of Modern Physics*, eight printing. (Macmillan, New York). First publ. 1927.
- Burke, W. L. (1980): *Spacetime, Geometry, Cosmology*. (University Science Books, Mill Valley, USA).
- Choquet-Bruhat, Y., DeWitt-Morette, C., Dillard-Bleick, M. (1996): *Analysis, Manifolds and Physics. Part I: Basics*, rev. ed. (Elsevier, Amsterdam). First publ. 1977.
- Cook, R. J. (2004): *Physical time and physical space in general relativity*. Am. J. Phys. **72**², 214–219.
- Curtis, W. D., Miller, F. R. (1985): *Differential Manifolds and Theoretical Physics*. (Academic Press, Orlando, USA).
- Dorgelo, H. B., Schouten, J. A. (1946): *On unities and dimensions. I*. Verh. Kon. Akad. Wetensch. Amsterdam **49**^{2, 3, 4}, 123–131, 282–291, 393–403.
- Fischer, A. E., Marsden, J. E. (1972): *The Einstein equations of evolution – a geometric approach*. J. Math. Phys. **13**⁴, 546–568.
- Flügge, S., ed. (1960): *Handbuch der Physik: Band III/1: Prinzipien der klassischen Mechanik und Feldtheorie* [Encyclopedia of Physics: Vol. III/1: Principles of Classical Mechanics and Field Theory]. (Springer, Berlin).
- Fock [Fok], V. A. (1964): *The Theory of Space, Time and Gravitation*, second revised ed. (Pergamon, Oxford). Transl. by N. Kemmer. First publ. in Russian 1955.
- Giacomo, P. (1984): *News from the bipm*. Metrologia **20**¹, 25–30.
- Gourgoulhon, É. (2012): *3+1 Formalism in General Relativity: Bases of Numerical Relativity*. (Springer, Heidelberg). First publ. 2007 as arXiv:gr-qc/0703035.
- Hartle, J. B. (2003): *Gravity: An Introduction to Einstein’s General Relativity*. (Addison-Wesley, San Francisco).
- Kilmister, C. W. (1973): *General Theory of Relativity*. (Pergamon, Oxford).
- Landau, L. D., Lifshitz [Lifšic], E. M. (1996): *The Classical Theory of Fields*, corr. repr. of the 4th English ed. (Butterworth-Heinemann, Oxford). Transl. from the 1987 seventh Russian edition by Morton Hamermesh. First publ. 1939.
- Marsden, J. E., Ratiu, T. (2007): *Manifolds, Tensor Analysis, and Applications*, 3rd ed. (Springer, New York). <http://www.math.cornell.edu/~web6520/Abraham-Marsden.pdf>. Written with the collaboration of Ralph Abraham. First publ. 1983.
- Misner, C. W., Thorne, K. S., Wheeler, J. A. (1973): *Gravitation*, repr. (W. H. Freeman and Company, New York). https://archive.org/details/Gravitation_201803.

- Penrose, R., Rindler, W. (2003): *Spinors and Space-Time. Vol. 1: Two-spinor calculus and relativistic fields*, corr. repr. (Cambridge University Press, Cambridge). First publ. 1984.
- Rindler, W. (1986): *Essential Relativity: Special, General, and Cosmological*, rev. second ed. (Springer, New York). First publ. 1969.
- Schouten, J. A. (1989): *Tensor Analysis for Physicists*, corr. second ed. (Dover, New York). First publ. 1951.
- Synge, J. L. (1960a): *Classical Dynamics*. In: Flügge (1960), I–VII, 1–225, 859–902.
- (1960b): *Relativity: The General Theory*. (North-Holland, Amsterdam).
- Truesdell III, C. A., Toupin, R. A. (1960): *The Classical Field Theories*. In: Flügge (1960), I–VII, 226–902. With an appendix on invariants by Jerald LaVerne Ericksen.
- Whitney, H. (1968a): *The mathematics of physical quantities. Part I: Mathematical models for measurement*. Am. Math. Monthly **75**², 115–138. See also Whitney (1968b).
- (1968b): *The mathematics of physical quantities. Part II: Quantity structures and dimensional analysis*. Am. Math. Monthly **75**³, 227–256. See also Whitney (1968a).