

Dimensional analysis in general relativity and in differential geometry

P.G.L. Porta Mana 

Kavli Institute, Trondheim, Norway <pgl@portamana.org>

13 December 2019; updated 19 June 2020

This note provides a short guide to performing dimensional analysis in general relativity and in differential geometry. It tries to revive Dorgelo and Schouten's notion of 'intrinsic' or 'absolute' dimension of a tensorial quantity. The intrinsic dimension is independent of the dimensions of the coordinates and expresses the physical or operational meaning of a tensor. The dimensional analysis of several important tensors and tensor operations is summarized. In particular it is shown that the components of a tensor need not have all the same dimension, and that the Riemann (once contravariant and twice covariant) and Ricci (fully covariant) curvature tensors are dimensionless. The relation between dimension and operational meaning for the metric and stress-energy-momentum tensors is discussed.

per la piccola Emma

1 Introduction

From the point of view of dimensional analysis, do all components of a tensor need to have the same dimension? What happens to these components if we choose coordinates that don't have dimensions of length or time? And if the components of a tensor have different dimensions, then does it make sense to speak of "the dimension of the tensor"? What are the dimensions of the metric and of the curvature tensors, in that case? What is the dimension of the constant in the Einstein equations?

A sense of insecurity often gets hold of many students (and possibly of some researchers) in relativity, when they have to discuss and answer questions like the above. This is evident in many question & answer websites and wiki pages, where some incorrect or unfounded statements about dimensional analysis in general relativity are in circulation. For example, the notion that the components of a tensor or the coordinate functions should all have the same dimension. As we'll see shortly, both notions are false.

Several factors contribute to these misconceptions and insecurity. Modern texts in Lorentzian and general relativity commonly use geometrized units. They say that, for finding the dimension of some constant in a tensorial equation, it's sufficient to compare the dimensions of the terms in the equation. But this procedure is sometimes not so immediate to apply, because some tensors don't have universally agreed dimensions – prime example the metric tensor. Older texts often use four coordinates with dimension of length, and base their dimensional analyses on that specific choice. They even multiply¹ coordinates or tensorial components having dimension of time by powers of c . A student thus gets the impression that coordinates ought to always be lengths, and that all components of a tensor ought to have the same dimension².

Dimensional analysis may thus not be as self-evident in general relativity and in differential geometry as authors in these subjects take it to be. The present note wants to provide a short but exhaustive guide to it. Some important dimensional-analysis questions in general relativity are also consistently settled in this note; for example the dimension of the Riemann curvature tensor, or the effect of the covariant or Lie derivatives on dimensions.

The application of dimensional analysis in general relativity seems to me most straightforward and self-evident if we rely on the coordinate-free or intrinsic approach to differential geometry, briefly recalled below, and if we adopt the perhaps overlooked notion of *intrinsic dimension* of a tensor.

The intrinsic dimension of a tensor was introduced under the name of 'absolute dimension' by Schouten and Dorgelo³ and used in Truesdell & Toupin⁴. As its name implies, this dimension is independent of the choice and dimensions of coordinate functions. It is distinct from the dimensions of the tensor's *components*, which instead depend on the dimensions of the coordinates. The intrinsic dimension of a tensor is determined by the latter's physical and operational⁵ meaning. It is therefore a natural notion for dimensional analysis in general relativity.

¹ e.g. Tolman 1949 p. 72 eq. (37.8); Landau & Lifshitz 1996 p. 80 eq. (32.15); Adler et al. 1975 p. 332 eq. (10.15). ² A recent work explicitly stating, if only in passing, that this needs not be the case is Kitano 2013 § X.

³ Dorgelo & Schouten 1946; Schouten 1989 ch. VI. ⁴ Truesdell & Toupin 1960 Appendix II. ⁵ Bridgman 1958.

A brief reminder of the intrinsic approach to differential geometry, with references, is given in the next section, together with some special notation necessary to our discussion. An introductory example of the basic way of reasoning about dimensional analysis in differential geometry is then presented in § 3. Sections 4–7 offer a more systematic discussion and a synopsis of dimensional analysis for the main tensorial operations. The notion of intrinsic dimension is explained in § 5. The intrinsic dimensions of the various curvature tensors, of the metric tensor, and of the stress-energy-momentum tensor are separately discussed in §§ 8–10. The (contravariant and thrice covariant) Riemann and (fully covariant) Ricci tensors, in particular, are found to have intrinsic dimension 1, that is, to be dimensionless. The operational motivation of the two standard choices for the dimension of the metric tensor are also discussed. The dimension of the constant in the Einstein equations is finally derived in § 11. This note obviously assumes familiarity with basic tensor calculus and related notions, for example of co- and contra-variance, tensor product, contraction. Some passages assume familiarity with the exterior calculus of differential forms. The general ideas, however, should be understandable even without such familiarity.

Finally, quoting Truesdell & Toupin⁶, “dimensional analysis remains a controversial and somewhat obscure subject. We do not attempt a complete presentation here”. References about recent developments in this subject are given in the final summary, § 12.

2 Intrinsic view of differential-geometric objects: brief reminder and notation

From the intrinsic point of view, a tensor is defined by its geometric properties. For example, a vector field v is an object that operates on functions defined on the (spacetime) manifold, yielding new functions, with the properties $v(af + bg) = av(f) + bv(g)$ and $v(fg) = v(f)g + fv(g)$ for all functions f, g and reals a, b . A covector field (1-form) ω is an object that operates on vector fields, yielding functions (‘duality’), with the property $\omega(fu + gv) = f\omega(u) + g\omega(v)$ for all vector fields u, v and functions f, g . The sum of vector or covector fields, and their products by functions – let’s call this ‘linearity’ – are defined in an obvious way. Tensors are constructed from these objects.

⁶ Truesdell & Toupin 1960 Appendix § 7 footnote 4.

A system of coordinates (x^i) is just a set of linearly independent functions. This set gives rise to a set of vectors fields $(\frac{\partial}{\partial x^i})$ and to a set of covector fields (dx^i) by the obvious requirements that $\frac{\partial}{\partial x^i}(x^j) = \delta_i^j$ and $dx^i(\frac{\partial}{\partial x^j}) = \delta_j^i$. These two sets can be used as bases to express all other vectors and covectors as linear combinations. A vector field v can thus be written as

$$v \equiv \sum_i v^i \frac{\partial}{\partial x^i} \equiv v^i \frac{\partial}{\partial x^i}, \quad (1)$$

where the *functions* $v^i := v(x^i)$ are its components with respect to the basis $(\frac{\partial}{\partial x^i})$. Analogously for a covector field.

For the presentation of the intrinsic view I recommend the excellent texts by Choquet-Bruhat et al. (1996), Boothby (2003), Abraham et al. (1988), Bossavit (1991), Burke (1987; 1980 ch. 2), and more on the general-relativity side Misner et al. (1973 ch. 9),ourgoulhon (2012 ch. 2), Penrose & Rindler (2003 ch. 4).

For the notation in dimensional analysis I use ISO conventions: $\dim(\mathbf{A})$ is the dimension of the quantity \mathbf{A} , and among the base quantities are mass M , length L , time T , temperature Θ , electric current I . Note that I don't discuss units – it doesn't matter here whether the unit for length is the metre or the centimetre, for example.

The number and ordering of a tensor's covariant and contravariant “slots”⁷ will often be important in our discussion. The traditional coordinate-free notation ‘ \mathbf{A} ’ unfortunately omits this information. We thus need a coordinate-free notation that makes it explicit. Penrose & Rindler⁸ propose an abstract-index notation where ‘ A_i^{jk} ’, for example, denotes a tensor covariant in its first slot and contravariant in its second and third slots. Every index in this notation is “a *label* whose sole purpose is to keep track of the type of tensor under discussion”⁹. So this notation doesn't stand for the set of *components* of the tensor. For the latter set, **bold** indices are used instead: ‘ \mathbf{A}_i^{jk} ’. In our discussion, however, where the difference between a tensor and its set of components will be crucial, this abstract-index notation unfortunately lends itself to conceptual and typographic misunderstanding.

I shall therefore use a notation such as $\mathbf{A}_{\bullet}^{\bullet\bullet}$ to indicate that \mathbf{A} is covariant in its first slot and contravariant in its second and third slots.

⁷ Misner et al. 1973 § 3.2.

⁸ Penrose & Rindler 2003 § 2.2.

⁹ Penrose & Rindler

2003 p. 75.

Its components would thus be (A_i^{jk}) . For brevity I'll call this a 'co-contra-contra-variant' tensor, with an obvious naming generalization for other tensor types.

The only weak points of this notation are the operations of transposition and contraction, which literal indices depict so well instead. Considering that transposition is a generalization of matrix transposition, and contraction is a generalization of trace, I'll use the following notation:

- $A^{\top\alpha\beta}$ is the transposition (swapping) of the α th and β th slots. Its coordinate-free definition is

$$(A^{\top\alpha\beta})(\dots, \overset{\alpha\text{th slot}}{\zeta}, \dots, \underset{\beta\text{th slot}}{\eta}, \dots) := A(\dots, \overset{\alpha\text{th slot}}{\eta}, \dots, \underset{\beta\text{th slot}}{\zeta}, \dots) \quad (2)$$

for each ζ, η of appropriate variance type.

- $\text{tr}_{\alpha\beta} A$ is the contraction of the α th and β th slots (which must have opposite variant types). Its coordinate-free definition is

$$(\text{tr}_{\alpha\beta} A)(\dots, \dots, \dots) := \sum_i A(\dots, \overset{\alpha\text{th slot}}{u_i}, \dots, \underset{\beta\text{th slot}}{\omega^i}, \dots) \quad (3)$$

for any arbitrary complete and linearly independent $\{u_i\}, \{\omega^j\}$ such that $\omega^j(u_i) = \delta_i^j$.

An analogous definition holds if the α th slot is covariant and the β th contravariant.

In index notation these operations are the familiar

$$\underset{\beta\text{th slot}}{A \dots \overset{\alpha\text{th slot}}{i} \dots j \dots} \mapsto \underset{\alpha\text{th slot}}{A \dots j \dots \overset{\beta\text{th slot}}{i} \dots} \quad \text{and} \quad \underset{\beta\text{th slot}}{A \dots \overset{\alpha\text{th slot}}{i} \dots i \dots}.$$

Contraction and transposition will be discussed only sparsely, so I hope you won't find the notation above too uncomfortable.

3 An introductory two-dimensional example

Let me first present a simple example of dimensional analysis in a two-dimensional spacetime. I provide very little explanation, letting the analysis speak for itself. The next sections will give a longer discussion of the general point of view, of the assumptions, and of cases with more elaborate geometric objects.

In a region of a two-dimensional spacetime we use coordinates (x, y) . These coordinates allow us to uniquely identify every event in the region (otherwise they wouldn't be coordinates). Let us say that coordinate x has dimension of temperature, and y of specific entropy:

$$\dim(x) = \Theta, \quad \dim(y) = s := \text{L}^2 \text{T}^{-2} \Theta^{-1}. \quad (4)$$

This choice may be possible for several reasons. For example, the region could be occupied by a heat-conducting material; in a specific spacetime foliation, its temperature increases along each 1-dimensional spacelike slice (constant y), and its entropy density is uniform on each slice but increases from slice to slice.¹⁰ The point here is that coordinates can have any dimensions owing to physical reasons.

From these coordinates we construct two covector fields (dx, dy) , and two vector fields $(\frac{\partial}{\partial x}, \frac{\partial}{\partial y})$ that serve as bases for the spaces of tangent covectors, vectors, and tensors. Their dimensions are

$$\begin{aligned} \dim(dx) &= \Theta & \dim(dy) &= s, \\ \dim\left(\frac{\partial}{\partial x}\right) &= \Theta^{-1} & \dim\left(\frac{\partial}{\partial y}\right) &= s^{-1}. \end{aligned} \quad (5)$$

Consider a contra-co-variant tensor field $\mathbf{A} \equiv \mathbf{A}^\bullet$, in this region. Using the basis fields above it can be written as

$$\mathbf{A} = A^x_x \frac{\partial}{\partial x} \otimes dx + A^x_y \frac{\partial}{\partial x} \otimes dy + A^y_x \frac{\partial}{\partial y} \otimes dx + A^y_y \frac{\partial}{\partial y} \otimes dy, \quad (6)$$

where $A^x_x := \mathbf{A}(dx, \frac{\partial}{\partial x})$ and so on are the components of the tensor in the coordinate system (x, y) .

By the rules of dimensional analysis, the two sides of the expansion above must have the same dimension. The four summands of the right side must also have the same dimension. Denoting $A := \dim(\mathbf{A})$, we thus have the four equations

$$\begin{aligned} A &= \dim(A^x_x) & A &= \dim(A^x_y) \Theta^{-1} s \\ A &= \dim(A^y_x) \Theta s^{-1} & A &= \dim(A^y_y), \end{aligned}$$

or

$$\begin{aligned} \dim(A^x_x) &= A & \dim(A^x_y) &= A \Theta s^{-1} \equiv A \text{L}^{-2} \text{T}^2 \Theta^2 \\ \dim(A^y_x) &= A \Theta^{-1} s & \dim(A^y_y) &= A. \end{aligned} \quad (7)$$

¹⁰ For general-relativistic thermomechanics see e.g. Eckart 1940; Maugin 1974; 1978a,b,c,d; Muschik & von Borzeszkowski 2014.

The intrinsic dimension of the tensor \mathbf{A} is A . The expansion (6) shows that this dimension is independent of the coordinate system, by construction – the expansion could be done in any other coordinate system, and the left side would be the same. The effect of coordinate transformations is examined more in detail in § 5. The intrinsic dimension is determined by the physical or operational meaning of the tensor. Together with the dimensions of the coordinates it determines the dimensions of the components, eq. (7), which need not be all equal.

This simple example should have disclosed the main points of dimensional analysis on manifolds, which will now be discussed in more generality.

In the derivation above we silently adopted a couple of natural conventions; for example, that the tensor product behaves similarly to multiplication with regard to dimensions. Such conventions are briefly discussed in § 12.

4 Coordinates

From a physical point of view, a coordinate is just a function that associates values of some physical quantity with the events in a region (the domain of the coordinate chart) of spacetime. Together with the other coordinates, such function allows us to uniquely identify every event within that region. Any physical quantity will do: the distance from something, the time elapsed since something, an angle, an energy density, the strength of a magnetic flux, a temperature, and so on. A coordinate can thus have any dimension: length L , time T , angle 1 , temperature Θ , magnetic flux $\Phi := ML^2T^{-2}I^{-1}$, and so on.

The functional relation between two sets of coordinates must of course be dimensionally consistent. For example, if $\dim(x^0) = T$ and $\dim(x^1) = L$, and we introduce a coordinate $\xi(x^0, x^1)$ with dimension 1 , additive in the previous two, then we must have $\xi = ax^0 + bx^1$ with $\dim(a) = T^{-1}$ and $\dim(b) = L^{-1}$.

5 Tensors: intrinsic dimension and components' dimensions

Consider a system of coordinates (x^i) with dimensions (X_i) , and the ensuing sets of covector fields (1-forms) dx^i and of vector fields $(\frac{\partial}{\partial x^i})$,

bases for the cotangent and tangent spaces. Their tensor products are bases for the tangent spaces of higher tensor types.

The differential dx^i traditionally has the same dimension as x^i : $\dim(dx^i) = X_i$, and the vector $\frac{\partial}{\partial x^i}$ traditionally has the inverse dimension: $\dim \frac{\partial}{\partial x^i} = X_i^{-1}$.

For our discussion let's take a concrete example: a contra-co-variant tensor field $\mathbf{A} \equiv \mathbf{A}^\bullet$. The discussion generalizes to tensors of other types in an obvious way.

The tensor \mathbf{A} can be expanded in terms of the basis vectors and covectors, as in § 2 and in the example of § 3:

$$\mathbf{A} = A^i_j \frac{\partial}{\partial x^i} \otimes dx^j \equiv A^0_0 \frac{\partial}{\partial x^0} \otimes dx^0 + A^0_1 \frac{\partial}{\partial x^0} \otimes dx^1 + \dots \quad (8)$$

Each function

$$A^i_j := \mathbf{A}\left(dx^i, \frac{\partial}{\partial x^j}\right) \quad (9)$$

is a component of the tensor in this coordinate system.

To make dimensional sense, all terms in the sum (8) must have the same dimension. This is possible only if the generic component A^i_j has dimension

$$\dim(A^i_j) = A X_i X_j^{-1}, \quad (10)$$

where A is common to all components. In fact, the $X_i X_j^{-1}$ term cancels the $X_i^{-1} X_j$ term coming from $\frac{\partial}{\partial x^i} \otimes dx^j$ in the sum (8), and each summand therefore has dimension A .

The generalization of the formula above to tensors of other types is obvious:

$$\dim(A^{ij\dots}_{kl\dots}) = A X_i X_j \dots X_k^{-1} X_l^{-1} \dots \quad (11)$$

where the ordering of the indices doesn't matter.

Clearly the components can have different dimensions. But this doesn't matter. What matters is that the sum (8) be dimensionally consistent. (Fokker¹¹, for example, uses a metric tensor with components having different dimensions.)

¹¹ Fokker 1965 § VII.1 p. 88.

The dimension A , which is also the dimension of the sum (8), I'll call the *intrinsic dimension* of the tensor \mathbf{A} , and we write

$$\dim(\mathbf{A}) = A. \quad (12)$$

This dimension is independent of any coordinate system. It reflects the physical or operational¹² meaning of the tensor. We shall see an example of such an operational analysis in §§ 9 and 10 for the metric and stress-energy-momentum tensors.

The notion of intrinsic dimension was introduced by Dorgelo and Schouten¹³ under the name 'absolute dimension'. I find the adjective 'intrinsic' more congruous to modern terminology (and less prone to suggest spurious connections with absolute values).

Different coordinate systems lead to different dimensions of the *components* of a tensor \mathbf{A} , but the absolute dimension of the tensor remains the same. Formula (11) for the dimensions of the components is consistent under changes of coordinates. For example, in new coordinates (\bar{x}^k) with dimensions (\bar{X}_k) , the new components of \mathbf{A} are

$$\bar{A}^k_l = A^i_j \frac{\partial \bar{x}^k}{\partial x^i} \frac{\partial x^j}{\partial \bar{x}^l} \quad (13)$$

and a quick check shows that $\dim(\bar{A}^k_l) = A \bar{X}_k \bar{X}_l^{-1}$, consistently with the general formula (11).

In the following I'll drop the adjective 'intrinsic' when it is clear from the context.

6 Tensor operations

By the reasoning of the previous section, which simply applies standard dimensional considerations to the basis expansion (8), it's easy to find the resulting intrinsic dimension of various operations and operators on tensors and tensor fields.

Here is a summary of the dimensional rules for the main differential-geometric operations and operators, except for the covariant derivative, the metric, and related tensors, discussed more in depth in §§ 8–9 below.

¹² Bridgman 1958; see also Synge 1960a § A.2; Truesdell & Toupin 1960 §§ A.3–4.

¹³ Dorgelo & Schouten 1946; Schouten 1989 ch. VI.

Some of these rules are actually definition or conventions, as briefly discussed in their description. The others can be proved; I only give a proof for one of them, leaving the other proofs as an exercise. For reference, in brackets I give the section of Choquet-Bruhat et al. (1996) where these operations are defined.

- The *tensor product* [III.B.5] multiplies dimensions:

$$\dim(\mathbf{A} \otimes \mathbf{B}) = \dim(\mathbf{A}) \dim(\mathbf{B}). \quad (14)$$

This is actually a definition or convention. We tacitly used this rule already in the example of § 3 and in § 5 for the coordinate expansion (8). It is a natural definition, because for tensors of order 0 (functions) the tensor product is just the ordinary product, and the dimension of a product is the product of the dimensions. This definition doesn't lead to inconsistencies.

- The *contraction* [III.B.5] or trace of the α th and β th slots of a tensor has the same dimension as the tensor:

$$\dim(\text{tr}_{\alpha\beta} \mathbf{A}) = \dim(\mathbf{A}). \quad (15)$$

Note that the formula above only holds *without raising or lowering indices*; see § 9 for those operations.

This operation can be traced back to the duality of vectors and covectors mentioned in § 2: a covector field ω operates linearly on a vector field v to yield a function $f = \omega(v)$. Also in this case we have that $\dim(f) = \dim(\omega) \dim(v)$ by definition or convention, and the rule (15) follows from this convention. Also in this case this convention seems very natural, owing to the linearity properties of the trace, and doesn't lead to inconsistencies.

- The *transposition*¹⁴ of the α th and β th slots of a tensor has the same dimension as the tensor:

$$\dim(\mathbf{A}^{\top\alpha\beta}) = \dim(\mathbf{A}). \quad (16)$$

¹⁴ called "building an isomer"¹⁴ by Schouten 1954 § I.3 p. 13; 1989 § II.4 p. 20.

- The *Lie bracket* [III.B.3] of two vectors has the product of their dimensions:

$$\dim([u, v]) = \dim(u) \dim(v). \quad (17)$$

In fact, in coordinates (x^i) the bracket can be expressed as

$$[u, v] = \left(u^j \frac{\partial v^i}{\partial x^j} - v^j \frac{\partial u^i}{\partial x^j} \right) \frac{\partial}{\partial x^i}, \quad (18)$$

and equating the dimensions of the left and right sides, considering that

$$\dim(u^i) = \dim(u) X_i, \quad \dim(v^i) = \dim(v) X_i, \quad (19)$$

we find again that all X terms cancel out, leaving the result (17).

- The *Lie derivative* [III.C.2] of a tensor with respect to a vector field has the product of the dimensions of the tensor and of the vector:

$$\dim(L_v \mathbf{A}) = \dim(v) \dim(\mathbf{A}). \quad (20)$$

Regarding operations and operators on differential forms:

- The *exterior product* [IV.A.1] of two differential forms multiplies their dimensions:

$$\dim(\omega \wedge \tau) = \dim(\omega) \dim(\tau). \quad (21)$$

- The *interior product* [IV.A.4] of a vector and a form multiplies their dimensions:

$$\dim(i_v \omega) = \dim(v) \dim(\omega). \quad (22)$$

- The *exterior derivative* [IV.A.2] of a form has the same dimension of the form:

$$\dim(d\omega) = \dim(\omega). \quad (23)$$

This can be proven using the identity $d i_v + i_v d = L_v$ or similar identities¹⁵ together with eqs (20) and (22).

- The *integral* [IV.B.1] of a form over a submanifold, or more generally a chain, M has the same dimension as the form:

$$\dim\left(\int_M \omega\right) = \dim(\omega). \quad (24)$$

¹⁵ Curtis & Miller 1985 ch. 9 p. 180 Theorem 9.78; Abraham et al. 1988 § 6.4 Theorem 6.4.8.

The reason is that the integral of a form over a chain ultimately rests on the standard definition of integration on the real line¹⁶, which satisfies the dimensional rule above. In fact, the integral is invariant with respect to reparameterizations of the chain; it depends only on its image (some texts¹⁷ even define chains as equivalence classes determined by their image.)

All rules above extend in obvious ways to inner-oriented forms¹⁸ (also called 'odd'¹⁹ or 'twisted'²⁰ forms) and to tensor densities.

7 Curves and integral curves

Consider a curve into spacetime, $C: s \mapsto P(s)$, with the parameter s having dimension $\dim(s) = S$.

If we consider the events of the spacetime manifold as dimensionless quantities, then the dimension of the tangent or velocity vector \dot{C} to the curve is

$$\dim(\dot{C}) = S^{-1}, \quad (25)$$

owing to the definition²¹

$$\dot{C} := \frac{\partial(x^i \circ C)}{\partial s} \frac{\partial}{\partial x^i}. \quad (26)$$

This has a quirky but interesting consequence. Given a vector field v we say that C is an integral curve for it if

$$v = \dot{C} \quad (27)$$

at all events $C(s)$ in the image of the curve (or more precisely $v_{C(s)} = \dot{C}_{C(s)}$ in usual differential-geometric notation²²). From the point of view of dimensional analysis this definition can only be valid if v has dimension S^{-1} . If v and s^{-1} have different dimensions – a case which could happen for physical reasons – the condition (26) must be modified into $v = k\dot{C}$, where k is a possibly dimensionful constant. This is equivalent to considering an affine and dimensional reparameterization of C .

¹⁶ e.g. Choquet-Bruhat et al. 1996 §§ IV.B.1–2; de Rham 1984 § 5 p. 21, § 6 p. 24; Abraham et al. 1988 § 7.1; Boothby 2003 § VI.2. ¹⁷ e.g. Martin 2004 § 10.4 p. 297; Fecko 2006 § 7.3.

¹⁸ Schouten 1989 ch. II. ¹⁹ De Rham 1984 ch. II. ²⁰ Burke 1983; 1995; Bossavit 1991 ch. 3. ²¹ Choquet-Bruhat et al. 1996 § III.B.1; Boothby 2003 § IV.(1.9).

²² Choquet-Bruhat et al. 1996 § III.B.1.

8 Connection, covariant derivative, curvature tensors

Consider an arbitrary connection²³ with covariant derivative ∇ . For the moment we don't assume the presence of any metric structure.

The covariant derivative of the product $f v$ of a function and a vector satisfies²⁴

$$\nabla(f v) = df \otimes v + f \nabla v. \quad (28)$$

The first summand, from formulae (23) and (14), has dimension $\dim(f) \dim(v)$; for dimensional consistency this must also be the dimension of the second summand. Thus

$$\dim(\nabla v) = \dim(v). \quad (29)$$

It follows that the *directional* covariant derivative ∇_u has dimension

$$\dim(\nabla_u v) = \dim(u) \dim(v), \quad (30)$$

and by its derivation properties²⁵ we see that formula (29) extends from vectors to tensors of arbitrary type.

In the coordinate system (x^i) , the action of the covariant derivative is carried by the *connection coefficients* or Christoffel symbols (Γ^i_{jk}) defined by

$$\nabla \frac{\partial}{\partial x^k} = \Gamma^i_{jk} dx^j \otimes \frac{\partial}{\partial x^i}. \quad (31)$$

From this equation and the previous ones it follows that these coefficients have dimensions

$$\dim(\Gamma^i_{jk}) = X_i X_j^{-1} X_k^{-1}. \quad (32)$$

The *torsion* $\tau^{\bullet} \dots$, *Riemann curvature* $R^{\bullet} \dots$, and *Ricci curvature* $Ric_{\bullet \bullet}$ tensors are defined by²⁶

$$\tau(u, v) := \nabla_u v - \nabla_v u - [u, v], \quad (33)$$

$$R(u, v; w) := \nabla_u \nabla_v w - \nabla_v \nabla_u w - \nabla_{[u, v]} w, \quad (34)$$

$$Ric_{\bullet \bullet} := \text{tr}_{13} R^{\bullet} \dots. \quad (35)$$

²³ Choquet-Bruhat et al. 1996 § V.B.

²⁵ Choquet-Bruhat et al. 1996 § V.B.1 p. 303.

²⁴ Choquet-Bruhat et al. 1996 § V.B.1.

²⁶ Choquet-Bruhat et al. 1996 § V.B.1.

From these definitions and the results of § 6 we find the dimensional requirements

$$\dim(\boldsymbol{\tau}^{\bullet..}) \dim(\boldsymbol{u}) \dim(\boldsymbol{v}) = \dim(\boldsymbol{u}) \dim(\boldsymbol{v}), \quad (36)$$

$$\dim(\boldsymbol{R}^{\bullet...}) \dim(\boldsymbol{u}) \dim(\boldsymbol{v}) \dim(\boldsymbol{w}) = \dim(\boldsymbol{u}) \dim(\boldsymbol{v}) \dim(\boldsymbol{w}), \quad (37)$$

$$\dim(\boldsymbol{Ric}_{..}) = \dim(\boldsymbol{R}^{\bullet...}), \quad (38)$$

which imply that *torsion*, *Riemann curvature*, and *Ricci curvature tensors* are *dimensionless*:

$$\dim(\boldsymbol{\tau}^{\bullet..}) = \dim(\boldsymbol{R}^{\bullet...}) = \dim(\boldsymbol{Ric}_{..}) = 1. \quad (39)$$

The exact contra- and co-variant type of these tensors is very important in the equations above. If we raise any of their indices using a metric, their dimensions will generally change.

Misner et al. (1973 p. 35) say that “curvature”, by which they seem to mean the Riemann tensor, has dimension L^{-2} . This statement is seemingly at variance with the dimensionless results (39). But I believe that Misner et al. refer to the *components* of the Riemann tensor, in specific coordinates of dimension L , and using geometrized units. In such specific coordinates every *component* R^i_{jkl} does indeed have dimension L^{-2} , according to the general formula (11), if and only if the intrinsic dimension of \boldsymbol{R} is unity, $\dim(\boldsymbol{R}) = 1$. So I believe that Misner et al.’s statement actually agrees with the results (39). This possible misunderstanding shows the importance of distinguishing between intrinsic dimensions, which don’t depend on any specific coordinate choice, and component dimensions, which do.

The formulae above are also valid if a metric is defined and the connection is compatible with it. The connection coefficients in this case are defined in terms of the metric tensor, but using the results of § 9 it’s easy to see that eqs (29), (30), (32), (39) still hold.

9 Metric and related tensors and operations

Let us now consider a metric tensor $\boldsymbol{g}_{..}$. What is its intrinsic dimension $\dim(\boldsymbol{g})$? There seem to be two choices in the literature; both can be derived from the operational meaning of the metric.

Consider a (timelike) worldline $s \mapsto C(s)$, $s \in [a, b]$, between events $C(a)$ and $C(b)$. The metric tells us the *proper time* Δt elapsed for an observer having that worldline, according to the formula

$$\Delta t = \int_a^b \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|} \, ds . \quad (40)$$

From the results of § 6 this formula implies that $T \equiv \dim(\Delta t) = \sqrt{\dim(\mathbf{g}_{..})}$, (independently of the dimension of s) and therefore

$$\dim(\mathbf{g}_{..}) = T^2. \quad (41)$$

Many authors²⁷, however, prefer to include a dimensional factor $1/c$ in front of the integral (40):

$$\Delta t = \frac{1}{c} \int_a^b \sqrt{|\mathbf{g}[\dot{C}(s), \dot{C}(s)]|} \, ds , \quad (42)$$

thus obtaining

$$\dim(\mathbf{g}_{..}) = L^2. \quad (43)$$

The choice (43) seems also supported by the traditional expression for the “line element ds^2 ” as it appears in many works:

$$ds^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2, \quad (44)$$

possibly with opposite signature (for an exception with dimension T^2 see Kilmister²⁸). If the coordinates (t, x, y, z) have the dimensions suggested by their symbols, this formula has dimension L^2 , so that if we interpret “ ds^2 ” as \mathbf{g} we find $\dim(\mathbf{g}) = L^2$. The line-element expression above often has an ambiguous differential-geometric meaning, however, because it may also represent the metric applied to some *unspecified* vector, that is, $\mathbf{g}(v, v)$, where v is left unspecified²⁹. In this case we have

$$L^2 = \dim(\mathbf{g}) \dim(v)^2 \quad (45)$$

and the dimension of \mathbf{g} is ambiguous or undefined, because the vector v could have any dimension.

²⁷ e.g. Fock 1964 § V.62 eq. (62.02); Curtis & Miller 1985 ch. 11 eq. (11.21); Rindler 1986 § 5.3 eq. (5.6); Hartle 2003 ch. 6 eq. (6.24). ²⁸ Kilmister 1973 ch. II p. 25. ²⁹ cf. Misner et al. 1973 Box 3.2 D p. 77.

The standard choices for $\dim \mathbf{g}$ are thus T^2 or L^2 . My favourite choice is the first, for reasons discussed by Synge and Bressan³⁰. Synge gives a vivid summary:³¹

We are now launched on the task of giving physical meaning to the Riemannian geometry [...]. It is indeed a Riemannian *chronometry* rather than *geometry*, and the word *geometry*, with its dangerous suggestion that we should go about measuring *lengths* with *yardsticks*, might well be abandoned altogether in the present connection

In fact, to measure the proper time Δt defined above we only need to ensure that a clock has the worldline C , and then take the difference between the clock's final and initial times. On the other hand, consider the case when the curve C is *spacelike*. Its proper length is still defined by the integral (40) apart from a dimensional constant. Its measurement, however, is more involved than in the timelike case. It requires dividing the curve into very short pieces, and having specially-chosen observers (orthogonal to the pieces) measure each piece. But the measurement of each piece actually relies on the measurement of *proper time*: each observer uses radar distance³², sending a lightlike signal which bounces back at the end of the piece, and measuring the time it takes to come back. Even if rigid rods are used, their calibration still relies on a measurement of time – this is also reflected in the current definition of the standard metre³³.

The metric \mathbf{g} can be considered as an operator mapping vectors to covectors, which we can compactly write as $\boldsymbol{\omega} = \mathbf{g}\mathbf{v}$ (instead of the cumbersome $\boldsymbol{\omega} = \text{tr}_{23}(\mathbf{g} \otimes \mathbf{v})$). The *inverse metric tensor* $\mathbf{g}^{-1\bullet}$ is then defined by the formula

$$\mathbf{g}^{-1}\mathbf{g} = \text{id}^\bullet, \quad \mathbf{g}\mathbf{g}^{-1} = \text{id}_\bullet, \quad (46)$$

so that

$$\dim(\mathbf{g}^{-1}) = \dim(\mathbf{g})^{-1}. \quad (47)$$

The *metric volume element*³⁴ in spacetime is a 4-form γ , equivalent to a completely antisymmetric tensor γ_{\dots} , such that $\gamma(e_0, e_1, e_2, e_3) = 1$ for every set of positively-oriented orthonormal vector fields (e_k) , that

³⁰ Synge 1960b §§ III.2–4; Bressan 1978 §§ 15, 18. ³¹ Synge 1960b § III.3 pp. 108–109. ³² Landau & Lifshitz 1996 § 84. ³³ BIPM 1983 p. 98; Giacomo 1984 p. 25.

³⁴ Abraham et al. 1988 § 6.2.

is, such that $\mathbf{g}(e_k, e_l) = \pm \delta_{kl}$ (remember that the orientation is not determined by the metric). It has only one non-zero component, given by the square root of the determinant of the (positively ordered) components (g_{ij}) of the metric:

$$\gamma = \sqrt{|\det(g_{ij})|} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3. \quad (48)$$

From this expression and the results of § 6 it can be shown that, in spacetime,

$$\dim(\gamma) = \dim(\mathbf{g})^2 \equiv \begin{cases} T^4 \\ L^4 \end{cases} \quad \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases}. \quad (49)$$

This is also the dimension of the *density* $|\gamma|$, which, as opposed to the volume element, has the property that $|\gamma|(e_0, e_1, e_2, e_3) = 1$ for all sets of orthonormal vector fields, not only positively-oriented ones (it is an inner-oriented 4-form).

The operation of *raising or lowering an index* of a tensor represents a contraction of the tensor product of that tensor with the metric or the metric inverse, for example $\mathbf{A}_{..} \equiv \text{tr}_{13}(\mathbf{A}^{\bullet} \otimes \mathbf{g}_{..})$ and similarly for tensors of other types. Therefore

$$\dim(\mathbf{A}_{...}) = \dim(\mathbf{A}^{\bullet} \dots) \dim(\mathbf{g}), \quad \dim(\mathbf{A}^{\bullet} \dots) = \dim(\mathbf{A}_{...}) \dim(\mathbf{g})^{-1}. \quad (50)$$

The formulae for the covariant derivative (29), connection coefficients (32), and curvature tensors (39) remain valid for a connection compatible with the metric. In this case the connection coefficients can be obtained from the metric by the formulae³⁵

$$\Gamma_{jk}^i = \frac{1}{2} \left(\frac{\partial}{\partial x^k} g_{jl} + \frac{\partial}{\partial x^j} g_{kl} - \frac{\partial}{\partial x^l} g_{jk} \right) g^{li}, \quad (51)$$

and it's easily verified that the dimensions (32) of these coefficients still hold. Also the result for the curvature tensors (39) still holds, since their expressions in terms of the connection coefficients is the same with or without a metric.

³⁵ Choquet-Bruhat et al. 1996 § V.B.2.

The scalar curvature ρ and the Einstein tensor \mathbf{G}_\bullet :

$$\rho := \text{tr } \mathbf{Ric}_\bullet \equiv \text{tr}_{23}(\mathbf{Ric} \otimes \mathbf{g}^{-1}), \quad \mathbf{G}_\bullet := \mathbf{Ric}_\bullet - \frac{1}{2}\rho \text{ id}_\bullet \quad (52)$$

have therefore dimension

$$\dim(\rho) = \dim(\mathbf{G}_\bullet) = \dim(\mathbf{g})^{-1} \equiv \begin{cases} \text{T}^{-2} \\ \text{L}^{-2} \end{cases} \quad \text{if } \dim(\mathbf{g}) := \begin{cases} \text{T}^2 \\ \text{L}^2 \end{cases} . \quad (53)$$

10 Stress-energy-momentum tensor

To find the dimension of the stress-energy-momentum \mathbf{T} , or ‘4-stress’ for short, let’s start with the analysis of the (3-)stress σ in Newtonian mechanics. The stress σ is the projection of the 4-stress \mathbf{T} onto a spacelike tangent plane with respect to some observer³⁶. If we assume that such spatial projection preserves the intrinsic dimension, then the 4-stress and the stress have the same intrinsic dimension.

In Newtonian mechanics the stress σ is an object that, integrated over the boundary of a body, gives the total surface force acting on the body³⁷ (such integration requires a flat connection). This means that it must be represented by a “force-valued” 2-form. Force, in turn, can be interpreted as an object that, integrated over a (spacelike) trajectory, gives an energy – the work done by the force along the trajectory. It’s therefore a 1-form. Putting these two requirements together, the stress turns out to be a covector-valued 2-form, equivalent to a tensor $\sigma_{\bullet\bullet}$ antisymmetric in its last two indices. Integrated over a surface, and then over a trajectory, it yields an energy. From § 6, integration of a form does not change the dimension of the form. Therefore

$$\dim(\sigma_{\bullet\bullet}) = \text{E} \equiv \text{ML}^2\text{T}^{-2} . \quad (54)$$

But usually the stress is represented by a co-contravariant tensor σ_\bullet . The latter is obtained by contracting the last two slots of $\sigma_{\bullet\bullet}$ with the inverse of the volume element of the 3-metric – this is the duality³⁸ between k -vectors and $(n - k)$ -covectors induced by the metric (and an orientation choice), where n is the geometric dimension of the manifold.

³⁶ Gourgoulhon 2012 § 3.4.1; Smarr & York 1978; York 1979; Smarr et al. 1980; Wilson & Mathews 2007 § 1.3; the projection doesn’t need to be orthogonal: Marsden & Hughes 1994 § 2.4; Hehl & Obukhov 2003 § B.1.4. ³⁷ Truesdell 1991 ch. III. ³⁸ Bossavit 1991 § 4.1.2.

If we assume the Newtonian 3-metric to have dimension L^2 , it can be shown similarly to § 9 that its volume element has dimension L^3 , and the inverse volume element has dimension L^{-3} . Thus we obtain

$$\dim(\sigma_{\bullet}) = EL^{-3} \equiv ML^{-1}T^{-2}, \quad (55)$$

an energy density (or ‘volumic energy’ according to ISO³⁹).

Since the stress σ_{\bullet} is the projection of T_{\bullet} and the projection preserves the intrinsic dimension, we finally find that T_{\bullet} also has the dimension of an energy density:

$$\dim(T_{\bullet}) = EL^{-3} \equiv ML^{-1}T^{-2}. \quad (56)$$

Note that other co- or contra-variant versions of the 4-stress have different intrinsic dimension, because they’re obtained by lowering or raising indices. For example, $\dim(T_{..}) = \dim(T_{\bullet}) \dim(g) = ML^{-1}$ if $\dim(g) := T^2$.

Let me add a passing remark. Even though in most texts the 4-stress is represented by a tensor of order 2, as above, its most fitting geometrical nature is still shrouded in mystery from the kinematic and the dynamical points of view. There are indications that it could be more properly represented by a covector-valued 3-form (equivalent to a tensor $T_{...}$ antisymmetric in the last three slots), or by a 3-vector-valued 3-form (equivalent to a tensor $T^{...}$ antisymmetric in the first three and last three slots), for reasons connected with integration, similar to those mentioned above for the stress $\sigma_{...}$. See for example the discussion about “ T ” by Misner et al.⁴⁰, the works by Segev⁴¹, the discussion by Burke⁴².

11 The constant in the Einstein equations

We finally arrive at the Einstein equations,

$$G = \kappa T \quad (57)$$

where κ (sometimes seen with a minus⁴³ depending on the signature of the metric or on the orientation of the stress) is Einstein’s constant. For

³⁹ ISO 2009 item A.6.2. ⁴⁰ Misner et al. 1973 ch. 15. ⁴¹ Segev 2002; 1986; Segev & Rodnay 1999; Segev 2000a,b. ⁴² Burke 1987 § 41. ⁴³ e.g. Tolman 1949 § 78 eq. (78.3); Fock 1964 § 52 eq. (52.06); Rindler 2006 § 14.2 eq. (14.8).

the dimension of κ we thus find

$$\dim(\kappa) = \dim(\mathbf{G}_\bullet) \dim(\mathbf{T}_\bullet)^{-1} \equiv \begin{cases} M^{-1}L & \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases} \\ M^{-1}L^{-1}T^2 & \end{cases} \quad (58)$$

This constant can be obtained from the dimensions of Newton's gravitational constant $\dim(G) = M^{-1}L^3T^{-2}$ (this is not the Einstein tensor \mathbf{G} !) and of the speed of light $\dim(c) = LT^{-1}$ only in the following ways, with an 8π factor coming from the Newtonian limit:

$$\kappa = \begin{cases} 8\pi G/c^2 & \text{if } \dim(\mathbf{g}) := \begin{cases} T^2 \\ L^2 \end{cases} \\ 8\pi G/c^4 & \end{cases} \quad (59)$$

The second choice is by far the most common, consistently with the most common choice of $\dim(\mathbf{g}) = L^2$ discussed before. The first choice appears for example in Fock⁴⁴ and Adler et al.⁴⁵.

12 Summary and conclusions

We have seen that dimensional analysis, with its familiar rules, can be seamlessly performed in general relativity and in differential geometry if we adopt the coordinate-free approach typical of modern texts. In this approach each tensor has an *intrinsic* dimension (a notion introduced by Schouten and Dorgelo). This dimension doesn't depend on the dimensions of the coordinates, and is determined by the physical or operational meaning of the tensor. It is therefore generally more profitable to focus on the intrinsic dimension of a tensor rather than on the dimensions of its components. The dimension of each specific component is easily found by formula (11): it's the product of the intrinsic dimension, times the dimension of the i th coordinate function for each contravariant index i , times the inverse of the dimension of the j th coordinate function for each covariant index j .

Dimensional analysis in differential geometry seems to rest on two main conventions: the tensor product and the action of covectors on vectors behave analogously to usual multiplication for the purposes of dimensional analysis. Alternative, equivalent sets of conventions can also be considered, for example involving the exterior derivative.

⁴⁴ Fock 1964 § 55 eqs (55.15) and (52.06).

⁴⁵ Adler et al. 1975 § 10.5 eq. (10.98).

We found or re-derived some essential results for general relativity, in particular that the Riemann $R^{\bullet}...$ and Ricci Ric_{\bullet} curvature tensors are dimensionless, and that the Einstein tensor G_{\bullet}^{\bullet} has the inverse dimension of the metric tensor. Maybe these results can be of importance for some current research involving scales and conformal factors⁴⁶. We also discussed the operational reasons behind two common choices of dimension for the metric tensor.

Since the dimensions of the components are usually different from the intrinsic dimension and depend on the coordinates, I recommend to avoid statements such as “the tensor A_i^{jk} has dimension X ”, because it leaves unclear whether “ A_i^{jk} ” is meant to represent the tensor in general (as in Penrose & Rindler’s notation), or to represent its set of components, or to represent just a specific component.

Dimensional analysis remains a controversial, obscure, but fascinating subject still today, 60 years from Truesdell & Toupin’s remark quoted in the Introduction. For an overview of some recent and creative approaches to it I recommend for example the works by Mari et al.⁴⁷, Domotor and Batitsky⁴⁸, Kitano⁴⁹, the extensive analysis by Dybkaer⁵⁰, the historical review by de Boer⁵¹, and references therein.

Thanks

... to Ingemar Bengtsson and Iván Davidovich for valuable comments on previous drafts. To the staff of the NTNU library for their always prompt support. To Mari, Miri, Emma for continuous encouragement and affection, and to Buster Keaton and Saitama for filling life with awe and inspiration. To the developers and maintainers of L^AT_EX, Emacs, AUCT_EX, Open Science Framework, R, Python, Inkscape, Sci-Hub for making a free and impartial scientific exchange possible. ☒

This work is financially supported by the Kavli Foundation and the Centre of Excellence scheme of the Research Council of Norway (Yasser Roudi group).

⁴⁶ e.g. Röhr & Uggla 2005; Cadoni & Tuveri 2019. ⁴⁷ Mari & Giordani 2012; Frigerio et al. 2010. ⁴⁸ Domotor 2017; Domotor & Batitsky 2016; Domotor 2012. ⁴⁹ Kitano 2013. ⁵⁰ Dybkaer 2010. ⁵¹ De Boer 1995.

Bibliography

(“de X ” is listed under D , “van X ” under V , and so on, regardless of national conventions.)

- Abraham, R., Marsden, J. E., Ratiu, T. (1988): *Manifolds, Tensor Analysis, and Applications*, 2nd ed. (Springer, New York). First publ. 1983.
- Adler, R., Bazin, M., Schiffer, M. (1975): *Introduction to General Relativity*, 2nd ed. (McGraw-Hill, Tokyo). First publ. 1965.
- BIPM (Bureau international des poids et mesures) (1983): 17^e conférence générale des poids et mesures (CGPM). (BIPM, Sèvres). <https://www.bipm.org/en/worldwide-metrology/cgpm/resolutions.html#comptes-rendus>.
- Boothby, W. M. (2003): *An Introduction to Differentiable Manifolds and Riemannian Geometry*, rev. 2nd ed. (Academic Press, Orlando, USA). First publ. 1975.
- Bossavit, A. (1991): *Differential Geometry: for the student of numerical methods in electromagnetism*. https://www.researchgate.net/publication/200018385_Differential_Geometry_for_the_student_of_numerical_methods_in_Electromagnetism.
- Bressan, A. (1978): *Relativistic Theories of Materials*. (Springer, Berlin).
- Bridgman, P. W. (1958): *The Logic of Modern Physics*, 8th pr. (Macmillan, New York). First publ. 1927.
- Burke, W. L. (1980): *Spacetime, Geometry, Cosmology*. (University Science Books, Mill Valley, USA).
- (1983): *Manifestly parity invariant electromagnetic theory and twisted tensors*. *J. Math. Phys.* **24**¹, 65–69.
- (1987): *Applied Differential Geometry*, repr. (Cambridge University Press, Cambridge). First publ. 1985.
- (1995): *Div, Grad, Curl Are Dead*. http://people.ucsc.edu/~rmont/papers/Burke_DivGradCurl.pdf; “preliminary draft II”. See also <http://www.uclick.org/~burke/>.
- Cadoni, M., Tuveri, M. (2019): *Galactic dynamics and long-range quantum gravity*. *Phys. Rev. D* **100**², 024029.
- Choquet-Bruhat, Y., DeWitt-Morette, C., Dillard-Bleick, M. (1996): *Analysis, Manifolds and Physics. Part I: Basics*, rev. ed. (Elsevier, Amsterdam). First publ. 1977.
- Curtis, W. D., Miller, F. R. (1985): *Differential Manifolds and Theoretical Physics*. (Academic Press, Orlando, USA).
- de Boer, J. (1995): *On the history of quantity calculus and the International System*. *Metrologia* **31**⁶, 405–429.
- de Rham, G. (1984): *Differentiable Manifolds: Forms, Currents, Harmonic Forms*. (Springer, Berlin). Transl. by F. R. Smith. First publ. in French 1955.
- Domotor, Z. (2012): *Algebraic frameworks for measurement in the natural sciences*. *Meas. Sci. Rev.* **12**⁶, 213–233.
- (2017): *Torsor theory of physical quantities and their measurement*. *Meas. Sci. Rev.* **17**⁴, 152–177.
- Domotor, Z., Batitsky, V. (2016): *An algebraic approach to unital quantities and their measurement*. *Meas. Sci. Rev.* **16**³, 103–126.
- Dorgelo, H. B., Schouten, J. A. (1946): *On unities and dimensions. [I, II, III]*. *Verh. Kon. Akad. Wetensch. Amsterdam* **49**^{2, 3, 4}, 123–131, 282–291, 393–403.
- Dybkaer, R. (2010): *An Ontology on Property: For Physical, Chemical and Biological Systems*, 2nd ed. (IUPAC, Research Triangle Park, USA). <https://ontology.iupac.org/>. First publ. 2004.

- Eckart, C. (1940): *The thermodynamics of irreversible processes. III. Relativistic theory of the simple fluid*. Phys. Rev. **58**¹⁰, 919–924.
- Fecko, M. (2006): *Differential Geometry and Lie Groups for Physicists*. (Cambridge University Press, Cambridge).
- Flügge, S., ed. (1960): *Handbuch der Physik: Band III/1: Prinzipien der klassischen Mechanik und Feldtheorie [Encyclopedia of Physics: Vol. III/1: Principles of Classical Mechanics and Field Theory]*. (Springer, Berlin).
- Fock [Fok], V. A. (1964): *The Theory of Space, Time and Gravitation*, 2nd rev. ed. (Pergamon, Oxford). Transl. by N. Kemmer. First publ. in Russian 1955.
- Fokker, A. D. (1965): *Time and Space, Weight and Inertia: A chronogeometrical introduction to Einstein's theory*. (Pergamon, Oxford). Transl. by D. Bijl, translation ed. by D. Field. First publ. in Dutch 1960.
- Frigerio, A., Giordani, A., Mari, L. (2010): *Outline of a general model of measurement*. Synthese **175**², 123–149.
- Giacomo, P. (1984): *News from the BIPM*. Metrologia **20**¹, 25–30.
- Gourgoulhon, É. (2012): *3+1 Formalism in General Relativity: Bases of Numerical Relativity*. (Springer, Heidelberg). First publ. 2007 as arXiv:gr-qc/0703035.
- Hartle, J. B. (2003): *Gravity: An Introduction to Einstein's General Relativity*. (Addison-Wesley, San Francisco).
- Hehl, F. W., Obukhov, Y. N. (2003): *Foundations of Classical Electrodynamics: Charge, Flux, and Metric*. (Birkhäuser, Boston).
- ISO (International Organization for Standardization) (2009): *ISO 80000-1:2009: Quantities and units 1: General*. International Organization for Standardization.
- Kilmister, C. W. (1973): *General Theory of Relativity*. (Pergamon, Oxford).
- Kitano, M. (2013): *Mathematical structure of unit systems*. J. Math. Phys. **54**⁵, 052901.
- Landau, L. D., Lifshitz [Lifšic], E. M. (1996): *The Classical Theory of Fields*, corr. repr. of the 4th English ed. (Butterworth-Heinemann, Oxford). Transl. from the 1987 seventh Russian edition by Morton Hamermesh. First publ. 1939.
- Mari, L., Giordani, A. (2012): *Quantity and quantity value*. Metrologia **49**⁶, 756–764.
- Marsden, J. E., Hughes, T. J. R. (1994): *Mathematical Foundations of Elasticity*, unabridged corr. republ. (Dover, New York). <http://resolver.caltech.edu/CaltechB00K:1983.002>. First publ. 1983.
- Martin, D. (2004): *Manifold Theory: Introduction for Mathematical Physicists*, corr. repr. (Horwood, Chichester). First publ. 1991.
- Mauguin, G. A. (1974): *Constitutive equations for heat conduction in general relativity*. J. Phys. A **7**⁴, 465–484.
- (1978a): *On the covariant equations of the relativistic electrodynamics of continua. I. General equations*. J. Math. Phys. **19**⁵, 1198–1205.
 - (1978b): *On the covariant equations of the relativistic electrodynamics of continua. II. Fluids*. J. Math. Phys. **19**⁵, 1206–1211.
 - (1978c): *On the covariant equations of the relativistic electrodynamics of continua. III. Elastic solids*. J. Math. Phys. **19**⁵, 1212–1219.
 - (1978d): *On the covariant equations of the relativistic electrodynamics of continua. IV. Media with spin*. J. Math. Phys. **19**⁵, 1220–1226.
- Misner, C. W., Thorne, K. S., Wheeler, J. A. (1973): *Gravitation*, repr. (W. H. Freeman and Company, New York). https://archive.org/details/Gravitation_201803.
- Muschik, W., von Borzeszkowski, H.-H. (2014): *Exploitation of the dissipation inequality in general relativistic continuum thermodynamics*. Arch. Appl. Mech. **84**⁹, 1517–1531.

- Penrose, R., Rindler, W. (2003): *Spinors and Space-Time. Vol. 1: Two-spinor calculus and relativistic fields*, corr. repr. (Cambridge University Press, Cambridge). First publ. 1984.
- Rindler, W. (1986): *Essential Relativity: Special, General, and Cosmological*, rev. 2nd ed. (Springer, New York). First publ. 1969.
- (2006): *Relativity: Special, General, and Cosmological*, 2nd ed. (Oxford University Press, Oxford). First publ. 2001.
- Röhr, N., Uggla, C. (2005): *Conformal regularization of Einstein's field equations*. *Class. Quantum Grav.* **22**¹⁷, 3775–3787.
- Schouten, J. A. (1954): *Ricci-Calculus: An Introduction to Tensor Analysis and Its Geometrical Applications*, 2nd ed. (Springer, Berlin). First publ. in German 1924.
- (1989): *Tensor Analysis for Physicists*, corr. 2nd ed. (Dover, New York). First publ. 1951.
- Segev, R. (1986): *Forces and the existence of stresses in invariant continuum mechanics*. *J. Math. Phys.* **27**¹, 163–170. <http://www.bgu.ac.il/~rsegev/Papers/JMP86S.pdf>.
- (2000a): *The geometry of Cauchy's fluxes*. *Arch. Rational Mech. Anal.* **154**³, 183–198. <http://www.bgu.ac.il/~rsegev/Papers/TheGeometryofCauchyFluxes.pdf>.
- (2000b): *Notes on stresses for manifolds*. *Rend. Sem. Mat. Univ. Pol. Torino* **58**², 199–206.
- (2001): *A correction of an inconsistency in my paper "Cauchy's theorem on manifolds"*. *J. Elasticity* **63**¹, 55–59. See Segev, Rodnay (1999).
- (2002): *Metric-independent analysis of the stress-energy tensor*. *J. Math. Phys.* **43**⁶, 3220–3231. <http://www.bgu.ac.il/~rsegev/Papers/MetricIndependent.pdf>.
- Segev, R., Rodnay, G. (1999): *Cauchy's theorem on manifolds*. *J. Elasticity* **56**², 129–144. <http://www.bgu.ac.il/~rsegev/Papers/CauchyTheoremOnManifolds.pdf>. See also erratum Segev (2001).
- Smarr, L., Taubes, C., Wilson, J. R. (1980): *General relativistic hydrodynamics: the comoving, Eulerian, and velocity potential formalisms*. In: Tipler (1980), ch. 11, 157–183.
- Smarr, L., York Jr., J. W. (1978): *Kinematical conditions in the construction of spacetime*. *Phys. Rev. D* **17**¹⁰, 2529–2551.
- Smarr, L. L., ed. (1979): *Sources of Gravitational Radiation*. (Cambridge University Press, Cambridge).
- Synge, J. L. (1960a): *Classical Dynamics*. In: Flügge (1960), I–VII, 1–225, 859–902.
- (1960b): *Relativity: The General Theory*. (North-Holland, Amsterdam).
- Tipler, F. J., ed. (1980): *Essays in General Relativity: A Festschrift for Abraham Taub*. (Academic Press, New York).
- Tolman, R. C. (1949): *Relativity, Thermodynamics and Cosmology*. (Oxford University Press). First publ. 1934.
- Truesdell III, C. A. (1991): *A First Course in Rational Continuum Mechanics. Vol. 1: General Concepts*, 2nd ed. (Academic Press, New York). First publ. 1977.
- Truesdell III, C. A., Toupin, R. A. (1960): *The Classical Field Theories*. In: Flügge (1960), I–VII, 226–902. With an appendix on invariants by Jerald LaVerne Ericksen.
- Wilson, J. R., Mathews, G. J. (2007): *Relativistic Numerical Hydrodynamics*, repr. (Cambridge University Press, Cambridge). First publ. 2003.
- York Jr., J. W. (1979): *Kinematics and dynamics of general relativity*. In: Smarr (1979), 83–126.