

Model-free prognosis and decision in personalized medicine: A case study on the conversion from Mild Cognitive Impairment to Alzheimer's Disease

P.G.L. Porta Mana^{1,2,*}, Ingrid Rye³, Alexandra Vik^{1,2}, Marek Kocinski^{2,4}, Arvid Lundervold^{2,4}, Astri J. Lundervold³, and Alexander S. Lundervold^{1,2}

¹*Department of Computer Science, Electrical Engineering and Mathematical Sciences, Western Norway University of Applied Sciences, Bergen, Norway*

²*Mohn Medical Imaging and Visualization Centre (MMIV), Department of Radiology, Haukeland University Hospital, Bergen, Norway*

³*Department of Biological and Medical Psychology, University of Bergen, Norway*

⁴*Department of Biomedicine, University of Bergen, Norway*

Correspondence*:
Corresponding Author
pgl@portamana.org

2 ABSTRACT

3 [Luca & Astri, draft]

4 Patients with Mild Cognitive Impairment (MCI) have an increased risk of a trajectory toward
5 Alzheimer's disease (AD). Early identification of patients with a high risk of AD is essential to
6 provide treatment before the disease is well-established in the brain. It is, therefore, of great
7 importance to study how well different kinds of predictors allow us to estimate a trajectory from
8 MCI towards AD in an individual patient.

9 But more is needed for a personalized approach to prognosis, prevention, and treatment, than
10 just the obvious requirement that prognoses be as best as they can be for each patient. Several
11 situational elements that can be different from patient to patient must be accounted for:

- 12 • the *kinds* of clinical data and evidence available for prognosis;
- 13 • the *outcomes* of the same kind of clinical data and evidence;
- 14 • the kinds of treatment or prevention strategies available, owing to different attitudes toward
15 life, different family networks and possibilities of familial support, different additional medical
16 factors such as physical disabilities, and different economic means;
- 17 • the advantages and disadvantages, benefits and costs of the same kinds of treatment or
18 prevention strategies; the patient has a major role in the quantification of such benefits and
19 costs;
- 20 • finally, the initial evaluation by the clinician – which often relies on too subtle clues (family
21 history, regional history, previous case experience) to be considered as measurable data.

22 Statistical decision theory is the normative quantification framework that takes into account these
23 fundamental differences. Medicine has the distinction of having been one of the first fields to
24 adopt this framework, exemplified in brilliant old and new textbooks on clinical decision-making.

25 Clinical decision-making makes allowance for these differences among patients through two
26 requirements. First, the quantification of prognostic evidence on one side, and of benefits and
27 costs of treatments and prevention strategies on the other, must be clearly separated and handled
28 in a modular way. Two patients can have the same prognostic evidence and yet very different
29 prevention options. Second, the quantification of independent prognostic evidence ought to be in
30 the form of *likelihoods about the health condition* (or equivalently of likelihood ratios, in a binary
31 case), that is, of the probabilities of the observed test outcomes given the hypothesized health
32 conditions. Likelihoods from independent clinical tests and predictors can then be combined
33 with a simple multiplication; for one patient, we could have three kinds of predictor available;
34 for another, we could have five. The clinician's pre-test assessment is included in the form of a
35 probability. These patient-dependent probabilities are combined with the patient-dependent costs
36 and benefits of treatment or prevention to arrive at the best course of action for that patient. The
37 main result underlying statistical decision theory is that decision-making *must* take this particular
38 mathematical form in order to be optimal and logically consistent.

39 The present work investigates the prognostic power of a set of neuropsychological and
40 Magnetic Resonance Imaging examinations, demographic data, and genetic information about
41 Apolipoprotein-E4 (APOE) status, for the prediction of the onset of Alzheimer's disease in patients
42 defined as mildly cognitively impaired at a baseline examination. The longitudinal data used come
43 from the ADNI database.

44 The prognostic power of these predictors is quantified in the form of a combined likelihood for
45 the onset of Alzheimer's disease. As a hypothetical example application of personalized clinical
46 decision making, three patient cases are considered where a clinician starts with prognostic
47 uncertainties, possibly coming from other tests, of 50%/50%, 25%/75%, 75%/25%. It is shown
48 how these pre-test probabilities are changed by the predictors.

49 This quantification also allows us to rank the relative prognostic power of the predictors. It is
50 found that several neuropsychological examinations have the highest prognostic power, much
51 higher than the genetic and imaging-derived predictors included in the present set.

52 Several additional advantages of this quantification framework are also exemplified and
53 discussed in the present work:

- 54 • missing data are automatically handled, and results having partial data are not discarded; this
55 quantification, therefore, also accounts for patient-dependent availability of *non-independent*
56 predictors;
- 57 • no modelling assumptions (e.g., linearity, gaussianity, functional dependence) are made;
- 58 • the prognostic power obtained is intrinsic to the predictors, that is, it is a bound for *any*
59 prognostic algorithm;
- 60 • variability ranges of the results owing to the finite size of the sample data are automatically
61 quantified.
- 62 • the values obtained, being probabilities, are more easily interpretable than scores of various
63 kinds.



64 Keywords: keyword, keyword, keyword, keyword, keyword, keyword, keyword, keyword

1 EACH PATIENT IS UNIQUE

65 Meet Olivia, Ariel, Bianca, Curtis.¹ These four persons don't know each other, but they have something in
66 common: they all suffer from a mild form of cognitive impairment, and are afraid that their impairment
67 will turn into Alzheimer's disease within a couple of years. In fact, this is why they recently underwent
68 some clinical analyses and cognitive tests. Today they got the results of their analyses. From these results
69 and other demographic factors their clinician is assessing their risk of developing Alzheimer, and will then
70 decide on possible preventive treatments together with the patients.

71 Besides this shared condition and worry, these patients have other things in common – but also some
72 differences. Let's take Olivia as reference and see the similarities and difference between her and the other
73 three:

- 74 • Olivia and Ariel turn out to have exactly identical clinical results, age, and geographical origin, and
75 very similar family histories. Ariel, however, is soon going to move to another country where a new
76 preventive treatment is available; this option is not open to Olivia.
- 77 • Olivia and Bianca also have exactly identical clinical results and age, and the same preventive options.
78 Bianca, however, comes from a different geographical region, having lower conversion rate, 30%, from
79 Mild Cognitive Impairment to Alzheimer. Moreover there is no history of Alzheimer in her family.
- 80 • Olivia and Curtis have different clinical results and age – in particular, Olivia has the risky APOE4
81 allele whereas Curtis hasn't, and Curtis is more than 20 years younger. But they otherwise come from
82 the same geographical region, have very similar family histories, and the same preventive options.

83 Figure  *** summarizes the similarity and differences between Olivia and the other three patients. Table 
84 *** reports the clinical results and demographic data common to Olivia, Ariel, Bianca; and those of Curtis.

¹ Fictive characters; any reference to real persons is purely coincidental

— Luca, old pieces of text —

Personalized diagnosis, prognosis, treatment, and prevention strategies must make allowance for several fundamental differences among patients:

- the *kinds* of clinical data and evidence available for diagnosis or prognosis can be different;
- the *values* of the same kind of clinical data and evidence can be different;
- the kinds of treatment or prevention options can be different;
- the advantages and disadvantages, benefits and costs of the same kinds of treatment or prevention can be different;
- finally, the evaluation of the clinician – which often relies on too subtle clues (family history, regional history, case experience) to be considered as measurable data – can be different.

Is there really a methodological framework that can take all these differences into account? Yes, there is, and Medicine has the distinction of having been one of the first fields to adopt it (Ledley and Lusted, 1959): Statistical Decision Theory. Its application in Medicine is explained and exemplified in several, brilliant, old and new textbooks (Weinstein and Fineberg, 1980; Sox et al., 2013; Hunink et al., 2014). This theory has mathematical and logical foundations and its principles constitute indeed the foundations for the definition and realization of Artificial Intelligence (Russell and Norvig, 2022) 🔧

The basics of clinical decision making 🔧 ..basics: each piece of evidence contributes with a likelihood or odds; they combine together and together with the clinician's pre-data evaluation. Then they are combined with the different benefits/costs of treatments or prevention strategies to find the optimal one. Decision trees can be necessary (but don't change this framework). Costs & benefits are evaluated by clinician & patient together.

$$\begin{array}{c}
 \overbrace{\text{post-test probability}} \\
 p(\text{health condition} \mid \text{results of all tests, prior info}) \propto \\
 \\
 \overbrace{\text{pre-test probability by clinician}} \\
 p(\text{health condition} \mid \text{prior info}) \times \\
 \\
 \text{likelihoods of tests} \left\{ \begin{array}{l} p(\text{result of 1st test} \mid \text{health condition, prior info}) \times \\ p(\text{result of 2nd test} \mid \text{health condition, prior info}) \times \\ \dots \end{array} \right. \quad (1)
 \end{array}$$

For Original Research Articles (Name et al., 1996), Clinical Trial Articles (LastName1 et al., 2013), and Technology Reports (Surname1, 2010), the introduction should be succinct, with no subheadings (Name, 1993). For Case Reports the Introduction should include symptoms at presentation (Surname, 2002), physical exams and lab results (LastName1 et al., 2011).

2 ARTICLE TYPES

For requirements for a specific article type please refer to the Article Types on any Frontiers journal page. Please also refer to Author Guidelines for further information on how to organize your manuscript in the required sections or their equivalents for your field

3 MANUSCRIPT FORMATTING

3.1 Heading Levels

3.2 Level 2

3.2.1 Level 3

3.2.1.1 Level 4

3.2.1.1.1 Level 5

3.3 Equations

Equations should be inserted in editable format from the equation editor.

$$\sum x + y = Z \quad (2)$$

3.4 Figures

Frontiers requires figures to be submitted individually, in the same order as they are referred to in the manuscript. Figures will then be automatically embedded at the bottom of the submitted manuscript. Kindly ensure that each table and figure is mentioned in the text and in numerical order. Figures must be of sufficient resolution for publication see here for examples and minimum requirements. Figures which are not according to the guidelines will cause substantial delay during the production process. Please see here for full figure guidelines. Cite figures with subfigures as figure 2a and 2b.

3.4.1 Permission to Reuse and Copyright

Figures, tables, and images will be published under a Creative Commons CC-BY licence and permission must be obtained for use of copyrighted material from other sources (including re-published/adapted/modified/partial figures and images from the internet). It is the responsibility of the authors to acquire the licenses, to follow any citation instructions requested by third-party rights holders, and cover any supplementary charges.

3.5 Tables

Tables should be inserted at the end of the manuscript. Please build your table directly in LaTeX. Tables provided as jpeg/tiff files will not be accepted. Please note that very large tables (covering several pages) cannot be included in the final PDF for reasons of space. These tables will be published as Supplementary Material on the online article page at the time of acceptance. The author will be notified during the typesetting of the final article if this is the case.

4 NOMENCLATURE

4.1 Resource Identification Initiative

To take part in the Resource Identification Initiative, please use the corresponding catalog number and RRID in your current manuscript. For more information about the project and for steps on how to search for an RRID, please click here.

4.2 Life Science Identifiers

Life Science Identifiers (LSIDs) for ZOOBANK registered names or nomenclatural acts should be listed in the manuscript before the keywords. For more information on LSIDs please see Inclusion of Zoological Nomenclature section of the guidelines.

5 ADDITIONAL REQUIREMENTS

For additional requirements for specific article types and further information please refer to Author Guidelines.

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

The Author Contributions section is mandatory for all articles, including articles by sole authors. If an appropriate statement is not provided on submission, a standard one will be inserted during the production process. The Author Contributions statement must describe the contributions of individual authors referred to by their initials and, in doing so, all authors agree to be accountable for the content of the work. Please see here for full authorship criteria.

FUNDING

Details of all funding sources should be provided, including grant numbers if applicable. Please ensure to add all necessary funding information, as after publication this is no longer possible.

ACKNOWLEDGMENTS

This is a short text to acknowledge the contributions of specific colleagues, institutions, or agencies that aided the efforts of the authors.

SUPPLEMENTAL DATA

Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures, please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be found in the Frontiers LaTeX folder.

DATA AVAILABILITY STATEMENT

The datasets [GENERATED/ANALYZED] for this study can be found in the [NAME OF REPOSITORY]
[LINK].

REFERENCES

- Hunink, M. G. M., Weinstein, M. C., Wittenberg, E., Drummond, M. F., Pliskin, J. S., Wong, J. B., et al. (2014). *Decision Making in Health and Medicine: Integrating Evidence and Values* (Cambridge: Cambridge University Press), 2 edn. doi:10.1017/CBO9781139506779. First publ. 2001
- [Dataset] LastName1, A., LastName2, A., and LastName3, A. (2011). Data title. doi:10.000/55555
- LastName1, A., LastName2, A., and LastName3, A. (2013). Article title. *Frontiers in Neuroscience* 30, 10127–10134. doi:10.3389/fnins.2013.12345
- Ledley, R. S. and Lusted, L. B. (1959). Reasoning foundations of medical diagnosis: Symbolic logic, probability, and value theory aid our understanding of how physicians reason. *Science* 130, 9–21. doi:10.1126/science.130.3366.9
- Name, A. (1993). *The title of the work* (The city: The name of the publisher)
- Name, C., Surname, D., and LastName, F. (1996). The title of the work. In *The title of the conference proceedings*, eds. E. Name1 and E. Name2 (The name of the publisher), 41–50
- Russell, S. J. and Norvig, P. (2022). *Artificial Intelligence: A Modern Approach*. Pearson series in artificial intelligence (Harlow, UK: Pearson), fourth global ed. edn. <http://aima.cs.berkeley.edu/global-index.html>, <https://archive.org/details/artificial-intelligence-a-modern-approach-4th-edition>. First publ. 1995
- Sox, H. C., Higgins, M. C., and Owens, D. K. (2013). *Medical Decision Making* (New York: Wiley), 2 edn. doi:10.1002/9781118341544. First publ. 1988
- Surname, B. (2002). The title of the work. In *The title of the book*, ed. E. Name (The city: The name of the publisher). 201–213
- Surname1, H. (2010). *The title of the work* (Patent country: Patent number)
- Weinstein, M. C. and Fineberg, H. V. (1980). *Clinical Decision Analysis* (Philadelphia: Saunders)

FIGURE CAPTIONS



Figure 1. Enter the caption for your figure here. Repeat as necessary for each of your figures



Figure 2a. This is Subfigure 1.



Figure 2b. This is Subfigure 2.

Figure 2. Enter the caption for your subfigure here. **(A)** This is the caption for Subfigure 1. **(B)** This is the caption for Subfigure 2.