

Model-free prognosis and decision in personalized medicine: A case study on the conversion from Mild Cognitive Impairment to Alzheimer's Disease

P.G.L. Porta Mana^{1,2,*}, I. Rye³, A. Vik^{1,2}, M. Kociński^{2,4}, A. Lundervold^{2,4}, A. J. Lundervold³, A. S. Lundervold^{1,2}

¹Department of Computer Science, Electrical Engineering and Mathematical Sciences, Western Norway University of Applied Sciences, Bergen, Norway

²Mohn Medical Imaging and Visualization Centre (MMIV), Department of Radiology, Haukeland University Hospital, Bergen, Norway

³Department of Biological and Medical Psychology, University of Bergen, Norway

⁴Department of Biomedicine, University of Bergen, Norway

Correspondence*:
Corresponding Author
pgl@portamana.org

2 ABSTRACT

3

4 Patients with Mild Cognitive Impairment have an increased risk of a trajectory toward Alzheimer's
5 Disease. Early identification of patients with a high risk of Alzheimer's Disease is essential to
6 provide treatment before the disease is well-established in the brain. great importance to study
7 how well different kinds of predictors allow us to prognose a trajectory from Mild Cognitive
8 Impairment towards Alzheimer's Disease in an individual patient.

9 But more is needed for a personalized approach to prognosis, prevention, and treatment, than
10 just the obvious requirement that prognoses be as best as they can be for each patient. Several
11 situational elements that can be different from patient to patient must be accounted for:


- 12 • the *kinds* of clinical data and evidence available for prognosis;
- 13 • the *outcomes* of the same kind of clinical data and evidence;
- 14 • the kinds of treatment or prevention strategies available, owing to different additional medical
15 factors such as physical disabilities, different attitudes toward life, different family networks
16 and possibilities of familial support, different economic means;
- 17 • the advantages and disadvantages, benefits and costs of the same kinds of treatment or
18 prevention strategies; the patient has a major role in the quantification of such benefits and
19 costs;
- 20 • finally, the initial evaluation by the clinician – which often relies on too subtle clues (family
21 history, regional history, previous case experience) to be considered as measurable data.


22 Statistical decision theory is the normative quantification framework that takes into account these
23 fundamental differences. Medicine has the distinction of having been one of the first fields to
24 adopt this framework, exemplified in brilliant old and new textbooks on clinical decision-making.

25 Clinical decision-making makes allowance for these differences among patients through two
26 requirements. First, the quantification of prognostic evidence on one side, and of benefits and
27 costs of treatments and prevention strategies on the other, must be clearly separated and handled
28 in a modular way. Two patients can have the same prognostic evidence and yet very different
29 prevention options. Second, the quantification of independent prognostic evidence ought to be in
30 the form of *likelihoods about the health condition* (or equivalently of likelihood ratios, in a binary
31 case), that is, of the probabilities of the observed test outcomes given the hypothesized health
32 conditions. Likelihoods from independent clinical tests and predictors can then be combined
33 with a simple multiplication; for one patient, we could have three kinds of predictor available;
34 for another, we could have five. The clinician's pre-test assessment is included in the form of a
35 probability. These patient-dependent probabilities are combined with the patient-dependent costs
36 and benefits of treatment or prevention to arrive at the best course of action for that patient. The
37 main result underlying statistical decision theory is that decision-making *must* take this particular
38 mathematical form in order to be optimal and logically consistent.

39 The present work investigates the prognostic power of a set of neuropsychological and Mag-
40 netic Resonance Imaging examinations, demographic data, and genetic information about
41 Apolipoprotein-E4. The present work investigates the prognostic power of a set of neuropsycholo-
42 gical and Magnetic Resonance Imaging examinations, demographic data, and genetic information
43 about Apolipoprotein-E4 (APOE) status, for the prediction of the onset of Alzheimer's Disease in
44 patients defined as mildly cognitively impaired at a baseline examination. The longitudinal data
45 used come from the ADNI database.

46 (APOE) status, for the prediction of the onset of Alzheimer's disease in patients defined as
47 mildly cognitively impaired at a baseline examination. The longitudinal data used come from the
48 ADNI database.

49 The prognostic power of these predictors is quantified in the form of a combined likelihood for
50 the onset of Alzheimer's disease. As a hypothetical example application of personalized clinical
51 decision making, three patient cases are considered where a clinician starts with prognostic
52 uncertainties, possibly coming from other tests, of 50%/50%, 25%/75%, 75%/25%. It is shown
53 how these pre-test probabilities are changed by the predictors.  [update this](#)

54  [rewrite following](#) This quantification also allows us to rank the relative prognostic power of the
55 predictors. It is found that several neuropsychological examinations have the highest prognostic
56 power, much higher than the genetic and imaging-derived predictors included in the present set.

57 Several additional advantages of this quantification framework are also exemplified and
58 discussed in the present work:

- 59 • missing data are automatically handled, and results having partial data are not discarded; this
60 quantification, therefore, also accounts for patient-dependent availability of *non-independent*
61 predictors;
- 62 • no modelling assumptions (e.g., linearity, gaussianity, functional dependence) are made;

- the prognostic power obtained is intrinsic to the predictors, that is, it is a bound for *any* prognostic algorithm;
- variability ranges of the results owing to the finite size of the sample data are automatically quantified.
- the values obtained, being probabilities, are more easily interpretable than scores of various kinds.

Keywords: Clinical decision making, Utility theory, Probability theory, Artificial Intelligence, Machine Learning, Base-rate fallacy

1 EACH PATIENT IS UNIQUE

Meet Olivia, Ariel, Bianca, Curtis.¹ These four persons don't know each other, but they have something in common: they all suffer from a mild form of cognitive impairment, and are afraid that their impairment will turn into Alzheimer's Disease within a couple of years. In fact, this is why they recently underwent some clinical analyses and cognitive tests. Today they got the results of their analyses. From these results and other relevant information, their clinician will assess their risk of developing Alzheimer. Clinician and patients will then decide among a set of possible preventive treatments.

Besides this shared condition and worry, these patients have other things in common – but also some differences. Let's take Olivia as reference and list the similarities and difference between her and the other three:

- Olivia and Ariel turn out to have exactly identical clinical results and age. They would also get similar benefits from the available preventive-treatment options. Ariel, however, comes from a different geographical region from Olivia, where the conversion rate from Mild Cognitive Impairment to Alzheimer is lower: about 30%, whereas in Olivia's it is around 45% (Petersen et al., 1999) Moreover there is no history of Alzheimer's Disease in Ariel's family.
- Olivia and Bianca also have exactly the same clinical results and age. They come from the same geographical region and have very similar family histories. In fact we shall see that they have the same probability of developing Alzheimer's disease. Bianca, however, suffers from several allergies and additional clinical conditions that would render some of the preventive options less beneficial to her.
- Olivia and Curtis have different clinical results – in particular, Olivia has the risky Apolipoprotein-E4 (APOE4) allele (?) whereas Curtis hasn't – and age, Olivia being more than 20 years older than Curtis. But they otherwise come from the same geographical region, have very similar family histories, and would get similar benefits from the preventive options.

We can categorize these differences as “difference in auxiliary information” (Olivia and Ariel), “difference in preventive benefits” (Olivia and Bianca), “difference in clinical factors” (Olivia and Curtis). Figure *** summarizes the similarity and differences between Olivia and the other three patients. Table *** reports the clinical results and demographic data common to Olivia, Ariel, Bianca, as well as those of Curtis.

Considering the similarities and differences among these patients, which treatments will be prescribed to them?

The purpose of the present work

¹ Fictive characters; any reference to real persons is purely coincidental




Figure 1. draft, needs better font sizes

Table 1. Clinical results & demographic data

Patient	Age	Sex	HC · 10 ^{−3}	APOE4	ANART	CFT	GDS	RAVLT-im	RAVLT-del	RAVLT-rec	TMTA	TMTB
Olivia, Ariel, Bianca	83.5	F	2.77	Y	14	18	0	29	12	3	126.5	117.0
Curtis	61.5	M	4.17	N	6	13	1	16	2	0	24.8	83.4

100 We will soon see that despite the many factors in common among these four patients – even despite the
101 identical clinical results for Olivia, Ariel, Bianca, and despite the identical probability of conversion for
102 Olivia and Bianca – *the optimal preventive option for each patient is different from those for the other three.*
103 This difference arises from the difference in clinical results, or auxiliary information, or preventive benefits.

— Luca, old pieces of text —

Personalized diagnosis, prognosis, treatment, and prevention strategies must make allowance for several fundamental differences among patients:

- the *kinds* of clinical data and evidence available for diagnosis or prognosis can be different;
- the *values* of the same kind of clinical data and evidence can be different;
- the kinds of treatment or prevention options can be different;
- the advantages and disadvantages, benefits and costs of the same kinds of treatment or prevention can be different;
- finally, the evaluation of the clinician – which often relies on too subtle clues (family history, regional history, case experience) to be considered as measurable data – can be different.

Is there really a methodological framework that can take all these differences into account? Yes, there is, and Medicine has the distinction of having been one of the first fields to adopt it (Ledley and Lusted, 1959): Statistical Decision Theory. Its application in Medicine is explained and exemplified in several, brilliant, old and new textbooks (Weinstein and Fineberg, 1980; Sox et al., 2013; Hunink et al., 2014). This theory has mathematical and logical foundations and its principles constitute indeed the foundations for the definition and realization of Artificial Intelligence (Russell and Norvig, 2022) 🔧

The basics of clinical decision making 🔧 ..basics: each piece of evidence contributes with a likelihood or odds; they combine together and together with the clinician's pre-data evaluation. Then they are combined with the different benefits/costs of treatments or prevention strategies to find the optimal one. Decision trees can be necessary (but don't change this framework). Costs & benefits are evaluated by clinician & patient together.

post-test probability

$p(\text{health condition} \mid \text{results of all tests, prior info}) \propto$

pre-test probability by clinician

$p(\text{health condition} \mid \text{prior info}) \times$

$$\text{likelihoods of tests} \left\{ \begin{array}{l} p(\text{result of 1st test} \mid \text{health condition, prior info}) \times \\ p(\text{result of 2nd test} \mid \text{health condition, prior info}) \times \\ \dots \end{array} \right. \quad (1)$$

For Original Research Articles (Name et al., 1996), Clinical Trial Articles (LastName1 et al., 2013), and Technology Reports (Surname1, 2010), the introduction should be succinct, with no subheadings (Name, 1993). For Case Reports the Introduction should include symptoms at presentation (Surname, 2002), physical exams and lab results (LastName1 et al., 2011).

2 ARTICLE TYPES

128 For requirements for a specific article type please refer to the Article Types on any Frontiers journal page.
129 Please also refer to Author Guidelines for further information on how to organize your manuscript in the
130 required sections or their equivalents for your field

3 MANUSCRIPT FORMATTING

131 3.1 Heading Levels

132 3.2 Level 2

133 3.2.1 Level 3

134 3.2.1.1 Level 4

135 3.2.1.1.1 Level 5

136 3.3 Equations

137 Equations should be inserted in editable format from the equation editor.

$$\sum x + y = Z \quad (2)$$

138 3.4 Figures

139 Frontiers requires figures to be submitted individually, in the same order as they are referred to in the
140 manuscript. Figures will then be automatically embedded at the bottom of the submitted manuscript. Kindly
141 ensure that each table and figure is mentioned in the text and in numerical order. Figures must be of
142 sufficient resolution for publication see here for examples and minimum requirements. Figures which are
143 not according to the guidelines will cause substantial delay during the production process. Please see here
144 for full figure guidelines. Cite figures with subfigures as figure ?? and ??.

145 3.4.1 Permission to Reuse and Copyright

146 Figures, tables, and images will be published under a Creative Commons CC-BY licence and
147 permission must be obtained for use of copyrighted material from other sources (including re-
148 published/adapted/modified/partial figures and images from the internet). It is the responsibility of the
149 authors to acquire the licenses, to follow any citation instructions requested by third-party rights holders,
150 and cover any supplementary charges.

151 3.5 Tables

152 Tables should be inserted at the end of the manuscript. Please build your table directly in LaTeX. Tables
153 provided as jpeg/tiff files will not be accepted. Please note that very large tables (covering several pages)
154 cannot be included in the final PDF for reasons of space. These tables will be published as Supplementary
155 Material on the online article page at the time of acceptance. The author will be notified during the
156 typesetting of the final article if this is the case.

4 NOMENCLATURE

4.1 Resource Identification Initiative

To take part in the Resource Identification Initiative, please use the corresponding catalog number and RRID in your current manuscript. For more information about the project and for steps on how to search for an RRID, please click [here](#).

4.2 Life Science Identifiers

Life Science Identifiers (LSIDs) for ZOOBANK registered names or nomenclatural acts should be listed in the manuscript before the keywords. For more information on LSIDs please see Inclusion of Zoological Nomenclature section of the guidelines.

5 ADDITIONAL REQUIREMENTS

For additional requirements for specific article types and further information please refer to Author Guidelines.

CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

The Author Contributions section is mandatory for all articles, including articles by sole authors. If an appropriate statement is not provided on submission, a standard one will be inserted during the production process. The Author Contributions statement must describe the contributions of individual authors referred to by their initials and, in doing so, all authors agree to be accountable for the content of the work. Please see [here](#) for full authorship criteria.

FUNDING

Details of all funding sources should be provided, including grant numbers if applicable. Please ensure to add all necessary funding information, as after publication this is no longer possible.

ACKNOWLEDGMENTS

This is a short text to acknowledge the contributions of specific colleagues, institutions, or agencies that aided the efforts of the authors.

SUPPLEMENTAL DATA

Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures, please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be found in the Frontiers LaTeX folder.

DATA AVAILABILITY STATEMENT

181 The datasets [GENERATED/ANALYZED] for this study can be found in the [NAME OF REPOSITORY]
182 [LINK].

REFERENCES

- 183 Hunink, M. G. M., Weinstein, M. C., Wittenberg, E., Drummond, M. F., Pliskin, J. S., Wong, J. B.,
184 et al. (2014). *Decision Making in Health and Medicine: Integrating Evidence and Values* (Cambridge:
185 Cambridge University Press), 2 edn. doi:10.1017/CBO9781139506779. First publ. 2001
186 [Dataset] LastName1, A., LastName2, A., and LastName3, A. (2011). Data title. doi:10.000/55555
187 LastName1, A., LastName2, A., and LastName3, A. (2013). Article title. *Frontiers in Neuroscience* 30,
188 10127–10134. doi:10.3389/fnins.2013.12345
189 Ledley, R. S. and Lusted, L. B. (1959). Reasoning foundations of medical diagnosis: Symbolic logic,
190 probability, and value theory aid our understanding of how physicians reason. *Science* 130, 9–21.
191 doi:10.1126/science.130.3366.9
192 Liu, C.-C., Kanekiyo, T., Xu, H., and Bu, G. (2013). Apolipoprotein E and Alzheimer disease: risk,
193 mechanisms and therapy. *Nat. Rev. Neurol.* 9, 106–118, 184. doi:10.1038/nrneurol.2012.263, doi:10.
194 1038/nrneurol.2013.32
195 Name, A. (1993). *The title of the work* (The city: The name of the publisher)
196 Name, C., Surname, D., and LastName, F. (1996). The title of the work. In *The title of the conference*
197 *proceedings*, eds. E. Name1 and E. Name2 (The name of the publisher), 41–50
198 Petersen, R. C., Smith, G. E., Waring, S. C., Ivnik, R. J., Tangalos, E. G., and Kokmen, E. (1999).
199 Mild cognitive impairment: Clinical characterization and outcome. *Arch. Neurol.* 56, 303–308, 760.
200 doi:10.1001/archneur.56.3.303, [https://jamanetwork.com/journals/jamaneurology/](https://jamanetwork.com/journals/jamaneurology/fullarticle/775068)
201 [fullarticle/775068](https://jamanetwork.com/journals/jamaneurology/fullarticle/775068)
202 Russell, S. J. and Norvig, P. (2022). *Artificial Intelligence: A Modern Approach*. Pear-
203 son series in artificial intelligence (Harlow, UK: Pearson), fourth global ed. edn.
204 <http://aima.cs.berkeley.edu/global-index.html>, [https://archive.org/](https://archive.org/details/details/artificial-intelligence-a-modern-approach-4th-edition)
205 [details/artificial-intelligence-a-modern-approach-4th-edition](https://archive.org/details/details/artificial-intelligence-a-modern-approach-4th-edition). First
206 publ. 1995
207 Sox, H. C., Higgins, M. C., and Owens, D. K. (2013). *Medical Decision Making* (New York: Wiley), 2 edn.
208 doi:10.1002/9781118341544. First publ. 1988
209 Surname, B. (2002). The title of the work. In *The title of the book*, ed. E. Name (The city: The name of the
210 publisher). 201–213
211 Surname1, H. (2010). *The title of the work* (Patent country: Patent number)
212 Weinstein, M. C. and Fineberg, H. V. (1980). *Clinical Decision Analysis* (Philadelphia: Saunders)

FIGURE CAPTIONS

”””