

Regression and exchangeability

Luca

<piero.mana@ntnu.no>

Draft of 24 December 2018 (first drafted 24 December 2018)

Exchangeability as a form of regression, regression as a form of exchangeability.

Note: Dear Reader & Peer, this manuscript is being peer-reviewed by you. Thank you.

1 Regression

‘Regression’, as usually intended, is the problem of inferring the value of a quantity x given that of another quantity t ; the two quantities having values in two specific sets. It is also assumed that in two or more observations with the same value of t the values of x may be different.

In this note I use ‘regression’ in a restricted sense: two or more observations with the same value of t have also the same value of x . So the second quantity is given by a function of the first. This is the case in many classification problems and similar inferences. The focus in this kind of problems is to infer x in an observation with a new value of t , given observations of the pair of quantities for different values of t .

If we assume for the moment that t and x assume finite sets of values, then also the set of functions between them is finite. Regression in the restricted sense above is then equivalent to inferring a function $f: t \mapsto x$:

$$p(x|t, I) = \int df \, p(x|t, f, I) p(f|I), \quad (1)$$

with

$$p(x|t, f, I) = \delta[x - f(t)]. \quad (2)$$

Thanks

... to Mari & Miri for continuous encouragement and affection, and to Buster Keaton and Saitama for filling life with awe and inspiration. To the developers and maintainers of L^AT_EX, Emacs, AUCT_EX, Open Science Framework, Python, Inkscape, Sci-Hub for making a free and unfiltered scientific exchange possible. 