

# Open questions in statistical mechanics [draft]

P.G.L. Porta Mana 

Western Norway University of Applied Sciences [<pgl@portamana.org>](mailto:pgl@portamana.org)

15 May 2021; updated 14 May 2021

Some unanswered questions in the foundations of statistical mechanics

## 1 Questions

### 1.1 Meaning and use of “average” in the canonical ensemble. Implications for the theory of fluctuations

In speaking of the canonical ensemble we say that the energy value  $U$  is only an “average”. Do we mean an average over time while we observe such a system? Or an average over identical preparations of such a system?

The difference is important because in the second case the system, after preparation, may have an energy  $U^*$ , with  $U^* \approx U$ , that is perfectly constant in time. It also means that such a system may be perfectly closed: we can use the canonical ensemble also for a perfectly closed system, but created with a protocol that does not always lead to the same energy.

From the second point of view the fluctuations of the system are not fluctuations in time while we observe it: they are differences among identical preparations of the system.

This also mean that the fluctuations that we observe in time in a system described by the canonical ensemble may be different from the fluctuations obtained across preparations, and may be absent altogether.

The Gaussian ensemble<sup>1</sup> allows for the exact specification of fluctuations across identical preparations.

---

<sup>1</sup> [johaletal2003](#); [hetherington1987](#).