

Enumerative Combinatorics

Lecture Notes

David Scholz

January 29, 2022

Abstract

Abstract

Contents

1	Introduction	1
1.1	Words	1
1.2	Permutations	2
1.3	Merry-go-rounds and Fermat's little theorem	3
1.4	Binomial coefficients	5
1.5	The Pascal triangle	6
1.6	Exercises	8
2	Permutations, binomial coefficients	9
2.1	The cab driver problem	9
2.2	Balls in boxes and multisets	10
2.3	Integer compositions	10
2.4	Principle of inclusion and exclusion	10
2.5	The derangement problem	10
2.6	Exercises	10
3	Linear recurrences - The Fibonacci sequence	12
3.1	Fibonacci's rabbit problem	12
3.2	Fibonacci numbers and the Pascal triangle	12
3.3	Domino tilings	12
3.4	Linear recurrence relations	12
3.5	The characteristic equation	12
3.6	Linear recurrence relations of order 2	12
3.7	The Binet formula	12
3.8	Linear recurrence relations of arbitrary order	12
3.9	The case of roots with multiplicities	12
3.10	Exercises	12
4	A nonlinear recurrence: many faces of Catalan numbers	13
4.1	Triangulations of a polygon	13
4.2	Recurrence relation for triangulations	13
4.3	The cashier problem	13
4.4	Dyck paths	13
4.5	Recurrence relations for Dyck paths	13
4.6	Reflection trick and a formula for Catalan numbers	13
4.7	Binary trees	13
4.8	Exercises	13

5	Generating functions: a unified approach to combinatorial problems - Solving linear recurrences	14
5.1	Generating functions: first examples	14
5.2	Formal power series	14
5.3	When are formal power series invertible?	14
5.4	Derivation of formal power series	14
5.5	Binomial theorem for negative integer exponents	14
5.6	Solving the Fibonacci recurrence relation	14
5.7	Generating functions of linear recurrence relations are rational	14
5.8	Solving linear recurrence relations: general case	14
5.9	Exercises	14
6	Generating function of the Catalan sequence	15
6.1	Composition of formal power series	15
6.2	Derivation and integration of formal power series	15
6.3	Chain rule - Inverse function theorem	15
6.4	Logarithm - Logarithmic derivative	15
6.5	Binomial theorem for arbitrary exponents	15
6.6	Generating function for Catalan numbers	15
6.7	Exercises	15
7	Partitions - Euler's generating function for partitions and pentagonal formula	16
7.1	Definition and first examples	16
7.2	Young diagrams	16
7.3	Generating function for partitions	16
7.4	Partition with odd and distinct summands	16
7.5	Sylvester's bijection	16
7.6	Euler's pentagonal theorem	16
7.7	Proof of Euler's pentagonal theorem	16
7.8	Computing the number of partitions via the pentagonal theorem	16
7.9	Exercises	16
8	Gaussian binomial coefficients	17
8.1	Generating function for partitions inside a rectangle	17
8.2	q -binomial coefficients: definition and first properties	17
8.3	Recurrence relation for q -binomial coefficients	17
8.4	Explicit formula for q -binomial coefficients	17
8.5	Euler's partition function	17
8.6	q -binomial coefficients in linear algebra	17

8.7	q -binomial theorem	17
8.8	Exercises	17

List of Figures

1	Initial train colouring.	3
2	Marry-go-round initial case.	3
3	Marry-go-round with only one colour.	4
4	The Cap Driver Problem.	9
5	Balls in boxes and mutlisets.	10

1 Introduction

We start by discussing fundamental combinatorial constructions such as words of a fixed alphabet, permutations of finite sets and the number of subsets with a given number of elements. While we give some examples of how some combinatorial problems naturally arise, our main goal in this first section is to provide an intuitive introduction to the topic of Enumerative Combinatorics.

1.1 Words

The first definition of this lecture is the definition of a word over a finite alphabet of symbols.

Definition 1.1 (word). Let A be a finite set, called *alphabet*. A *word* is a finite sequence of elements of A . The *size* of A is its cardinality $|A|$. The *size* or *length* of a word is its number of elements from A .

We might abuse language to a certain extent and call the elements of A *letters*.

Example 1.2. Let $A = \{1, 2, 3, 4\}$, then 112, 132, 1234, 111, 2345, and so on, are all words over A . Notice that the length of the words is not fixed.

In the study of combinatorics a very classical question is as follows. Given an alphabet A of length $|A| = n$, how many words over A of length k exist? This question leads us to the following theorem.

Theorem 1.1. The number of words of length k in an alphabet consisting of n letters is equal to n^k .

Proof. There are n possibilities for the first letter. The same holds for all other letters. Thus, the total number of words equals

$$\underbrace{n \cdot n \cdot n \cdots n}_{k\text{-times}} = n^k$$

□

The above proof is fairly simple, but there is a different possible interpretation of this problem. Let X and Y be sets. We might ask: what is the total number of maps of the form $X \rightarrow Y$ without restrictions (an element in Y can be hit multiple times)? It appears that this question can be solved using Theorem 1.1. For seeing this, we number the elements of X and Y using natural numbers. Given a map $f : X \rightarrow Y$, the sequence $f(1), f(2), \dots, f(k)$ forms a word of length k . The number of words is therefore equivalent to the number of such maps.

1.2 Permutations

A notion that is closely related to the definition of a word, is the definition of a *permutation*. First of all, the basic idea is to count maps, but usually with certain properties. Let X and Y be sets with $|X| = k$ and $|Y| = n$. We might ask: how many bijections $f : X \rightarrow Y$ are there? Recall that a map is bijective if it is surjective and injective. In other words, given two elements $x, y \in X$ such that $f(x) = f(y)$, then it is implied that $x = y$ (the injective property), and every element of Y appears in the preimage of a certain element of X (the surjective property). It follows that the cardinalities of X and Y must be equal. These observations lead us to the definition of a permutation.

Definition 1.3 (permutation). A *permutation* is a bijection from $\{1, 2, 3, \dots, n\}$ to itself.

In that sense the number of permutations is the number of bijections from $\{1, 2, \dots, n\}$ to itself. Equivalently, a permutation might be interpreted as a total linear ordering of the set $\{1, 2, 3, \dots, n\}$. If so, we need to arrange them in a certain way. How many ways to do this are there? The following theorem provides one possible solution.

Theorem 1.2. The number of permutations of $\{1, 2, \dots, n\}$ is equal to $n! = 1 \cdot 2 \cdot 3 \cdots n$.¹

Proof. There are n choices for the first element. For the second element there are $n - 1$ choices since one element is already occupied. For the k -th element there are $n - k + 1$ choices. For the last element there is one choice. Thus, the total number of permutations is given by $n(n - 1)(n - 2) \cdots 2 \cdot 1 = n!$. This completes the proof. \square

In the following, we will generalize the above result to injective maps. Let $k \leq n$. In how many ways can we produce a collection of k elements out of n ? Such a collection is called a k -permutation.

Definition 1.4 (k -permutation). A k -permutation is a total linear ordering on a k -element subset of $\{1, 2, \dots, n\}$.

Example 1.5. Let $\{1, 2, 3\}$. A 1-permutation is just a 1-element subset, thus there is nothing order. The set of 2-permutations is given by $\{\{1, 2\}, \{1, 3\}, \{2, 1\}, \{2, 3\}, \{3, 1\}, \{3, 2\}\}$. Thus, there are six 2-permutations in total.

The general case of the number of k -permutations is examined in the following theorem.

Theorem 1.3. The number of k -permutations of $\{1, 2, \dots, n\}$ is equal to

$$n(n - 1) \cdots (n - k + 1).$$

¹Recall that $0! = 1$.

Proof. There are n possible choices for the first entry of the subset. There are $(n - 1)$ possible choices for the second element because the second element can occupy any entry except the one of the first element. For the k -th entry there are $n - k + 1$ possible choices. All these possibilities are independent, hence the total number of k -permutations is equal to their product $n(n - 1) \cdots (n - k + 1)$. \square

The number $n(n - 1) \cdots (n - k + 1)$ is sometimes called the k decreasing power and is denoted by $n^{\downarrow k}$.

1.3 Merry-go-rounds and Fermat's little theorem

The problem which we are going to discuss next will illustrate a connection of combinatorics to algebra and possibly number theory.

Consider a train Q consisting of p carriages. Suppose that we have n colours. In how many ways can we paint Q , if we can paint each carriage with one of the n colours? Consider figure 1 as an example. There is a train with 5 carriages which are coloured in a certain way using 3 different colours.




Figure 1: Initial train colouring.

Now, this is exactly the word problem, thus the solution is given by n^p .

Question: Suppose that instead of a train we have a merry-go-round and suppose further, for some reason, that p is prime. In how many ways can we paint each carriage of a merry-go-round using n colours?




Figure 2: Merry-go-round initial case.

In figure 2 a merry-go-round with 5 carriages is visualized. They are again coloured using 3 colours.

What is the difference to the problem we have considered before? In this case we can rotate the carriages without changing its colours, but these colourings are considered to be identical.

Idea: We cut the merry-go-round (consider the two lines in figure 2 as the "cutting point") and make a train out of it and there are exactly p places on which we could make a cut.

The first naive guess would be that the solution to our problem is equal to $\frac{n^p}{p}$, since we can get p trains from each merry-go-round. This is wrong for an obvious reason. The number $\frac{n^p}{p}$ is by no means always integer. So let us try to understand what is wrong with this reasoning. First, if we cut at two different places in our merry-go-round, then we could get the same train. An easy way for seeing this is to consider a merry-go-round in which each carriage is coloured using the same colour. There are still p ways to make a train out of it but each of them is identical. Compare this reasoning with figure 3.




Figure 3: Merry-go-round with only one colour.

To say it in different words, if we rotate a merry-go-round by an arbitrary number of times, it will eventually coincide with itself, thus if a merry-go-round goes to itself after some nontrivial rotation, then all carriages have the same colour. Now, notice that this is not the case if not all carriages have the same colour, since then we get exactly p different trains, but observe that this is only true provided that p is prime. In order to see this, consider a merry-go-round with 4 carriages, and assume that we have two colours. black and white. Then, there are only two different trains which we can get from this merry-go-round. Colour the merry-go-round as follows: black - white - black - white. Then after two rotations, we get the same train. Thus, instead of four trains we get two. However, if p is prime, then our claim is indeed true.

If our merry-go-round has carriages of different colours, then we can compute the number of such merry-go-rounds. It is the number of trains divided by p . What is the number of trains which have different colors? This is n to the power of p , which is the total number of trains, minus the number of trains of the same color, which is just n and for each color, there is only one train for which all carriages have this given colour. Hence, our final result is given by the following equation.

$$\frac{n^p - n}{p} + p$$

It is again surprising that the above number is always an integer. This is not obvious at all and proving this using number theory is quite more involved. In general, we have obtained a very nice result given by the following theorem.

Theorem 1.4. Given $n \in \mathbb{N}$ and a prime p , then $n^p - n \mid p$ ($n^p - n$ is divisible by p).

The above is Fermat's little Theorem and we have proven it in a completely combinatorial way.

1.4 Binomial coefficients

Next, we consider the following important definition.

Definition 1.6 (binomial coefficient). Let $k \leq n$. The number of k -element subsets of $\{1, 2, \dots, n\}$ is called a *binomial coefficient* $\binom{n}{k}$ (" n choose k ").

How do we interpret this number? Suppose that we are hiring people to certain positions and suppose that we have n candidates and we have k identical positions. The number of selecting k people out of n is called $\binom{n}{k}$. Notice that this is different than the problem from problem about k -permutations. In our current case, the order of the k elements that we choose does not matter. Thus, choosing all k element subsets of a set and then ordering them in all possible ways is equal to the k -permutations that we have seen above. How do we count the number of k -element subsets of a set?

Theorem 1.5. The number of k -element subsets of a set with cardinality n is equal to

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1) \cdots (n-k+1)}{k!}.$$

Proof. We consider ordered collections of distinct elements of length k . There are n ways of choosing the first element, and $n - k + 1$ ways of choosing the k -th element. Hence there are

$$n(n-1) \cdots (n-k+1) = \frac{n!}{(n-k)!}$$

ways such ordered collections. Since we are concerned with the number of unordered subsets and each subset has cardinality k , each subset can be rearranged in $k!$ ways. Therefore, the number of ways to choose a k -element subset of a set with cardinality n is given by

$$\frac{\frac{n!}{(n-k)!}}{k!} = \frac{n!}{k!(n-k)!}.$$

This completes the proof. \square

With the above theorem, we have shown something additionally that is not obvious at all and not necessarily easy to prove either. We have also shown that the number $\frac{n(n-1)\cdots(n-k+1)}{k!}$ is integer.

At this point we provide some properties of binomial coefficients. We leave the proof as an easy exercise to the reader.

- (i) $\binom{n}{0} = \binom{n}{n} = 1.$
- (ii) $\binom{n}{1} = \binom{n}{n-1} = n$
- (iii) $\binom{n}{1} = \binom{n}{n-k}.$
- (iv) $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$

Let us examine property (iv) in more detail. Despite its obvious algebraic proof, there is an intuitive combinatorial version as well. We provide a proof sketch.

Theorem 1.6. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Proof-sketch. Given a set $A = \{1, \dots, n\}$ with n elements. The number of k -element subsets of A is $\binom{n}{k}$ according to theorem 1.5. Suppose that we pick the element 1 of A , then all k -element subsets of A are divided into two parts, some of them contain the element 1, and the others do not. Since we have already picked one element, we need to pick $k-1$ elements out of the remaining $n-1$ elements. There are $\binom{n-1}{k-1}$ subsets containing 1. Thus, the subsets of cardinality k that do not contain 1 is equal to $\binom{n-1}{k}$, because we want to pick a k -element subset out of $n-1$ elements since 1 element is forbidden. Adding up the two parts are equal to $\binom{n}{k}$. This completes the proof. \square

1.5 The Pascal triangle

Binomial coefficients can be arranged in a triangle which is called the Pascal triangle. Its rows are indexed by integers from 0 up to infinity and in each row we write down all binomial coefficients. For example, for $\binom{n}{k}$, $n = 4$ represents the fifth row of the Pascal triangle

and since $k \leq n$ there are 5 binomial coefficients, namely $\binom{4}{0} = 1$, $\binom{4}{1} = 4$, $\binom{4}{2} = 6$, $\binom{4}{3} = 4$ and $\binom{4}{4} = 1$.

The relation given In Theorem 1.6 provides a way of constructing the Pascal triangle. It says that the sum of two neighbors, namely $\binom{n-1}{k-1}$ and $\binom{n-1}{k}$ is equal to $\binom{n}{k}$. Thus, taking two neighbors in one row and adding them up gives as the element below them (e.g. $3+3$ in row $n=3$ provides the element 6 in the row $n=4$).

$$\begin{array}{cccccccc}
n = 0 & & & & & & & 1 \\
n = 1 & & & & 1 & & 1 & \\
n = 2 & & & 1 & & 2 & & 1 \\
n = 3 & & 1 & & 3 & & 3 & 1 \\
n = 4 & & 1 & 4 & & 6 & 4 & 1 \\
n = 5 & 1 & 5 & 10 & & 10 & 5 & 1 \\
n = 6 & 1 & 6 & 15 & 20 & 15 & 6 & 1
\end{array}$$

Proposition 1.1. $\binom{n}{0} + \binom{n}{k} + \cdots + \binom{n}{n} = 2^n$.

Proof. The binomial coefficient $\binom{n}{k}$ is the number of k -element subsets of a n -element set $A = \{1, \dots, n\}$. Hence,

$$\sum_{k=0}^n \binom{n}{k}$$

is equal to the number of all possible subsets of A . We can associate a subset of A with a binary number of length n whereby each digit in this binary number represents a specific element in A . Now, setting a digit to 1 means that we pick the element from A at this position for our subset. Setting it to 0 means that we do not pick it. This is equal to the number of maps $\{1, \dots, n\} \rightarrow \{0, 1\}$ and the total number of such maps is equal to 2^n or in other words, the number of binary numbers that can be represented with n digits is equal to 2^n . This completes the proof. \square

We call the above binomial coefficients, because they arise as coefficients in the binomial formulas $(a + b)^n$. Let us prove this.

Theorem 1.7. (Newton's Binomial Theorem)

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} \cdot b^k = a^n + n \cdot a^{n-1} \cdot b + \frac{n(n-1)}{2} a^{n-2} \cdot b^2 + \dots + n \cdot a \cdot b^{n-1} + b^n$$

Proof. We expand the left-hand-side.

$$(a+b)^n = \underbrace{(a+b)(a+b)(a+b)\cdots(a+b)}_{n\text{-times}}$$

Every term in the above sum is the result of choosing either a or b from each of the n factors. We want to calculate the coefficient for $a^{n-k}b^k$. Since b^k , we must choose b from exactly k of the factors and we choose a from the remaining $n - k$ factors. Thus, the coefficients for $a^{n-k}b^k$ is equal to $\binom{n}{k}$. This completes the proof. \square

Corollary 1.1. $\binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \binom{n}{3} + \cdots + (-1)^n \binom{n}{n} = 0$

Proof. Consider the binomial formula as provided in Theorem 1.7. Now, let $a = 1$ and b (or vice versa), it follows that

$$(1 - 1)^n = \sum (-1)^k \binom{n}{k}.$$

\square

Notice, that Newton's binomial Theorem provides an easy proof for proposition 1.1 as well by considering $(1 + 1)^n = \sum \binom{n}{k}$.

1.6 Exercises

1. Find the number of subsets of a 10-element set with an odd number of elements.
2. Compute the sum $\sum_{k=0}^n 2^k \binom{n}{k}$.
3. Find the number of ways to rearrange the letters of the word *EULER*.
4. Find the number of ways to rearrange the letters of the word *CATALAN*.
5. How many necklaces can be formed with 7 distinct beads? Necklaces can be rotated and/or flipped over. All beads must be used.

2 Permutations and binomial coefficients continued

The following lecture discusses some more problems in which binomial coefficients arise and will lead us to the famous principle of exclusion and inclusion.

2.1 The cab driver problem

We will begin with the so called 'cab driver problem'. Suppose that we have a city with streets arranged in a regular grid (like Manhattan) and suppose that our city contains n blocks from north to south and m blocks from west to east. Now, suppose that we have a cab driver that wants to drive from one point on the grid to another point, say from A to B . In figure 4 the general setup is visualized.




Figure 4: The Cap Driver Problem.

Question: In how many ways can we get from A to B ? We are only allowed to go east or north and we are not allowed to go backwards.

Let $P_{m,n}$ be the number of paths from A to B . We notice first, that we can solve this problem by considering a sequence of letters. Let us denote "going north" by the letter "N" and "going east" by the letter "E", then a path from A to B corresponds to a sequence of N's and E's of length $m + n$ with n N's and m E's. The total number of such words is equal to the total number of paths from A to B .

There is also another solution. Starting from any point in the above grid, there are only two choices: go north or go east. Let A be the starting point and denote "going north in the first step" by A' and "going east in the second step" by A'' , then the total number of paths from A to B is equal to the total number of paths from A' to B plus the total number of paths from A'' to B . The result is a so called recurrence relation given by the following equation.

$$P_{A \rightarrow B} = P_{A' \rightarrow B} + P_{A'' \rightarrow B}$$

In general, if we go one step north the first time, then there are $P_{m,n-1}$ paths left from this position to B . Similarly, if we go east first, then there are $P_{m-1,n}$ paths left from this position to B . This results in the following relation.

$$P_{m,n} = P_{m,n-1} + P_{m-1,n}$$

Given the initial conditions $P_{0,n} = P_{m,0} = 1$, and $P_{m,1} = m + 1$ and $P_{1,n} = n + 1$, our final solution can be found quickly.

$$P_{m,n} = \binom{m+n}{m} = \binom{m+n}{n}$$

2.2 Balls in boxes and multisets

Suppose that we have k balls and n boxes. In how many ways can we put k balls into n boxes? If the maximal capacity of each box is 1, the answer would be $\binom{n}{k}$ if $n \geq k$ and 0 otherwise. What happens if there are no restrictions on the capacity of boxes?

This question leads us to the definition of a multiset. Let X be a set.

Definition 2.1. A multiset is a function $\mu : X \rightarrow \mathbb{Z}_{\geq 0}$. The size of a multiset is $\sum_{x \in X} \mu(x)$.

q

Example 2.2. Assume that we have 5 boxes, that is $X = \{1, 2, 3, 4, 5\}$, and $\mu(1) = 1, \mu(2) = 0, \mu(3) = 3$ and $\mu(4) = 1$. Compare this with figure 5. Notice that box 3 contains 3 balls. The size of μ is 5.




Figure 5: Balls in boxes and multisets.

2.3 Integer compositions

2.4 Principle of inclusion and exclusion

2.5 The derangement problem

2.6 Exercises

1. Find the number of lattice paths from $(-2, -2)$ to $(3, 3)$ passing through the point $(0, 0)$ (each segment of a path goes north or east).

2. In how many ways is it possible to permute the letters in the word *EUCLID* in such a way that the order of the vowels (*E, U, I*) is unchanged?
3. Find the number of merry-go-rounds formed by 10 carriages of two different colours.
4. Find the number of positive integers not exceeding 300 and **not** divisible by any of 2, 3, and 5.

3 Linear recurrences - The Fibonacci sequence

Hallo

- 3.1 Fibonacci's rabbit problem
- 3.2 Fibonacci numbers and the Pascal triangle
- 3.3 Domino tilings
- 3.4 Linear recurrence relations
- 3.5 The characteristic equation
- 3.6 Linear recurrence relations of order 2
- 3.7 The Binet formula
- 3.8 Linear recurrence relations of arbitrary order
- 3.9 The case of roots with multiplicities
- 3.10 Exercises

4 A nonlinear recurrence: many faces of Catalan numbers

Hallo

4.1 Triangulations of a polygon

4.2 Recurrence relation for triangulations

4.3 The cashier problem

4.4 Dyck paths

4.5 Recurrence relations for Dyck paths

4.6 Reflection trick and a formula for Catalan numbers

4.7 Binary trees

4.8 Exercises

5 Generating functions: a unified approach to combinatorial problems - Solving linear recurrences

Hallo

- 5.1 Generating functions: first examples
- 5.2 Formal power series
- 5.3 When are formal power series invertible?
- 5.4 Derivation of formal power series
- 5.5 Binomial theorem for negative integer exponents
- 5.6 Solving the Fibonacci recurrence relation
- 5.7 Generating functions of linear recurrence relations are rational
- 5.8 Solving linear recurrence relations: general case
- 5.9 Exercises

6 Generating function of the Catalan sequence

Hallo

6.1 Composition of formal power series

6.2 Derivation and integration of formal power series

6.3 Chain rule - Inverse function theorem

6.4 Logarithm - Logarithmic derivative

6.5 Binomial theorem for arbitrary exponents

6.6 Generating function for Catalan numbers

6.7 Exercises

7 Partitions - Euler's generating function for partitions and pentagonal formula

Hallo

7.1 Definition and first examples

7.2 Young diagrams

7.3 Generating function for partitions

7.4 Partition with odd and distinct summands

7.5 Sylvester's bijection

7.6 Euler's pentagonal theorem

7.7 Proof of Euler's pentagonal theorem

7.8 Computing the number of partitions via the pentagonal theorem

7.9 Exercises

8 Gaussian binomial coefficients

Hallo

- 8.1 Generating function for partitions inside a rectangle
- 8.2 q -binomial coefficients: definition and first properties
- 8.3 Recurrence relation for q -binomial coefficients
- 8.4 Explicit formula for q -binomial coefficients
- 8.5 Euler's partition function
- 8.6 q -binomial coefficients in linear algebra
- 8.7 q -binomial theorem
- 8.8 Exercises