

1 Taylor Series

1.1 Important Maclaurin Series

Trigonometric Functions

$$\sin(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

Hyperbolic Functions

$$\sinh(x) = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!} = x + \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cosh(x) = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

Exponential Function

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$$

Natural Logarithm (for $|x| < 1$)

$$\log(1-x) = -\sum_{n=1}^{\infty} \frac{x^n}{n} = -x - \frac{x^2}{2} - \frac{x^3}{3} - \dots$$

$$\log(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Geometric Series (for $|x| < 1$)

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots$$

Binomial Series (for $|x| < 1, \alpha \in \mathbb{C}$)

$$(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n, \quad \binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$$

This includes the square root series for $\alpha = \frac{1}{2}$ and the infinite geometric series for $\alpha = -1$.

$$(1+x)^{\frac{1}{2}} = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \dots$$

2 Linear Algebra Fundamentals

2.1 Matrix Adjoints

A is Hermitian if $A^* = A$ and symmetric if $A^T = A$

- Eigenvectors of A for distinct eigenvalues are orthogonal
- The algebraic and geometric multiplicities are the same.
- A has a spectral decomposition

$$A = U\Lambda U^{-1} = U\Lambda U^*$$

2.2 Norms

2.2.1 Vector Norms $\|\cdot\|_p$

$$\|x\|_p = \left(\sum |x_i|^p \right)^{\frac{1}{p}}$$

Sum Norm, $p = 1$

$$\|x\|_{sum} = \|x\|_1 = \sum |x_i|$$

Euclidean Norm, $p = 2$

$$\|x\|_{Euclidean} = \|x\|_2 = \sqrt{\sum |x_i|^2}$$

Maximum Norm, $p = \infty$

$$\|x\|_{max} = \|x\|_{\infty} = \max \{|x_i|\}$$

All finite dimensional norms behave similarly with respect to convergence.

Examples: $\|x\|_2 \leq \|x\|_1$ and $\|x\|_1 \leq \sqrt{n}\|x\|_2$.

2.2.2 Matrix Norms

Matrix norms have the property $\|AB\| \leq \|A\| \|B\|$.

The first variety is the Induced Norms

$$\|A\|_p = \sup_{x \neq 0, x \in \mathbb{C}^n} \frac{\|Ax\|_p}{\|x\|_p} = \sup_{\|x\|_p=1, x \in \mathbb{C}^n} \|Ax\|_p$$

- $p = 1$ leads to the maximum absolute column sum

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^m |a_{ij}|$$

- $p = 2$ leads to the largest singular value of A , ie the square root of the largest eigenvalue of the positive semi-definite matrix A^*A

$$\|A\|_2 = \sqrt{\lambda_{\max}(A^*A)} = \sigma_{\max}(A)$$

- $p = \infty$ leads to the maximum absolute row sum

$$\|A\|_{\infty} = \max_{1 \leq i \leq m} \sum_{j=1}^n |a_{ij}|$$

Fun facts

- For any induced norm, $\|A^r\|_p \geq \rho(A)^r$, where equality holds for $p = 2$.

2.2.3 Schatten Norms

The Schatten norms arise when applying the p -norm to the vector of singular values of a matrix. Let $\{\sigma_i\}$ be the singular values of A .

$$\|A\|_p = \left(\sum_{i=1}^{\min(m,n)} \sigma_i^p \right)^{\frac{1}{p}}$$

Schatten norms are unitarily invariant. That is, for any unitary matrices U, V ,

$$\|A\|_p = \|UAV\|_p$$

- For $p = 1$, we have the **Nuclear norm** also known as the **Trace norm**

$$\|A\|_* = \sum_{i=1}^{\min(m,n)} \sigma_i$$

- $p = 2$. The Frobenius norm is the finite dimensional analog of the Hilbert Schmidt norm

$$\|A\|_F = \|A\|_2 = \left(\sum_{j=1}^n \sum_{i=1}^m |a_{ij}|^2 \right)^{\frac{1}{2}}$$

with the interesting property

$$\|A\|_F = \sqrt{\text{tr}(A^*A)} = \sqrt{\sum_{i=1}^{\min(m,n)} \sigma_i^2}$$

- For $p = \infty$, we have the **spectral norm**

$$\|A\|_* = \max_{1 \leq i \leq \min(m,n)} \sigma_i$$

2.2.4 Matrix Determinants

Useful properties of matrix determinants are

- $\det(AB) = \det(A) \det(B)$
- $\det(A^{-1}) = \det(A)^{-1}$

2.2.5 Condition Numbers

The condition number κ measures the sensitivity of a function with respect to an argument. That it is the maximum relative change.

In general, this is

$$\kappa(f) = \frac{(f(x + \Delta x) - f(x))}{f(x)} \frac{x}{\Delta x} = \frac{(f(x + \Delta x) - f(x))}{\Delta x} \frac{x}{f(x)}$$

For a differentiable function, this is

$$\kappa(f) = \frac{x f'(x)}{f(x)}$$

For $Ax = b$,

$$\kappa(A) = \frac{\|A^{-1} \delta b\|}{\|\delta b\|} \frac{\|b\|}{\|A^{-1} b\|}$$

2.3 Positive Definite Matrices

Which for any consistent norm (ie, norms with $\kappa(A) \geq 1$) becomes

$$\kappa(A) = \|A^{-1}\| \|A\|$$

- If $\|\cdot\| = \|\cdot\|_2$, then

$$\kappa(A) = \frac{\sigma_{max}(A)}{\sigma_{min}(A)}$$

If A is normal, then

$$\kappa(A) = \frac{\lambda_{max}(A)}{\lambda_{min}(A)}$$

If A is unitary, then

$$\kappa(A) = 1$$

In general, for a differentiable function $f(x)$,

$$\kappa(f) = \frac{\|J(x)\|}{\|f(x)\|} \|x\|$$

2.3 Positive Definite Matrices

For a matrix A , the following are equivalent

- All of its eigenvalues are positive
- It may form an inner product such as $\langle \vec{x}, A\vec{y} \rangle$
- It is the Gram matrix of linearly independent vectors
- It's leading principle minors are all positive
- It has a unique Cholesky decomposition

2.4 Matrix Calculus

If f is a scalar function of many variables, then

$$\partial f = \frac{\partial f}{\partial \vec{x}} = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \frac{\partial f}{\partial x_3} \right)$$

One could also take the directional derivative in the direction \vec{u}

$$\partial_{\vec{u}} f = \partial f \cdot \vec{u}$$

The derivative of a vector function is written as

$$\frac{\partial \vec{y}}{\partial \vec{x}} = (J(\vec{y}))(\vec{x}) \quad J_{ij} = \frac{\partial y_i}{\partial x_j}$$

- $\frac{\partial \vec{x}}{\partial \vec{x}} = I$
- $\frac{\partial A\vec{x}}{\partial \vec{x}} = A$
- $\frac{\partial \vec{x}^T A}{\partial \vec{x}} = A^T$
- $\frac{\partial \vec{u}^T v}{\partial \vec{x}} = \frac{\partial \vec{u}}{\partial \vec{x}} \vec{v} + \frac{\partial \vec{v}}{\partial \vec{x}} \vec{u}$

- $\frac{\partial \vec{x}^T x}{\partial \vec{x}} = 2\vec{x}$
- $\frac{\partial \vec{u}^T A \vec{v}}{\partial \vec{x}} = \frac{\partial \vec{u}}{\partial \vec{x}} A \vec{v} + \frac{\partial \vec{v}}{\partial \vec{x}} A^T \vec{u}$
- $\frac{\partial \vec{x}^T A \vec{x}}{\partial \vec{x}} = A \vec{x} + A^T \vec{x}$
- $\frac{\partial^2 \vec{x}^T A \vec{x}}{(\partial \vec{x})^2} = A + A^T$

3 Common Theorems

3.1 Reminders

- The mean value theorem for a differentiable $f(x)$ is

$$f(x) - f(y) = f'(\theta x + (1 - \theta)y)(x - y)$$

for some $\theta \in [0, 1]$.

- For polynomial division, dividing a polynomial of degree n $p(x)$ by the monomial $(x - a)$ leaves you with a polynomial of degree $n - 1$ and

$$p(x) = (x - a)q(x) + r(x)$$

That is, $q(x), r(x) \in \mathbb{P}_{n-1}$. $q(x)$ and $r(x)$ are unique. Note that $p(a) = r(a)$.

3.2 Fredholm Alternative , Linear Algebra Version

For $A \in \mathbb{C}^{n \times m}$ and $b \in \mathbb{C}^{m \times 1}$,

- Either $A\vec{x} = \vec{b}$ has a solution \vec{x}
- OR: $A^T \vec{y} = 0$ has a solution \vec{y} with $\vec{y}^T \vec{b} \neq 0$.

That is, $A\vec{x} = \vec{b}$ has a solution if and only if for any \vec{y} s.t. $A^T \vec{y} = 0$, $\vec{y}^T \vec{b} = 0$.

4 Polynomials

4.1 Special Polynomials

4.1.1 Legendre Polynomials

Defined by

$$\frac{d}{dx} \left((1 - x^2) \frac{d}{dx} P_n(x) \right) + n(n + 1) P_n(x) = 0$$

Also by Rodrigues' formula

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{(dx)^n} ((x^2 - 1)^n)$$

or by the recurrence relation

$$(n + 1)P_{n+1}(x) = (2n + 1)xP_n(x) - nP_{n-1}(x)$$

They are orthogonal in the sense that

$$\int_{-1}^1 P_m(x)P_n(x)dx = \frac{2}{2n+1}\delta_{mn}$$

- The first few are

$$P_0 = 1$$

$$P_1 = x$$

$$P_2 = \frac{1}{2}(3x^2 - 1)$$

$$P_3 = \frac{1}{2}(5x^3 - 3x)$$

- These arise when you carry out the GS process on the canonical polynomial basis $1, x, x^2, \dots$
- They are antisymmetric

$$P_n(-x) = (-1)^n P_n(x)$$

4.1.2 Chebyshev Polynomials

The Chebyshev polynomials of the first kind are defined as solutions to

$$(1-x^2)y'' - xy' + n^2y = 0$$

$$(1-x^2)y'' - 3xy' + n(n+2)y = 0$$

by the recurrence relation

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x) \quad T_0(x) = 1, T_1(x) = x$$

These also have the important trigonometric definition

$$T_n(x) = \cos(n \arccos(x)) = \cosh(n \operatorname{arccosh}(x))$$

In other words, these satisfy

$$T_n(\cos(\theta)) = \cos(n\theta)$$

They are orthogonal in the sense

$$\int_{-1}^1 \frac{T_n(x)T_m(x)}{\sqrt{1-x^2}}dx = \begin{cases} 0 & n \neq m \\ \pi & n = m = 0 \\ \frac{\pi}{2} & n = m > 0 \end{cases}$$

The Chebyshev polynomials of the second kind are defined by

$$U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x) \quad U_0(x) = 1, U_1(x) = 2x$$

These satisfy

$$U_n(\cos(\theta)) = \frac{\sin((n+1)\theta)}{\sin(\theta)}$$

They are orthogonal in the sense

$$\int_{-1}^1 U_n(x)U_m(x)\sqrt{1-x^2}dx = \begin{cases} 0 & n \neq m \\ \frac{\pi}{2} & n = m \end{cases}$$

4.1.3 Hermite Polynomials

The physicists' Hermite Polynomials are defined by

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2} = \left(2x - \frac{d}{dx}\right)^n 1$$

With the recursion relations

$$H_{n+1}(x) = 2xH_n(x) - H'_n(x)$$

and

$$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x)$$

These are orthogonal in the sense that

$$\int_{-\infty}^{\infty} H_m(x) H_n(x) e^{-x^2} dx = \sqrt{\pi} 2^n n! \delta_{nm}$$

4.1.4 Bernstein Polynomials

Let the Bernstein basis polynomials be defined by

$$b_{\nu,n}(x) = \binom{n}{\nu} x^{\nu} (1-x)^{n-\nu} \quad \nu = 0, \dots, n$$

with coefficients β_{ν} . So the Bernstein polynomial of degree n is defined as

$$B_n(x) = \sum_{\nu=0}^n \beta_{\nu} b_{\nu,n}(x)$$

Now consider

$$B_n(f)(x) = \sum_{\nu=0}^n f\left(\frac{\nu}{n}\right) b_{\nu,n}(x)$$

It can be shown that

$$\lim_{n \rightarrow \infty} B_n(f)(x) = f(x)$$

and the convergence is uniform on $[0, 1]$.

- These are used in the proof of the Stone-Weierstrass approximation theorem.

4.1.5 Lagrange Polynomials

Given a distinct set $\{x_j\}$, the Lagrange basis polynomials are defined by

$$\ell_j(x) = \prod_{0 \leq m \leq k, m \neq j} \frac{x - x_m}{x_j - x_m} \quad 0 \leq j \leq k$$

We see that

$$\ell_j(x_m) = \delta_{jm}$$

- These interpolate a function $f(x)$ exactly when

$$L(x) = \sum_{j=0}^n \prod_{0 \leq m \leq k, m \neq j} \frac{x - x_m}{x_j - x_m} f(x_j)$$

4.2 Polynomial Interpolation

Interpolation refers to approximations that are exact for certain parts of the data.

4.2.1 Using Lagrange Polynomials

In general, for n data points, polynomial interpolation has the form

$$p_n(x) = \sum_{i=0}^n f(x_i) \left(\prod_{0 \leq j \leq n, j \neq i} \frac{x - x_j}{x_i - x_j} \right)$$

which comes from solving the Vandermonde matrix problem

$$\begin{pmatrix} x_0^n & \dots & x_0^0 \\ \vdots & \ddots & \vdots \\ x_n^n & \dots & x_n^0 \end{pmatrix} \begin{pmatrix} a_n \\ \vdots \\ a_0 \end{pmatrix} = \begin{pmatrix} x_0^n & \dots & 1 \\ \vdots & \ddots & \vdots \\ x_n^n & \dots & 1 \end{pmatrix} \begin{pmatrix} a_n \\ \vdots \\ a_0 \end{pmatrix} = \begin{pmatrix} f_0 \\ \vdots \\ f_n \end{pmatrix}$$

These interpolants are exact at the data points.

4.2.2 Using Hermite Polynomials

Hermite interpolation involves a function and its first m derivatives at n data points.

$$\|f(x) - H(x)\| = \frac{f^{(K)}(\eta)}{K!} \prod_{i=1}^n (x - x_i)^{k_i}$$

where $K = mn$ is the total number of data points. and $k_i = m - 1$.

4.2.3 Using Newton's Method

The Newton basis polynomials are defined as

$$N(x) = \sum_{j=0}^k a_j \eta_j(x)$$

With basis polynomials

$$\eta_0(x) = 1 \quad \eta_j(x) = \prod_{i=0}^{j-1} (x - x_i) \quad j > 0$$

And coefficients defined as the forward differences

$$a_j = [y_0, \dots, y_j] = \frac{[y_1, \dots, y_j] - [y_0, \dots, y_{j-1}]}{x_j - x_0}$$

$$\begin{array}{ll} x_0 & y_0 = [y_0] \\ & [y_0, y_1] \\ x_1 & y_1 = [y_1] \quad [y_0, y_1, y_2] \\ & [y_1, y_2] \quad [y_0, y_1, y_2, y_3] \\ x_2 & y_2 = [y_2] \quad [y_1, y_2, y_3] \\ & [y_2, y_3] \\ x_3 & y_3 = [y_3] \end{array}$$

So then

$$N(x) = [y_k] + [y_k, y_{k-1}] (x - x_k) + \dots + [y_k, y_0] (x - x_k)(x - x_{k-1}) \dots (x - x_1)$$

If x_i are equally spaces with $x_i = x_0 + ih$, $x = s + x_0$, then $x - x_i = s - ih$

$$N(x) = [y_k] + [y_k, y_{k-1}] (s - kh) + \dots + [y_k, y_0] (s - kh)(s - (k-1)h) \dots (s - h)$$

4.3 Approximation by Polynomial

The Stone-Weierstrass Theorem (also known as Weierstrass Approximation Theorem) states that every continuous function on a closed interval can be uniformly approximated by a polynomial function. That is, for every $\epsilon > 0$, there exists an n and a n th degree polynomial $p_n(x)$ such that

$$\|f - p\|_{\infty} < \epsilon$$

Consider the Bernstein polynomials on $[0, 1]$. Notice that

$$f(x) = f(x)(x + (1-x))^n = f(x) \sum_{i=0}^n \binom{n}{i} x^i (1-x)^{n-i}$$

So then

$$f(x) - (B_n f)(x) = \sum_{i=0}^n \left(f(x) - f\left(\frac{i}{n}\right) \right) \binom{n}{i} x^i (1-x)^{n-i}$$

and thus

$$|f(x) - (B_n f)(x)| \leq \sum_{i=0}^n \left| f(x) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} x^i (1-x)^{n-i}$$

This is a continuous function on a compact interval, thus

- It's maximum will be achieved, say at $x = \eta$
- The function is uniformly continuous on the interval, so for all $\epsilon > 0$, there exists a $\delta > 0$ such that for all $x, |x - y| < \delta$ implies $|f(x) - f(y)| < \epsilon$

Let δ be such that $|f(\eta) - f(x)| < \frac{\epsilon}{2}$. Now let N denote the index subset such that $\frac{i}{n}$ is within δ of η . So then

$$\begin{aligned} |f(x) - (B_n f)(x)| &\leq \sum_{i \in N} \frac{\epsilon}{2} \binom{n}{i} \eta^i (1-\eta)^{n-i} + \sum_{i \notin N} \left| f(\eta) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} \eta^i (1-\eta)^{n-i} \\ &< \frac{\epsilon}{2} + \sum_{i \notin N} \left| f(\eta) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} \eta^i (1-\eta)^{n-i} \end{aligned}$$

Now notice that

$$\begin{aligned} \sum_{i \notin N} \left| f(\eta) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} \eta^i (1-\eta)^{n-i} &\leq \sum_{i \notin N} \left| f(\eta) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} \eta^i (1-\eta)^{n-i} \frac{(\eta - \frac{i}{n})^2}{\delta^2} \\ &\leq \sum_{i \notin N} \left(|f(\eta)| + \left| f\left(\frac{i}{n}\right) \right| \right) \binom{n}{i} \eta^i (1-\eta)^{n-i} \frac{(\eta - \frac{i}{n})^2}{\delta^2} \end{aligned}$$

Let M be such that $|f(x)| < M$, so

$$\sum_{i \notin N}^n \left| f(\eta) - f\left(\frac{i}{n}\right) \right| \binom{n}{i} \eta^i (1-\eta)^{n-i} \leq 2M\delta^{-2} \sum_{i=1}^n \binom{n}{i} \eta^i (1-\eta)^{n-i} \left(\eta - \frac{i}{n}\right)^2$$

Somehow we get this down to

$$|f(x) - (B_n f)(x)| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

4.4 Best Approximation

Let V be a Banach space and let $T \subset V$, an approximation p is a best approximation if for a given f ,

$$\|f - p\| \leq \|f - \hat{p}\| \quad \forall \hat{p} \in T$$

In other words, if

$$\|f - p\| = E_T(f) = \inf_{\hat{p} \in T} \|f - \hat{p}\|$$

- A best approximation exists if T is compact

Since finite dimensional linear spaces are compact, at least one best approximation is guaranteed to exist. This amounts to finding the coefficients $\{\alpha_i\}$ to minimize the distance

$$d(\alpha_1, \dots, \alpha_n) = \|f - (\alpha_1 v_1 + \dots + \alpha_n v_n)\|$$

4.4.1 pre-Hilbert Case

In the case that V is a pre-Hilbert space and $f \in U$, $p \in U$ the best approximation from U if and only if

$$\langle f - p, v \rangle = 0 \quad \forall v \in U$$

This can be used to compute the best approximation using the normal equations. Pick $\{g_k\}$ such that $\text{span } \{g_k\} = U$.

$$\left\langle f - \sum_{j=1}^n \alpha_j g_j, g_k \right\rangle = 0 \implies \sum_{j=1}^n \alpha_j \langle g_j, g_k \rangle = \langle f, g_k \rangle \quad \forall k$$

If $\{g_k\}$ is orthonormal,

$$\alpha_k = \langle f, g_k \rangle \quad \forall k$$

- Zero Theorem: If the set of polynomials $\{\psi_1, \dots, \psi_n\}$ forms an orthonormal set on $[a, b]$, with respect to a weight function $\omega(x)$, then each of these polynomials has only simple real zeros, all of which lie in (a, b)

4.5 Least Squares

Given an inner product $\langle \cdot, \cdot \rangle$ and a set of functions $\{\phi_i\}_{i=0,\dots,n}$, we see the best approximation of $f(x)$ in the norm given by this inner product. That is, we seek to choose $\{c_i\}$ such that for any other choice of sets $\{\hat{c}_i\}$,

$$\left\| f(x) - \sum_{i=0}^n c_i \phi_i(x) \right\|_{\mathcal{H}} \leq \left\| f(x) - \sum_{i=0}^n \hat{c}_i \phi_i(x) \right\|_{\mathcal{H}}$$

This is achieved by solving the system

$$\begin{pmatrix} \langle \phi_0, \phi_0 \rangle & \langle \phi_0, \phi_1 \rangle & \dots & \langle \phi_0, \phi_n \rangle \\ \langle \phi_1, \phi_0 \rangle & \langle \phi_1, \phi_1 \rangle & & \\ \vdots & & \ddots & \\ \langle \phi_n, \phi_0 \rangle & \langle \phi_n, \phi_1 \rangle & & \langle \phi_n, \phi_n \rangle \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \\ \vdots \\ c_n \end{pmatrix} = \begin{pmatrix} \langle f, \phi_0 \rangle \\ \langle f, \phi_1 \rangle \\ \vdots \\ \langle f, \phi_n \rangle \end{pmatrix}$$

In the discrete case, we instead consider the system of equations

$$\begin{pmatrix} \phi_0(x_0) & \phi_1(x_0) & \dots & \phi_n(x_0) \\ \phi_0(x_1) & \phi_1(x_1) & & \\ \vdots & & \ddots & \\ \phi_0(x_m) & \phi_1(x_m) & & \phi_n(x_m) \end{pmatrix}_{m \times n} \begin{pmatrix} c_0 \\ c_1 \\ \vdots \\ c_n \end{pmatrix}_{n \times 1} = \begin{pmatrix} f(x_0) \\ f(x_1) \\ \vdots \\ f(x_m) \end{pmatrix}_{m \times 1}$$

Which has the form $A\vec{c} = \vec{f}$. This has the solution

$$A^* A \vec{c} = A^* \vec{f} \implies \vec{c} = (A^* A)^{-1} A^* \vec{f}$$

5 Numerical Integration

5.1 Connection to Interpolation

Let $\{p_k(x)\}_{k=0}^n$ be a set of orthogonal polynomials on $[a, b]$ with respect to the inner product

$$\langle f, g \rangle = \int_a^b f(x)g(x)\omega(x)dx$$

The quadrature rule given by

$$I(f) = \int_a^b f(x)\omega(x)dx \approx Q(f) = \int_a^b \omega(x) \left(\sum_{j=1}^n f(x_j)L_j(x) \right) dx = \sum_{j=1}^n w_j f(x_j)$$

$$w_j = \int_a^b L_j(x)\omega(x)dx \quad L_j(x) = \prod_{k=1, k \neq j}^n \frac{x - x_k}{x_j - x_k}$$

is exact for polynomials of degree at most $n - 1$.

If $x_j, j = 1, \dots, n$ are the zeros of the highest degree polynomial $p_n(x)$, then the quadrature rule given above has exact degree $2n - 1$. This is because

$$f(x) = p_n(x)q(x) + r(x) \quad q(x), r(x) \in \mathbb{P}_{n-1}(x)$$

So then

$$I(f) = I(p_n q + r) = \int_a^b q(x)p_n(x)\omega(x)dx + \int_a^b r(x)\omega(x)dx = \int_a^b r(x)\omega(x)dx = I(r) = Q(r)$$

since our formula is exact for $r(x)$. So then since $p_n(x_j) = 0$,

$$Q(r) = \sum_{j=1}^n w_j r(x_j) = \sum_{j=1}^n w_j (p_n(x_j)q(x_j) + r(x_j)) = Q(f)$$

Thus this method works for polynomials of degree $2n-1$. If we pick $f \in \mathbb{P}_{2n}$, then $q(x) \in \mathbb{P}_n$ and $r(x) \in \mathbb{P}_{n-1}$, so $\langle q, p_n \rangle \neq 0$, so

$$I(f) - Q(f) = I(qp_n + r) - Q(p_nq + r) = Q(r) + \langle q, p_n \rangle - Q(r) = \langle q, p_n \rangle \neq 0$$

5.2 Newton Cotes Formulas

The Newton Cotes formulas assume $n+1$ equidistant sampling points where n is the degree of the method. The closed formulas include the endpoints:

$$x_i = a + \frac{i}{n}(b-a) \quad f_i = f(x_i) \quad i = 0, \dots, n$$

while the open formulas use

$$x_i = a + \frac{i}{n}(b-a) \quad f_i = f(x_i) \quad i = 1, \dots, n-1$$

The methods are derived via the Lagrange Basis polynomials.

$$\int_a^b f(x)dx \approx I = \int_a^b L(x)dx = \int_a^b \left(\sum_{i=0}^n f(x_i)l_i(x) \right) dx = \sum_{i=0}^n f(x_i) \int_a^b l_i(x)dx$$

or for open rules

$$\int_a^b f(x)dx \approx I = \sum_{i=1}^{n-1} f(x_i) \int_a^b l_i(x)dx$$

These formulas can suffer from instability for large n unless they are converted into a composite rule.

5.2.1 Closed Schemes

- Trapezoid Rule:

A degree 1 closed scheme is

$$I = \frac{b-a}{2} (f_0 + f_1) \quad h = b-a$$

With error term

$$E = -\frac{(b-a)^3}{12} f^{(2)}(\xi)$$

- Simpson's Rule:

A degree 2 closed scheme is

$$I = \frac{b-a}{6} (f_0 + 4f_1 + f_2) \quad h = \frac{(b-a)}{2}$$

With error term

$$E = -\frac{(b-a)^5}{2880} f^{(4)}(\xi)$$

- Simpson's $\frac{3}{8}$ Rule:
A degree 3 scheme is

$$I = \frac{b-a}{8} (f_0 + 3f_1 + 3f_2 + f_3) \quad h = \frac{(b-a)}{3}$$

With error term

$$E = -\frac{(b-a)^5}{6480} f^{(4)}(\xi)$$

- Boole's Rule :
A degree 4 scheme is

$$I = \frac{b-a}{90} (7f_0 + 32f_1 + 12f_2 + 32f_3 + 7f_4) \quad h = \frac{(b-a)}{4}$$

With error term

$$E = -\frac{(b-a)^7}{1935360} f^{(6)}(\xi)$$

5.2.2 Open Schemes

- Midpoint Rule:
A degree 2 open scheme is

$$I = b-a (f_1) \quad h = \frac{(b-a)}{2}$$

With error term

$$E = \frac{(b-a)^3}{24} f^{(2)}(\xi)$$

- Trapezoid Method:
A degree 3 open scheme is

$$I = \frac{b-a}{2} (f_1 + f_2) \quad h = \frac{(b-a)}{3}$$

With error term

$$E = \frac{(b-a)^3}{36} f^{(2)}(\xi)$$

- Milne's Rule:
A degree 4 open scheme is

$$I = \frac{b-a}{4} (2f_1 - f_2 + 2f_3) \quad h = \frac{(b-a)}{4}$$

With error term

$$E = \frac{7(b-a)^5}{23040} f^{(4)}(\xi)$$

6 Iterative Schemes for $Ax = b$

Given an iterative scheme where $A = L + D + R = M + N$

$$M\vec{x}^{(k+1)} = N\vec{x}^{(k)} + b$$

Thus if we have the amplification matrix $G = M^{-1}N$,

$$\vec{e}^{(k+1)} = G\vec{e}^{(k)}$$

By induction, $\vec{e}^{(k+1)} = G^k \vec{e}^{(0)}$. Perform the spectral decomposition of G ,

$$\vec{e}^{(k)} = R\Gamma^K R^{-1} \vec{e}^{(0)}$$

If $\Gamma = \text{diag}(\gamma_1, \dots, \gamma_n)$ with $|\gamma_1| \geq \dots \geq |\gamma_n|$ then we have convergence if $|\gamma_1| < 1$. since then $G^K \rightarrow 0$ as $k \rightarrow \infty$.

6.1 Jacobi Iteration

$$M = D \quad N = L + R = D - A \implies G = I - D^{-1}A$$

or alternatively,

$$Du_{k+1} = -(L + R)u_k + b$$

This method converges when $\rho(D^{-1}(L + R)) < 1$

- This method is guaranteed to converge when is strictly or irreducibly diagonally dominant. That is, when $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|$

6.2 Gauss Seidel

$$M = L + D \quad N = -R \implies G = -(L + D)^{-1}R$$

or alternatively,

$$(L + D)u_{k+1} = -Ru_k + b$$

This method converges when $\rho((L + D)^{-1}R) < 1$

- This method is guaranteed to converge when is strictly or irreducibly diagonally dominant.
- This method is guaranteed to converge when A is SPD

6.3 Successive Over-Relaxation (SOR)

Let ω be a parameter, and then

$$A = D + L + R \quad M = \frac{1}{\omega}(D + \omega L) \quad N = \frac{1}{\omega}((1 - \omega)D - \omega R)$$

so then

$$G = \omega(D + \omega L)^{-1} \frac{1}{\omega}((1 - \omega)D - \omega R) = (D + \omega L)^{-1}((1 - \omega)D - \omega R)$$

That is,

$$(D + \omega L)u_{k+1} = ((1 - \omega)D - \omega R)u_k + f$$

- If $A \in \mathbb{R}^{n \times n}$ is SPD, and $D + \omega L$ is nonsingular, then SOR converges for all $0 < \omega < 2$
- If $\omega = 1$, this method reduces to the Gauss Seidel method.

7 Matrix Decompositions


7.1 Singular Value Decomposition

The SVD of a $m \times n$ matrix is a factorization of the form

$$M = U\Sigma V^*$$

where

- U is an $m \times m$ unitary matrix made of the left-singular vectors of M , the eigenvectors of MM^*
- Σ is an $m \times n$ rectangular diagonal matrix with non-negative real numbers on the diagonal, corresponding to the square roots of the non-zero eigenvalues of both M^*M and MM^*
- V is an $n \times n$ unitary matrix made of the right-singular vectors of M , the eigenvectors of M^*M



8 Nonlinear Solvers

8.1 Newton's Method

$$x_{k+1} = x_k - ((Jf)(x))^{-1} f(x_k)$$

8.2 Secant Method

$$x_{k+1} = x_k - f(x_k) \frac{x_k - x_{k-1}}{f(x_k) - f(x_{k-1})}$$

Letting $x_{k+1} = x^* + e_{k+1}$ for the solution x^* ,

$$e_{k+1} = e_k - f(x^* + e_k) \frac{e_k - e_{k-1}}{f(x^* + e_k) - f(x^* + e_{k-1})}$$

We see that

$$e_{k+1} = \frac{f(x^* + e_k)e_k - f(x^* + e_{k-1})e_k - f(x^* + e_k)(e_k - e_{k-1})}{f(x^* + e_k) - f(x^* + e_{k-1})}$$

Which simplifies to

$$e_{k+1} = \frac{f(x^* + e_k)e_{k-1} - f(x^* + e_{k-1})e_k}{f(x^* + e_k) - f(x^* + e_{k-1})}$$

We see that since $f(x^*) = 0$,

$$\begin{aligned} f(x^* + e_k) &= f'(x^*)e_k + \frac{1}{2}f''(x^*)(e_k)^2 + \frac{1}{6}f'''(\eta)(e_k)^3 \\ f(x^* + e_{k-1}) &= f'(x^*)e_{k-1} + \frac{1}{2}f''(x^*)(e_{k-1})^2 + \frac{1}{6}f'''(\eta)(e_{k-1})^3 \end{aligned}$$

So then

$$f(x^* + e_k) - f(x^* + e_{k-1}) = (e_k - e_{k-1}) \left(f'(x^*) + \frac{1}{2}f''(x^*)(e_k + e_{k-1}) + \frac{1}{6}f'''(\eta)q(e_k, e_{k-1}) \right)$$

And

$$\begin{aligned} f(x^* + e_k)e_{k-1} - f(x^* + e_{k-1})e_k &= \frac{1}{2}f''(x^*)(e_k^2e_{k-1} - e_ke_{k-1}^2) + \frac{1}{6}f'''(\eta)(e_k^3e_{k-1} - e_ke_{k-1}^3) \\ &= e_ke_{k-1}(e_k - e_{k-1}) \left(\frac{1}{2}f''(x^*) + \frac{1}{6}f'''(\eta)(e_k + e_{k-1}) \right) \end{aligned}$$

So

$$e_{k+1} = e_ke_{k-1} \frac{\left(\frac{1}{2}f''(x^*) + \frac{1}{6}f'''(\eta)(e_k + e_{k-1}) \right)}{\left(f'(x^*) + \frac{1}{2}f''(x^*)(e_k + e_{k-1}) + \frac{1}{6}f'''(\eta)q(e_k, e_{k-1}) \right)}$$

assuming $f''(x^*) \neq 0$

$$|e_{k+1}| \leq |e_k| |e_{k-1}| \left| \frac{\left(\frac{1}{2}f''(x^*) + \frac{1}{6}f'''(\eta)(e_k + e_{k-1}) \right)}{\left(f'(x^*) + \frac{1}{2}f''(x^*)(e_k + e_{k-1}) + \frac{1}{6}f'''(\eta)q(e_k, e_{k-1}) \right)} \right|$$

So if we assume small initial errors,

$$|e_{k+1}| \leq M |e_k| |e_{k-1}| \quad M = \frac{1}{2} \frac{|f''(x^*)|}{|f'(x^*)|}$$

So assume $|e_{k+1}| \approx C |e_k|^p$, and $|e_k| \approx D |e_{k-1}|^p$. Then $|e_{k+1}| \approx CD^p |e_{k-1}|^{p^2}$. So then

$$CD^p |e_{k-1}|^{p^2} \approx D |e_{k-1}|^p |e_{k-1}|$$

So we want to solve

$$p^2 - p - 1 = 0 \implies p = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 \pm \sqrt{5}}{2}$$

So our method is of order $\frac{1+\sqrt{5}}{2}$.

9 ODE Solvers

We consider an ODE of the form

$$u'(t) = f(t, u)$$

9.1 Single Step Methods

Single step methods have the form

$$y_{i+1} = y_i + h\Phi(t, y; h), h = \delta t > 0$$

Given a solution $u(t)$, we define the truncation error of the method to be the difference between the exact and approximate solutions over a step of size h :

$$T(t, y; h) = \frac{1}{h} (y_{i+1} - u(t+h))$$

or alternatively,

$$T(t, y; h) = \Phi(t, y; h) - \frac{1}{h} (u(t+h) - u(t))$$

- A method is consistent if $T(t, y; h) \rightarrow 0$ uniformly as $h \rightarrow 0$. That is if

$$\lim_{h \rightarrow 0} \|T(t, y; h)\|_{\infty} \rightarrow 0$$

We have consistency if and only if $T(t, y; 0) = f(t, y)$.

- The method is accurate of order p if for some vector norm, we have

$$\|T(t, y; h)\| \leq Ch^p$$

for some C . That is, as $h \rightarrow 0$,

$$T(t, y; h) = \mathcal{O}(h^p)$$

$p > 0$ implies consistency, and usually $p \geq 1$.

- The method is zero stable if there exists a constant C such that for any two sequences y_k and \hat{y}_k with $y_0 \neq \hat{y}_0$,

$$|y_k - \hat{y}_k| \leq C \max_{0 \leq i \leq k-1} \{|y_i - \hat{y}_i|\} \text{ as } h \rightarrow 0$$

Sufficient condition: A method is zero stable for any ode $y' = f(t, y)$ for Lipschitz f if and only if the spectral radius of the iteration is bounded by 1 with equality holding only for simple eigenvalues.

9.1.1 Single Step Methods

Euler's method has the form

$$y_{i+1} = y_i + hf(t_i, y_i)$$

And is first order unless $f_t + f_y f = 0$

One can use Taylor expansion methods to improve the order of accuracy. We calculate the total derivatives of f ,

$$\begin{aligned} f^{(0)} &= f(t, y) \\ f^{(1)} &= f_t(t, y) + f_y(t, y)f(t, y) \\ &\vdots \\ f^{(k+1)} &= f_t^{(k)}(t, y) + f_y^{(k)}(t, y)f^{(k)}(t, y) \end{aligned}$$

Then using $u^{(k+1)}(t) = f^{(k)}(t, u(t))$, we can form an approximation of order p via

$$y_{i+1} = y_i + h \left(f^{(0)}(t, y) + \frac{1}{2}hf^{(1)}(t, y) + \dots + \frac{1}{p!}h^{p-1}f^{(p-1)}(t, y) \right)$$

It can be shown that for this method,

$$\|T(t, y, h)\| \leq \frac{C_p}{(p+1)!}h^p$$

Again this may perform better if $f^{(p)} = 0$.

9.2 Runge-Kutta Methods

We could use multiple stages in our calculation, using

$$\begin{aligned} k_1(t, y) &= f(t, y) \\ k_2(t, y) &= f(t + \mu h, y + \mu h k_1) \\ y_{i+1} &= y_i + h(\alpha_1 k_1 + \alpha_2 k_2) \end{aligned}$$

In general, two stages offer at best an order 2 method, achieved by using the family

$$\alpha_1 + \alpha_2 = 1 \quad \alpha_2 \mu = \frac{1}{2} \quad \alpha_2 \neq 0 \text{ (arbitrary)}$$

To extend the idea of the two-stage methods to r stage methods, we use

$$\begin{aligned} k_1(t, y) &= f(t, y) \\ k_s(t, y; h) &= f\left(t + \mu_s h, y + h \sum_{j=1}^{s-1} \lambda_{sj} k_j\right) \\ y_{i+1} &= y_i + h \left(\sum_{s=1}^r \alpha_s k_s \right) \end{aligned}$$

For consistency we require $\sum_{s=1}^r \alpha_s = 1$, and it is natural to impose

$$\mu_s = \sum_{j=1}^{s-1} \lambda_{sj} \quad s = 2, 3, \dots, r$$

These methods can be summarize in the form of a Butcher Tableau as seen here:

$$\begin{array}{c|ccc}
 \mu_1 = 0 & & & \\
 \mu_2 & \lambda_{21} & & \\
 \mu_3 & \lambda_{31} & \lambda_{32} & \\
 \vdots & \vdots & & \ddots \\
 \mu_s & \lambda_{s1} & \lambda_{s2} & \dots \lambda_{s,s-1} \\
 \hline
 & \alpha_1 & \alpha_2 & \dots \alpha_s
 \end{array}$$

So if $f(u) = Cu$ is linear,

$$\begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ -\lambda_{s1} & \dots & 1 \end{pmatrix} \begin{pmatrix} k_1 \\ \vdots \\ k_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} Cy_i$$

and so

$$\begin{aligned}
 y_{i+1} &= y_i + h \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_s \end{pmatrix}^T \begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ -\lambda_{s1} & \dots & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} Cy_i \\
 y_{i+1} &= \left(1 + Ch \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_s \end{pmatrix}^T \begin{pmatrix} 1 & & \\ \vdots & \ddots & \\ -\lambda_{s1} & \dots & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right) y_i
 \end{aligned}$$

9.2.1 Stability

Runge Kutta Methods are absolute stable if the spectral radius of the amplification factor is strictly bounded by 1. That is, for linear scalar problems $u'(t) = Cu(t)$, if

$$|1 + ChG| < 1$$

This leads to a region of Ch for which the method is stable.

$$-1 < 1 + ChG < 1 \implies -2G^{-1} < Ch < 0$$

9.3 0

Linear Multistep Methods

We could develop a recurrence - like method in the form

$$\sum_{j=0}^k \alpha_j y_{n+j} = h \sum_{j=0}^k \beta_j f(t_{n+j}, y_{n+j})$$

With real $\{\alpha_i\}, \{\beta_i\}$, $\alpha_k \neq 0$ and α_0, β_0 are not both zero. If $\beta_k = 0$, then this method is explicit. Otherwise it is implicit. This is linear since it involves linear combinations of $\{y_i\}$.

- Implicit Euler Method

$$y_{n+1} = y_n + \frac{1}{2}h(f_{n+1})$$

- The Implicit Trapezium method

$$y_{n+1} = y_n + \frac{1}{2}h(f_{n+1} + f_n)$$

- The Explicit 4-step Adams-Bashforth Method

$$y_{n+4} = y_{n+3} + \frac{1}{24}h(55f_{n+3} - 59f_{n+2} + 37f_{n+1} - 9f_n)$$

- For a linear multi-step method that is consistent with the ODE, zero stability is necessary and sufficient for convergence.

9.4 Absolute Stability

Apply a given method

A linear multi-step method is absolutely stable for a given Ch if and only if for that Ch , all of the roots of the stability polynomial $p(z, Ch)$ satisfy $|r_i| < 1$, $i = 1, \dots, k$.

- Absolutely stable methods are also zero stable.

9.5 Implicit Methods

We can also consider implicit Runge-Kutta methods

$$k_s = f\left(t + \mu_s h, y + h \sum_{j=1}^r \lambda_{sj} k_j(t, y; h)\right) \quad s = 1, 2, \dots, r$$

$$y_{i+1} = y_i + h \left(\sum_{s=1}^r \alpha_s k_s \right)$$

For consistency we require $\sum_{s=1}^r \alpha_s = 1$, and it is natural to impose

$$\mu_s = \sum_{j=1}^{s-1} \lambda_{sj} \quad s = 2, 3, \dots, r$$

The maximum order attainable by any Runge Kutta method is equal to the number of stages used for methods of less than order 4.