

Principles of Pharmacogenomics: Pharmacokinetic, Pharmacodynamic, and Clinical Implications

Y. W. Francis Lam^{1,2}

¹Department of Pharmacology, University of Texas Health Science Center at San Antonio, San Antonio, TX, United States; ²College of Pharmacy, University of Texas at Austin, Austin, TX, United States

OUTLINE

Objectives	2	Polymorphisms in Drug-Transporter Genes	21
Introduction	2	<i>The ABC-Efflux Transporters</i>	21
Polymorphisms in Cytochrome P450 Enzymes	3	ABCB1	21
CYP2D6	4	ABCC1 and ABCC2	23
CYP2C19	9	ABCG2	23
CYP2C9	11	<i>The SLC-Uptake Transporters</i>	24
CYP2C8	13	Organic Anion Transporting Polypeptides	24
CYP3A4/5/7	13	OAT1B1	24
CYP4F2	15	OATP2B1	26
CYP2B6	15	OATP1B3	26
CYP2A6	17	Organic Cation Transporters	26
CYP1A2	17	Organic Anion Transporters	27
Polymorphisms in Non-CYP450 Drug-Metabolizing Enzymes	18	Polymorphisms in Drug-Target Genes	27
UDP-Glucuronosyl Transferase	18	<i>Drug Target Receptor Genes in Oncology</i>	28
Thiopurine-S-methyltransferase	19	<i>Drug Target Receptor Genes in Cardiology</i>	30
Dihydropyrimidine Dehydrogenase (DYPD)	19	<i>Drug Target Genes in Psychiatry</i>	32
N-acetyltransferase	20	<i>Drug Target Genes in Pain Management</i>	32
Glutathione-S-transferase	20	Signal Transduction Proteins	33
		Enzyme Genes	34
		Ion Channel Genotype	36

Conclusion	37	References	37
Questions for Discussion	37		

OBJECTIVES

1. Describe how a single-nucleotide polymorphism (SNP) can affect protein function or expression, and consequently, influence drug response.
2. Explain how genetic polymorphisms for drug metabolism or drug-transporter proteins may influence drug pharmacokinetics.
3. Contrast phenotypic responses to genetic variation for drug metabolism versus drug-target proteins.
4. Describe novel drug developed based on an understanding of genes involved in disease pathophysiology.
5. Explain how genetic polymorphisms at the drug-target site may influence drug pharmacodynamics.

INTRODUCTION

Significant interpatient variability in drug response is largely attributed to innate differences among individuals in their capacity to process and respond to medications. Pharmacogenomics involves incorporating information about a person’s genotype into drug therapy decisions, with the goal of providing the most effective and safest therapy for that individual. Over the last decade, there have been significant advances in our understanding of the contribution of genetic differences in pharmacokinetics and pharmacodynamics toward interindividual variability in drug response. Not only may pharmacogenomics lead to improved use of existing therapies, but it may also lead to novel drugs developed based on an improved understanding of genetic control of cellular functions.

The human genome comprises approximately 20,000 protein-coding genes. By far the most common variation is the single-nucleotide polymorphism (SNP), which is defined as single-base differences that exist between individuals. Over 22 million SNPs have been reported in the human genome [1]. SNPs that result in amino acid substitution are termed nonsynonymous. Nonsynonymous SNPs occurring in coding regions of the gene (e.g., exons) can impact protein activity and have significant consequences on responses to medications that depend on the protein for metabolism, transport, or eliciting cellular effects. Synonymous polymorphisms do not result in amino acid substitution; however, those occurring in a gene regulatory region (e.g., promoter region, intron) may alter gene expression and the amount of protein that is produced. Two or more SNPs are often inherited together more frequently than would be expected based on chance alone. This is referred to as linkage disequilibrium (LD). A haplotype refers to a set of SNPs that are in LD. Other types of variation that can affect gene expression or protein conformation include insertion–deletion polymorphisms (indels), short tandem repeats, and copy number variants (CNVs). A CNV represents a DNA segment (≥1 kb) with a variable number of copies of that segment, because of duplications, deletions, or rearrangement, and constitutes a major source of interindividual variation in the human genome. A unique reference SNP identifier (rs number) is assigned for each genetic variant, and exists as an SNP data repository, the National Center for Biotechnology Information (NCBI) Single-Nucleotide Polymorphism Database (dbSNP).

Polymorphisms commonly occur for genes encoding drug metabolism, drug transporter,




FIGURE 1.1 Location of genetic variations affecting drug response. Those occurring in genes for drug metabolism or transport can affect drug pharmacokinetics, whereas SNPs in genes encoding for drug-target proteins can impact drug pharmacodynamics.

and drug-target proteins (Fig. 1.1). Drug metabolism and transporter genotypes can affect drug availability at the target site, whereas drug-target genotype can affect a patient's sensitivity to a drug. In many instances, genes for proteins involved in drug disposition, together with genes for proteins at the drug-target site, jointly influence drug response. In addition, genetic polymorphisms in absorption, distribution, metabolism, and excretion (ADME) and target genes also contribute to ethnic heterogeneity in drug response [2]. Research advances have resulted in continued identification of association between genetic polymorphisms and response, with recent focus on genome-wide association studies (GWAS) in populations worldwide.

The terms pharmacogenetics and pharmacogenomics are often used interchangeably. Because drug responses are mostly determined by multiple, rather than single, proteins, recent trends of investigations on determinants of drug response have shifted from pharmacogenetics to pharmacogenomics. However, for simplicity, this chapter treats pharmacogenetics and pharmacogenomics as synonymous.

Despite the scientific advances made, personalized medicine envisioned many years ago has in many cases yet to become a reality. Exceptions to this largely exists in oncology and more recently in cardiology, in which genotyping to determine clopidogrel effectiveness is starting to become routine at some large academic medical centers [3,4]. Examples of genotype-guided therapies are beginning to emerge in other therapeutic areas, which are discussed in detail throughout this book. However, significant challenges still exist in ethical, socioeconomic, regulatory, legislative, drug development, and educational issues that need to be addressed and resolved before personalized medicine can be practically and satisfactorily implemented in clinical practice on a broader scale. The goal of this chapter is to review the pharmacokinetic and pharmacodynamics basis of individualized therapy, and briefly discuss the challenges of implementing pharmacogenomics in clinical practice. Further indepth discussion of specific therapeutic areas and/or disease states, as well as ethical, socioeconomic, regulatory, legislative, drug development, technological, and educational issues will be the focus of subsequent chapters.

POLYMORPHISMS IN CYTOCHROME P450 ENZYMES

The cytochrome P450 (CYP) superfamily of isoenzymes represents the most important and studied metabolic enzymes that exhibit clinically relevant genetic polymorphisms. Within this superfamily of isoenzymes, 57 different CYP genes and 58 pseudogenes have been identified, and, based on the similarity in their amino acid sequences, are grouped into 18 families and 44 subfamilies with increasing extent of sequence similarity. Of these genes and pseudogenes, 42 are involved in the metabolism of exogenous xenobiotics and endogenous substances, such as steroids and prostaglandins, and 15 are known to be

involved in the metabolism of drugs in humans [5]. Information regarding *CYP* allele nomenclature and specific genetic variations defining different metabolic phenotypes had been available at the Karolinska Institute website: www.cypalleles.ki.se, for more than a decade, and recently moved to the new Pharmacogene Variation (PharmVar) Consortium, which serves as a new hub for pharmacogene nomenclature [6].

The genes encoding *CYP*s are highly polymorphic, with SNPs in the *CYP* gene locus accounting for most of the variations in *CYP* activity, resulting in functional genetic polymorphism for several isoenzymes, including *CYP2A6*, *CYP2B6*, *CYP2C9*, *CYP2C19*, *CYP2D6*, and *CYP3A4/5*. Additional types of *CYP* polymorphisms cause gene deletions, deleterious mutations resulting in premature stop codon or splicing defects, amino acid changes, gene duplications, and CNV. Different alleles or functional variants of these polymorphisms for individual drug metabolizing genes are defined with a “star” (*) designation. A combination of two *alleles, for example, *CYP2D6**1/*1, is used to classify individuals into several genetically defined metabolic phenotypes with different expressions of enzyme activity. In general, the poor metabolizers (PMs) inherit two defective or deleted alleles and exhibit abolished-enzyme activity; the intermediate metabolizers (IMs) carry either one functional and one defective allele, or two partially defective alleles, and, in both cases, have reduced activity of the enzyme. The normal metabolizers are typically known as the extensive metabolizers (EMs) with two functional alleles and normal enzyme activity; and the ultrarapid metabolizers (UMs) carry a duplicated or amplified gene variant, resulting in two or multiple copies of the functional allele and very high enzyme activity.

In general, the clinical consequences of genetically altered-enzyme activity would depend on whether the pharmacological activity resides with the parent compound or the metabolite, and the relative contribution of the polymorphic

isoenzyme to the overall metabolism of the drug. For the majority of the drugs, PMs would exhibit a higher risk of adverse drug reactions (ADRs), whereas UMs would experience lower efficacy when administered standard-dosage regimen of a drug that is mostly dependent on the polymorphic enzyme for elimination. In the case of a prodrug, the UMs exhibit higher incidence of ADRs, and the PMs experience lower efficacy, reflecting a difference in the extent of therapeutically active metabolite formed between the two metabolic genotypes.

Among the different *CYP* gene polymorphisms, those affecting *CYP2D6*, *CYP2C19*, and *CYP2C9* are currently the most relevant with also the most abundant data, as well as representing most of the revised regulatory labeling information. Their potential role in translating the expanding pharmacogenomic knowledge into dose requirements and therapeutic decisions will be discussed first. An overview of the other major *CYP* isoenzymes will also be presented.

CYP2D6

CYP2D6 is the only drug-metabolizing *CYP* enzyme that is not inducible, and the significant interindividual differences in enzyme activity are largely attributed to genetic variations. *CYP2D6* is located on chromosome 22 and consists of 4382 nucleotides. The *CYP2D6* gene, which codes for the *CYP2D6* enzyme, is composed of 497 amino acids. In addition, the *CYP2D6* gene polymorphisms are also the best characterized among all of the *CYP* variants, with at least 100 alleles identified. Nevertheless, Sistonen et al. [7] demonstrated that, even with the extensive number of alleles, determining 20 different haplotypes by genotyping 12 SNPs could predict the real phenotype with 90%–95% accuracy.

Among the multiple *CYP2D6* alleles, *CYP2D6**1, *CYP2D6**2, *CYP2D6**33, and *CYP2D6**35 are active alleles with normal enzyme activity, whereas the two most

important null variants are *CYP2D6**4 (c.1846G>A, rs3892097) and *CYP2D6**5 (gene deletion), resulting in an inactive enzyme and absence of enzyme, respectively. Significant reduction in enzyme activity is commonly associated with *CYP2D6**10 (c.100C>T, rs1065852), *CYP2D6**17 (c.1023C>T, rs28371706, c.2850C>T, rs16947), and *CYP2D6**41 (c.2988G>A, rs28371725), and phenotypically expressed as IM. In addition, to these reduced function alleles, the IM phenotype has also been associated with the *CYP2D6**9, *29, and *36 variants [5]. Additional loss-of-function alleles include *CYP2D6**3, *6–*8, *11–*16, *19–*21, *38, *40, and *42. *CYP2D6* is also the first CYP isoenzyme for which CNVs were reported [8]. Individuals carrying up to 13 functional copies of the *CYP2D6**2 allele [9] have been reported to exhibit variation in response to different drugs [10,11]. After these initial reports, gene duplication has also been documented for the *CYP2D6**1, *4, *6, *10, *17, *29, *35, *41, *43, and *45 variants [12]. Therefore, although UMs can result from duplication or multiduplication of the active *CYP2D6* gene, duplication of partially functional and nonfunctional genes

can also occur, resulting in different levels of gene expression and phenotypes of metabolic importance (Table 1.1). A CYP activity score has also been recommended for use in classifying the different 2D6 phenotypic groups [13]. More recently, a software tool (originally named “Constellation” and subsequently renamed as “Astrolabe”) capable of allowing rapid, automated phenotype assignment has been made available for academic research at no cost [14].

Significant interethnic variations in *CYP2D6* allele and phenotype distributions have also been well documented. The normal function *CYP2D6**2 has been reported in approximately 25% of Caucasians, 31% of Africans, and 10%–12% of East Asians [15]. *CYP2D6**4 and *CYP2D6**5 (allelic frequency of about 20%–25% and 4%–6%, respectively) are predominantly found in Caucasian PMs, whereas the predominant variants in people of Asian and African heritage are *CYP2D6**10 (allelic frequency of about 50%) and *CYP2D6**17 (allelic frequency of about 20%–34%), respectively, both resulting in the IM phenotype. Therefore, even though the classic PM phenotypic frequencies determined

TABLE 1.1 Functional *CYP2D6* Polymorphisms, Expected Enzyme Activity, and Predicted Metabolic Phenotypes for Selected Common Variants

Allelic Variants and Polymorphism	Functional Effect on Enzyme Activity	Predicted Metabolic Phenotypes
Active: *1, *2, *2A, *33, *35	Normal activity	Extensive metabolizers: <ul style="list-style-type: none"> • Homozygous carriers of two active alleles
Partially active: *9, *10 (P34S), *17 (T107I, R296C), *29, *36, *41 (splicing defect)	Reduced activity	Heterozygous carriers of an active and a partially active allele Intermediate metabolizers:
Inactive: *3 (<i>frame shift</i>) *4 (<i>splicing defect</i>), *5 (<i>gene deletion</i>) *6 (<i>frame shift</i>), *7, *8, *11, *12, *13, *14, *15, *16, *19, *20, *21, *38, *40, *42 Gene duplication have been reported for both *4 and *6	Loss-of-function	<ul style="list-style-type: none"> • Heterozygous carriers of an active and a loss-of-function allele • Homozygous carriers of two reduced activity alleles • Heterozygous carriers of a partially active allele and a loss-of-function allele
Gene duplications/copy number variants *1, *2, *10, *17, *29, *35, *41, *43, *45	Enhanced activity	Poor metabolizers: <ul style="list-style-type: none"> • Homozygous carriers of two loss-of-function alleles Ultrarapid metabolizers: <ul style="list-style-type: none"> • Carriers of ≥3 active alleles

in Asians (about 0%–1% of population) and Africans (0%–5% of population) are lower than that reported for the Caucasians (5%–14% of population), the high prevalence of *CYP2D6**10 and *CYP2D6**17 in these two IM populations provides a biologic and molecular explanation for reported higher drug concentrations and/or the practice of prescribing lower dosage requirements in people of Asian and African heritage [16–19]. On the other hand, the UM phenotypic frequency is much higher in Northeast Africa and Oceania, including the Saudi Arabian (20%) and black Ethiopian (29%) populations when compared to Caucasians (1%–10%) and East Asians (0%–2%).

Even though accounting for a small percent of total CYP content in the liver, *CYP2D6* mediates the metabolism of approximately 20%–30% of currently marketed drugs, and *CYP2D6* polymorphism affects significantly the elimination of 50% of these drugs [20], which include antidepressants, antipsychotics, analgesics, antiarrhythmics, antiemetics, and anticancer drugs. Although differences in pharmacokinetic parameters (elimination half-lives, clearances, and areas under the plasma concentration–time curves) for *CYP2D6* substrates could be demonstrated among the different metabolic phenotypes, the significant overlap in *CYP2D6* activities in EMs and IMs result in therapeutic implication mostly for the PM and UM phenotypes. In the past, the clinical relevance of *CYP2D6* polymorphism primarily concerned the increased prevalence of ADRs in PMs administered standard doses of drugs that rely significantly on *CYP2D6* for elimination. These drugs include the antianginal agent perhexiline (neuropathy) [21], the antiarrhythmic agent propafenone (proarrhythmic events) [22], and neuroleptic agents such as perphenazine (sedation and parkinsonism) [23,24].

More recently, occurrences of ADRs have also been highlighted in UMs, primarily a result of a 10–30-fold increase in metabolite concentrations. The most cited example is that of codeine, which

is converted by *CYP2D6* to the pharmacologically more active metabolite morphine. UMs administered the usual therapeutic dose of codeine have been reported to exhibit symptoms of narcotic overdose associated with significantly elevated morphine concentrations. This toxicity potential had been highlighted in several case reports [25–29], including a fatal case of a breast-fed infant that was attributed to extensive formation of morphine from codeine taken by the mother who is a UM [26]. (Table 1.2) Prior to this unfortunate case, codeine has been considered safe for managing pain associated with childbirth, as literature reported low amounts of codeine are usually found in breast milk. Therefore, this fatal case underscores the importance of understanding how genes can affect pharmacological and therapeutic outcome associated with exposure to drug and/or active metabolite.

Given the high incidence of codeine use in postgestational women, Madadi et al. subsequently performed a case-control study in breast-fed infants with or without central nervous system depression signs and symptoms after exposed to codeine during breast feedings. They reported that breast-fed infants from mothers who are *CYP2D6* UMs and homozygous carriers of *UGT2B7**2 (rs7439366; *UGT2B7* is a phase 2 enzyme involved in codeine glucuronidation) have an increased risk of potentially life-threatening central nervous system depression [30]. Since 2007, the Food and Drug Administration (FDA) had issued several warning in revised prescribing information for codeine label. Citing the risk of morphine overdose in children and breast-fed infants and warnings from the FDA, the World Health organization, Health Canada, and the European Medicine Agency, the Academy of Pediatrics had recently cautioned the use of codeine in children, regardless of age [31].

Samer et al. reported higher incidence of oxycodone toxicity in UMs that could be partially related to *CYP2D6*-mediated metabolism to oxymorphone. The toxicity incidence is especially higher in those with concurrent

TABLE 1.2 Summary of Selected Literature on Impact of CYP2D6 Genotype and/or Drug Interaction on Opioid Safety

	Allelic Variants Reported	Adverse Events	References
Codeine	<i>CYP2D6</i> *1 ×3	Life-threatening opioid intoxication exacerbated by drug interaction (with erythromycin and voriconazole) and renal insufficiency	[25]
	<i>CYP2D6</i> *2A/*2×2	Fatality in a breast-fed baby whose mother is a UM	[26]
	<i>CYP2D6</i> *1 ×N	Fatality in a two-yr-old child due to respiratory arrest	[27]
	<i>CYP2D6</i> ×2	Occurrence of apnea and brain injury in a 29-mo old child	[28]
	<i>CYP2D6</i> gene duplication <i>CYP2D6</i> *1 ×N	Fatality in two children who are UMs. Respiratory depression in an <i>CYP2D6</i> EM who survived	[29]
Hydrocodone	<i>CYP2D6</i> *2A/*41	Fatality in a child who also received concurrent clarithromycin	[33]
Oxycodone	<i>CYP2D6</i> UMs	Greater toxicity, especially in those administered ketoconazole	[32]
Tramadol	Heterozygous carrier of a wild-type allele duplication	22-yr with a near-fatal case of cardiac arrest and high concentration of tramadol metabolite	[34]
	<i>CYP2D6</i> gene duplication	Tramadol-related respiratory depression	[35]

ketoconazole administration [32]. Similarly, drug interaction with clarithromycin might have played a role in the fatal case after hydrocodone exposure experienced by a 5-yr-old developmentally delayed child with a *CYP2D6**2A/*41 genotype [33]. In addition, tramadol cardiotoxicity and respiratory depression have been reported in UMs [34,35] with high level of the active O-desmethyltramadol [34], which has been reported to exhibit a high correlation with increased plasma epinephrine level [36]. The FDA also recently updated its safety warning for tramadol.

In addition to implications for ADR, the efficacy of prodrugs (such as codeine and hydrocodone) would also be reduced in PMs because less parent drug is converted by CYP2D6 to its respective active metabolite: morphine or hydromorphone, resulting in little analgesic relief [37]. However, despite strong evidence of a genotype effect on the pharmacokinetics of codeine and hydrocodone, the impact on dosage requirement is much less obvious. In this

regard, the value of CYP2D6 genotype lies more with guiding the choice of the appropriate analgesic rather than genotype-based dosage recommendation [13,38]. In particular, avoidance of codeine, the only opioid analgesic with a Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline, is recommended for PMs and UMs. In addition, hydrocodone may not be a good alternative analgesic agent to codeine in these patient populations [13].

There are similar reports of lower efficacy in PMs with venlafaxine [39]. Another example is tamoxifen, in which CYP2D6 plays a major role in the formation of the abundant and pharmacologically more active metabolite, endoxifen [40]. Because endoxifen possesses greater affinity for the estrogen receptor than tamoxifen, PMs with the *CYP2D6**4/*4 genotype have been shown to have an increased risk of breast cancer recurrence and worse relapse-free survival, as well as a much lower incidence of moderate or severe hot flashes [40]. The results of Kiyotani et al. [41] showed that the association between tamoxifen

response and CYP2D6 genotype in Japanese breast cancer patients was only evident for those patients receiving tamoxifen monotherapy, and underscores the importance of considering concomitant drug therapy in pharmacogenomics association study with tamoxifen and possibly other drugs. Conflicting data and continued debate complicate the adoption of CYP2D6 genotyping in the therapeutic use of tamoxifen currently for patients with estrogen-receptor positive breast cancer. Nevertheless, available evidence strongly supports a role for CYP2D6 in pharmacological activation of tamoxifen [42] and possibly a likelihood of lesser therapeutic benefit in PMs [43], with the ultimate impact on patient outcome to be tested in prospective clinical studies.

Inadequate therapeutic response with implications for dosage adjustment had also been demonstrated for UMs administered CYP2D6 substrates. The best evidence described two patients with multiple copies of *CYP2D6*2* requiring the tricyclic antidepressant nortriptyline 500 mg daily (vs. usual recommended daily dose of 100–150 mg) in one patient [44] and clomipramine 300 mg/day (vs. 25–150 mg) in another patient [9] to achieve adequate therapeutic response. Similarly, lower efficacy in UMs has been reported with other antidepressants [45,46] and antiemetics such as ondansetron [47].

Nevertheless, the therapeutic significance of CYP2D6 is not only impacted by genetic polymorphism but also by the potential of CYP2D6-mediated drug–drug interaction, with clinical implications in patients with different metabolic phenotypes resulting from competitive inhibition of CYP2D6. As shown by Hamelin and colleagues [48], the pharmacological consequences of drug–drug interaction via CYP2D6 inhibition are of greater magnitude in EMs, with pronounced and prolonged hemodynamic responses to metoprolol, than in PM.

Potent CYP2D6 inhibitors had been shown to reduce the metabolic capacity of EMs significantly so that individual EM could appear

metabolically as PM during concurrent administration [49] which could have therapeutic significance in patients taking multiple drugs. For example, it is not uncommon that tamoxifen-treated patients are also taking antidepressants such as selective serotonin reuptake inhibitors (SSRIs), for both their antidepressant effect as well as their offlabel use to manage hot flashes. In view of the abundance and greater antiestrogenic activity of endoxifen, concurrent administration of SSRIs that are potent inhibitors of CYP2D6 (such as fluoxetine and paroxetine) should best be avoided, and SSRIs with a lesser extent of CYP2D6 inhibition (such as citalopram and venlafaxine) would be better alternate antidepressants if there is a need for concurrent antidepressant therapy with tamoxifen.

Interestingly, Gryn et al. described significant reduction in endoxifen concentration in a patient with *CYP2D6*1/*41* genotype. The reported endoxifen concentration was described as “well below levels seen in most CYP2D6 poor metabolizers.” Although the case report did not investigate the mechanism for the altered level, the authors suggested it could be secondary to the patient’s concurrent treatment with phenytoin. Phenytoin is a potent inducer of multiple drug-metabolizing enzymes as well as the efflux drug-transporter ABCB1 (also known as multidrug resistance transporter, and described in more details in later sections) [50], which mediates the efflux transport of endoxifen [51]. Although the clinical outcome was not described, this case underscores the importance of evaluating the modulating effect of drug interaction when utilizing genotyping in individualized therapy [52]. Similar modulating effects on other genes encoding different metabolizing enzymes are described in later sections.

In addition, it is important to realize that the potential for drug interaction via CYP2D6 inhibition could also be affected by the basal metabolic activity of the individual patient. We have shown that the UM phenotype could affect the potential for drug interaction with paroxetine, a

CYP2D6 substrate as well as a potent CYP2D6 inhibitor, whence a UM with three functional CYP2D6 copies had undetectable paroxetine concentration with standard dosing and showed no inhibitory effect at CYP2D6 [53].

CYP2C19

CYP2C19 is located on chromosome 10q23.33 and is a large gene consisting of 90,209 nucleotides and yet coding for CYP2C19 that contains only 490 amino acids. Compared to the CYP2D6 polymorphism, polymorphisms in the CYP2C19 gene do not affect as many drugs, and their clinical implication has not been extensively evaluated. However, studies involving the proton pump inhibitors (PPIs) provide extensive pharmacokinetic and clinical evidence, as well as the economic impact of the importance of taking into consideration of CYP2C19 polymorphism in the management of gastroesophageal diseases.

Over the years, as many as 30 CYP2C19 alleles, including those with no functional activity (*2, *3, *4, *6, *7) and those associated with reduced catalytic activity (*5 and *8), have been identified (www.cypalleles.ki.se/cyp2c19.htm). The principal null alleles are *2 (c.681G>A, rs4244285) and *3 (c.636 G>A, rs4986893), resulting in an inactive CYP2C19 enzyme, and accounting for the vast majority of the PM phenotype in Caucasians (1%–6%), black Africans (1%–7.5%), and Asians (10%–25%). Genotyping these two defective alleles has been shown to detect about 84%, greater than 90%, and about 100%, of PMs in Caucasians, Africans, and Asians, respectively. The detection rate for Caucasians PMs could be increased to about 92% by including the less common CYP2C19*4 (rs28399504) and CYP2C19*6 (rs72552267) in the genotyping assay. Similar to other CYP2C19 rare variants, *5 (rs56337013), *7 (rs72558186), and *8 (rs41291556) have <1% allele frequency. Individuals carrying at least one functional allele are referred as EMs, whereas those with one functional and one loss-of-function allele are IMs. Of interest

is that, similar to CYP2D6 polymorphism, a “gain of function” CYP2C19*17 allele (c.-806C>T rs12248560) was identified in the 5'-flanking region of CYP2C19, with increased gene transcription associated with high enzyme activity and an EM phenotype [54].

CYP2C19*2 and *3 are commonly found in Asians, with allele frequencies of about 30% and approximately 10%, respectively. In contrast, the allele frequency of *3 is <1% in Caucasians and African Americans, even though the *2 occurs at a frequency of about 13% and approximately 18%, respectively, in these two ethnic groups. About 50% of the Chinese population possess either the *1/*2 and *1/*3 genotypes, and 24% have the *2/*2, *2/*3, or *3/*3 genotypes [55]. In contrast, only about 2%–5% and 30%–40% of the Caucasian population, respectively, have the *2/*2 and *1/*2 genotypes. Similar frequencies of the heterozygous and homozygous variant genotypes are reported in persons of African descent. The higher prevalence of PMs and heterozygote EMs carrying defective CYP2C19 alleles in Asians likely account for reports of slower rates of metabolism of CYP2C19 substrates and the practice of prescribing lower diazepam dosages for patients of Chinese heritage [56,57]. An opposite direction in ethnic variation was observed in the prevalence of CYP2C19*17 (18% in Swedes and Ethiopians vs. 4% in Chinese), with the *1/*17 and *17/*17 genotypes occurring in more Caucasians and Ethiopians (up to 36%) than Asians (8% of Chinese and 1% of Japanese) [54].

CYP2C19 accounts for about 3% of total hepatic CYP content, and CYP2C19 polymorphism affects the metabolism of PPIs (omeprazole, lansoprazole, pantoprazole, rabeprazole), antidepressants (citalopram, sertraline, moclobemide, amitriptyline, clomipramine), the antiplatelet agent clopidogrel, the antifungal drug voriconazole, the benzodiazepine diazepam, and the anticancer drug cyclophosphamide. Similar to CYP2D6, CYP2C19 is also susceptible to inhibition by drugs such as cimetidine, fluoxetine, and diazepam. The inhibition occurs in a gene

dose-dependent manner in which carriers of two *CYP2C19**17 alleles exhibit the greatest extent of inhibition compared to little to no inhibition for patients with *CYP2C19* PM phenotype.

The PPIs and clopidogrel provide the best examples of clinical relevance of *CYP2C19* polymorphism. When compared to EMs, PMs showed 5- to 12-fold increases in the area under the curve (AUC) of omeprazole, lansoprazole, and pantoprazole [58,59], whereas homozygous carriers of the *CYP2C19**17 were shown to have a modest 2.1-fold lower AUC than EMs [60]. In addition, the *CYP2C19* genotype significantly affects the achievable intragastric pH with PPI therapy. In subjects who took a single 20-mg dose of omeprazole, Furuta et al. showed a good relationship not only between *CYP2C19* genotype and AUC, but also between the genotype and achievable intragastric pH: 4.5 in PMs, 3.3 in heterozygous EMs, and 2.1 in homozygous EMs [61]. Given the smaller dependency of esomeprazole and rabeprazole on *CYP2C19* for metabolism, the pharmacological action of these two PPIs is less affected by the *CYP2C19* polymorphism [62,63].

An important treatment strategy in the management of patients with peptic ulcer disease is eradication of *Helicobacter pylori* with a regimen of PPI and antibiotics. *CYP2C19* genotype-related pharmacological effects have also been associated with improved eradication rate of *H. pylori* after dual [64] or triple therapy including omeprazole [65], lansoprazole [66], or pantoprazole [67]. The cure rate achieved with dual- and triple-therapy regimens was 100% in PMs compared with 29%–84% in EMs [64–67]. Furuta et al. also reported a much higher eradication rate of 97% in EMs who failed initial triple therapy (lansoprazole, clarithromycin, and amoxicillin) and subsequently were retreated with high-dose lansoprazole (30 mg four times daily) and amoxicillin [68]. In addition, to showing a gene-dose effect in achieving desirable ranges of intragastric pH and *H. pylori* cure rates for lansoprazole, Furuta et al. also demonstrated the cost

effectiveness of pharmacogenomics-guided dosing when compared to conventional dosing [69]. On the other hand, despite increased metabolism of PPI in carriers of *CYP2C19**17 and the potential of therapeutic failure [54,70], eradication rates of *H. pylori* have so far not to be shown to be associated with the *CYP2C19**17 allele, at least for patients with peptic ulcer disease and receiving the triple regimen of pantoprazole, amoxicillin, and metronidazole [67,71].

In healthy volunteers given a single 200-mg dose of voriconazole, Wang et al. demonstrated a 48% lower AUC in heterozygous carriers of the *CYP2C19**17 allele as compared to homozygous carriers of *CYP2C19**1 [72]. This finding is consistent with data that is more recent showing correlation between *CYP2C19* polymorphism and target voriconazole concentrations, with an increased risk of subtherapeutic trough concentration in patients with the *CYP2C19* UM phenotype [73–75]. Investigators have also shown 42% lower escitalopram concentrations and 21% lower AUC in patients who are homozygous carriers of *CYP2C19**17 when compared to *CYP2C19**1 homozygotes [76]. Clearly, *CYP2C19**17 homozygotes might require higher doses of most *CYP2C19* substrates, including PPIs [60,70], antidepressants, and voriconazole [72]. However, despite the presence of pharmacokinetic differences, the impact of *CYP2C19**17 on therapeutic outcomes with these *CYP2C19* substrates have not been evaluated extensively or confirmed.

Clopidogrel is an antiplatelet prodrug that requires *CYP2C19*-mediated conversion to its active metabolite for therapeutic effect [77], with most of pharmacokinetic and pharmacodynamic evidence related to the *CYP2C19**2 allele [77–82]. Shuldiner et al. conducted a GWAS in which *ex vivo* adenosine diphosphate (ADP)-induced platelet aggregation at baseline and after 7 days of clopidogrel were measured in a genetically homogenous cohort of 429 healthy Amish subjects. In addition, 400,230 SNPs were evaluated in each subject for association with

platelet activity. They reported that the SNP rs12777823 on chromosome 10q24 with the greatest association signal is in strong LD with *CYP2C19**2, accounting for 12% of the interindividual variation in platelet aggregation during clopidogrel treatment. As importantly, there was no association between the *CYP2C19* polymorphism and baseline platelet aggregation [83]. The results from this GWAS confirmed results from previous candidate gene studies regarding the role of *CYP2C19* as a major genetic determinant of clopidogrel response [78–82]. In a follow-up study of 227 patients undergoing percutaneous coronary intervention (PCI), the investigators also reported a higher incidence of cardiovascular death in carriers of the *2 allele (20.9% vs. 10%) at 1-yr follow-up. No association with response was found for other *CYP2C19* alleles, including *3, *5 (rs56337013), and *17, that were also genotyped in the study [83]. A recent meta-analysis confirms the association of the *CYP2C19* nonfunctional allele and high-risk of adverse cardiovascular events in patients who underwent PCI [84].

Although the increased production of the active clopidogrel metabolite in carriers of the *17 allele has been associated with greater inhibition of platelet aggregation [85,86] and better clinical outcomes [87], there is also the potential of increased bleeding risk [88]. In addition, the increased response of the *17 allele has been suggested not as a direct effect, but rather attributed to that of the *1 allele [89]. Given this consideration, there is no specific therapeutic recommendation for this gain-of-function allele in the most recent practice guideline for *CYP2C19* genotyping [90].

Even with involvement of other non-genetic factors [91], the increased risks of major adverse cardiovascular events and stent thrombosis in carriers of at least one *CYP2C19**2 allele were confirmed in two meta-analyses that included almost 22,000 patients [88,92]. Differences in patient selection for analysis likely account for the lack of association reported in two other recent meta-analyses, which included a

significant number of low-risk patients, such as those with acute coronary syndrome managed medically or patients with atrial fibrillation [93,94]. The meta-analysis of Hulot et al. [92] also evaluated the drug interaction potential of PPIs because of their inhibitory effect toward *CYP2C19*, resulting in a metabolic phenotype of *CYP2C19* PM similar to that of carriers of the *2 allele. Both Hulot et al. and another study [92,95] suggest that the detrimental effects of PPIs on cardiovascular outcomes with clopidogrel likely occur at a higher frequency in high-risk patients receiving both drugs. Current data do not provide sufficient information to determine whether the observed adverse effects of PPI usage in high-risk patients (e.g., patients undergoing PCI) are related to *CYP2C19* inhibition or yet-to-be-discovered mechanisms.

Based on the increasing amount of literature data supporting an association between *CYP2C19**2 and poor clopidogrel response, the FDA has made several revisions to the approved product label of clopidogrel. Although the March 2010 version specifically addresses the implication for homozygotes, there is no guidance on the implication for heterozygotes. In addition, as with other revised labels with additional genetic information, there is little guidance on clinical management of carriers of *CYP2C19**2. The September 2016 label warns of diminished effectiveness in *CYP2C19* poor metabolizers and suggests the use of different platelet P2Y₁₂ inhibitors in those patients. In light of the scientific and clinical evidences as well as the regulatory decision, several recent clinical studies addressing alternative antiplatelet agents have been initiated and are discussed in Chapter 6.

CYP2C9

In addition to *CYP2C19*, another important member of the *CYP2C* subfamily of enzymes is *CYP2C9* containing 490 amino acids. It is encoded by *CYP2C9* consisting of 50,708 nucleotides and located on chromosome 10q24.1 in close

proximity to *CYP2C19*. To date, approximately 60 *CYP2C9* alleles (www.cypalleles.ki.se/cyp2c9.htm) have been identified in the regulatory and coding regions of *CYP2C9*, with *CYP2C9**2 (c.430C>T, rs1799853) and *CYP2C9**3 (c.1075A>C, rs1057910) being the most common in persons of European descent and the most extensively studied. Both reduced-function alleles exhibit single amino-acid substitutions (p.R144C and p.I359L, respectively) in the coding region, accounting for lower enzyme activity by approximately 30% for *2 and 80% for *3 [96]. Other reduced-function alleles of potential importance included *5 (rs28371686), *6 (rs9332131), *8 (rs7900194), and *11 (rs28371685). [97–100] In addition, a “gain-of-function” *CYP2C9* (rs7089580) variant in intron 3 has been identified [97].

Significant variations in *CYP2C9* alleles and genotype frequencies exist among different ancestry groups. Both *CYP2C9**2 and *CYP2C9**3 are more common in Caucasians (11% and 7%, respectively) than in Asians and Africans. In fact, *CYP2C9**2 has not been detected in Asians, in whom *CYP2C9**3 is the most common allele. On the other hand, *CYP2C9**8, as well as *5, *6, and *11 (albeit all at a lower frequency than 8), are present almost exclusively in African Americans. The novel *CYP2C9* c.18786A>T variant (rs7089580) was reported to occur in about 40% of the African American population, and *CYP2C9**8 (c.449G>A, rs7900194) appears to be a major contributor to *CYP2C9* expression in this ethnic group [97]. Approximately 1% and 0.4% of Caucasians have the *2/*2 and *3/*3 genotypes, respectively. The *1/*3 genotype occurs at a frequency of 4% in the Chinese and Japanese populations, with almost complete absence of the other genotypes (*2/*2, *2/*3, *1/*2, and *3/*3).

CYP2C9 accounts for about 20% of total hepatic CYP content and is involved in the metabolism of about 10% of currently marketed drugs. These *CYP2C9* substrates include the nonsteroidal antiinflammatory drugs such as celecoxib, ibuprofen, and flurbiprofen; oral anticoagulants such as acenocoumarol, and phenprocoumon,

and the S-isomer of warfarin; oral antidiabetic agents such as glibenclamide, glimepiride, glipizide, glyburide and tolbutamide; antiepileptic agents such as phenytoin, and antihypertensive agents such as candesartan, irbesartan, and losartan. The enzyme reduction associated with the *3 allele is greater than that with the *2 allele, with a 5- to 10-fold reduction in homozygous *3 carriers and two-fold reduction in heterozygous *3 carriers, when compared to homozygous *1 carriers. For example, clearance of warfarin is reduced by 90%, 75%, and 40% in subjects with the corresponding *CYP2C9* genotypes of *3/*3, *1/*3, and *1/*2 [101]. respectively. Interestingly, the effects of several reduced-function alleles appear to be substrate dependent. For the *2 allele, a significant effect was shown for clearances of acenocoumarol, tolbutamide, and warfarin but not for other substrates. On the other hand, nonsteroidal anti-inflammatory drug (NSAID)-associated gastrointestinal bleeding was shown to be related to the *3 but not the *2 variant [102]. Similarly, although the *8 allele has no effect on clearance of losartan, it decreases enzyme activity of warfarin and phenytoin, and exhibits an increased activity toward tolbutamide [103].

Of all of the *CYP2C9* substrates, warfarin is the most extensively studied with dosing implications for different metabolic phenotypes. *CYP2C9* polymorphism, together with the literature information regarding the gene that encodes the warfarin target, vitamin K epoxide reductase complex (VKORC1) [104], provide promising translational use of the pharmacogenomic data [105,106], with revised language regarding their impact incorporated into the drug label [107]. *CYP2C9* mediates the conversion of the active S-enantiomer of warfarin to an inactive metabolite. Most of the data document that the *2 and *3 alleles are associated with greater difficulty with warfarin induction therapy, increased time to achieve stable dosing, lower mean-dose requirement (e.g., as low as ≤ 1.5 mg/day with *3/*3), as well as increased risks of elevated, international normalized ratios (INRs) and bleeding [105,108,109]. Giving the 30%

and 80% difference in enzyme activity reduction between the *2 and *3 alleles, the warfarin-dose requirements differ between carriers of these two alleles. Compared to homozygous carriers of the *1 allele, data suggest a dose reduction of 30% and 47% for patients with the heterozygous genotypes of *CYP2C9**1/*2 and *CYP2C9**1/*3, respectively, and up to 80% for patients with the homozygous *CYP2C9**3/*3 genotype [106,108,110,111].

In addition, with the difference in allele prevalence among different ancestral groups, the strength of association between the *2 and *3 alleles and genotypes is stronger in Caucasians [112,113]. Other recently identified alleles (*5, *6, *8, and *11) have been reported to better predict dose requirement (20% lower for *8 carrier) and adverse outcomes in African Americans [97–99,103,112,114]. On the other hand, the “gain-of-function” *CYP2C9* c.18786A>T allele was reported to contribute a higher-dose requirement (3.7 mg/week/allele) [97]. Finally, concurrent drugs with significant modulating effect on *CYP2C9* activity would also have an impact on the association between *CYP2C9* genotypes and warfarin-dose requirement [115]. The effect of *CYP4F2* and *VKORC1* genotypes on warfarin pharmacokinetics and pharmacodynamics will be discussed in later sections of this chapter.

CYP2C8

In addition to *CYP2C9* and *CYP2C19*, the other clinically relevant members of the highly homologous genes (*CYP2C18*–*CYP2C19*–*CYP2C9*–*CYP2C8*) that cluster on chromosome 10q24 [83] is *CYP2C8*. To date, several SNPs within the coding region of the *CYP2C8* gene have been identified (www.cypalleles.ki.se/cyp2c8.htm). The more common variants are *2 (c.805A>T, rs11572103, resulting in p.I269F), *3 with two amino acid substitutions (c.416G>A, rs11572080 with p.R139K, and c.1196A>G, rs10509681 with p.K399R) reportedly to be in total LD, and *4 (c.792C>G, rs1058930, p.I264M). Both *3 and *4 alleles are more common in

Caucasians (with the *4 variant reportedly only found in Caucasians). On the other hand, *2 and a rare allele, *5 (rs72558196, frame-shift deletion) are exclusively found in Africans and Japanese, respectively [116,117].

Accounting for about 7% of total hepatic content, the hepatic expression level of *CYP2C8* lies between that of *CYP2C19* and *CYP2C9* [118], and it plays an important role in the metabolism of different drugs, primarily the antidiabetic agents (pioglitazone, repaglinide, rosiglitazone, and troglitazone), the anticancer agents (paclitaxel), the antiarrhythmic drug amiodarone, and the anti-malarial agents amodiaquine and chloroquine. The smaller number of substrates as compared to *CYP2C9* and *CYP2C19* presumably leads to the lesser interest in studying *CYP2C8* polymorphism. As a result, the molecular mechanisms underlying interindividual variations in *CYP2C8* activity remain unclear. Decreased elimination of R-ibuprofen has been reported in carriers of *CYP2C8**3 [119,120]. However, with the presence of a strong LD between *CYP2C8**3 and *CYP2C9**2 [119,121], the individual contribution of *CYP2C8**3 remains to be elucidated. In contrast, increased metabolism of repaglinide was reported in heterozygous carriers of *CYP2C8**3 when compared to carriers of either *1 or *4 [122]. Although this finding is interesting, other reports showed that genetic polymorphism of the hepatic uptake transporter plays a more important role in determining repaglinide pharmacokinetics [123]. The identification of two *CYP2C8* haplotypes: a high-activity allele associated with *CYP2C8**1B and a low activity associated with *CYP2C8**4 [124], further highlights the need to characterize the different *CYP2C8* variants, including their functional relevance.

CYP3A4/5/7

A total of four *CYP3A* genes have been described in humans: *CYP3A4*, *CYP3A5*, *CYP3A7*, and *CYP3A43*; with *CYP3A7* primarily important in fetal *CYP3A* metabolism and

CYP3A4 exhibiting little functional or clinical relevance. More than 20 variants in the coding region of *CYP3A4*, most of them associated with reduced catalytic activity of the enzyme, have been identified to date. [125] The significance of the reduced-function allele *CYP3A4**22 C>T SNP (rs35599367) in intron 6, which results in 20% decrease in enzyme activity, has been extensively evaluated recently, especially in conjunction with *CYP3A5* SNP [126–129]. *CYP3A5* expression is highly polymorphic with the loss-of-function *3 allele (c.6986A>G, rs776746) in intron 3 as the most common variant, which results in a splicing defect and absence of enzyme activity. Other loss-of-function and reduced-function *CYP3A5* variants include the *2 (rs28365083; g.27289C>A; T398N), *6 (14690G>A; rs10264272), and *7 (rs41303343; 27131_27132ins T) alleles [130,131].

In general, *CYP3A4* polymorphism is more common in Caucasians, with *2 and *7 being the more prevalent alleles, whereas Asians have higher frequencies of *16 and *18 variants. Of note, is that *CYP3A4**22 is absent in both Asian and African populations. Carriers of the wild-type *CYP3A5**1 allele (also known as *CYP3A5* “expressors”) are more common in Asians (up to 50%) and Blacks (up to 90%) than in Caucasians (about 15%). The allele frequency of *CYP3A5**3 is much higher in Caucasians and Asians, occurring in 90% and 75% of the populations, respectively, *versus* a relatively low frequency of 20% in Africans. On the other hand, both *CYP3A5**6 and *7 are absent in Caucasians and Asians but present in Africans with frequencies up to 17% [130,132,133]. The *CYP3A5**2 allele has a frequency of less than 1% in Caucasians and is mostly absent in other ethnic populations.

CYP3A4 accounts for about 40% of the total hepatic CYP content and mediates the metabolism of more than 50% of currently used drugs with many examples from the pharmacological classes of macrolide antibiotics, antidepressants, antipsychotics, anxiolytics, calcium channel blockers, immunosuppressants, opiates, and the statins. The

current consensus is that *CYP3A4* polymorphisms are mostly of minor clinical relevance, and unlikely responsible for the 10- to 40-fold interindividual variations in *CYP3A4* activities. This is likely a result of low variant allele frequencies, only small changes in enzyme activity in the presence of a variant allele, as well as the overlapping substrate specificity between *CYP3A4* and *CYP3A5*. The significant variability in *CYP3A4* activity is more likely related to a large number of drugs capable of altering the enzyme through induction or inhibition in the liver and the gastrointestinal tract. Therefore, there is currently no uniform agreement on metabolizer subgroups for *CYP3A4*.

On the other hand, the clinical relevance of *CYP3A* genetic polymorphism is primarily associated with *CYP3A5*. The pharmacokinetics of the immunosuppressive agent tacrolimus is dependent on the *CYP3A5* genotype, with a higher-dosage requirement in homozygous or heterozygous carriers of *CYP3A5**1 [134,135]. In addition, results from a randomized controlled trial showed that pharmacogenetic-guided dosing based on *CYP3A5* genotype was associated with greater achievement of target tacrolimus concentrations when compared to standard dosing based on body weight [136]. Nevertheless, the overall clinical relevance of *CYP3A5* polymorphism is limited by its small contribution (2%–3%) to the total *CYP3A* metabolism. [137,138], and reportedly impacted by timing of tacrolimus therapy. In a meta-analysis of tacrolimus-dose requirement and rejection rate, Tang et al. indicated that the effect of *CYP3A5* polymorphism (*CYP3A5**3) is most prominent during the first month of tacrolimus therapy, suggesting that *CYP3A5* genotyping might be useful to guide initial dosing of tacrolimus for prevention of early graft rejection [139]. Inclusion of both *CYP3A4**22 and *CYP3A5**3 status have been shown in many recent studies to significantly improve tacrolimus dose prediction [126–129,140]. Therapeutic and pharmacogenomic recommendations for tacrolimus were included in the recent CPIC guideline [141].

On the other hand, despite significant effect of *CYP3A4**1G (g.20230G>A, rs2242480) and *CYP3A5**3 on ticagrelor pharmacokinetics in a recent study of healthy Chinese subjects, there was no association on the extent of inhibition of platelet aggregation. Therefore, the investigators concluded that no dosage adjustment based on *CYP3A4* and *CYP3A5* genotypes is necessary [142].

CYP4F2

There are six members within the *CYP4F* gene subfamily residing on chromosome 19p13.1-2: *CYP4F2*, *CYP4F3*, *CYP4F8*, *CYP4F11*, *CYP4F12*, and *CYP4F22*. The importance of *CYP4F2*, a vitamin-K oxidase, is related to the recent report of its role in mediating the conversion of vitamin K₁ to hydroxyvitamin K₁. Increased *CYP4F2* activity causes decreased activation of vitamin K-dependent clotting factors, reflecting the consequence of reduced availability and reduction of vitamin K₁ to vitamin KH₂ necessary for carboxylation and activation of the clotting factors. On the other hand, the g.7253233C>T (rs2108622, p.V433M) SNP in exon 2 of the *CYP4F2* gene results in lower protein expression and enzyme activity, and consequently greater vitamin K₁ availability [143,144]. The T allele at rs2108622 confers the *CYP4F2**3 designation. Some ethnic differences in the V433M SNP has been reported, with the M433 allele occurring at a much lower frequency in African Americans [114], which contrast with its high occurrence in Indonesians and Egyptians [145,146].

Although genome-wide association studies enable detection of weaker genetic signals [144], *CYP4F2* genotype nevertheless only accounts for 1%–3% of the overall variability of warfarin-dose requirement [144,147], in contrast to *CYP2C9* genotype that accounts for approximately 10%–12% of the variability. Homozygous carriers of the M allele of the p.V433M SNP had been shown to require an approximate 1 mg/day higher dose of warfarin than homozygous

carriers of the V allele [148]. However, additional studies demonstrated the association between the *CYP4F2* genotypes and dose requirements in Caucasians and Asians [144,147,149,150] but not in African Americans, Egyptians, or Indonesians [114,145,146]. This could reflect ethnic differences in *CYP4F2* allele and genotype frequencies distribution, the minor contribution of *CYP4F2* [151], as well as the modulating effects of other more important dose-requirement variables such as *CYP2C9* and *VKORC1*.

CYP2B6

Although several variant alleles with low enzyme expression, including *CYP2B6**6 and *18, have been identified, to date there have not been any reports of the presence of an important loss-of-function allele. Among the variant alleles, the *CYP2B6**6 haplotype carrying two nonsynonymous SNPs (c.516G>T, rs3745274 and c.785A>G, rs2279343 causing two amino acid changes: p.Q172H and p.K262R, respectively) in exon 4 is the most common and occurs commonly in Caucasians and Asians (16%–26% allele frequency), whereas *18 (c.983T>C, rs28399499, I328T) is more common in Black subjects with allele frequencies of 7%–9% [152]. Interestingly, the 785A>G SNP resulting in the K262R amino acid change also occurs as a separate allele, *CYP2B6**4 (rs2279343 without rs3745274), and results in increased expression and enzyme activity [153,154]. Whether the 516G>T and 785A>G mutations are linked to additional mutations creating specific haplotypes causing either high or low *CYP2B6* activities is not known. Gatanaga et al. also reported a new *26 allele containing 499G for the c.499C>G SNP (rs3826711), and 499G always coexists with 516G>T and 785A>G, thus representing a novel haplotype containing the 499C>G, 516G>T and 785A>G SNPs [155].

CYP2B6 accounts for up to 6%–10% of total CYP content in the liver [156,157], and known substrates include anticancer drugs such as

cyclophosphamide and ifosfamide, the smoking cessation agent bupropion, the antiretroviral agents efavirenz and nevirapine, as well as methadone. In addition to reduced activation of cyclophosphamide leading to lower antitumor efficacy, *CYP2B6* gene variants play a significant role in determining bupropion and methadone pharmacokinetic variabilities, in particular with *CYP2B6**6 (decreased clearance) [158–160] and *CYP2B6**4 (increased clearance) [159]. Levran et al. reported that the mean daily methadone dose in heroin addicts was 88 and 96 mg, respectively, for homozygous carriers of variant alleles 785A>G and 516G>T; as compared to 133 and 129 mg, respectively, for heterozygous carriers of the two variant alleles; and 150 and 151 mg, respectively, for wild-type homozygotes [161]. In individuals whose death was attributed to methadone poisoning, *CYP2B6**4, *6, and *9 alleles were associated with higher postmortem methadone blood concentrations ($P \leq .05$) [162]. However, despite report of longer corrected QT interval (QTc) interval in *CYP2B6* slow metabolizers [163], a clear relationship between *CYP2B6* genotype and risk of cardiac arrhythmia and sudden death remains to be determined.

The potential clinical relevance of *CYP2B6* has been evaluated primarily with the nonnucleoside reverse transcriptase inhibitors efavirenz and nevirapine. Increased central nervous system side effects associated with variable systemic exposure of efavirenz could be the result of patients being carriers of the *6 or *18 alleles [155,164]. Incorporating determination of additional less-frequent alleles such as *26 and *29 could further improve the prediction of elevated plasma efavirenz concentrations [155,164,165]. Altered concentrations of, and clinical outcome associated with, nevirapine have also been associated with *CYP2B6* rs3745274 SNP.

In a prospective study of the effect of *CYP2B6* polymorphism on efavirenz concentrations and exposure, 456 patients infected with the human immunodeficiency virus type 1 (HIV-1) were

genotyped for different SNPs, including the 499C>G, 15631G>T and 18053A>G polymorphisms [155]. All patients received the standard-dosage regimen of 600 mg/day, and extremely high concentrations ($9,500 \pm 2,580$ ng/mL) were obtained in all 14 patients with the *CYP2B6**6/*6 genotype and in both patients with the *CYP2B6**6/*26 genotype. In contrast, only two patients with other *CYP2B6* genotypes had similarly high efavirenz concentrations, and both were heterozygous carrier of either the *6 allele (7,140 ng/mL) or *26 allele (9,710 ng/mL). Therefore, the *6 and *26 alleles were both associated with high efavirenz concentrations, and patients with the *CYP2B6**6/*6 or the *CYP2B6**6/*26 genotype had the highest concentrations with standard-dosage regimen of 600 mg/day.

To investigate the feasibility of dose reduction in patients with high efavirenz concentrations secondary to *CYP2B6* polymorphism, the investigators then reduced the efavirenz-dosage regimen to 400-mg/day in five patients and to 200 mg/day in another seven patients. The genotypes in these 12 patients included nine *6/*6 homozygotes, two *6/*26 heterozygotes, and one *1/*26 heterozygote. The plasma concentrations decreased proportionally with the dose reductions. Despite receiving the lower-dosage regimens for more than 6 months, the 12 patients were able to maintain therapeutically effective anti-HIV-1 activity with HIV-1 load continuously less than 50 copies/mL. Central nervous system side effects were reported to be much less frequent at the lower-dosage regimens. Similar therapeutic success with persistent suppressed HIV-1 load was also demonstrated in efavirenz-naïve patients (*6/*6 and *6/*26), who were administered the lower-dosage regimen of 400-mg/day. The overall study results demonstrated the feasibility of genotype-based efavirenz-dose reduction in patients with *CYP2B6* *6/*6 and *6/*26 genotypes, with additional advantages of less central nervous system side effects and lower treatment cost.

CYP2A6

CYP2A6 only accounts for about 4% of total CYP450 content, and significant variations in CYP2A6 activity are primarily a result of genetic influence. The *CYP2A6* gene is located on chromosome 19 and codes for the protein CYP2A6 consisting of 494 amino acids. With more than 40 variants identified, the primary variants for CYP2A6 polymorphism (www.cypalleles.ki.se) include CYP2A6*2 (rs1801272, g.1799T>A), CYP2A6*4 (gene deletion), CYP2A6*5 (rs5031017 g.6582G>T), and CYP2A6*20 (rs28399444, frame shift), all of which are associated with abolished enzyme activity. Additional alleles associated with reduced enzyme activity include *7 (rs5031016, g.6558 T>C), *10 (rs28399468, g.6600G>T) *11 (rs111033610, g.3391T>C), *17 (rs28399454, g.5065G>A), *18 (rs1809810 g.5668A>T), and *19 (rs5031016 g.6558T>C). As with other CYP polymorphisms, there are substantial inter-ethnic differences in allele frequency. Deletion of the *CYP2A6* gene is very common in Asian patients [166], which likely accounts for the dramatic difference in the high occurrence of PMs in Asian (20%) versus Caucasian populations ($\leq 1\%$).

Nicotine is metabolized by CYP2A6 to cotinine, and the clinical relevance of the *CYP2A6* polymorphism has been primarily investigated in managing patients with tobacco abuse. Nonsmokers were found to be more likely to carry defective *CYP2A6* alleles such as *7 and *9 than were smokers. In addition, smokers with defective *CYP2A6* alleles smoked fewer cigarettes and were more likely to quit. These results likely reflect higher nicotine concentrations, enhanced nicotine tolerance and increased adverse effects from nicotine in CYP2A6 poor metabolizers. Based on these observations, CYP2A6 inhibition may have a role in the management of tobacco dependency [166].

CYP1A2

Located on chromosome 15, *CYP1A2* consists of 7,758 nucleotides and encodes the enzyme CYP1A2 that contains 516 amino acids. Polymorphisms of the CYP1 family of genes have been studied for association with cancer susceptibility. Several *CYP1A2* SNPs have been identified, including *CYP1A2**1C (rs2069514, -3860G>A) and the haplotype *CYP1A2**1K containing three variants: -739T>G (rs2069526), -729C>T (rs12720461), and -163C>A (rs762551). However, to date there has been no consistent report of any functional *CYP1A2* alleles that result in important changes in gene expression and enzyme activity. Therefore, in contrast to other CYP isoenzymes such as CYP2C19 and CYP2C9, there is less agreement in the literature regarding acceptable method of defining *CYP1A2* metabolic phenotype by *CYP1A2* genotype.

Nevertheless, a unique aspect of the *CYP1A2* gene is that a specific allele, *CYP1A2**1F (rs762551) containing a c.-163C>A mutation in intron 1, has been shown to affect CYP1A2 inducibility [167] and the magnitude of increased caffeine metabolism in smokers [168,169]. However, conflicting reports have been reported for other *CYP1A2* substrates [170–172]. This gene–environment interaction makes genotype–phenotype prediction of phenotype much more difficult. Finally, promoter variation is less likely to result in substrate-dependent effects, and the functional importance of increased CYP1A2 inducibility is currently unknown.

CYP1A2 contributes up to 10% of the total hepatic P450 content. However, unlike other CYP isoenzymes, it only mediates the metabolism of several commonly used drugs such as olanzapine, clozapine, duloxetine, and theophylline [173,174]. Although pharmacokinetic studies evaluating CYP1A2 inducibility by smoking or omeprazole have been performed, none of the studies have produced consensus information.

POLYMORPHISMS IN NON-CYP450 DRUG-METABOLIZING ENZYMES

Genetic polymorphisms in many non-P450 enzymes also play a role in influencing metabolism and elimination of many drugs. Among these enzymes, UDP-glucuronosyl transferase (UGT), thiopurine-S-methyltransferase (TPMT), dihydropyrimidine dehydrogenase (DPD), N-acetyltransferase (NAT), and glutathione-S-transferase (GST) have been characterized and their clinical relevance studied.

UDP-Glucuronosyl Transferase

The uridine diphosphate (UDP)-glucuronosyl transferase (UGT) enzymes are divided into two distinct subfamilies: UGT1 and UGT2. UGT1A1 has been the most extensively investigated among the UGT1A enzymes. A polymorphism with an extra thymine-adenine (TA) repeat (TA insertion) in the 5'-promoter region of the *UGT1A1* gene results in the (TA)⁷TAA allele or *UGT1A1**28 (rs8175347), with a 35% decrease in transcriptional activity of *UGT1A1* and lower enzyme activity than the wild-type (TA)⁶TAA allele (*UGT1A1**1) [175]. Another UGT1A1 polymorphism, *UGT1A1**6 (rs4148323) carrying the c.211G>A SNP and p.G71R substitution in exon 1, has also been associated with lower enzymatic activity [176]. Although the *28 variant is more common in Caucasians (29%–40%) and Africans (36%–43%) than in Asians (13%–16%), the *6 is found only in Asians with a frequency of 16%–23%.

UGT contributes about 35% of phase II drug metabolism and is involved in glucuronidation of endogenous compounds and xenobiotics. For UGT1A1, the substrates include bilirubin, SN-38 (active and toxic metabolite of the anticancer drug irinotecan), raltegravir (inhibitor of the HIV integrase enzyme), clozapine, bazedoxifene (an investigational selective estrogen receptor modulator for prevention and treatment of postmenopausal osteoporosis), and eltrombopag (a

thrombopoietin receptor agonist for the management of thrombocytopenia).

Irinotecan, also known as CPT-11, is a pro-drug that requires metabolic activation via carboxylesterase to SN-38, a potent inhibitor of topoisomerase I. SN-38 is inactivated via glucuronidation by the polymorphic *UGT1A1* enzyme. Both *UGT1A1**28 and *6 had been associated with impaired SN-38 glucuronidation, especially in patients who are homozygous carriers (*UGT1A1**28 TA7/TA7) [176,177]. The ensuing high SN-38 concentrations lead to increased SN-38 excretion into the gut lumen, predisposing patients to severe diarrhea even with standard irinotecan-dosage regimens. Abnormally high SN-38 concentrations have also been reported in patients with severe neutropenia [178]. These pharmacogenetic-related adverse reactions have also been demonstrated in prospective clinical trial [179] that led to FDA approval of the Invader UGT1A1 Molecular Assay (Third Wave Technologies) for genotyping *UGT1A1* alleles and revision of the irinotecan product label to include consideration of lower initial dose requirement for individuals who are homozygous for *UGT1A1**28, although the genetic testing is not a requirement. The predictive value of *UGT1A1**28 polymorphism has recently been confirmed in a meta-analysis of 58 studies [180]. With the involvement of UGT1A1 in bilirubin glucuronidation confirmed by three meta-analyses [181–183] and the prevalence of *UGT1A1**6 among Asian populations, UGT1A1 may play a role in the high incidence of neonatal hyperbilirubinemia in those populations [184].

A different UGT enzyme, UGT2B7, plays an important role in mediating the conversion of morphine to the pharmacologically active metabolite, morphine-6-glucuronide. Darbari et al. reported that homozygous and heterozygous carriers of the G variant for the -840 G>A SNP (rs7438135) had significant higher parent to metabolite concentration ratio compared to individuals with the A/A genotype ($P = .03$) [185]. A second *UGT2B7* SNP (rs7439366, C802T) was

implicated for morphine toxicity in a patient with the *T/T* genotype that resulted in increased morphine-6-glucuronide formation [186].

Thiopurine-S-methyltransferase

Thiopurine-S-methyltransferase (TPMT) is encoded by the *TPMT* gene that has a nonsynonymous SNP and resulting in reduced TPMT enzymatic activity. Although more than 31 variants of the *TPMT* gene have been identified, the five most studied one are *TPMT**2 (rs1800462, G238>C, reduced activity), *TPMT**3A (a haplotype consisting of the two nonsynonymous SNPs, G460>A and A719>G, abolished activity), *TPMT**3B (rs1800460, G460>A, reduced activity) *TPMT**3C (rs1142345, A719>G, reduced activity), and *TPMT**4 (rs1800584, G626>A, very low activity). About 95% of intermediate or low TPMT activity in affected patients are associated with *TPMT**2, *TPMT**3A, or *TPMT**3C. Approximately 10% and 0.3% of the Caucasian population is heterozygous and homozygous, respectively, of these mutant alleles.

TPMT mediates the inactivation of thiopurine drugs, including thioguanine, 6-mercaptopurine, and its precursor, azathioprine. Compared to patients who possess the wild-type alleles, homozygotes or heterozygotes for the *TPMT* mutant alleles have much higher levels of the cytotoxic thiopurine nucleotides and are at higher risk for developing serious hematological toxicities during treatment with standard-dosage regimens of the thiopurine drugs [187]. As a result, patients with absent and low TPMT activity can only tolerate 5 and 50% of the standard 6-mercaptopurine regimen.

The TPMT metabolism represents one of the most-investigated drug metabolic pathways that demonstrates the clinical relevance of genetic polymorphism. Currently, TPMT is the only drug-metabolizing enzyme with significant acceptance and widespread testing in clinical practice, with genotyping or phenotyping (determination of TPMT activity in red blood

cells) recommended by the FDA, and the thiopurine drugs are one of several medications that have clinical guidelines available through the CPIC [188,189].

Dihydropyrimidine Dehydrogenase (DYPD)

In addition to being the initial and rate-limiting enzyme that catalyzes pyrimidines such as uracil and thymine, dihydropyrimidine dehydrogenase (DYPD), a minor phase I metabolizing enzyme, also mediates the metabolism of 5-fluorouracil (5-FU) and capecitabine. Genetic polymorphisms in the *DYPD* gene that encodes DYPD result in DYPD-deficiency phenotypes with an overall frequency in the general population of about 3%–5%, which varies significantly among many ethnic groups [190].

With more than 30 SNPs in *DYPD*, the most-relevant decreased functional variants associated with grade 3- and grade 4-toxicities in 5-FU-treated patients include c.1905+1G>A, also known in the literature as *DYPD**2A or *DYPD*:IVS14+1G>A (rs3918290); c.1679 T>G, also known as *DYPD**13 (rs55886062, p.I560S); c.2846A>T (rs67376798 p.D949V); and c.1129-5923C>G (rs75017182) [191,192]. Other variants such as c.85T>C (*DYPD**9A, rs1801265, p.C29R) do not result in altered DYPD activity [193]. Among the four functional SNPs, the G>A mutation within intron 14 results in a protein with no catalytic activity and is found in approximately 40%–50% of patients who have either a partial or complete DYPD deficiency. Homozygous and heterozygous carriers of the variant IVS14+1G>A allele have complete absence and 50%, respectively, of normal DYPD activity, and significant, sometimes life-threatening 5-FU-related toxicities [194]. However, the risk of severe toxicity is not necessarily related to *DYPD* genotypes [195]. This may be due to sensitivity of the *DYPD* genetic testing being dependent on which *DYPD* variants are included in specific test panels [193].

N-acetyltransferase

Genetic polymorphism in acetylation capacity was reported more than 50 yrs ago, when two distinct phenotypes of rapid acetylator (RA) and slow acetylator (SA) were noted in patients enrolled in a clinical trial of the antituberculosis drug isoniazid [196]. Subsequently, the phenotype differences were associated with enzyme activities of two cytosolic enzymes N-acetyltransferase-1 (NAT1) and N-acetyltransferase-2 (NAT2), which are encoded by the *NAT1* and *NAT2* genes, respectively. The NAT2 enzyme is primarily responsible for acetylation of aromatic amines and hydrazines. Polymorphism in *NAT2* results in more than 10 *NAT2* alleles, with *NAT2**4 reported as the wild-type allele, and *NAT2**5 (rs1801280) carrying the c.341T>C SNP that results in the p.I114T amino acid change, *NAT2**6 (rs1799930) with c.590G>A SNP and p.R197Q substitution, as well as *NAT2**7 (rs1799931) with c.857G>A SNP and corresponding p.G286E substitution as the primary variant alleles [197]. These three variant alleles account for the majority of the SA phenotype. The prevalence of SAs varies significantly in different ethnic groups: 90% of Arab populations, 40%–60% of Caucasians, and 5%–25% of East Asians.

Substrates for NAT include numerous arylamine- and hydrazine-containing drugs such as sulfamethoxazole, hydralazine, isoniazid, and procainamide. High blood levels of these and similarly, acetylated drugs in SAs have been associated with lupus-like syndrome (hydralazine and procainamide), peripheral neuropathy (isoniazid), and liver damage (sulfapyridine). In addition, to drug therapy, *NAT2* polymorphism has also been implicated in susceptibility to developing different types of cancer, with SA having an increased risk after prolonged exposure to carcinogenic arylamines and other industrial chemicals [198].

Glutathione-S-transferase

The human glutathione-S-transferase (GST) family of cytosolic enzymes contains at least 17 genes divided into seven classes: α , μ , π , σ , θ , ζ , and ω . Of these, the most important genes are *GSTM1* of the μ class, *GSTT1* of the θ class, *GSTP1* of the π class, and *GSTA1* of the α class. Both deletion polymorphisms and SNPs exist for *GST* genes. Gene deletion results in the null variants, *GSTM1**0 and *GSTT1**0 and loss of *GSTM1* and *GSTT1* enzyme function, respectively. Two polymorphisms of the *GSTP1* gene have been described: rs947894 carrying the exon 5 c.A1404G SNP, and p.I105V substitution at codon 105, and rs1799811 carrying the exon 6 c.C2294T SNP and p.A114V substitution at codon 114. Four different haplotypes have been described for the population: *GSTP1**A (¹⁰⁵Ile-¹¹⁴Ala), *GSTP1**B (¹⁰⁵Val-¹¹⁴Ala), *GSTP1**C (¹⁰⁵Val-¹¹⁴Val), and *GSTP1**D (¹⁰⁵Ile-¹¹⁴Val) [199]. A point mutation in the promoter of the *GSTA1* gene results in lower promoter activity associated with the *GSTA1**B allele. In contrast to deleted *GST* genotypes, the *GSTP1* and *GSTA1* polymorphisms result in genotypes with low-activity enzymes.

The frequency of occurrence of the two *GST* null alleles varies significantly among different populations. Between 42% and 58% of Caucasians and 27%–41% of Africans are reported to be lacking the *GSTM1* gene. For the *GSTT1* gene, the null-allele frequency ranges from 2% to 42% for Caucasians, 50%–60% in Asians, 15%–20% of African Americans, and less than 10% in Hispanics [200,201]. The *GSTP1* and *GSTA1* polymorphisms have been reported to occur in up to 40% of Caucasians and 54% of Africans, and 40% of Caucasians and 41% of Africans, respectively.

GSTs are detoxification enzymes that mediate the conjugation of reduced glutathione with different substrates that include carcinogens and chemotherapeutic agents, such as oxaliplatin-based chemotherapy and chlorambucil

[200,202–204], with poorer response and reduced overall survival in patients with *GSTM1*0* or *GSTT1*0* genotypes, and treated with oxaliplatin-based chemotherapy or anthracycline-based induction therapy [204–207]. Because GSTs are detoxification enzymes, the shortened survival in patients with reduced GST activity might be related to severe drug-related toxicity, as evidenced by a higher frequency of grade 4 neutropenia in homozygous carriers of *GSTM1*0* and treated with oxaliplatin-based chemotherapy for their metastatic colorectal cancer [208]. The important detoxification role of GST has also been reported in a recent study in which pediatric patients with *GSTM1* and *CYP2C9* variants have a higher risk of developing hemorrhagic cystitis when treated with the combined regimen of busulfan and cyclophosphamide [209].

POLYMORPHISMS IN DRUG-TRANSPORTER GENES

Membrane transporters are present at many endothelial and epithelial barriers, including the blood brain barrier (BBB), the intestinal epithelial cells, the hepatocytes, and the renal tubular cells. By facilitating drug excretion into the gastrointestinal tract and bile, from the liver and kidney, as well as limiting the amount of drug crossing the BBB, they provide an important physiological role of protecting humans against toxic xenobiotics, and have recently been recognized as important determinants of drug disposition and response [210]. These drug transporters can be broadly classified into two groups: the efflux adenosine triphosphate-binding cassette (ABC, and formerly known as multidrug resistance [MDR]) family of transporters, and the uptake solute carrier (SLC) family of transporters. In all, 49 members are present within the human ABC-transporter family. Based on homology of their amino acid sequences, they are further classified into seven subfamilies. Of all the ABC

transporters, the better-known examples are ABCB1 (P-glycoprotein [Pgp] or MDR1), ABCC1 (multidrug resistance 1 [MRP1]), ABCC2 (multidrug resistance [MRP2]), and ABCG2 (breast cancer resistance protein [BCRP]). For the SLC family, there are 360 members that are subdivided into 46 subfamilies. The better-known SLC transporters are organic anion-transporting polypeptide (OATP), organic cation transporter (OCT), and organic anion transporter (OAT). Genetic variants of the genes encoding these drug-transport proteins (<http://www.pharmGKB.org>) have been discovered that affect their expression, substrate specificity, and/or intrinsic transport activity, and ultimately disposition, efficacy, and safety of many drug substrates.

The ABC-Efflux Transporters

ABCB1

ABCB1 was the first recognized and the most-studied ABC transporter. It is encoded by the highly polymorphic *ABCB1*, with more than 50 SNPs and three insertion/deletion polymorphisms reported. The most common studied SNPs are the c.C1236T (rs1128503) silent polymorphism in exon 12, the c.G2677A/T (rs2032582) polymorphism in exon 21, and the c.C3435T (rs1045642) silent polymorphism in exon 26. The c.G2677A/T polymorphism results in a change in amino acid sequence p.A893S (G2677T) SNP or p. A893T (G2677A) SNP. Ethnic variations in allelic variant distribution are well known [211,212]. In addition, strong LD among these SNPs had been reported to create haplotypes consisting of 1236C>T, 2677G>A/T, and 3435C>T. The three *ABCB1* SNPs and their haplotypes (Table 1.3) are important in expression and function of ABCB1.

The functional and clinical implication of the *ABCB1* polymorphism was first evaluated for the C3435T SNP with digoxin as the substrate, demonstrating a relationship between lower

TABLE 1.3 Selected ABC Transporters Polymorphisms Indicating Allele Variants and Frequency, and Drug Substrates

Genes	Allele Variants, Amino Acid Change	Frequency (%)	Drug Substrate Examples
<i>ABCB1</i>	C3435T	48%–59% in Caucasians 37%–66% in Asians 10%–27% in Africans	Protease inhibitors (ritonavir, saquinavir, nelfinavir) Anticancer drugs (anthracyclines, taxanes, vinca alkaloids, imatinib)
	C1236T	34%–42% in Caucasians 60%–72% in Asians 15%–21% in Africans	Immunosuppressants (cyclosporine, tacrolimus) Antibiotics (erythromycin, levofloxacin) Calcium channel blockers (diltiazem, verapamil) Digoxin, pivalastatin, simvastatin
	G2677T, A893S	38%–47% in Caucasians 32%–62% in Asians ≤15% in Africans	
	G2677A, A893T	1%–10% in Caucasians 3%–22% in Asians	
	1236C>T/2677G>T/3435C>T haplotype	23%–42% in Caucasians 28%–56% in Asians 4.5%–8.7% in Africans	
<i>ABCC2</i>	1249G>A, V417I	22%–26% in Caucasians 13%–19% in Asians 14% in Africans	Reverse transcriptase inhibitors (tenofovir), Anticancer drugs (anthracyclines, vinca alkaloids, methotrexate, SN-38 glucuronide), pravastatin, rifampin
<i>ABCG2</i>	421C>A, Q141K	6%–14% in Caucasians 15%–36% in Asians 0%–5% in Africans	Anticancer drugs (methotrexate, imatinib, gefitinib, SN-38, SN-38 glucuronide, topotecan), Apixaban, atorvastatin, rosuvastatin, glyburide, dolutegravir

expression of *ABCB1* and increased digoxin bioavailability and plasma concentration after oral administration in TT homozygotes with reduced *ABCB1* activity [213]. In two separate studies, investigators showed that CC genotype of the C3435T SNP (increased Pgp expression) is associated with reduced efficacy and a higher risk of myalgia after treatment with atorvastatin [214] and increased statin-associated increase in serum creatine kinase [215], presumably due to lower intracellular concentration and higher plasma concentration of statin.

The polymorphism also affects plasma concentrations and clinical effects of protease inhibitors. After 6-mo therapy with efavirenz or nelfinavir, patients with the TT genotype had a greater rise in cluster of differentiation 4 (CD4) cell counts than patients with the CC genotype

[216]. Therefore, *ABCB1* genotyping may have a role in predicting responses to protease inhibitors. The *ABCB1* haplotype TTT (rs1128503, rs2032582, rs1045642) was reported to be responsible for increased morphine exposure and the exhibition of morphine sensitivity in a patient [186]. In addition, Sadhasivam et al. reported an association between another *ABCB1* variant (rs9282564) and increased risk of morphine-induced respiratory depression in patients with the GG and GA genotypes of this SNP [217].

Nevertheless, conflicting results have been reported regarding the functional and clinical significance of the polymorphism for different substrates including psychotropics (see Chapter 7), antiretroviral protease inhibitors, immunosuppressants, and anticancer drugs. This may be due to the use of different assays

and study designs to identify ABCB1 substrates, the overlapping substrate specificity between ABCB1 and other enzymes and transporters; e.g., CYP3A4 for cyclosporine and OATP transporters for fexofenadine [218], the existence of strong LD necessitating a haplotype approach rather than individual SNPs in association studies or clinical evaluations. In addition, the 1236C>T/2677G>T/3435C>T haplotype was shown to affect the inhibition of substrate transport and not the transport process per se [219]. Thus, the functional effect of *ABCB1* polymorphism may be more modest than previously thought. Whether additional mutations resulting in loss of function, significant change in substrate specificity or functionality would have a bigger impact is not known, and awaits further studies for clarification [219,220].

ABCC1 and ABCC2

Both *ABCC1* and *ABCC2* are involved in the biliary excretion of conjugated drugs such as glucuronides or sulfates of tamoxifen and SN-38 glucuronide [221,222], organic anions, and some nonconjugated drugs such as methotrexate and pravastatin (Table 1.3), and exhibit overlapping substrate specificities for a variety of drugs. Genetic variation in *ABCC1* gene is rare, whereas polymorphisms of *ABCC2* gene are more common, including the c.1249G>A SNP (rs2273697) in exon 10 resulting in a p.V417I substitution and lower protein expression. Another identified polymorphism is the c.3972C>T SNP (rs3740066) in exon 28 with a p.I324I amino acid substitution [223].

Patients with the 1249G>A variant and receiving tenofovir were reported to have higher risk of drug-induced renal proximal tubulopathy, possibly a result of reduced renal drug excretion [224]. In an exploratory study of an association between *ABCC2* polymorphisms and haplotypes with irinotecan disposition in a cohort of 167 Caucasian patients with solid tumors, a total of 15 *ABCC2* haplotypes were constructed from six variants of *ABCC2* gene. The *ABCC2**2




FIGURE 1.2 Schematic representation of potential protective effect of *ABCC2* polymorphism against irinotecan-induced diarrhea.

haplotype (low activity) was found to be associated with lower irinotecan clearance of 28.3L/h in 48 patients compared with 31.6L/h in 75 patients not carrying the haplotype ($P=.02$). Interestingly, patients carrying the *ABCC2**2 haplotype but not the *UGT1A1**28 allele experienced lower incidence of severe grade 3–4 diarrhea (odds ratio of 0.15) compared to patients carrying at least one *UGT1A1**28 allele (odds ratio of 1.87), suggesting a protective effect of *ABCC2**2 haplotype against diarrhea occurrence [225]. Because *ABCC2* mediates the secretion of SN-38 glucuronide into the bile, the protective effect might reflect a lower exposure of intestinal epithelial cells to SN-38 that is formed after cleavage of SN-38 glucuronide by β -glucuronidase within the intestine (Fig. 1.2).

ABCG2

The *ABCG2* gene encodes the BCRP, which is also known as mitoxantrone resistant protein (MXR), or placenta-specific ATP binding cassette transporter (ABCP). More than 80 polymorphisms in *ABCG2* have been reported, with the most studied being the c.421C>A SNP (rs2231142, p.Gln141Lys) in exon 5 that results in a p.Q141K substitution and lower protein expression (Table 1.3) [226]. The c.421C>A

variant with K141 is commonly present in different ethnic groups, being more common in Asians and Caucasians than in sub-Saharan Africans [211,212,227].

Patients carrying the c.421C>A SNP were reported to have increased concentrations of gefitinib and topotecan [228,229], resulting in higher incidence of gefitinib-induced diarrhea [230]. Increased risk of diarrhea was also associated with the *ABCG2* polymorphism in patients with cancer and receiving rituximab plus cyclophosphamide/doxorubicin/vincristine/prednisone (R-CHOP) therapy [231]. *ABCG2* also plays a role in disposition of other drugs, with the c.421C>A variant reducing biliary excretion of apixaban [232], dolutegravir [233], and rosuvastatin [234]. In 305 Chinese patients with hypercholesterolemia treated with 10mg of rosuvastatin per day, a gene dose-dependent reduction in low-density lipoprotein cholesterol levels was observed in a carrier of the C421A variant [235]. In both the JUPITER trial, with more than 4,000 patients, and another study with a cohort of 291 Chinese patients, a strong association at the genome-wide level significance has been reported between the C421A variant and altered statin efficacy [236,237].

The SLC-Uptake Transporters

Organic Anion Transporting Polypeptides

In contrast to *ABCB1*, OATPs are influx or uptake transporters. In addition, to facilitating hepatic uptake of drugs such as statins and antidiabetic agents from the blood into hepatocytes for further metabolism or biliary secretion, OATP also mediates the transport of several endogenous compounds, including bile salts, across the cell membrane. A total of 11 OATP transporters have been identified and classified into six families [238]. Of the human OATPs, OATP1A2, OATP1B1, OATP1B3, and OATP2B1 (Table 1.4) are the best characterized.

OAT1B1

The human *SLCO1B1* gene encodes OATP1B1, which is also known as OATP-C. Since the discovery of the first c.521T>C SNP [239], multiple SNPs have been reported for *SLCO1B1*, with 17 different *SLCO1B1* alleles identified [240]. The 521T>C SNP (rs4149056) with p.V174A substitution results in lower expression of the OATP1B1 protein and reduced transport activity. The 521T>C SNP is more common in Caucasians and Asians than in Africans (Table 1.4). Another very common mutation in all investigated ethnic groups is the c.388A>G SNP (rs2306283) resulting in p.N130D substitution, although conflicting results exist regarding associated changes in transport activity. More importantly, though, the 521T>C SNP and 388A>G SNP are in LD, resulting in several functionally distinct haplotypes, e.g., *OATB1B1**5 carrying the 388A/521C, *OATB1B1**15 carrying the 388G/521C, and *OATB1B1**17 carrying the 388G/521C with -11187A/-10499A of two additional SNPs in the promoter region of *SLCO1B1* [240,241]. Both *5 and *15 haplotypes contain the 521C allele and have been associated with reduced activity.

OATP1B1 plays an important role in hepatic uptake of the 3-hydroxy-3-methylglutaryl-Coenzyme A (HMG-CoA) reductase inhibitors such as pravastatin and rosuvastatin, as well as simvastatin acid, the active metabolite of simvastatin. The 521T>C variant has been associated with altered pharmacokinetics of simvastatin acid, with the CC homozygotes having more than two–three-fold increased systemic exposure compared to the other two genotypes [242], potentially resulting in increased toxicity [243], and with decreased intracellular concentration of simvastatin acid for inhibiting HMG-CoA reductase in hepatocytes, a lower efficacy for cholesterol reduction [244]. In a GWAS, 316,184 SNPs were compared between 96 patients treated with 80mg/day of simvastatin and suffering from myopathy and 96 control subjects without the adverse drug effect. A noncoding rs4363657

TABLE 1.4 Selected SLC Transporters Polymorphisms Indicating Allele Variants and Frequency, and Drug Substrates

Genes	Allele Variants, Amino Acid Change	Frequency (%)	Drug Substrate Examples
<i>SLCO1B1</i>	521T>C, V174A	8%–22% in Caucasians 1%–19% in Asians 1%–5% in Africans	HMG-CoA reductase inhibitors (atorvastatin, simvastatin acid, pravastatin, rosuvastatin), Anticancer drugs (SN-38, methotrexate),
	388A>G, N130D	30%–46% in Caucasians 54%–84% in Asians 72%–81% in Africans	Antibacterials (rifampicin, cefazolin), repaglinide, valsartan
<i>SLCO2B1</i>	1457C>T, S486F	1%–6% in Caucasians 25%–36% in Asians 10%–41% in Africans	HMG-CoA reductase inhibitors (atorvastatin, fluvastatin, pravastatin, rosuvastatin), glibenclamide, fexofenadine, motelukast
	935G>A, R312Q	2%–14% in Caucasians 21%–40% in Asians 7%–15% in Africans	
<i>SLCO1B3</i>	334T>G, S112A	74%–89% in Caucasians 64%–83% in Asians 35%–41% in Africans	Anticancer drugs (docetaxel, paclitaxel), digoxin
	699G>A, M233I	71%–90% in Caucasians 64%–84% in Asians 34%–48% in Africans	
<i>SLC22A1</i>	1201G>A, G401S	1% in Caucasians 0% in Asians 1% in Africans	Metformin
	1393G>A, G465R	4% in Caucasians 0% in Asians, Africans	
	1256delATG, M420del	60% in Caucasians 74%–81% in Asians 74% in Africans	
<i>SLC22A2</i>	808G>T, A270S	16% in Caucasians 14%–17% in Asians 11% in Africans	Metformin

SNP within intron 11 of *SLCO1B1*, found to be in nearly complete LD with the rs4149056 polymorphism (521T>C, V174A) ($r^2 > 0.95$), was identified as the only strong SNP marker associated with simvastatin-induced myopathy. The odds ratio (OR) for myopathy was reported as 4.3 per copy of the C allele, and 17.4 in CC homozygotes compared with TT homozygotes [245].

In a more recent study, carriers of the T521C SNP were shown to have an OR of 8.86 ($P < .01$) for statin-induced serum creatine kinase elevation,

whereas the impact of the A388G SNP was much smaller (OR of 0.24, $P < .05$) [215]. The magnitude of the clinical significance shown in these two studies [215,245] suggests potential value of genotyping to screen out patients with abnormal OATP1B1 activity to improve the therapeutic index of simvastatin, and may be for other HMG-CoA reductase inhibitors such as pravastatin that are also OATP1B1 substrates [246–248]. Indeed, both the 521T>C SNP and *SLCO1B1**17 haplotype had been shown to be associated with increased

pravastatin concentrations and decreased efficacy [249–251]. Based on the results of these and other studies, the CPIC has made recommendations for genotype-based dosing of statins in its 2014 guideline update [252]. In addition, both *SLCO1B1* drug–drug interaction with concurrent drug therapy and drug–gene interaction with *SLCO1B1* variants can alter statin transport and subsequent metabolism and, hence, the risk for statin-related ADR such as rhabdomyolysis [253].

OATP2B1

OATP2B1, also known as OATP-B, possesses substrate selectivity similar to that of OATP1B1 [254]. OATP2B1 has also been found to be expressed in the luminal membrane of the small intestinal enterocytes [255], and hence would have a role in drug absorption. Since the first discovery of genetic polymorphism, several sequence mutations of *OATP2B1* have been described, including the c.1457C>T SNP (rs2306168), c.601G>A SNP (rs35199625), c.935G>A SNP (rs12422149), c.43C>T SNP (rs56837383), and a nine-nucleotide deletion of three amino acids 26–28 (26–28, p.QNT) of *OATP2B1* [256]. Although decreased transport activity had been shown mostly in vitro for most of these SNPs, the results are not consistent among all studies. In addition, significant ethnic variabilities exist in allele frequency of these SNPs; for example, the allelic frequency of c.1457C>T SNP is higher in Asians (31%) compared to Caucasians (3%)

A recent study evaluated the impact of the 1457C>T SNP on fexofenadine pharmacokinetics in Japanese subjects, and found similar pharmacokinetic parameters among the three genotype groups [257]. Although the same SNP did not affect the absorption of the leukotriene receptor antagonist motelukast, patients who carry the 935A variant allele of the c.935G>A SNP was reported to show lower plasma concentration and lesser pharmacological response [258]. Yet a separate study reported the lack of an association between motelukast and the

c.935G>A SNP. Although this might suggest that the effect of *SLCO2B1* SNP on drug absorption could be substrate dependent, more importantly, additional studies with other substrates would need to be performed for clarification of the effect of *SLCO2B1* on drug disposition.

OATP1B3

In humans, the *SLCO1B3* gene encodes OATP1B3, which was previously also known as OATP8 and LST-2. Several sequence variations exist for the *SLCO1B3* gene. The c.334T>G SNP (rs4149117) and the c.699G>A SNP (rs7311358) occur at a high frequency in Caucasian populations. Although OATP1B3 mediates the hepatic uptake of several drugs, including taxanes [259], a study in 90 patients with cancer from six different ethnic groups reported that there were no associations between paclitaxel clearance and the two *OATP1B3* SNPs [260]. Similarly, no associations were found between docetaxel pharmacokinetics and *OATP1B3* SNPs [261,262]. The role of *OATP1B3* polymorphisms in drug disposition and response await further clarification from future studies.

In summary, OATP polymorphisms can affect disposition and possibly response for a large number of drugs. Current evidence strongly suggests a vital role of specific SNPs of *SLCO1B1* gene (e.g., 521T>C) for statin efficacy and adverse effects. Similar data for other OATP1B1 substrates from future clinical studies would provide further evidence of the value of prospective genotyping for *SLCO1B1* variants in individualizing drug therapy. In contrast, the data for validating the functional role of *SLCO2B1* and *SLCO1B3* polymorphisms are not as clear. Much more work is needed for clarifying the clinical significance of these SNPs for predicting pharmacokinetic profile for, and clinical response to, OATP2B1 and OATP1B3 substrates.

Organic Cation Transporters

Three organic cation transporters (OCTs) have been identified in humans: OCT1, OCT2,

and OCT3, all of which are members of the SLC22A family, and are encoded by the corresponding *SLC22A1*, *SLC22A2*, and *SLC22A3* genes, respectively (Table 1.4). OCT1 is primarily expressed in the hepatocytes and mediates cellular uptake of drugs into the liver. OCT2 is primarily expressed in the proximal tubules of the kidney. In contrast, OCT3 is more broadly distributed in the body.

Located on chromosome 6, *SLC22A1* is highly polymorphic with reduced or loss of transporter functional activity secondary to four coding polymorphisms: c.181C>T (rs12208357, p.Arg61Cys), c.1393G>A (rs34059508, p.Gly465Arg), c.1201G>A (rs34130495, p.Gly401Ser), and OCT1 Met420 deletion of three bases ATG at codon 420 of exon 7 and collectively designated as rs72552763 [263]. The Met420 deletion variant commonly occurs in Caucasians and African Americans with a frequency of 18.5% and 5%, respectively. OCT1 genotypes have been shown to contribute to interindividual variability in disposition of several drugs, including ondansetron, metformin, morphine, and tramadol [264–267].

Of the different SNPs that have been identified for the *SLC22A2* gene, the most relevant one is the c.808G>T SNP (rs316019) that results in the p.A270S substitution. The antidiabetic drug metformin is primarily renally eliminated by active tubular secretion via OCT2. Homozygotes of the low-activity 270S variant had been shown to have lower renal clearance and higher plasma concentrations of metformin when compared to homozygous carriers of the wild-type 270A [268,269]. Interestingly, Tzvetkov et al. demonstrated that OCT1 is also expressed in the distal tubule and may play a role in tubular reabsorption of metformin. They reported that homozygous and heterozygous carriers of various haplotypes of low-activity alleles of several *SLC22A1* polymorphisms (c.1201G>A SNP with p.G401S substitution, c.1393G>A SNP with p.G465R substitution, and a deletion resulting in M420del) were associated with increases in metformin renal clearance

by about 20%–30% [267]. Nevertheless, OCT1 is primarily expressed in the hepatocyte, the major site of action of metformin. The same low-activity OCT1 variant alleles of these polymorphisms have also been reported to decrease hepatic uptake of metformin with resultant lower blood glucose response [270], and more recently for fenoterol, resulting in increased systemic exposure and drug-related toxicities [271]. The effects of genetic polymorphisms in other transporters such as the multidrug and toxin extrusion transporters as well as pharmacological targets for metformin is further discussed in Chapter 9.

Organic Anion Transporters

In contrast to the OCT belonging to the same SLC22 family, the organic anion transporters (OATs) primarily mediates the transport of organic anions. Four OATs have been studied regarding their tissue location: OAT1, OAT2, and OAT3 are primarily expressed in the basolateral membrane of the renal proximal tubule, whereas OAT4 is located at the apical side. Therefore, OAT1, OAT2, and OAT3 are responsible for uptake of drug substrates into the tubular cells and OAT4 mediates their secretion into the renal tubule. Although several polymorphisms have been reported for *SLC22A6* encoding OAT1, *SLC22A7* encoding OAT2, *SLC22A8* encoding OAT3, and *SLC22A11* encoding OAT4, the allele frequency of these SNPs are all $\leq 1\%$ and their functional significance have not been clarified [272,273].

POLYMORPHISMS IN DRUG-TARGET GENES

The study of pharmacodynamics encompasses the biochemical and physiological effects of drugs on the body and the relationship between drug concentration and drug effect. Drugs exert their effects through interaction with numerous protein types, including cell

surface receptors (e.g., β -adrenergic, 5-hydroxytryptamine receptors, and μ -opioid receptors), enzymes (e.g., vitamin K epoxide reductase complex 1, adenosine monophosphate-activated protein kinase, and catecho-O-methyltransferase), and ion channel proteins (e.g., sodium and potassium channels, epithelial sodium channel). Additionally, numerous intracellular signaling proteins downstream from the target protein are involved in eliciting drug response. Genetic variation affecting either the activity or expression of a drug-target or intracellular signaling protein can have significant consequences for pharmacodynamic drug response.

Phenotypic response to genetic variation for drug-target proteins generally differs from that of drug-metabolizing enzymes and drug transporters (Table 1.5). As illustrated in Fig. 1.3, variation in drug-metabolizing enzymes results in distinct phenotypes (e.g., PMs, EMs, or UMs), as described in the earlier section. With precision oncology the expression of drug-target receptor gene for tumor cells predicts drug efficacy. In addition, there are a limited number of examples of genetic variants in drug-target proteins in germline cells that result in distinct pharmacodynamic effects. One of these examples involves mutations in the vitamin K epoxide complex subunit 1 (*VKORC1*) gene, in which rare non-synonymous mutations result in warfarin resistance, and exceptional high doses (30 mg/day or higher) are required to achieve therapeutic anticoagulation. Most polymorphisms that impact drug pharmacodynamics tend to be much more subtle and help explain response variability across a single distribution curve. For example, commonly occurring variations in the *VKORC1* regulatory regions help explain the significant interpatient variability in the warfarin dose required to produce optimal anticoagulation, as described in detail in Chapter 6. The remainder of this section discusses examples of genes for various types of drug-target proteins that contribute to the interpatient variability in pharmacodynamic drug responses.

Drug Target Receptor Genes in Oncology

Several cancer chemotherapy agents have been developed based on findings that overexpression of certain tumor cell surface receptors drives tumor cell growth and which are further described in Chapters 3 and 5. Expression of the epidermal growth factor receptor type 2 (HER2), also known as Her2/neu and ErbB2, is one such example that influences disease prognosis and predicts drug. Overexpression of HER2 occurs in approximately 20% of metastatic breast cancers and is associated with more aggressive cancer and poor prognosis [274]. Trastuzumab is a recombinant monoclonal antibody that was developed to target HER2 and block growth and survival of HER2-dependent tumors. The addition of trastuzumab to chemotherapeutic regimens to treat breast cancer significantly slows the progression of breast cancer in women with HER2-positive tumors, with treatment effects positively correlated with the degree of HER2 overexpression [275]. Thus, testing to confirm HER2 overexpression is necessary before trastuzumab use. Although the data are conflicting and require further confirmation, a genetic association between HER2 655 A > G (Ile/Val) polymorphism (rs1136201) and trastuzumab cardiotoxicity has also been suggested [276].

The epidermal growth factor receptor (EGFR), also known as HER1 or ErbB1, is overexpressed in head and neck, colon, and rectal cancer. EGFR overexpression is associated with cancer growth and invasion and portends a poor clinical prognosis. The discovery of the *EGFR* gene and its role in cancer prognosis led to the development of EGFR antagonists, including cetuximab, panitumumab, erlotinib, and gefitinib. Cetuximab is a recombinant monoclonal antibody that binds to the extracellular domain of the EGFR, thus preventing epidermal growth factor and other ligands from activating the receptor. Cetuximab is indicated in the treatment of metastatic

TABLE 1.5 Consequences of Selected Genetic Variation in Drug Disposition and Drug Target Proteins

Gene	Drug Examples	Clinical Consequence
<i>CYP2D6</i>	Atomoxetine	PMs may have 10-fold greater atomoxetine exposure
	Codeine	UMs are at increased risk for morphine toxicity (details in Table 1.2)
<i>CYP2C9</i>	Warfarin	CYP2C9 deficiency increases bleeding risk
<i>CYP2C19</i>	Clopidogrel	CYP2C19 deficiency reduces drug effectiveness
<i>G6PD</i>	Rasburicase	G6PD deficiency increases risk for hemolytic anemia
<i>TPMT</i>	Azathioprine, 6-mercaptopurine, Thioguanine	Nonfunctional genotype increases the risk of serious, life-threatening myelosuppression with conventional drug doses
<i>UGT1A1</i>	Irinotecan	Reduced function genotype increases risk for drug-induced neutropenia
<i>DPD</i>	Capecitabine, 5-fluorouracil	DPD deficiency may lead to severe diarrhea, neutropenia, neurotoxicity)
<i>SLCO1B1</i>	Simvastatin	Increased risk for myopathy
DRUG-TARGET GENES		
<i>EGFR</i>	Cetuximab, Panitumumab	Determines drug effectiveness
<i>HER2</i>	Trastuzumab	Determines drug effectiveness
<i>ADRB1</i>	β -blockers	Influences variability in blood pressure response and possibly mortality reduction
<i>VKORC1</i>	Warfarin	Determines dose needed for optimal anticoagulation
<i>KCNJ11</i> and <i>ABCC8</i>	Sulfonylureas	Drug effectiveness
<i>KCNMB1</i>	Verapamil	Possibly determines reduction in blood pressure
<i>DRD3</i>	Antipsychotics	Risk for tardive dyskinesia
<i>GRK5</i>	β -blockers	Drug effectiveness on clinical outcomes in heart failure
<i>ATM</i>	Metformin	Antidiabetic response
<i>SLC6A4</i>	SSRIs	Drug effectiveness
<i>HTR2A</i>	Clozapine	Drug effectiveness

colorectal cancers that overexpress EGFR, in which it has been shown to improve survival [277,278]. Similar to cetuximab, panitumumab is a monoclonal antibody that blocks activation of the EGFR and is indicated in metastatic colorectal cancer that progresses despite chemotherapy with fluoropyrimidine-, oxaliplatin-, and irinotecan-containing regimens. Erlotinib and

gefitinib also target the EGFR and are indicated in nonsmall cell lung cancer.

Other examples of targeted chemotherapy developed based on genetic abnormalities include:

- rituximab, a monoclonal antibody used to treat CD20-positive, B-cell non-Hodgkin's lymphoma and chronic lymphocytic leukemia;




FIGURE 1.3 Many drug-metabolizing enzyme polymorphisms are inactivating resulting in distinct phenotypes, such as the poor metabolizer and extensive metabolizer phenotypes. In contrast, drug-receptor polymorphisms tend to be more subtle and help explain variability across single distribution curve.

- imatinib mesylate, a kinase inhibitor developed to block the product of a reciprocal translocation between chromosomes 9 and 22, occurring in 95% of patients with chronic myeloid leukemia; and
- crizotinib, an anaplastic lymphoma kinase (ALK) and c-ros oncogene1, receptor tyrosine kinase (ROS-1) inhibitor that targets the echinoderm microtubule associated protein like 4–anaplastic lymphoma kinase (EML4-ALK) gene fusion product in nonsmall-cell lung cancer.

Drug Target Receptor Genes in Cardiology

β_1 -receptors are located in the heart and kidney, in which they are involved in the regulation of heart rate, cardiac contractility, and plasma renin release. β_1 -receptor mediated effects contribute importantly to the pathophysiology of numerous cardiovascular diseases, including hypertension, coronary artery disease, and heart failure. In particular, plasma renin release and activation of the renin-angiotensin-aldosterone system lead to increased blood volume and

vasoconstriction in hypertension. Increases in heart rate and cardiac contractility increases myocardial oxygen demand, thus contributing to myocardial ischemia in patients with coronary heart disease. Furthermore, increased sympathetic nervous system activity is one of the primary mechanisms contributing to cardiac remodeling and heart failure progression. Consequently, β -blockers exert beneficial effects across cardiovascular diseases, resulting in blood pressure reduction in hypertension, lowering of myocardial oxygen demand in ischemic heart disease, and attenuation of cardiac remodeling in heart failure. There is evidence that genetic variation for the adrenoceptor β_1 (ADRB1) may influence the effectiveness of β -blocker therapy.

The ADRB1 is encoded by an intronless gene, located on chromosome 10q24-26. Two common nonsynonymous SNPs in the ADRB1, p.S49G and p.R389G are in strong LD. The S49G SNP is located in the extracellular region of the receptor near the amino terminus, and the R389G variant is located in the cytoplasmic tail in the G-protein coupling domain of the ADRB1. In vitro studies show lesser receptor downregulation with the S49 form of the receptor and both greater receptor coupling to the G-protein and greater adenylyl cyclase activity with the R389 form [279,280]. There are ethnic differences in the S49G and R389G allele frequencies, with a G49 frequency of 12%–16% in Caucasians and 23%–28% in African Americans and a G389 frequency of 24%–34% in Caucasians and 39%–46% in African Americans [281].

The ADRB1 gene has been the primary focus of research into genetic determinants of responses to β -blockers in hypertension, coronary heart disease, and heart failure [282–284]. In each case, the R389 allele or S49-R389 haplotype has been associated with greater response to β -blockade presumable because of greater adrenergic activity with this allele and haplotype. For example, treatment of hypertension with metoprolol produced greater blood pressure reduction in patients who were homozygous for the S49-R389

haplotype than in carriers of the G49 or G389 allele [285]. Among patients with coronary heart disease, the S49-R389 haplotype was associated with an increased risk for death compared to other haplotypes, an effect negated by treatment with atenolol [286]. In patients with heart failure, the homozygous R389 genotype was associated with greater improvements in left ventricular ejection fraction with carvedilol or metoprolol and greater survival benefits with bucindolol [287–289]. These clinical data are consistent with the in vitro data implying greater agonist-mediated effects (e.g., greater sympathetic nervous system-driven hemodynamic effects) with the S49 and R389 alleles and suggest that *ADRB1* genotype is an important determinant of blood pressure and cardiac responses to β -blockers.

The *ADRB1* genotype is also associated with β -blocker tolerability in heart failure. β -blockers are indicated for patients with heart failure because they attenuate the detrimental effects of the sympathetic nervous system on heart failure progression. However, because β -blockers have negative inotropic effects (i.e., reduced cardiac contractility), they can worsen heart failure when first started. For this reason, they must be started in very low doses with careful up-titration. Although most heart failure patients tolerate β -blocker initiation at low doses and slow up-titration, some experience significant heart failure exacerbation. The influence of *ADRB1* genotype on tolerability to β -blocker initiation and up-titration has been examined, and it was found that carriers of the Gly389 allele or the 49Ser/Ser genotype more frequently require increases in concomitant heart failure therapy (predominately diuretics) for management of symptoms of worsening heart failure during β -blocker titration than patients with other genotypes [290].

Stimulation of the presynaptic adrenergic α_2C -receptor (ADRA2C) results in inhibition of norepinephrine release, and has been correlated with β -blocker response. The ADRA2C Del322-325 ($\alpha_{2C\text{del}322-325}$) polymorphism causes

an inframe deletion of 12 nucleic acids, resulting in the loss of four amino acids in the ADRA2C protein and loss of protein function. Loss of ADRA2C function would be expected to result in less inhibition of norepinephrine release, and consequently increased norepinephrine levels and sympathetic tone. The frequency of the Del322-325 variant exhibits marked variability by ancestry, with a frequency of approximately 0.40 in African Americans and <0.05 in those of European descent [291], and homozygosity for Del322-325 variant has been associated with higher risk of heart failure in African Americans compared to Caucasians.

In a large, multicenter, randomized, placebo-controlled heart failure trial, investigators found that individuals with the Del322-325 allele had greater reductions in sympathetic activity with bucindolol, a nonselective β -blocker with α_1 -receptor blocker properties. However, individuals with the wild-type (Ins322-325) ADRA2C genotype derived significant survival benefits from bucindolol, whereas Del322-325 allele carriers did not [292]. The mechanism underlying this association was not determined. However, it was hypothesized that the significant sympatholytic activity with bucindolol in Del322-325 allele carriers caused detrimental clinical effects. These findings might explain the negative association between bucindolol use and heart failure survival in the study population overall. Specifically, whereas carvedilol, metoprolol, and bisoprolol were all shown to improve survival in heart failure, bucindolol was not [293,294]. However, compared to other β -blocker trials, the trial with bucindolol enrolled a large number of African Americans, in whom the Del322-325 allele, associated with lack of benefit with bucindolol, is ten-times more common.

With its pharmacological targets at the *ADRB1* on myocytes and on the adrenergic neurons, the interaction of both *ADRB1* and *ADRA2C* SNPs could further modulate the drug response. A recent clinical trial identified subsets of populations with different responses based

on evaluation of both genotypes. Enhanced bucindolol efficacy was associated with *ADRB1* homozygotes, whereas intermediate efficacy was observed in patients with Gly variant of the *ADRB1* SNP and homozygote carriers of the wild-type α_{2C} 322-325. In contrast, a lack of efficacy was reported in carriers of the *ADRB1* Gly variant and homozygous carriers of the Del322-325 allele [295]. Similarly, Reddy et al. recently reported that in children with dilated cardiomyopathy, β -blockers produced better hemodynamics and preservation of cardiac function in those with high-risk genotypes, including α_{2Cdel} 322-325 and β_1 Arg389 [296]. Similar combinatorial pharmacogenomic approaches have also been investigated in psychopharmacology and is discussed in Chapter 7.

Drug Target Genes in Psychiatry

Antidepressants target 5-hydroxytryptamine (5-HT) receptors, and a number of studies have examined the association between antidepressant treatment response and 5-HT genotype, as described in more details in Chapter 7. However, results of these studies are largely inconsistent and even conflicting. For example, in a large-scale association study of 68 candidate genes, only the synonymous IVS2 A/G (rs7997012) SNP within intron 2 of the *HTR2A* gene, which codes for the postsynaptic 5-HT_{2A} receptor, was associated with response to citalopram [297]. Although a large study in European Caucasians confirmed the association between the rs7997012 SNP and antidepressant response, the findings were opposite of those in the initial study [298].

The majority of drug-target genetic associations discussed so far related to drug effectiveness. Variation in the *DRD3* gene, encoding for the dopamine D3 receptor, is an example of drug-target genotype linked to adverse drug effects. Specifically, the *DRD3* p.S9G variant has been implicated in risk for developing

tardive dyskinesia, an irreversible movement disorder that develops after long-term antipsychotic treatment, particularly with typical antipsychotics. In a meta-analysis, the G9 allele was significantly overrepresented among 317 patients with tardive dyskinesia compared to 463 patients without this adverse drug effect [299]. Furthermore, G9 allele homozygotes had higher abnormal involuntary movement scores compared to both heterozygotes and S9 allele homozygotes. This association was confirmed in another meta-analysis [300].

Drug Target Genes in Pain Management

Over the years, several SNPs have been discovered for genes that encode different analgesic drug targets, and association studies had been carried out in various pain phenotypes. Not surprisingly, nonreplication of findings is common. The μ -opioid receptor (MOR) is the primary drug target for endogenous opioid peptides and the opioid analgesics. With more than 100 variants of the μ -opioid receptor gene (*OPRM1*) identified [301], the most studied polymorphism is the c.118A>G (rs1799971) SNP in exon 1 of *OPRM1* that results in p.N40D, and lower mRNA expression and protein amount associated with the G allele. This functional difference between the two alleles is reflected in stronger binding by the G allele to the endogenous opiate β -endorphins, thereby affecting opioid action at the receptor site, with decreased opioid potency by a factor of two to three [302]. This is evident by the report of Oertel et al., who showed that despite a stronger binding, the signal efficacy is weaker in regions of the brain that are important to pain perception and experience [303].

Decreased clinical response to opioids had been shown in carriers of the G allele. Klepstad et al. reported morphine-dosage requirements in 207 cancer patients differed among carriers of the wild type versus that of the variant allele. Four homozygous carriers of the G allele

required 225 ± 143 mg/day for effective pain control compared to 97 ± 89 mg/day in 78 wild-type homozygotes ($P = .006$). However, dosage requirement for heterozygote was 66 ± 50 mg/day, so there was no evidence of a gene-dose effect [304]. Chou et al. also reported similar findings of different dosage requirements for postoperative pain with patient-controlled analgesia, at 24 and 48 h after total knee arthroplasty respectively, of 22.3 ± 10 mg and 40.4 ± 21 mg in homozygous carriers of the G allele versus 16 ± 8 mg and 25.3 ± 15.5 mg in wild-type homozygotes [305]. Two patients identified as “low” responders of morphine requiring 1.8 and 2 gm/day were identified as a carrier of the G allele [306,307]. These and other clinical trial results [308–312] suggest that *OPRM1* genotype and haplotype analyses could have clinical implication for pain control in a variety of patients. Chidambaran et al. also reported a higher risk of morphine-induced respiratory depression associated with the A118G SNP [312].

In addition to morphine, a significant association has also been shown between the A118G SNP and decreased potency of morphine-6-glucuronide (M6G), the pharmacologically active metabolite of morphine. Using pharmacokinetic–pharmacodynamic modeling, the study showed that the effector site EC_{50} for M6G was 714 ± 197 nmol/L in six homozygous carriers of the wild-type, $1,475 \pm 424$ nmol/L in five heterozygotes, and 3,140 nmol/L in a homozygous carrier of the G allele [313]. Additional studies have reported decreased effect and higher-dosage requirement for other opioid agonists, including fentanyl and alfentanil, in carriers of the G allele [309,314–318]. However, negative association with A118G SNP has been reported for fentanyl, which was attributed by the investigators to the small sample size of the study [319]. In view of the low frequency of A118G SNP, the issue of small sample size with associated low statistical power to detect difference in analgesic doses and/or outcome is important.

Signal Transduction Proteins

Signal transduction encompasses the cascade of events following drug binding to a receptor that ultimately leads to a change in cellular response. G-protein receptor kinase 5 (GRK5) is an example of a signal transduction protein linked to drug response. The ADBR1 and other adrenergic receptors are coupled to GTP-binding proteins also called G-proteins. Upon ligand binding, the receptor couples to the intracellular G-protein to elicit a cellular response. GRKs phosphorylate cardiac receptors, essentially inhibiting receptor-mediated signaling and, thus, serving in a manner analogous to natural β -blockade. The *GRK5* p.Q41L polymorphism occurs commonly in African Americans, with an allele frequency $>30\%$. However, it rarely occurs in Caucasians. The L41 allele has been found to more effectively uncouple agonist-mediated receptor signaling and has been associated with increased transplant-free survival in African Americans with heart failure [320]. Patients with the L41 allele derived no benefit from β -blocker therapy, presumably, because they already have inherent downregulation of ADRB1 receptor signaling [321]. However, in patients with the *GRK5* 41QQ genotype, which is associated with a poor prognosis, treatment with β -blocker therapy significantly improved transplant-free survival [320].

The dopamine and serotonin receptors targeted by antipsychotics are also G-protein-coupled receptors (GPCRs) and signal to effector proteins through intracellular G-protein subunits. Regulators of G-protein signaling shorten the duration of neurotransmitter-mediated receptor signaling through the GPCRs. The regulator of G-protein signaling 4 (RGS4) is one such regulator, and it regulates the activity of the GPCRs. The gene that encodes RSG4 had been identified as a vulnerability gene for schizophrenia [322,323], and variants of *RSG4* have been studied as predictors for antipsychotic treatment response.

Three SNPs of *RSG4* have been reported to confer differential treatment responses in three ethnic groups. In patients of African descent, those with the CC genotype of the rs951439 SNP had longer (391 days) and better (21% improvement based on the Positive and Negative Syndrome Scale [PANSS]) response to perphenazine than ziprasidone (124 days and 5% worsening, respectively). On the other hand, the same patient population with the TT genotype of the rs2842030 SNP responded better to perphenazine (24% improvement in the PANSS) than to quetiapine, risperidone, and ziprasidone. A sharp contrast in association was shown in patients with European descent, in which risperidone treatment resulted in better response with the TT genotype of the rs951349 SNP and GG genotype of the rs2842030 SNP [324]. In 120 schizophrenic patients of Chinese descent, rs2661319 of *RSG4* was found to predict response to risperidone treatment [325]. These data with *RSG4* polymorphisms underscore the importance of patient population stratification by ethnicity in pharmacogenomics investigations. It is also noteworthy that the investigators of the Chinese study also had reported in other studies that polymorphisms affecting the dopamine D₂ receptor (Ser311Cys), D₃ receptor (Ser9Gly), and 5-HT_{2A} receptor (102-T/C) predict treatment response to risperidone [326–328]. Whether evaluating a combination of SNPs could result in better response prediction remains to be investigated.

The alpha adducin (ADD1) gene encodes for α -adducin, a cytoskeletal protein involved in signal transduction and renal sodium transport. The *ADD1* p.G460W variant is associated with greater renal sodium–potassium pump activity, renal sodium retention, and salt-sensitive hypertension [329,330]. Given its role in regulating sodium reabsorption and potentially mediating increased hypertension risk, the *ADD1* gene has been studied for its contribution to diuretic response. Although the 460W allele has been linked to greater blood

pressure reduction with thiazide diuretics, the data are inconsistent [329,331]. The *ADD1* gene appears to interact with other genes involved in renal sodium reabsorption, including the neural precursor cell expressed, developmentally downregulated 4-like (NEDD4L) and lysine-deficient protein kinase 1 (WNK) genes [329]. This may explain the inconsistencies in the data when *ADD1* is analyzed alone rather than in the context of other genes involved in renal sodium handling, and illustrates the likely contribution of multiple genes to the efficacy of many drugs.

Enzyme Genes

VKORC1 is the target site for warfarin. Specifically, warfarin inhibits VKORC1 to prevent regeneration of a reduced form of vitamin K necessary for clotting factor activation. Two common variants, -1639G>A (rs9923231) and -1173C>T (rs9934438) in the *VKORC1* regulatory regions, are associated with reduced gene expression [332]. The frequency of 1639A allele of rs9923231 is highest in Asians (~90%) and lowest in persons of African descent (10%), with an intermittent frequency in populations of European descent (~40%) [333].

Numerous studies have documented the association between *VKORC1* genotype and warfarin-dose requirements. The -1639AA, AG, and GG genotypes are associated with average warfarin-dose requirements of approximately 3, 5, and 6 mg/day, respectively. The two SNPs are equally predictive for predicting warfarin-dose requirement. Recent data suggest that dosing based on one of the *VKORC1* SNP, in addition, to *CYP2C9* genotype, leads to more accurate dose prediction and may reduce the risk for adverse clinical outcomes early in the course of warfarin therapy [334,335]. The *VKORC1* genotype is described in detail in Chapter 6.

The angiotensin-converting enzyme (ACE) gene has been widely studied for its effects on ACE inhibitor response. An insertion/deletion

(I/D) polymorphism in intron 16 of the *ACE* gene results in the presence or absence of a 287-base-pair fragment. The *ACE* D allele has been linked consistently to higher plasma concentrations of ACE, the enzyme responsible for the conversion of angiotensin I to the potent vasoconstrictor angiotensin II [336]. Given its association with ACE concentrations, a number of investigators have examined whether the I/D polymorphism contributes to the interpatient variability in ACE inhibitor response. However, much of the data with the I/D polymorphism and blood pressure response to ACE inhibitors are inconsistent and even conflicting, with some studies demonstrating greater response with the D/D genotype, whereas others have shown greater response with the I/I genotype, and further studies showing no association. In one of the largest pharmacogenetic studies to date, including nearly 38,000 patients, there was no association between the *ACE* I/D genotype and either blood pressure response or cardiovascular or renal outcomes with antihypertensive therapy [337].

Numerous polymorphic proteins are involved in the complex signaling pathway of the renin-angiotensin system, including renin, angiotensinogen, the angiotensin II type 1 receptor, bradykinin, and aldosterone synthase. Thus, a likely explanation for the inconsistent data with the *ACE* gene and ACE inhibitor response in hypertension is that a single polymorphism provides minimal contribution to ACE inhibitor response. Rather, ACE inhibitor response may be best determined by a combination of multiple polymorphisms occurring in multiple genes involved in the renin-angiotensin pathway.

The data with the *ACE* I/D genotype and ACE inhibitor response in patients with heart failure are more compelling. In this population, the *ACE* D allele has been associated with an increased risk for cardiac transplant or death [338–342]. As described in detail in Chapter 6, the detrimental effect of the *ACE* D allele on transplant-free survival appears greatest among patients who are

taking lower than recommended doses of ACE inhibitors. These data suggest that maximizing the ACE inhibitor dose may be necessary in *ACE* D allele carriers to attenuate the harmful effects of this allele [342,343].

Metformin is an antidiabetic drug that works in part by activating adenosine monophosphate-activated protein kinase (AMPK), which is a master regulator of cell and body energy homeostasis and glucose uptake in skeletal muscle [344]. The ataxia-telangiectasia mutated (ATM) is a DNA repair gene that acts upstream of AMPK and appear necessary for metformin action [345]. A GWAS identified a significant association between metformin response in type 2 diabetic patients and a polymorphism in a locus containing the *ATM* gene [345]. The potential significance of ATM for metformin response is further described in Chapter 9.

The catecho-O-methyltransferase (COMT) enzyme is a key modulator of the adrenergic and dopaminergic systems, which play a role in pain modulation. A functional SNP in the *COMT* gene, c.472G>A (rs4680), results in a p.V158M substitution with three- to four-fold decrease in enzyme activity. The reduced enzyme activity leads to decreased dopamine degradation and subsequent increases in norepinephrine and epinephrine levels that may be associated with exaggerated levels of pain [346]. Down regulation of endorphins with compensatory upregulation of MOR has also been suggested to be a result of the SNP [347,348]. Cancer patients who are homozygous carriers of the M variant (high pain-sensitivity patients) reportedly required more morphine (155 ± 160 mg/day) than heterozygotes (117 ± 10 mg/day) and homozygous carriers of the wild-type V allele (95 ± 99 mg/day) [349]. These results were replicated in a later study [350]. Reyes-Gibby et al. also reported higher morphine-dosage requirement: 63% and 23%, respectively, for satisfactory pain control in patients with the *COMT* Val/Val and Val/Met genotypes compared to carriers of

Met/Met genotype ($P = .02$) [311]. The increased morphine toxicity reported in the patient by Madadi et al. had a *COMT* haplotype CCG (rs4633, rs4818, rs4680) and a *G/G* genotype for the *OPRM1* (rs1799971, A118G) SNP, in addition, to the *ABCB1* haplotype [186]. This again highlights the importance of multiple genes in mediating drug disposition and effect and the need of genotyping multiple functional polymorphisms in pharmacogenomic studies. Interestingly, in addition to reporting negative association for the *OPRM1* A118G (rs1799971) polymorphism that was discussed in the section on drug-target genes in patient management, Landau et al. also did not find a correlation between fentanyl therapeutic outcomes with *COMT* genotype [319].

Ion Channel Genotype

One of the most often cited examples of ion channel genes with consequences for drug response are genes for the pore-forming channel proteins that affect potassium and sodium transport across the cardiac cell membrane. Mutations in cardiac ion channel genes predispose individuals to congenital long-QT syndrome. Moreover, there is evidence that these mutations may increase the risk for drug-induced torsades de pointes [351,352]. This is discussed in detail in Chapter 6.

The large-conductance calcium and voltage-dependent potassium (BK) channel is another example of an ion channel with genetic contributions to drug response. The BK channel is found in vascular smooth muscle and consists of pore-forming- α and regulatory- $\beta 1$ subunits. The $\beta 1$ subunit enhances calcium sensitivity and decreases smooth muscle cell excitability, thus attenuating smooth muscle contraction. The *KCNMB1* gene encodes for the BK channel $\beta 1$ subunit. A common SNP in the *KCNMB1* gene, Glu65Lys, results in a gain of function of the channel and increased calcium sensitivity compared to the wild type [353]. Given its role in mediating calcium sensitivity, the *KCNMB1*

gene was examined for its effect on response to the calcium channel blocker, verapamil. Among patients with hypertension and coronary heart disease who were started on verapamil, 65Lys allele carriers achieved blood pressure control more rapidly than 65Glu homozygotes, suggesting that the Glu65Lys SNP enhances response to calcium channel blockers and contributes to the interpatient variability in blood pressure reduction during calcium channel blocker therapy [354].

The epithelial sodium channel (ENaC) is another example of an ion channel with genetic contributions to drug response. The ENaC is located in the distal renal tubule and collecting duct of the nephron and serves as the final site for sodium reabsorption. The channel is composed of α , β , and γ subunits, encoded by the *SCNN1A*, *SCNN1B*, and *SCNN1G* genes, respectively. In a healthy volunteer study, SNPs in the *SCNN1B* and *SCNN1G* genes were associated with natriuretic and diuretic responses to single oral doses of loop diuretics. Loop diuretics are commonly prescribed for managing symptoms of fluid overload in heart failure. Whether genes encoding for ENaC subunits influence response to loop diuretics in heart failure remains to be determined. But, given the significant consequences of under- or overdosing loop diuretics in this disease, such information could have significant clinical value.

The potassium inwardly rectifying channel, subfamily J, member 11 gene (*KCNJ11*) and the sulfonylurea receptor gene (*ABCC8*) encode the Kir6.2 and sulfonylurea receptor-1 (SUR1) subunits of pancreatic ATP-sensitive potassium (K_{ATP}) channels, respectively. Activating mutations in the *KCNJ11* and *ABCC8* cause K_{ATP} channels to remain open, which promotes hyperpolarization of the pancreatic β cell membrane and impaired insulin release [355,356]. Sulfonylurea drugs promote K_{ATP} channel closure, thereby attenuating the effects of activating mutations in *KCNJ11* and *ABCC8*. As such, sulfonylureas are especially

effective in patients with *KCNJ11* or *ABCC8* activating mutations [356,357]. Chapter 9 provides a more detailed discussion of these genetic variations and their effects on response to sulfonylurea agents.

CONCLUSION

Variations in genes influencing drug pharmacokinetics and pharmacodynamics often jointly influence drug response, as is the case with warfarin, the dose requirements for which are influenced by both the *CYP2C9* and *VKORC1* genotypes. Thus, when taking a candidate gene approach to discovery of variants impacting drug response, genes encoding proteins involved in determining drug bioavailability (transporter proteins, drug metabolizing enzymes) and response (receptor, enzyme, ion channel, and/or intracellular signaling proteins) should be considered. Genome-wide approaches to identifying determinants of drug response may reveal previously unknown proteins involved in eliciting drug response that represent potential biomarkers for predicting drug effectiveness or risk for toxicity. In addition, proteins involved in disease pathophysiology may represent attractive targets for drug development, as most often demonstrated in the area of oncology.

QUESTIONS FOR DISCUSSION

1. What are examples of drug metabolism and drug-transporter genotypes that affect drug response?
2. What are examples of drugs developed based on an understanding of genes involved in disease pathophysiology?
3. What are examples of drug-target genes with implications for drug response?
4. How might genes for drug metabolism, drug transport, and/or drug-target sites jointly influence drug response?

References

- [1] Database of single nucleotide polymorphisms (dbSNP). National Center for Biotechnology Information, National Library of Medicine, Bethesda (MD). Nucleic Acid Res 2000;28:352–5 (dbSNP Build ID: 137).
- [2] Li J, Zhang L, Zhou H, Stoneking M, Tang K. Global patterns of genetic diversity and signals of natural selection for human ADME genes. *Human Molecular Genetics* 2011;20:528–40.
- [3] Baralt C. UF delivers promise of personalized medicine to heart patients. 2012. UF&Shands News 1992. <http://news.ufl.edu/archive/2012/06/uf-delivers-promise-of-personalized-medicine-to-heart-patients.html>.
- [4] Pulley JM, Denny JC, Peterson JF, Bernard GR, Vnencak-Jones CL, Ramirez AH, et al. Operational implementation of prospective genotyping for personalized medicine: the design of the Vanderbilt PREDICT project. *Clinical Pharmacology and Therapeutics* 2012;92:87–95.
- [5] Zanger UM, Turpeinen M, Klein K, Schwab M. Functional pharmacogenetics/genomics of human cytochromes P450 involved in drug biotransformation. *Analytical and Bioanalytical Chemistry* 2008;392:1093–108.
- [6] Gaedigk A, Ingelman-Sundberg M, Miller NA, Leeder JS, Whirl-Carrillo M, Klein TE, et al. The pharmacogene variation (PharmVar) consortium: incorporation of the human cytochrome P450 (cyp) allele nomenclature database. *Clinical Pharmacology and Therapeutics* 2018;103:399–401.
- [7] Sistonen J, Sajantila A, Lao O, Corander J, Barbuiani G, Fuselli S. CYP2D6 worldwide genetic variation shows high frequency of altered activity variants and no continental structure. *Pharmacogenetics and Genomics* 2007;17:93–101.
- [8] Beoris M, Amos Wilson J, Garces JA, Lukowiak AA. CYP2D6 copy number distribution in the US population. *Pharmacogenetics and Genomics* 2016;26:96–9.
- [9] Bertilsson L, Dahl ML, Sjoqvist F, Aberg-Wistedt A, Humble M, Johansson I, et al. Molecular basis for rational megaprescribing in ultrarapid hydroxylators of debrisoquine. *Lancet* 1993;341:63.
- [10] Dalen P, Dahl ML, Bernal Ruiz ML, Nordin J, Bertilsson L. 10-Hydroxylation of nortriptyline in white persons with 0, 1, 2, 3, and 13 functional CYP2D6 genes. *Clinical Pharmacology and Therapeutics* 1998;63:444–52.
- [11] Johansson I, Lundqvist E, Bertilsson L, Dahl ML, Sjoqvist F, Ingelman-Sundberg M. Inherited amplification of an active gene in the cytochrome P450 CYP2D locus as a cause of ultrarapid metabolism of debrisoquine. *Proceedings of the National Academy of Sciences of the United States of America* 1993;90:11825–9.
- [12] Gaedigk A, Ndjountche L, Divakaran K, Dianne Bradford L, Zineh I, Oberlander TF, et al. Cytochrome P4502D6 (CYP2D6) gene locus heterogeneity: characterization

- of gene duplication events. *Clinical Pharmacology and Therapeutics* 2007;81:242–51.
- [13] Crews KR, Gaedigk A, Dunnenberger HM, Leeder JS, Klein TE, Caudle KE, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for cytochrome P450 2D6 genotype and codeine therapy: 2014 update. *Clinical Pharmacology and Therapeutics* 2014;95:376–82.
 - [14] Twist GP, Gaedigk A, Miller NA, Farrow EG, Willig LK, Dinwiddie DL, et al. Constellation: a tool for rapid, automated phenotype assignment of a highly polymorphic pharmacogene, CYP2D6, from whole-genome sequences. *NPJ Genomic Medicine* 2016;1:15007.
 - [15] Man M, Farmen M, Dumaual C, Teng CH, Moser B, Irie S, et al. Genetic variation in metabolizing enzyme and transporter genes: comprehensive assessment in 3 major East Asian subpopulations with comparison to Caucasians and Africans. *The Journal of Clinical Pharmacology* 2010;50:929–40.
 - [16] Lin KM, Finder E. Neuroleptic dosage for Asians. *American Journal of Psychiatry* 1983;140:490–1.
 - [17] Mihara K, Otani K, Tybring G, Dahl ML, Bertilsson L, Kaneko S. The CYP2D6 genotype and plasma concentrations of mianserin enantiomers in relation to therapeutic response to mianserin in depressed Japanese patients. *Journal of Clinical Psychopharmacology* 1997;17:467–71.
 - [18] Droll K, Bruce-Mensah K, Otton SV, Gaedigk A, Sellers EM, Tyndale RF. Comparison of three CYP2D6 probe substrates and genotype in Ghanaians, Chinese and Caucasians. *Pharmacogenetics* 1998;8:325–33.
 - [19] Masimirembwa C, Persson I, Bertilsson L, Hasler J, Ingelman-Sundberg M. A novel mutant variant of the CYP2D6 gene (CYP2D6*17) common in a black African population: association with diminished debrisoquine hydroxylase activity. *British Journal of Clinical Pharmacology* 1996;42:713–9.
 - [20] Zanger UM, Schwab M. Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation. *Pharmacology and Therapeutics* 2013;138:103–41.
 - [21] Barclay ML, Sawyers SM, Begg EJ, Zhang M, Roberts RL, Kennedy MA, et al. Correlation of CYP2D6 genotype with perhexiline phenotypic metabolizer status. *Pharmacogenetics* 2003;13:627–32.
 - [22] Lee JT, Kroemer HK, Silberstein DJ, Funck-Brentano C, Lineberry MD, Wood AJ, et al. The role of genetically determined polymorphic drug metabolism in the beta-blockade produced by propafenone. *New England Journal of Medicine* 1990;322:1764–8.
 - [23] Dahl-Puustinen ML, Liden A, Alm C, Nordin C, Bertilsson L. Disposition of perphenazine is related to polymorphic debrisoquin hydroxylation in human beings. *Clinical Pharmacology and Therapeutics* 1989;46:78–81.
 - [24] Spina E, Ancione M, Di Rosa AE, Meduri M, Caputi AP. Polymorphic debrisoquine oxidation and acute neuroleptic-induced adverse effects. *European Journal of Clinical Pharmacology* 1992;42:347–8.
 - [25] Gasche Y, Daali Y, Fathi M, Chiappe A, Cottini S, Dayer P, et al. Codeine intoxication associated with ultrarapid CYP2D6 metabolism. *New England Journal of Medicine* 2004;351:2827–31.
 - [26] Koren G, Cairns J, Chitayat D, Gaedigk A, Leeder SJ. Pharmacogenetics of morphine poisoning in a breast-fed neonate of a codeine-prescribed mother. *Lancet* 2006;368:704.
 - [27] Ciszkowski C, Madadi P, Phillips MS, Lauwers AE, Koren G. Codeine, ultrarapid-metabolism genotype, and postoperative death. *New England Journal of Medicine* 2009;361:827–8.
 - [28] Voronov P, Przybylo HJ, Jagannathan N. Apnea in a child after oral codeine: a genetic variant - an ultra-rapid metabolizer. *Paediatric Anaesthesia* 2007;17:684–7.
 - [29] Kelly LE, Rieder M, van den Anker J, Malkin B, Ross C, Neely MN, et al. More codeine fatalities after tonsillectomy in North American children. *Pediatrics* 2012;129:e1343–7.
 - [30] Madadi P, Ross CJ, Hayden MR, Carleton BC, Gaedigk A, Leeder JS, et al. Pharmacogenetics of neonatal opioid toxicity following maternal use of codeine during breastfeeding: a case-control study. *Clinical Pharmacology and Therapeutics* 2009;85:31–5.
 - [31] Tobias JD, Green TP, Cote CJ. Section on a, pain M, committee on D. Codeine: time to say “No”. *Pediatrics* 2016;138:e20162396.
 - [32] Samer CF, Daali Y, Wagner M, Hopfgartner G, Eap CB, Rebsamen MC, et al. Genetic polymorphisms and drug interactions modulating CYP2D6 and CYP3A activities have a major effect on oxycodone analgesic efficacy and safety. *British Journal of Pharmacology* 2010;160:919–30.
 - [33] Madadi P, Hildebrandt D, Gong IY, Schwarz UI, Ciszkowski C, Ross CJ, et al. Fatal hydrocodone overdose in a child: pharmacogenetics and drug interactions. *Pediatrics* 2010;126:e986–9.
 - [34] Elkalioubie A, Allorge D, Robriquet L, Wiart JF, Garat A, Broly F, et al. Near-fatal tramadol cardiotoxicity in a CYP2D6 ultrarapid metabolizer. *European Journal of Clinical Pharmacology* 2011;67:855–8.
 - [35] Stamer UM, Stuber F, Muders T, Musshoff F. Respiratory depression with tramadol in a patient with renal impairment and CYP2D6 gene duplication. *Anesthesia and Analgesia* 2008;107:926–9.
 - [36] Garcia-Quetglas E, Azanza JR, Sadaba B, Munoz MJ, Gil I, Campanero MA. Pharmacokinetics of tramadol enantiomers and their respective phase I metabolites in relation to CYP2D6 phenotype. *Pharmacological Research* 2007;55:122–30.

- [37] Sindrup SH, Brosen K, Bjerring P, Arendt-Nielsen L, Larsen U, Angelo HR, et al. Codeine increases pain thresholds to copper vapor laser stimuli in extensive but not poor metabolizers of sparteine. *Clinical Pharmacology and Therapeutics* 1990;48:686–93.
- [38] Crews KR, Gaedigk A, Dunnenberger HM, Klein TE, Shen DD, Callaghan JT, et al. Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines for codeine therapy in the context of cytochrome P450 2D6 (CYP2D6) genotype. *Clinical Pharmacology and Therapeutics* 2012;91:321–6.
- [39] Wijnen PA, Limantoro I, Drent M, Bekers O, Kuijpers PM, Koek GH. Depressive effect of an antidepressant: therapeutic failure of venlafaxine in a case lacking CYP2D6 activity. *Annals of Clinical Biochemistry* 2009;46:527–30.
- [40] Goetz MP, Rae JM, Suman VJ, Safgren SL, Ames MM, Visscher DW, et al. Pharmacogenetics of tamoxifen biotransformation is associated with clinical outcomes of efficacy and hot flashes. *Journal of Clinical Oncology* 2005;23:9312–8.
- [41] Kiyotani K, Mushiroda T, Hosono N, Tsunoda T, Kubo M, Aki F, et al. Lessons for pharmacogenomics studies: association study between CYP2D6 genotype and tamoxifen response. *Pharmacogenetics and Genomics* 2010;20:565–8.
- [42] Schroth W, Winter S, Murdter T, Schaeffeler E, Eccles D, Eccles B, et al. Improved prediction of endoxifen metabolism by CYP2D6 genotype in breast cancer patients treated with tamoxifen. *Frontiers in Pharmacology* 2017;8:582.
- [43] Goetz MP, Sangkuhl K, Guchelaar HJ, Schwab M, Province M, Whirl-Carrillo M, et al. Clinical pharmacogenetics implementation consortium (CPIC) guideline for CYP2D6 and tamoxifen therapy. *Clinical Pharmacology and Therapeutics* 2018;103:770–7.
- [44] Bertilsson L, Aberg-Wistedt A, Gustafsson LL, Nordin C. Extremely rapid hydroxylation of debrisoquine: a case report with implication for treatment with nortriptyline and other tricyclic antidepressants. *Therapeutic Drug Monitoring* 1985;7:478–80.
- [45] Kawanishi C, Lundgren S, Agren H, Bertilsson L. Increased incidence of CYP2D6 gene duplication in patients with persistent mood disorders: ultrarapid metabolism of antidepressants as a cause of nonresponse. A pilot study. *European Journal of Clinical Pharmacology* 2004;59:803–7.
- [46] Rau T, Wohlleben G, Wuttke H, Thuerauf N, Lunkenheimer J, Lanczik M, et al. CYP2D6 genotype: impact on adverse effects and nonresponse during treatment with antidepressants—a pilot study. *Clinical Pharmacology and Therapeutics* 2004;75:386–93.
- [47] Candiotti KA, Birnbach DJ, Lubarsky DA, Nhuch F, Kamat A, Koch WH, et al. The impact of pharmacogenomics on postoperative nausea and vomiting: do CYP2D6 allele copy number and polymorphisms affect the success or failure of ondansetron prophylaxis? *Anesthesiology* 2005;102:543–9.
- [48] Hamelin BA, Bouayad A, Methot J, Jobin J, Desgagnés P, Poirier P, et al. Significant interaction between the nonprescription antihistamine diphenhydramine and the CYP2D6 substrate metoprolol in healthy men with high or low CYP2D6 activity. *Clinical Pharmacology and Therapeutics* 2000;67:466–77.
- [49] Alfaro CL, Lam YW, Simpson J, Ereshefsky L. CYP2D6 status of extensive metabolizers after multiple-dose fluoxetine, fluvoxamine, paroxetine, or sertraline. *Journal of Clinical Psychopharmacology* 1999;19:155–63.
- [50] Gryn SE, Teft WA, Kim RB. Profound reduction in the tamoxifen active metabolite endoxifen in a patient on phenytoin for epilepsy compared with a CYP2D6 genotype matched cohort. *Pharmacogenetics and Genomics* 2014;24:367–9.
- [51] Teft WA, Mansell SE, Kim RB. Endoxifen, the active metabolite of tamoxifen, is a substrate of the efflux transporter P-glycoprotein (multidrug resistance 1). *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2011;39:558–62.
- [52] Shah RR, Gaedigk A, A LL, Eichelbaum M, Stingl J, Smith RL. CYP450 genotype and pharmacogenetic association studies: a critical appraisal. *Pharmacogenomics* 2016;17:259–75.
- [53] Lam YW, Gaedigk A, Ereshefsky L, Alfaro CL, Simpson J. CYP2D6 inhibition by selective serotonin reuptake inhibitors: analysis of achievable steady-state plasma concentrations and the effect of ultrarapid metabolism at CYP2D6. *Pharmacotherapy* 2002;22:1001–6.
- [54] Sim SC, Risinger C, Dahl ML, Aklilu E, Christensen M, Bertilsson L, et al. A common novel CYP2C19 gene variant causes ultrarapid drug metabolism relevant for the drug response to proton pump inhibitors and antidepressants. *Clinical Pharmacology and Therapeutics* 2006;79:103–13.
- [55] Yamada S, Onda M, Kato S, Matsuda N, Matsuhisa T, Yamada N, et al. Genetic differences in CYP2C19 single nucleotide polymorphisms among four Asian populations. *Journal of Gastroenterology* 2001;36:669–72.
- [56] Ghoneim MM, Korttila K, Chiang CK, Jacobs L, Schoenwald RD, Mewaldt SP, et al. Diazepam effects and kinetics in caucasians and orientals. *Clinical Pharmacology and Therapeutics* 1981;29:749–56.
- [57] Kumana CR, Lauder IJ, Chan M, Ko W, Lin HJ. Differences in diazepam pharmacokinetics in Chinese and white Caucasians—relation to body lipid stores. *European Journal of Clinical Pharmacology* 1987;32:211–5.

- [58] Furuta T, Sugimoto M, Shirai N, Ishizaki T. CYP2C19 pharmacogenomics associated with therapy of *Helicobacter pylori* infection and gastro-esophageal reflux diseases with a proton pump inhibitor. *Pharmacogenomics* 2007;8:1199–210.
- [59] Andersson T, Holmberg J, Rohss K, Walan A. Pharmacokinetics and effect on caffeine metabolism of the proton pump inhibitors, omeprazole, lansoprazole, and pantoprazole. *British Journal of Clinical Pharmacology* 1998;45:369–75.
- [60] Baldwin RM, Ohlsson S, Pedersen RS, Mwinyi J, Ingelman-Sundberg M, Eliasson E, et al. Increased omeprazole metabolism in carriers of the CYP2C19*17 allele; a pharmacokinetic study in healthy volunteers. *British Journal of Clinical Pharmacology* 2008;65:767–74.
- [61] Furuta T, Ohashi K, Kosuge K, Zhao XJ, Takashima M, Kimura M, et al. CYP2C19 genotype status and effect of omeprazole metabolism in carriers of the CYP2C19*17 allele; a pharmacokinetic study in healthy volunteers. *Clinical Pharmacology and Therapeutics* 1999;65:552–61.
- [62] Lou HY, Chang CC, Sheu MT, Chen YC, Ho HO. Optimal dose regimens of esomeprazole for gastric acid suppression with minimal influence of the CYP2C19 polymorphism. *European Journal of Clinical Pharmacology* 2009;65:55–64.
- [63] Qiao HL, Hu YR, Tian X, Jia LJ, Gao N, Zhang LR, et al. Pharmacokinetics of three proton pump inhibitors in Chinese subjects in relation to the CYP2C19 genotype. *European Journal of Clinical Pharmacology* 2006;62:107–12.
- [64] Furuta T, Ohashi K, Kamata T, Takashima M, Kosuge K, Kawasaki T, et al. Effect of genetic differences in omeprazole metabolism on cure rates for *Helicobacter pylori* infection and peptic ulcer. *Annals of Internal Medicine* 1998;129:1027–30.
- [65] Tanigawara Y, Aoyama N, Kita T, Shirakawa K, Komada F, Kasuga M, et al. CYP2C19 genotype-related efficacy of omeprazole for the treatment of infection caused by *Helicobacter pylori*. *Clinical Pharmacology and Therapeutics* 1999;66:528–34.
- [66] Kawabata H, Habu Y, Tomioka H, Kutsumi H, Kobayashi M, Oyasu K, et al. Effect of different proton pump inhibitors, differences in CYP2C19 genotype and antibiotic resistance on the eradication rate of *Helicobacter pylori* infection by a 1-week regimen of proton pump inhibitor, amoxicillin and clarithromycin. *Alimentary Pharmacology and Therapeutics* 2003;17:259–64.
- [67] Gawronska-Szklarz B, Siuda A, Kurzawski M, Bielicki D, Marlicz W, Drozdziak M. Effects of CYP2C19, MDR1, and interleukin 1-B gene variants on the eradication rate of *Helicobacter pylori* infection by triple therapy with pantoprazole, amoxicillin, and metronidazole. *European Journal of Clinical Pharmacology* 2010;66:681–7.
- [68] Furuta T, Shirai N, Takashima M, Xiao F, Hanai H, Sugimura H, et al. Effect of genotypic differences in CYP2C19 on cure rates for *Helicobacter pylori* infection by triple therapy with a proton pump inhibitor, amoxicillin, and clarithromycin. *Clinical Pharmacology and Therapeutics* 2001;69:158–68.
- [69] Furuta T, Shirai N, Kodaira M, Sugimoto M, Nogaki A, Kuriyama S, et al. Pharmacogenomics-based tailored versus standard therapeutic regimen for eradication of *H. pylori*. *Clinical Pharmacology and Therapeutics* 2007;81:521–8.
- [70] Hunfeld NG, Mathot RA, Touw DJ, van Schaik RH, Mulder PG, Franck PF, et al. Effect of CYP2C19*2 and *17 mutations on pharmacodynamics and kinetics of proton pump inhibitors in Caucasians. *British Journal of Clinical Pharmacology* 2008;65:752–60.
- [71] Kurzawski M, Gawronska-Szklarz B, Wrzesniewska J, Siuda A, Starzynska T, Drozdziak M. Effect of CYP2C19*17 gene variant on *Helicobacter pylori* eradication in peptic ulcer patients. *European Journal of Clinical Pharmacology* 2006;62:877–80.
- [72] Wang G, Lei HP, Li Z, Tan ZR, Guo D, Fan L, et al. The CYP2C19 ultra-rapid metabolizer genotype influences the pharmacokinetics of voriconazole in healthy male volunteers. *European Journal of Clinical Pharmacology* 2009;65:281–5.
- [73] Hamadeh IS, Klinker KP, Borgert SJ, Richards AI, Li W, Mangal N, et al. Impact of the CYP2C19 genotype on voriconazole exposure in adults with invasive fungal infections. *Pharmacogenetics and Genomics* 2017;27:190–6.
- [74] Lamoureux F, Duflo T, Woillard JB, Metsu D, Pereira T, Compagnon P, et al. Impact of CYP2C19 genetic polymorphisms on voriconazole dosing and exposure in adult patients with invasive fungal infections. *International Journal of Antimicrobial Agents* 2016;47:124–31.
- [75] Wang T, Zhu H, Sun J, Cheng X, Xie J, Dong H, et al. Efficacy and safety of voriconazole and CYP2C19 polymorphism for optimised dosage regimens in patients with invasive fungal infections. *International Journal of Antimicrobial Agents* 2014;44:436–42.
- [76] Rudberg I, Mohebi B, Hermann M, Refsum H, Molden E. Impact of the ultrarapid CYP2C19*17 allele on serum concentration of escitalopram in psychiatric patients. *Clinical Pharmacology and Therapeutics* 2008;83:322–7.
- [77] Gurbel PA, Bergmeijer TO, Tantry US, ten Berg JM, Angiolillo DJ, James S, et al. The effect of CYP2C19 gene polymorphisms on the pharmacokinetics and pharmacodynamics of prasugrel 5-mg, prasugrel 10-mg and clopidogrel 75-mg in patients with coronary artery disease. *Thrombosis and Haemostasis* 2014;112:589–97.

- [78] Collet JP, Hulot JS, Pena A, Villard E, Esteve JB, Silvain J, et al. Cytochrome P450 2C19 polymorphism in young patients treated with clopidogrel after myocardial infarction: a cohort study. *Lancet* 2009;373:309–17.
- [79] Hulot JS, Bura A, Villard E, Azizi M, Remones V, Goyenvalle C, et al. Cytochrome P450 2C19 loss-of-function polymorphism is a major determinant of clopidogrel responsiveness in healthy subjects. *Blood* 2006;108:2244–7.
- [80] Kim KA, Park PW, Hong SJ, Park JY. The effect of CYP2C19 polymorphism on the pharmacokinetics and pharmacodynamics of clopidogrel: a possible mechanism for clopidogrel resistance. *Clinical Pharmacology and Therapeutics* 2008;84:236–42.
- [81] Mega JL, Close SL, Wiviott SD, Shen L, Hockett RD, Brandt JT, et al. Cytochrome p-450 polymorphisms and response to clopidogrel. *New England Journal of Medicine* 2009;360:354–62.
- [82] Umemura K, Furuta T, Kondo K. The common gene variants of CYP2C19 affect pharmacokinetics and pharmacodynamics in an active metabolite of clopidogrel in healthy subjects. *Journal of Thrombosis and Haemostasis* 2008;6:1439–41.
- [83] Shuldiner AR, O'Connell JR, Bliden KP, Gandhi A, Ryan K, Horenstein RB, et al. Association of cytochrome P450 2C19 genotype with the antiplatelet effect and clinical efficacy of clopidogrel therapy. *Journal of the American Medical Association* 2009;302:849–57.
- [84] Sorich MJ, Rowland A, McKinnon RA, Wiese MD. CYP2C19 genotype has a greater effect on adverse cardiovascular outcomes following percutaneous coronary intervention and in Asian populations treated with clopidogrel: a meta-analysis. *Circulation Cardiovascular Genetics* 2014;7:895–902.
- [85] Frere C, Cuisset T, Gaborit B, Alessi MC, Hulot JS. The CYP2C19*17 allele is associated with better platelet response to clopidogrel in patients admitted for non-ST acute coronary syndrome. *Journal of Thrombosis and Haemostasis* 2009;7:1409–11.
- [86] Sibbing D, Koch W, Gebhard D, Schuster T, Braun S, Stegherr J, et al. Cytochrome 2C19*17 allelic variant, platelet aggregation, bleeding events, and stent thrombosis in clopidogrel-treated patients with coronary stent placement. *Circulation* 2010;121:512–8.
- [87] Tiroch KA, Sibbing D, Koch W, Roosen-Runge T, Mehilli J, Schomig A, et al. Protective effect of the CYP2C19*17 polymorphism with increased activation of clopidogrel on cardiovascular events. *American Heart Journal* 2010;160:506–12.
- [88] Mega JL, Simon T, Collet JP, Anderson JL, Antman EM, Bliden K, et al. Reduced-function CYP2C19 genotype and risk of adverse clinical outcomes among patients treated with clopidogrel predominantly for PCI: a meta-analysis. *Journal of the American Medical Association* 2010;304:1821–30.
- [89] Gurbel PA, Tantry US, Shuldiner AR. Letter by Gurbel et al regarding article, "Cytochrome 2C19*17 allelic variant, platelet aggregation, bleeding events, and stent thrombosis in clopidogrel-treated patients with coronary stent placement". *Circulation* 2010;122:e478. author reply e9.
- [90] Scott SA, Sangkuhl K, Stein CM, Hulot JS, Mega JL, Roden DM, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for CYP2C19 genotype and clopidogrel therapy: 2013 update. *Clinical Pharmacology and Therapeutics* 2013;94:317–23.
- [91] Geisler T, Schaeffeler E, Dippon J, Winter S, Buse V, Bischofs C, et al. CYP2C19 and nongenetic factors predict poor responsiveness to clopidogrel loading dose after coronary stent implantation. *Pharmacogenomics* 2008;9:1251–9.
- [92] Hulot JS, Collet JP, Silvain J, Pena A, Bellemain-Appaix A, Barthelemy O, et al. Cardiovascular risk in clopidogrel-treated patients according to cytochrome P450 2C19*2 loss-of-function allele or proton pump inhibitor coadministration: a systematic meta-analysis. *Journal of the American College of Cardiology* 2010;56:134–43.
- [93] Bauer T, Bouman HJ, van Werkum JW, Ford NF, ten Berg JM, Taubert D. Impact of CYP2C19 variant genotypes on clinical efficacy of antiplatelet treatment with clopidogrel: systematic review and meta-analysis. *BMJ* 2011;343:d4588.
- [94] Holmes MV, Perel P, Shah T, Hingorani AD, Casas JP. CYP2C19 genotype, clopidogrel metabolism, platelet function, and cardiovascular events: a systematic review and meta-analysis. *Journal of the American Medical Association* 2011;306:2704–14.
- [95] Kwok CS, Loke YK. Meta-analysis: the effects of proton pump inhibitors on cardiovascular events and mortality in patients receiving clopidogrel. *Alimentary Pharmacology and Therapeutics* 2010;31:810–23.
- [96] Stubbins MJ, Harries LW, Smith G, Tarbit MH, Wolf CR. Genetic analysis of the human cytochrome P450 CYP2C9 locus. *Pharmacogenetics* 1996;6:429–39.
- [97] Perera MA, Gamazon E, Cavallari LH, Patel SR, Poindexter S, Kittles RA, et al. The missing association: sequencing-based discovery of novel SNPs in VKORC1 and CYP2C9 that affect warfarin dose in African Americans. *Clinical Pharmacology and Therapeutics* 2011;89:408–15.
- [98] Allabi AC, Gala JL, Horsmans Y, Babaoglu MO, Bozkurt A, Heusterspreute M, et al. Functional impact of CYP2C95, CYP2C96, CYP2C98, and CYP2C911 in vivo among black Africans. *Clinical Pharmacology and Therapeutics* 2004;76:113–8.
- [99] Dickmann LJ, Rettie AE, Kneller MB, Kim RB, Wood AJ, Stein CM, et al. Identification and functional characterization of a new CYP2C9 variant (CYP2C9*5) expressed among African Americans. *Molecular Pharmacology* 2001;60:382–7.

- [100] Scott SA, Jaremko M, Lubitz SA, Kornreich R, Halperin JL, Desnick RJ. CYP2C9*8 is prevalent among African-Americans: implications for pharmacogenetic dosing. *Pharmacogenomics* 2009;19:1243–55.
- [101] Takahashi H, Kashima T, Nomoto S, Iwade K, Tainaka H, Shimizu T, et al. Comparisons between in-vitro and in-vivo metabolism of (S)-warfarin: catalytic activities of cDNA-expressed CYP2C9, its Leu359 variant and their mixture versus unbound clearance in patients with the corresponding CYP2C9 genotypes. *Pharmacogenetics* 1998;8:365–73.
- [102] Figueiras A, Estany-Gestal A, Aguirre C, Ruiz B, Vidal X, Carvajal A, et al. CYP2C9 variants as a risk modifier of NSAID-related gastrointestinal bleeding: a case-control study. *Pharmacogenetics and Genomics* 2016;26:66–73.
- [103] Liu Y, Jeong H, Takahashi H, Drozda K, Patel SR, Shapiro NL, et al. Decreased warfarin clearance associated with the CYP2C9 R150H (*8) polymorphism. *Clinical Pharmacology and Therapeutics* 2012;91:660–5.
- [104] Rieder MJ, Reiner AP, Gage BF, Nickerson DA, Eby CS, McLeod HL, et al. Effect of VKORC1 haplotypes on transcriptional regulation and warfarin dose. *New England Journal of Medicine* 2005;352:2285–93.
- [105] Anderson JL, Horne BD, Stevens SM, Grove AS, Barton S, Nicholas ZP, et al. Randomized trial of genotype-guided versus standard warfarin dosing in patients initiating oral anticoagulation. *Circulation* 2007;116:2563–70.
- [106] Sconce EA, Khan TI, Wynne HA, Avery P, Monkhouse L, King BP, et al. The impact of CYP2C9 and VKORC1 genetic polymorphism and patient characteristics upon warfarin dose requirements: proposal for a new dosing regimen. *Blood* 2005;106:2329–33.
- [107] Kim MJ, Huang SM, Meyer UA, Rahman A, Lesko LJ. A regulatory science perspective on warfarin therapy: a pharmacogenetic opportunity. *The Journal of Clinical Pharmacology* 2009;49:138–46.
- [108] Higashi MK, Veenstra DL, Kondo LM, Wittkowsky AK, Srinouanprachanh SL, Farin FM, et al. Association between CYP2C9 genetic variants and anticoagulation-related outcomes during warfarin therapy. *Journal of the American Medical Association* 2002;287:1690–8.
- [109] Aithal GP, Day CP, Kesteven PJ, Daly AK. Association of polymorphisms in the cytochrome P450 CYP2C9 with warfarin dose requirement and risk of bleeding complications. *Lancet* 1999;353:717–9.
- [110] Lindh JD, Holm L, Andersson ML, Rane A. Influence of CYP2C9 genotype on warfarin dose requirements—a systematic review and meta-analysis. *European Journal of Clinical Pharmacology* 2009;65:365–75.
- [111] Steward DJ, Haining RL, Henne KR, Davis G, Rushmore TH, Trager WF, et al. Genetic association between sensitivity to warfarin and expression of CYP2C9*3. *Pharmacogenetics* 1997;7:361–7.
- [112] Limdi NA, Arnett DK, Goldstein JA, Beasley TM, McGwin G, Adler BK, et al. Influence of CYP2C9 and VKORC1 on warfarin dose, anticoagulation attainment and maintenance among European-Americans and African-Americans. *Pharmacogenomics* 2008;9:511–26.
- [113] Limdi NA, McGwin G, Goldstein JA, Beasley TM, Arnett DK, Adler BK, et al. Influence of CYP2C9 and VKORC1 1173C/T genotype on the risk of hemorrhagic complications in African-American and European-American patients on warfarin. *Clinical Pharmacology and Therapeutics* 2008;83:312–21.
- [114] Cavallari LH, Langae TY, Momary KM, Shapiro NL, Nutescu EA, Coty WA, et al. Genetic and clinical predictors of warfarin dose requirements in African Americans. *Clinical Pharmacology and Therapeutics* 2010;87:459–64.
- [115] Zhong SL, Yu XY, Liu Y, Xu D, Mai LP, Tan HH, et al. Integrating interacting drugs and genetic variations to improve the predictability of warfarin maintenance dose in Chinese patients. *Pharmacogenetics and Genomics* 2012;22:176–82.
- [116] Cavaco I, Stromberg-Norklit J, Kaneko A, Msellem MI, Dahoma M, Ribeiro VL, et al. CYP2C8 polymorphism frequencies among malaria patients in Zanzibar. *European Journal of Clinical Pharmacology* 2005;61:15–8.
- [117] Soyama A, Saito Y, Komamura K, Ueno K, Kamakura S, Ozawa S, et al. Five novel single nucleotide polymorphisms in the CYP2C8 gene, one of which induces a frame-shift. *Drug Metabolism Pharmacokinetics* 2002;17:374–7.
- [118] Edwards RJ, Adams DA, Watts PS, Davies DS, Boobis AR. Development of a comprehensive panel of antibodies against the major xenobiotic metabolising forms of cytochrome P450 in humans. *Biochemical Pharmacology* 1998;56:377–87.
- [119] Garcia-Martin E, Martinez C, Tabares B, Frias J, Agundez JA. Interindividual variability in ibuprofen pharmacokinetics is related to interaction of cytochrome P450 2C8 and 2C9 amino acid polymorphisms. *Clinical Pharmacology and Therapeutics* 2004;76:119–27.
- [120] Martinez C, Garcia-Martin E, Blanco G, Gamito FJ, Ladero JM, Agundez JA. The effect of the cytochrome P450 CYP2C8 polymorphism on the disposition of (R)-ibuprofen enantiomer in healthy subjects. *British Journal of Clinical Pharmacology* 2005;59:62–9.

- [121] Lundblad MS, Stark K, Eliasson E, Oliw E, Rane A. Biosynthesis of epoxyeicosatrienoic acids varies between polymorphic CYP2C enzymes. *Biochemical and Biophysical Research Communications* 2005;327:1052–7.
- [122] Niemi M, Leathart JB, Neuvonen M, Backman JT, Daly AK, Neuvonen PJ. Polymorphism in CYP2C8 is associated with reduced plasma concentrations of repaglinide. *Clinical Pharmacology and Therapeutics* 2003;74:380–7.
- [123] Niemi M, Backman JT, Kajosaari LI, Leathart JB, Neuvonen M, Daly AK, et al. Polymorphic organic anion transporting polypeptide 1B1 is a major determinant of repaglinide pharmacokinetics. *Clinical Pharmacology and Therapeutics* 2005;77:468–78.
- [124] Rodriguez-Antona C, Niemi M, Backman JT, Kajosaari LI, Neuvonen PJ, Robledo M, et al. Characterization of novel CYP2C8 haplotypes and their contribution to paclitaxel and repaglinide metabolism. *The Pharmacogenomics Journal* 2008;8:268–77.
- [125] Miyazaki M, Nakamura K, Fujita Y, Guengerich FP, Horiuchi R, Yamamoto K. Defective activity of recombinant cytochromes P450 3A4.2 and 3A4.16 in oxidation of midazolam, nifedipine, and testosterone. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2008;36:2287–91.
- [126] Andreu F, Colom H, Elens L, van Gelder T, van Schaik RHN, Hesselink DA, et al. A new CYP3A5*3 and CYP3A4*22 cluster influencing tacrolimus target concentrations: a population approach. *Clinical Pharmacokinetics* 2017;56:963–75.
- [127] de Jonge H, Elens L, de Loor H, van Schaik RH, Kuypers DR. The CYP3A4*22 C>T single nucleotide polymorphism is associated with reduced midazolam and tacrolimus clearance in stable renal allograft recipients. *The Pharmacogenomics Journal* 2015;15:144–52.
- [128] Elens L, van Gelder T, Hesselink DA, Haufroid V, van Schaik RH. CYP3A4*22: promising newly identified CYP3A4 variant allele for personalizing pharmacotherapy. *Pharmacogenomics* 2013;14:47–62.
- [129] Gijzen VM, van Schaik RH, Elens L, Soldin OP, Soldin SJ, Koren G, et al. CYP3A4*22 and CYP3A combined genotypes both correlate with tacrolimus disposition in pediatric heart transplant recipients. *Pharmacogenomics* 2013;14:1027–36.
- [130] Kuehl P, Zhang J, Lin Y, Lamba J, Assem M, Schuetz J, et al. Sequence diversity in CYP3A promoters and characterization of the genetic basis of polymorphic CYP3A5 expression. *Nature Genetics* 2001;27:383–91.
- [131] Lamba J, Hebert JM, Schuetz EG, Klein TE, Altman RB. PharmGKB summary: very important pharmacogene information for CYP3A5. *Pharmacogenetics and Genomics* 2012;22:555–8.
- [132] Lamba JK, Lin YS, Thummel K, Daly A, Watkins PB, Strom S, et al. Common allelic variants of cytochrome P4503A4 and their prevalence in different populations. *Pharmacogenetics* 2002;12:121–32.
- [133] Sata F, Sapone A, Elizondo G, Stocker P, Miller VP, Zheng W, et al. CYP3A4 allelic variants with amino acid substitutions in exons 7 and 12: evidence for an allelic variant with altered catalytic activity. *Clinical Pharmacology and Therapeutics* 2000;67:48–56.
- [134] Hesselink DA, van Schaik RH, van der Heiden IP, van der Werf M, Gregoor PJ, Lindemans J, et al. Genetic polymorphisms of the CYP3A4, CYP3A5, and MDR-1 genes and pharmacokinetics of the calcineurin inhibitors cyclosporine and tacrolimus. *Clinical Pharmacology and Therapeutics* 2003;74:245–54.
- [135] Zheng H, Zeevi A, Schuetz E, Lamba J, McCurry K, Griffith BP, et al. Tacrolimus dosing in adult lung transplant patients is related to cytochrome P4503A5 gene polymorphism. *The Journal of Clinical Pharmacology* 2004;44:135–40.
- [136] Thervet E, Lorient MA, Barbier S, Buchler M, Fichoux M, Choukroun G, et al. Optimization of initial tacrolimus dose using pharmacogenetic testing. *Clinical Pharmacology and Therapeutics* 2010;87:721–6.
- [137] Stevens JC, Hines RN, Gu C, Koukouritaki SB, Manro JR, Tandler PJ, et al. Developmental expression of the major human hepatic CYP3A enzymes. *Journal of Pharmacology and Experimental Therapeutics* 2003;307:573–82.
- [138] Westlind-Johnsson A, Malmbo S, Johansson A, Otter C, Andersson TB, Johansson I, et al. Comparative analysis of CYP3A expression in human liver suggests only a minor role for CYP3A5 in drug metabolism. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2003;31:755–61.
- [139] Tang HL, Xie HG, Yao Y, Hu YF. Lower tacrolimus daily dose requirements and acute rejection rates in the CYP3A5 nonexpressers than expressers. *Pharmacogenetics and Genomics* 2011;21:713–20.
- [140] Lloberas N, Elens L, Llaudo I, Padullés A, van Gelder T, Hesselink DA, et al. The combination of CYP3A4*22 and CYP3A5*3 single-nucleotide polymorphisms determines tacrolimus dose requirement after kidney transplantation. *Pharmacogenetics and Genomics* 2017;27:313–22.
- [141] Birdwell KA, Decker B, Barbarino JM, Peterson JF, Stein CM, Sadee W, et al. Clinical pharmacogenetics implementation consortium (CPIC) guidelines for CYP3A5 genotype and tacrolimus dosing. *Clinical Pharmacology and Therapeutics* 2015;98:19–24.
- [142] Liu S, Shi X, Tian X, Zhang X, Sun Z, Miao L. Effect of CYP3A4(*)1G and CYP3A5(*)3 polymorphisms on pharmacokinetics and pharmacodynamics of ticagrelor in healthy Chinese subjects. *Frontiers in Pharmacology* 2017;8:176.

- [143] McDonald MG, Rieder MJ, Nakano M, Hsia CK, Rettie AE. CYP4F2 is a vitamin K1 oxidase: an explanation for altered warfarin dose in carriers of the V433M variant. *Molecular Pharmacology* 2009;75:1337–46.
- [144] Takeuchi F, McGinnis R, Bourgeois S, Barnes C, Eriksson N, Soranzo N, et al. A genome-wide association study confirms VKORC1, CYP2C9, and CYP4F2 as principal genetic determinants of warfarin dose. *PLoS Genetics* 2009;5:e1000433.
- [145] Shahin MH, Khalifa SI, Gong Y, Hammad LN, Sallam MT, El Shafey M, et al. Genetic and nongenetic factors associated with warfarin dose requirements in Egyptian patients. *Pharmacogenetics and Genomics* 2011;21:130–5.
- [146] Suriapranata IM, Tjong WY, Wang T, Utama A, Raharjo SB, Yuniadi Y, et al. Genetic factors associated with patient-specific warfarin dose in ethnic Indonesians. *BMC Medical Genetics* 2011;12:80.
- [147] Chan SL, Suo C, Lee SC, Goh BC, Chia KS, Teo YY. Translational aspects of genetic factors in the prediction of drug response variability: a case study of warfarin pharmacogenomics in a multi-ethnic cohort from Asia. *The Pharmacogenomics Journal* 2012;12:312–8.
- [148] Caldwell MD, Awad T, Johnson JA, Gage BF, Falkowski M, Gardina P, et al. CYP4F2 genetic variant alters required warfarin dose. *Blood* 2008;111:4106–12.
- [149] Cha PC, Mushiroda T, Takahashi A, Kubo M, Minami S, Kamatani N, et al. Genome-wide association study identifies genetic determinants of warfarin responsiveness for Japanese. *Human Molecular Genetics* 2011;19:4735–44.
- [150] Choi JR, Kim JO, Kang DR, Yoon SA, Shin JY, Zhang X, et al. Proposal of pharmacogenetics-based warfarin dosing algorithm in Korean patients. *Journal of Human Genetics* 2011;56:290–5.
- [151] Danese E, Montagnana M, Johnson JA, Rettie AE, Zamboni CF, Lubitz SA, et al. Impact of the CYP4F2 p.V433M polymorphism on coumarin dose requirement: systematic review and meta-analysis. *Clinical Pharmacology and Therapeutics* 2012;92:746–56.
- [152] Mehlotra RK, Bockarie MJ, Zimmerman PA. CYP2B6 983T>C polymorphism is prevalent in West Africa but absent in Papua New Guinea: implications for HIV/AIDS treatment. *British Journal of Clinical Pharmacology* 2007;64:391–5.
- [153] Jinno H, Tanaka-Kagawa T, Ohno A, Makino Y, Matsushima E, Hanioka N, et al. Functional characterization of cytochrome P450 2B6 allelic variants. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2003;31:398–403.
- [154] Kirchheiner J, Klein C, Meineke I, Sasse J, Zanger UM, Mordt TE, et al. Bupropion and 4-OH-bupropion pharmacokinetics in relation to genetic polymorphisms in CYP2B6. *Pharmacogenetics* 2003;13:619–26.
- [155] Gatanaga H, Hayashida T, Tsuchiya K, Yoshino M, Kuwahara T, Tsukada H, et al. Successful efavirenz dose reduction in HIV type 1-infected individuals with cytochrome P450 2B6 *6 and *26. *Clinical Infectious Diseases* 2007;45:1230–7.
- [156] Hofmann MH, Blievernicht JK, Klein K, Saussele T, Schaeffeler E, Schwab M, et al. Aberrant splicing caused by single nucleotide polymorphism c.516G>T [Q172H], a marker of CYP2B6*6, is responsible for decreased expression and activity of CYP2B6 in liver. *Journal of Pharmacology and Experimental Therapeutics* 2008;325:284–92.
- [157] Wang H, Tompkins LM. CYP2B6: new insights into a historically overlooked cytochrome P450 isozyme. *Current Drug Metabolism* 2008;9:598–610.
- [158] Gadel S, Friedel C, Kharasch ED. Differences in methadone metabolism by CYP2B6 variants. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2015;43:994–1001.
- [159] Kharasch ED, Regina KJ, Blood J, Friedel C. Methadone pharmacogenetics: CYP2B6 polymorphisms determine plasma concentrations, clearance, and metabolism. *Anesthesiology* 2015;123:1142–53.
- [160] Benowitz NL, Zhu AZ, Tyndale RF, Dempsey D, Jacob 3rd P. Influence of CYP2B6 genetic variants on plasma and urine concentrations of bupropion and metabolites at steady state. *Pharmacogenetics and Genomics* 2013;23:135–41.
- [161] Levran O, Peles E, Hamon S, Randesi M, Adelson M, Kreek MJ. CYP2B6 SNPs are associated with methadone dose required for effective treatment of opioid addiction. *Addiction Biology* 2013;18:709–16.
- [162] Bunten H, Liang WJ, Pounder DJ, Seneviratne C, Osselt D. OPRM1 and CYP2B6 gene variants as risk factors in methadone-related deaths. *Clinical Pharmacology and Therapeutics* 2010;88:383–9.
- [163] Eap CB, Crettol S, Rougier JS, Schlafper J, Sintra Grilo L, Deglon JJ, et al. Stereoselective block of hERG channel by (S)-methadone and QT interval prolongation in CYP2B6 slow metabolizers. *Clinical Pharmacology and Therapeutics* 2007;81:719–28.
- [164] Rotger M, Tegude H, Colombo S, Cavassini M, Furrer H, Decosterd L, et al. Predictive value of known and novel alleles of CYP2B6 for efavirenz plasma concentrations in HIV-infected individuals. *Clinical Pharmacology and Therapeutics* 2007;81:557–66.
- [165] Rotger M, Saumoy M, Zhang K, Flepp M, Sahli R, Decosterd L, et al. Partial deletion of CYP2B6 owing to unequal crossover with CYP2B7. *Pharmacogenetics and Genomics* 2007;17:885–90.
- [166] Malaiyandi V, Sellers EM, Tyndale RF. Implications of CYP2A6 genetic variation for smoking behaviors and nicotine dependence. *Clinical Pharmacology and Therapeutics* 2005;77:145–58.

- [167] Thorn CF, Aklillu E, Klein TE, Altman RB. PharmGKB summary: very important pharmacogene information for CYP1A2. *Pharmacogenetics and Genomics* 2012;22:73–7.
- [168] Ghotbi R, Christensen M, Roh HK, Ingelman-Sundberg M, Aklillu E, Bertilsson L. Comparisons of CYP1A2 genetic polymorphisms, enzyme activity and the genotype-phenotype relationship in Swedes and Koreans. *European Journal of Clinical Pharmacology* 2007;63:537–46.
- [169] Sachse C, Brockmoller J, Bauer S, Roots I. Functional significance of a C→A polymorphism in intron 1 of the cytochrome P450 CYP1A2 gene tested with caffeine. *British Journal of Clinical Pharmacology* 1999;47:445–9.
- [170] Mihara K, Kondo T, Suzuki A, Yasui-Furukori N, Ono S, Otani K, et al. Effects of genetic polymorphism of CYP1A2 inducibility on the steady-state plasma concentrations of trazodone and its active metabolite m-chlorophenylpiperazine in depressed Japanese patients. *Pharmacology and Toxicology* 2001;88:267–70.
- [171] Shimoda K, Someya T, Morita S, Hirokane G, Yokono A, Takahashi S, et al. Lack of impact of CYP1A2 genetic polymorphism (C/A polymorphism at position 734 in intron 1 and G/A polymorphism at position -2964 in the 5'-flanking region of CYP1A2) on the plasma concentration of haloperidol in smoking male Japanese with schizophrenia. *Progress In Neuro-Psychopharmacology and Biological Psychiatry* 2002;26:261–5.
- [172] van der Weide J, Steijns LS, van Weelden MJ. The effect of smoking and cytochrome P450 CYP1A2 genetic polymorphism on clozapine clearance and dose requirement. *Pharmacogenetics* 2003;13:169–72.
- [173] Czerwensky F, Leucht S, Steimer W. CYP1A2*1D and *1F polymorphisms have a significant impact on olanzapine serum concentrations. *Therapeutic Drug Monitoring* 2015;37:152–60.
- [174] Perera V, Gross AS, Polasek TM, Qin Y, Rao G, Forrest A, et al. Considering CYP1A2 phenotype and genotype for optimizing the dose of olanzapine in the management of schizophrenia. *Expert Opinion on Drug Metabolism and Toxicology* 2013;9:1115–37.
- [175] Beutler E, Gelbart T, Demina A. Racial variability in the UDP-glucuronosyltransferase 1 (UGT1A1) promoter: a balanced polymorphism for regulation of bilirubin metabolism? *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:8170–4.
- [176] Jinno H, Tanaka-Kagawa T, Hanioka N, Saeki M, Ishida S, Nishimura T, et al. Glucuronidation of 7-ethyl-10-hydroxycamptothecin (SN-38), an active metabolite of irinotecan (CPT-11), by human UGT1A1 variants, G71R, P229Q, and Y486D. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2003;31:108–13.
- [177] Ando Y, Saka H, Asai G, Sugiura S, Shimokata K, Kamataki T. UGT1A1 genotypes and glucuronidation of SN-38, the active metabolite of irinotecan. *Annals of Oncology* 1998;9:845–7.
- [178] Wasserman E, Myara A, Lokiec F, Goldwasser F, Trivin F, Mahjoubi M, et al. Severe CPT-11 toxicity in patients with Gilbert's syndrome: two case reports. *Annals of Oncology* 1997;8:1049–51.
- [179] Iyer L, Das S, Janisch L, Wen M, Ramirez J, Karrison T, et al. UGT1A1*28 polymorphism as a determinant of irinotecan disposition and toxicity. *The Pharmacogenomics Journal* 2002;2:43–7.
- [180] Liu XH, Lu J, Duan W, Dai ZM, Wang M, Lin S, et al. Predictive value of UGT1A1*28 polymorphism in irinotecan-based chemotherapy. *Journal of Cancer* 2017;8:691–703.
- [181] Chen G, Ramos E, Adeyemo A, Shriner D, Zhou J, Doumatey AP, et al. UGT1A1 is a major locus influencing bilirubin levels in African Americans. *European journal of human genetics. European Journal of Human Genetics* 2012;20:463–8.
- [182] Johnson AD, Kavousi M, Smith AV, Chen MH, Dehghan A, Aspelund T, et al. Genome-wide association meta-analysis for total serum bilirubin levels. *Human Molecular Genetics* 2009;18:2700–10.
- [183] Sanna S, Busonero F, Maschio A, McArdle PF, Usala G, Dei M, et al. Common variants in the SLCO1B3 locus are associated with bilirubin levels and unconjugated hyperbilirubinemia. *Human Molecular Genetics* 2009;18:2711–8.
- [184] Kadakol A, Ghosh SS, Sappal BS, Sharma G, Chowdhury JR, Chowdhury NR. Genetic lesions of bilirubin uridine-diphosphoglucuronate glucuronosyltransferase (UGT1A1) causing Crigler-Najjar and Gilbert syndromes: correlation of genotype to phenotype. *Human Mutation* 2000;16:297–306.
- [185] Darbari DS, van Schaik RH, Capparelli EV, Rana S, McCarter R, van den Anker J. UGT2B7 promoter variant -840G>A contributes to the variability in hepatic clearance of morphine in patients with sickle cell disease. *American Journal of Hematology* 2008;83:200–2.
- [186] Madadi P, Sistonen J, Silverman G, Gladly R, Ross CJ, Carleton BC, et al. Life-threatening adverse events following therapeutic opioid administration in adults: is pharmacogenetic analysis useful? *Pain Research and Management* 2013;18:133–6.
- [187] Relling MV, Hancock ML, Rivera GK, Sandlund JT, Ribeiro RC, Krynetski EY, et al. Mercaptopurine therapy intolerance and heterozygosity at the thio-purine S-methyltransferase gene locus. *Journal of the National Cancer Institute* 1999;91:2001–8.

- [188] Relling MV, Gardner EE, Sandborn WJ, Schmiegelow K, Pui CH, Yee SW, et al. Clinical pharmacogenetics implementation consortium guidelines for thiopurine methyltransferase genotype and thiopurine dosing: 2013 update. *Clinical Pharmacology and Therapeutics* 2013;93:324–5.
- [189] Relling MV, Gardner EE, Sandborn WJ, Schmiegelow K, Pui CH, Yee SW, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for thiopurine methyltransferase genotype and thiopurine dosing. *Clinical Pharmacology and Therapeutics* 2011;89:387–91.
- [190] Lu Z, Zhang R, Diasio RB. Dihydropyrimidine dehydrogenase activity in human peripheral blood mononuclear cells and liver: population characteristics, newly identified deficient patients, and clinical implication in 5-fluorouracil chemotherapy. *Cancer Research* 1993;53:5433–8.
- [191] Morel A, Boisdron-Celle M, Fey L, Soulie P, Craipeau MC, Traore S, et al. Clinical relevance of different dihydropyrimidine dehydrogenase gene single nucleotide polymorphisms on 5-fluorouracil tolerance. *Molecular Cancer Therapeutics* 2006;5:2895–904.
- [192] Vreken P, Van Kuilenburg AB, Meinsma R, Smit GP, Bakker HD, De Abreu RA, et al. A point mutation in an invariant splice donor site leads to exon skipping in two unrelated Dutch patients with dihydropyrimidine dehydrogenase deficiency. *Journal of Inherited Metabolic Disease* 1996;19:645–54.
- [193] Amstutz U, Henricks LM, Offer SM, Barbarino J, Schellens JHM, Swen JJ, et al. Clinical pharmacogenetics implementation consortium (CPIC) guideline for dihydropyrimidine dehydrogenase genotype and fluoropyrimidine dosing: 2017 update. *Clinical Pharmacology and Therapeutics* 2018;103:210–6.
- [194] Van Kuilenburg AB, Vreken P, Beex LV, Meinsma R, Van Lenthe H, De Abreu RA, et al. Heterozygosity for a point mutation in an invariant splice donor site of dihydropyrimidine dehydrogenase and severe 5-fluorouracil related toxicity. *European Journal of Cancer* 1997;33:2258–64.
- [195] Yen JL, McLeod HL. Should DPD analysis be required prior to prescribing fluoropyrimidines? *European Journal of Cancer* 2007;43:1011–6.
- [196] Evans DA, Manley KA, Mc KV. Genetic control of isoniazid metabolism in man. *British Medical Journal* 1960;2:485–91.
- [197] Hein DW. Molecular genetics and function of NAT1 and NAT2: role in aromatic amine metabolism and carcinogenesis. *Mutation Research* 2002;506–507:65–77.
- [198] Cartwright RA, Glashan RW, Rogers HJ, Ahmad RA, Barham-Hall D, Higgins E, et al. Role of N-acetyltransferase phenotypes in bladder carcinogenesis: a pharmacogenetic epidemiological approach to bladder cancer. *Lancet* 1982;2:842–5.
- [199] Ali-Osman F, Akande O, Antoun G, Mao JX, Buolamwini J. Molecular cloning, characterization, and expression in *Escherichia coli* of full-length cDNAs of three human glutathione S-transferase Pi gene variants. Evidence for differential catalytic activity of the encoded proteins. *Journal of Biological Chemistry* 1997;272:10004–12.
- [200] Davies SM, Robison LL, Buckley JD, Tjoa T, Woods WG, Radloff GA, et al. Glutathione S-transferase polymorphisms and outcome of chemotherapy in childhood acute myeloid leukemia. *Journal of Clinical Oncology* 2001;19:1279–87.
- [201] Navarro SL, Chang JL, Peterson S, Chen C, King IB, Schwarz Y, et al. Modulation of human serum glutathione S-transferase A1/2 concentration by cruciferous vegetables in a controlled feeding study is influenced by GSTM1 and GSTT1 genotypes. *Cancer Epidemiology, Biomarkers and Prevention* 2009;18:2974–8.
- [202] Ciaccio PJ, Tew KD, LaCreta FP. Enzymatic conjugation of chlorambucil with glutathione by human glutathione S-transferases and inhibition by ethacrynic acid. *Biochemical Pharmacology* 1991;42:1504–7.
- [203] Petros WP, Hopkins PJ, Spruill S, Broadwater G, Vredenburgh JJ, Colvin OM, et al. Associations between drug metabolism genotype, chemotherapy pharmacokinetics, and overall survival in patients with breast cancer. *Journal of Clinical Oncology* 2005;23:6117–25.
- [204] Goekkurt E, Al-Batran SE, Hartmann JT, Mogck U, Schuch G, Kramer M, et al. Pharmacogenetic analyses of a phase III trial in metastatic gastroesophageal adenocarcinoma with fluorouracil and leucovorin plus either oxaliplatin or cisplatin: a study of the arbeitsgemeinschaft internistische onkologie. *Journal of Clinical Oncology* 2009;27:2863–73.
- [205] Voso MT, D'Alo F, Putzulu R, Mele L, Scardocci A, Chiusolo P, et al. Negative prognostic value of glutathione S-transferase (GSTM1 and GSTT1) deletions in adult acute myeloid leukemia. *Blood* 2002;100:2703–7.
- [206] Xiao Z, Yang L, Xu Z, Zhang Y, Liu L, Nie L, et al. Glutathione S-transferases (GSTT1 and GSTM1) genes polymorphisms and the treatment response and prognosis in Chinese patients with de novo acute myeloid leukemia. *Leukemia Research* 2008;32:1288–91.
- [207] Kap EJ, Richter S, Rudolph A, Jansen L, Ulrich A, Hoffmeister M, et al. Genetic variants in the glutathione S-transferase genes and survival in colorectal cancer patients after chemotherapy and differences according to treatment with oxaliplatin. *Pharmacogenetics and Genomics* 2014;24:340–7.
- [208] McLeod HL, Sargent DJ, Marsh S, Green EM, King CR, Fuchs CS, et al. Pharmacogenetic predictors of adverse events and response to chemotherapy in metastatic colorectal cancer: results from North American Gastrointestinal Intergroup Trial N9741. *Journal of Clinical Oncology* 2010;28:3227–33.

- [209] Uppugunduri CRS, Storelli F, Mlakar V, Huezio-Diaz Curtis P, Rezgui A, Theoret Y, et al. The association of combined GSTM1 and CYP2C9 genotype status with the occurrence of hemorrhagic cystitis in pediatric patients receiving myeloablative conditioning regimen prior to allogeneic hematopoietic stem cell transplantation. *Frontiers in Pharmacology* 2017;8:451.
- [210] Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, et al. Membrane transporters in drug development. *Nature Reviews Drug Discovery* 2010;9:215–36.
- [211] Gradhand U, Kim RB. Pharmacogenomics of MRP transporters (ABCC1-5) and BCRP (ABCG2). *Drug Metabolism Reviews* 2008;40:317–54.
- [212] Marzolini C, Paus E, Buclin T, Kim RB. Polymorphisms in human MDR1 (P-glycoprotein): recent advances and clinical relevance. *Clinical Pharmacology and Therapeutics* 2004;75:13–33.
- [213] Sakaeda T, Nakamura T, Horinouchi M, Kakumoto M, Ohmoto N, Sakai T, et al. MDR1 genotype-related pharmacokinetics of digoxin after single oral administration in healthy Japanese subjects. *Pharmaceutical Research* 2001;18:1400–4.
- [214] Hoenig MR, Walker PJ, Gurmsey C, Beadle K, Johnson L. The C3435T polymorphism in ABCB1 influences atorvastatin efficacy and muscle symptoms in a high-risk vascular cohort. *Journal of Clinical Lipidology* 2011;5:91–6.
- [215] Ferrari M, Guasti L, Maresca A, Mirabile M, Contini S, Grandi AM, et al. Association between statin-induced creatine kinase elevation and genetic polymorphisms in SLCO1B1, ABCB1 and ABCG2. *European Journal of Clinical Pharmacology* 2014;70:539–47.
- [216] Fellay J, Marzolini C, Meaden ER, Back DJ, Buclin T, Chave JP, et al. Response to antiretroviral treatment in HIV-1-infected individuals with allelic variants of the multidrug resistance transporter 1: a pharmacogenetics study. *Lancet* 2002;359:30–6.
- [217] Sadhasivam S, Chidambaran V, Zhang X, Meller J, Esslinger H, Zhang K, et al. Opioid-induced respiratory depression: ABCB1 transporter pharmacogenetics. *The Pharmacogenomics Journal* 2015;15:119–26.
- [218] Wachter VJ, Wu CY, Benet LZ. Overlapping substrate specificities and tissue distribution of cytochrome P450 3A and P-glycoprotein: implications for drug delivery and activity in cancer chemotherapy. *Molecular Carcinogenesis* 1995;13:129–34.
- [219] Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, et al. A “silent” polymorphism in the MDR1 gene changes substrate specificity. *Science* 2007;315:525–8.
- [220] Fung KL, Pan J, Ohnuma S, Lund PE, Pixley JN, Kimchi-Sarfaty C, et al. MDR1 synonymous polymorphisms alter transporter specificity and protein stability in a stable epithelial monolayer. *Cancer Research* 2014;74:598–608.
- [221] Lien EA, Solheim E, Lea OA, Lundgren S, Kvinnsland S, Ueland PM. Distribution of 4-hydroxy-N-desmethyltamoxifen and other tamoxifen metabolites in human biological fluids during tamoxifen treatment. *Cancer Research* 1989;49:2175–83.
- [222] Sugiyama Y, Kato Y, Chu X. Multiplicity of biliary excretion mechanisms for the camptothecin derivative irinotecan (CPT-11), its metabolite SN-38, and its glucuronide: role of canalicular multispecific organic anion transporter and P-glycoprotein. *Cancer Chemotherapy and Pharmacology* 1998;(42 Suppl.):S44–9.
- [223] Itoda M, Saito Y, Soyama A, Saeki M, Murayama N, Ishida S, et al. Polymorphisms in the ABCC2 (cMOAT/MRP2) gene found in 72 established cell lines derived from Japanese individuals: an association between single nucleotide polymorphisms in the 5′-untranslated region and exon 28. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2002;30:363–4.
- [224] Izzedine H, Hulot JS, Villard E, Goyenvallée C, Dominguez S, Ghosn J, et al. Association between ABCC2 gene haplotypes and tenofovir-induced proximal tubulopathy. *The Journal of Infectious Diseases* 2006;194:1481–91.
- [225] de Jong FA, Scott-Horton TJ, Kroetz DL, McLeod HL, Friberg LE, Mathijssen RH, et al. Irinotecan-induced diarrhea: functional significance of the polymorphic ABCC2 transporter protein. *Clinical Pharmacology and Therapeutics* 2007;81:42–9.
- [226] Imai Y, Nakane M, Kage K, Tsukahara S, Ishikawa E, Tsuruo T, et al. C421A polymorphism in the human breast cancer resistance protein gene is associated with low expression of Q141K protein and low-level drug resistance. *Molecular Cancer Therapeutics* 2002;1:611–6.
- [227] Bosch TM, Kjellberg LM, Bouwers A, Koeleman BP, Schellens JH, Beijnen JH, et al. Detection of single nucleotide polymorphisms in the ABCG2 gene in a Dutch population. *American Journal of Pharmacogenomics* 2005;5:123–31.
- [228] Li J, Cusatis G, Brahmer J, Sparreboom A, Robey RW, Bates SE, et al. Association of variant ABCG2 and the pharmacokinetics of epidermal growth factor receptor tyrosine kinase inhibitors in cancer patients. *Cancer Biology and Therapy* 2007;6:432–8.
- [229] Sparreboom A, Loos WJ, Burger H, Sissung TM, Verweij J, Figg WD, et al. Effect of ABCG2 genotype on the oral bioavailability of topotecan. *Cancer Biology and Therapy* 2005;4:650–8.
- [230] Cusatis G, Gregorc V, Li J, Spreafico A, Ingersoll RG, Verweij J, et al. Pharmacogenetics of ABCG2 and adverse reactions to gefitinib. *Journal of the National Cancer Institute* 2006;98:1739–42.
- [231] Kim IS, Kim HG, Kim DC, Eom HS, Kong SY, Shin HJ, et al. ABCG2 Q141K polymorphism is associated

- with chemotherapy-induced diarrhea in patients with diffuse large B-cell lymphoma who received frontline rituximab plus cyclophosphamide/doxorubicin/vincristine/prednisone chemotherapy. *Cancer Science* 2008;99:2496–501.
- [232] Ueshima S, Hira D, Fujii R, Kimura Y, Tomitsuka C, Yamane T, et al. Impact of ABCB1, ABCG2, and CYP3A5 polymorphisms on plasma trough concentrations of apixaban in Japanese patients with atrial fibrillation. *Pharmacogenetics and Genomics* 2017;27:329–36.
- [233] Tsuchiya K, Hayashida T, Hamada A, Oki S, Oka S, Gatanaga H. High plasma concentrations of dolutegravir in patients with ABCG2 genetic variants. *Pharmacogenetics and Genomics* 2017;27:416–9.
- [234] Zhang W, Yu BN, He YJ, Fan L, Li Q, Liu ZQ, et al. Role of BCRP 421C>A polymorphism on rosuvastatin pharmacokinetics in healthy Chinese males. *Clinica Chimica Acta* 2006;373:99–103.
- [235] Tomlinson B, Hu M, Lee VW, Lui SS, Chu TT, Poon EW, et al. ABCG2 polymorphism is associated with the low-density lipoprotein cholesterol response to rosuvastatin. *Clinical Pharmacology and Therapeutics* 2010;87:558–62.
- [236] Chasman DI, Giulianini F, MacFadyen J, Barratt BJ, Nyberg F, Ridker PM. Genetic determinants of statin-induced low-density lipoprotein cholesterol reduction: the justification for the use of statins in prevention: an intervention trial evaluating rosuvastatin (JUPITER) trial. *Circulation Cardiovascular Genetics* 2012;5:257–64.
- [237] Lee HK, Hu M, Lui S, Ho CS, Wong CK, Tomlinson B. Effects of polymorphisms in ABCG2, SLCO1B1, SLC10A1 and CYP2C9/19 on plasma concentrations of rosuvastatin and lipid response in Chinese patients. *Pharmacogenomics* 2013;14:1283–94.
- [238] Hagenbuch B, Meier PJ. Organic anion transporting polypeptides of the OATP/SLC21 family: phylogenetic classification as OATP/SLCO superfamily, new nomenclature and molecular/functional properties. *Pflügers Archiv* 2004;447:653–65.
- [239] Tamai I, Nezu J, Uchino H, Sai Y, Oku A, Shimane M, et al. Molecular identification and characterization of novel members of the human organic anion transporter (OATP) family. *Biochemical and Biophysical Research Communications* 2000;273:251–60.
- [240] Tirona RG, Leake BF, Merino G, Kim RB. Polymorphisms in OATP-C: identification of multiple allelic variants associated with altered transport activity among European- and African-Americans. *Journal of Biological Chemistry* 2001;276:35669–75.
- [241] Pasanen MK, Backman JT, Neuvonen PJ, Niemi M. Frequencies of single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide 1B1 SLCO1B1 gene in a Finnish population. *European Journal of Clinical Pharmacology* 2006;62:409–15.
- [242] Pasanen MK, Neuvonen M, Neuvonen PJ, Niemi M. SLCO1B1 polymorphism markedly affects the pharmacokinetics of simvastatin acid. *Pharmacogenetics and Genomics* 2006;16:873–9.
- [243] Voora D, Shah SH, Spasojevic I, Ali S, Reed CR, Salisbury BA, et al. The SLCO1B1*5 genetic variant is associated with statin-induced side effects. *Journal of the American College of Cardiology* 2009;54:1609–16.
- [244] Meyer zu Schwabedissen HE, Albers M, Baumeister SE, Rimbach C, Nauck M, Wallaschofski H, et al. Function-impairing polymorphisms of the hepatic uptake transporter SLCO1B1 modify the therapeutic efficacy of statins in a population-based cohort. *Pharmacogenetics and Genomics* 2015;25:8–18.
- [245] Link E, Parish S, Armitage J, Bowman L, Heath S, Matsuda F, et al. SLCO1B1 variants and statin-induced myopathy—a genomewide study. *New England Journal of Medicine* 2008;359:789–99.
- [246] Niemi M, Pasanen MK, Neuvonen PJ. SLCO1B1 polymorphism and sex affect the pharmacokinetics of pravastatin but not fluvastatin. *Clinical Pharmacology and Therapeutics* 2006;80:356–66.
- [247] Nishizato Y, Ieiri I, Suzuki H, Kimura M, Kawabata K, Hirota T, et al. Polymorphisms of OATP-C (SLC21A6) and OAT3 (SLC22A8) genes: consequences for pravastatin pharmacokinetics. *Clinical Pharmacology and Therapeutics* 2003;73:554–65.
- [248] Pasanen MK, Fredrikson H, Neuvonen PJ, Niemi M. Different effects of SLCO1B1 polymorphism on the pharmacokinetics of atorvastatin and rosuvastatin. *Clinical Pharmacology and Therapeutics* 2007;82:726–33.
- [249] Niemi M, Neuvonen PJ, Hofmann U, Backman JT, Schwab M, Lutjohann D, et al. Acute effects of pravastatin on cholesterol synthesis are associated with SLCO1B1 (encoding OATP1B1) haplotype *17. *Pharmacogenetics and Genomics* 2005;15:303–9.
- [250] Niemi M, Schaeffeler E, Lang T, Fromm MF, Neuvonen M, Kyrklund C, et al. High plasma pravastatin concentrations are associated with single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide-C (OATP-C, SLCO1B1). *Pharmacogenetics* 2004;14:429–40.
- [251] Tachibana-Iimori R, Tabara Y, Kusuhabara H, Kohara K, Kawamoto R, Nakura J, et al. Effect of genetic polymorphism of OATP-C (SLCO1B1) on lipid-lowering response to HMG-CoA reductase inhibitors. *Drug Metabolism Pharmacokinetics* 2004;19:375–80.
- [252] Ramsey LB, Johnson SG, Caudle KE, Haidar CE, Voora D, Wilke RA, et al. The clinical pharmacogenetics implementation consortium guideline for SLCO1B1 and simvastatin-induced myopathy: 2014 update. *Clinical Pharmacology and Therapeutics* 2014;96:423–8.

- [253] Tamraz B, Fukushima H, Wolfe AR, Kaspera R, Totah RA, Floyd JS, et al. OATP1B1-related drug-drug and drug-gene interactions as potential risk factors for cerivastatin-induced rhabdomyolysis. *Pharmacogenetics and Genomics* 2013;23:355–64.
- [254] Tamai I, Nozawa T, Koshida M, Nezu J, Sai Y, Tsuji A. Functional characterization of human organic anion transporting polypeptide B (OATP-B) in comparison with liver-specific OATP-C. *Pharmaceutical Research* 2001;18:1262–9.
- [255] Kobayashi D, Nozawa T, Imai K, Nezu J, Tsuji A, Tamai I. Involvement of human organic anion transporting polypeptide OATP-B (SLC21A9) in pH-dependent transport across intestinal apical membrane. *Journal of Pharmacology and Experimental Therapeutics* 2003;306:703–8.
- [256] Nozawa T, Nakajima M, Tamai I, Noda K, Nezu J, Sai Y, et al. Genetic polymorphisms of human organic anion transporters OATP-C (SLC21A6) and OATP-B (SLC21A9): allele frequencies in the Japanese population and functional analysis. *Journal of Pharmacology and Experimental Therapeutics* 2002;302:804–13.
- [257] Imanaga J, Kotegawa T, Imai H, Tsutsumi K, Yoshizato T, Ohyama T, et al. The effects of the SLCO2B1 c.1457C>T polymorphism and apple juice on the pharmacokinetics of fexofenadine and midazolam in humans. *Pharmacogenetics and Genomics* 2011;21:84–93.
- [258] Mougey EB, Feng H, Castro M, Irvin CG, Lima JJ. Absorption of montelukast is transporter mediated: a common variant of OATP2B1 is associated with reduced plasma concentrations and poor response. *Pharmacogenetics and Genomics* 2009;19:129–38.
- [259] Smith NF, Acharya MR, Desai N, Figg WD, Sparreboom A. Identification of OATP1B3 as a high-affinity hepatocellular transporter of paclitaxel. *Cancer Biology and Therapy* 2005;4:815–8.
- [260] Smith NF, Marsh S, Scott-Horton TJ, Hamada A, Mielke S, Mross K, et al. Variants in the SLCO1B3 gene: interethnic distribution and association with paclitaxel pharmacokinetics. *Clinical Pharmacology and Therapeutics* 2007;81:76–82.
- [261] Baker SD, Verweij J, Cusatis GA, van Schaik RH, Marsh S, Orwick SJ, et al. Pharmacogenetic pathway analysis of docetaxel elimination. *Clinical Pharmacology and Therapeutics* 2009;85:155–63.
- [262] Chew SC, Singh O, Chen X, Ramasamy RD, Kulkarni T, Lee EJ, et al. The effects of CYP3A4, CYP3A5, ABCB1, ABCC2, ABCC2 and SLCO1B3 single nucleotide polymorphisms on the pharmacokinetics and pharmacodynamics of docetaxel in nasopharyngeal carcinoma patients. *Cancer Chemotherapy and Pharmacology* 2011;67:1471–8.
- [263] Goswami S, Gong L, Giacomini K, Altman RB, Klein TE. PharmGKB summary: very important pharmacogene information for SLC22A1. *Pharmacogenetics and Genomics* 2014;24:324–8.
- [264] Tzvetkov MV, dos Santos Pereira JN, Meineke I, Saadatmand AR, Stingl JC, Brockmoller J. Morphine is a substrate of the organic cation transporter OCT1 and polymorphisms in OCT1 gene affect morphine pharmacokinetics after codeine administration. *Biochemical Pharmacology* 2013;86:666–78.
- [265] Tzvetkov MV, Saadatmand AR, Bokelmann K, Meineke I, Kaiser R, Brockmoller J. Effects of OCT1 polymorphisms on the cellular uptake, plasma concentrations and efficacy of the 5-HT(3) antagonists tropisetron and ondansetron. *The Pharmacogenomics Journal* 2012;12:22–9.
- [266] Tzvetkov MV, Saadatmand AR, Lotsch J, Tegeder I, Stingl JC, Brockmoller J. Genetically polymorphic OCT1: another piece in the puzzle of the variable pharmacokinetics and pharmacodynamics of the opioidergic drug tramadol. *Clinical Pharmacology and Therapeutics* 2011;90:143–50.
- [267] Tzvetkov MV, Vormfelde SV, Balen D, Meineke I, Schmidt T, Seht D, et al. The effects of genetic polymorphisms in the organic cation transporters OCT1, OCT2, and OCT3 on the renal clearance of metformin. *Clinical Pharmacology and Therapeutics* 2009;86:299–306.
- [268] Song IS, Shin HJ, Shim EJ, Jung IS, Kim WY, Shon JH, et al. Genetic variants of the organic cation transporter 2 influence the disposition of metformin. *Clinical Pharmacology and Therapeutics* 2008;84:559–62.
- [269] Wang ZJ, Yin OQ, Tomlinson B, Chow MS. OCT2 polymorphisms and in-vivo renal functional consequence: studies with metformin and cimetidine. *Pharmacogenetics and Genomics* 2008;18:637–45.
- [270] Shu Y, Sheardown SA, Brown C, Owen RP, Zhang S, Castro RA, et al. Effect of genetic variation in the organic cation transporter 1 (OCT1) on metformin action. *Journal of Clinical Investigation* 2007;117:1422–31.
- [271] Tzvetkov MV, Matthaei J, Pojar S, Faltraco F, Vogler S, Prukop T, et al. Increased systemic exposure and stronger cardiovascular and metabolic adverse reactions to fenoterol in individuals with heritable OCT1 deficiency. *Clinical Pharmacology and Therapeutics* 2018;103:868–78.
- [272] Bhatnagar V, Xu G, Hamilton BA, Truong DM, Eraly SA, Wu W, et al. Analyses of 5' regulatory region polymorphisms in human SLC22A6 (OAT1) and SLC22A8 (OAT3). *Journal of Human Genetics* 2006;51:575–80.
- [273] Xu G, Bhatnagar V, Wen G, Hamilton BA, Eraly SA, Nigam SK. Analyses of coding region polymorphisms in apical and basolateral human organic anion transporter (OAT) genes [OAT1 (NKT), OAT2, OAT3, OAT4, URAT (RST)]. *Kidney International* 2005;68:1491–9.
- [274] Hudis CA. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine* 2007;357:39–51.

- [275] Goldenberg MM. Trastuzumab, a recombinant DNA-derived humanized monoclonal antibody, a novel agent for the treatment of metastatic breast cancer. *Clinical Therapeutics* 1999;21:309–18.
- [276] Gomez Pena C, Davila-Fajardo CL, Martinez-Gonzalez LJ, Carmona-Saez P, Soto Pino MJ, Sanchez Ramos J, et al. Influence of the HER2 Ile655Val polymorphism on trastuzumab-induced cardiotoxicity in HER2-positive breast cancer patients: a meta-analysis. *Pharmacogenetics and Genomics* 2015;25:388–93.
- [277] Jonker DJ, O'Callaghan CJ, Karapetis CS, Zalcberg JR, Tu D, Au HJ, et al. Cetuximab for the treatment of colorectal cancer. *New England Journal of Medicine* 2007;357:2040–8.
- [278] Bokemeyer C, Bondarenko I, Makhson A, Hartmann JT, Aparicio J, de Braud F, et al. Fluorouracil, leucovorin, and oxaliplatin with and without cetuximab in the first-line treatment of metastatic colorectal cancer. *Journal of Clinical Oncology* 2009;27:663–71.
- [279] Mason DA, Moore JD, Green SA, Liggett SB. A gain-of-function polymorphism in a G-protein coupling domain of the human beta1-adrenergic receptor. *Journal of Biological Chemistry* 1999;274:12670–4.
- [280] Levin MC, Marullo S, Muntaner O, Andersson B, Magnusson Y. The myocardium-protective Gly-49 variant of the beta 1-adrenergic receptor exhibits constitutive activity and increased desensitization and down-regulation. *Journal of Biological Chemistry* 2002;277:30429–35.
- [281] Pacanowski MA, Johnson JA. PharmGKB submission update: IX. ADRB1 gene summary. *Pharmacological Reviews* 2007;59:2–4.
- [282] Cooper-DeHoff RM, Johnson JA. Hypertension pharmacogenomics: in search of personalized treatment approaches. *Nature Reviews Nephrology* 2016;12:110–22.
- [283] Si D, Wang J, Xu Y, Chen X, Zhang M, Zhou H. Association of common polymorphisms in beta1-adrenergic receptor with antihypertensive response to carvedilol. *Journal of Cardiovascular Pharmacology* 2014;64:306–9.
- [284] Wu D, Li G, Deng M, Song W, Huang X, Guo X, et al. Associations between ADRB1 and CYP2D6 gene polymorphisms and the response to beta-blocker therapy in hypertension. *Journal of International Medical Research* 2015;43:424–34.
- [285] Johnson JA, Zineh I, Puckett BJ, McGorray SP, Yarandi HN, Pauly DF. Beta 1-adrenergic receptor polymorphisms and antihypertensive response to metoprolol. *Clinical Pharmacology and Therapeutics* 2003;74:44–52.
- [286] Pacanowski MA, Gong Y, Cooper-Dehoff RM, Schork NJ, Shriver MD, Langae TY, et al. beta-adrenergic receptor gene polymorphisms and beta-blocker treatment outcomes in hypertension. *Clinical Pharmacology and Therapeutics* 2008;84:715–21.
- [287] Chen L, Meyers D, Javorsky G, Burstow D, Lolekha P, Lucas M, et al. Arg389Gly-beta1-adrenergic receptors determine improvement in left ventricular systolic function in nonischemic cardiomyopathy patients with heart failure after chronic treatment with carvedilol. *Pharmacogenetics and Genomics* 2007;17:941–9.
- [288] Terra SG, Hamilton KK, Pauly DF, Lee CR, Patterson JH, Adams KF, et al. Beta1-adrenergic receptor polymorphisms and left ventricular remodeling changes in response to beta-blocker therapy. *Pharmacogenetics and Genomics* 2005;15:227–34.
- [289] Liggett SB, Mialet-Perez J, Thaneemit-Chen S, Weber SA, Greene SM, Hodne D, et al. A polymorphism within a conserved beta(1)-adrenergic receptor motif alters cardiac function and beta-blocker response in human heart failure. *Proceedings of the National Academy of Sciences of the United States of America* 2006;103:11288–93.
- [290] Terra SG, Pauly DF, Lee CR, Patterson JH, Adams KF, Schofield RS, et al. beta-Adrenergic receptor polymorphisms and responses during titration of metoprolol controlled release/extended release in heart failure. *Clinical Pharmacology and Therapeutics* 2005;77:127–37.
- [291] Small KM, Forbes SL, Rahman FF, Bridges KM, Liggett SB. A four amino acid deletion polymorphism in the third intracellular loop of the human alpha 2C-adrenergic receptor confers impaired coupling to multiple effectors. *Journal of Biological Chemistry* 2000;275:23059–64.
- [292] Bristow MR, Murphy GA, Krause-Steinrauf H, Anderson JL, Carlquist JF, Thaneemit-Chen S, et al. An alpha2C-adrenergic receptor polymorphism alters the norepinephrine-lowering effects and therapeutic response of the beta-blocker bucindolol in chronic heart failure. *Circulation Heart Failure* 2010;3:21–8.
- [293] Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, et al. 2009 focused update incorporated into the ACC/AHA 2005 guidelines for the diagnosis and management of heart failure in adults a report of the American College of Cardiology Foundation/American Heart Association task force on practice guidelines developed in collaboration with the International Society for Heart and Lung Transplantation. *Journal of the American College of Cardiology* 2009;53:e1–90.
- [294] The Beta-Blocker Evaluation of Survival Trial Investigators. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. *New England Journal of Medicine* 2001;344:1659–67.
- [295] O'Connor CM, Fiuzat M, Carson PE, Anand IS, Plehn JF, Gottlieb SS, et al. Combinatorial pharmacogenetic interactions of bucindolol and beta1, alpha2C adrenergic receptor polymorphisms. *PLoS One* 2012;7:e44324.
- [296] Reddy S, Fung A, Manlihot C, Tierney ES, Chung WK, Blume E, et al. Adrenergic receptor genotype

- influences heart failure severity and beta-blocker response in children with dilated cardiomyopathy. *Pediatric Research* 2015;77:363–9.
- [297] Peters EJ, Slager SL, Jenkins GD, Reinalda MS, Garriock HA, Shyn SI, et al. Resequencing of serotonin-related genes and association of tagging SNPs to citalopram response. *Pharmacogenetics and Genomics* 2009;19:1–10.
- [298] Horstmann S, Lucae S, Menke A, Hennings JM, Ising M, Roeske D, et al. Polymorphisms in GRIK4, HTR2A, and FKBP5 show interactive effects in predicting remission to antidepressant treatment. *Neuropsychopharmacology* 2010;35:727–40.
- [299] Lerer B, Segman RH, Fangerau H, Daly AK, Basile VS, Cavallaro R, et al. Pharmacogenetics of tardive dyskinesia: combined analysis of 780 patients supports association with dopamine D3 receptor gene Ser9Gly polymorphism. *Neuropsychopharmacology* 2002;27:105–19.
- [300] Bakker PR, van Harten PN, van Os J. Antipsychotic-induced tardive dyskinesia and the Ser9Gly polymorphism in the DRD3 gene: a meta analysis. *Schizophrenia Research* 2006;83:185–92.
- [301] Ikeda K, Ide S, Han W, Hayashida M, Uhl GR, Sora I. How individual sensitivity to opiates can be predicted by gene analyses. *Trends in Pharmacological Sciences* 2005;26:311–7.
- [302] Bond C, LaForge KS, Tian M, Melia D, Zhang S, Borg L, et al. Single-nucleotide polymorphism in the human mu opioid receptor gene alters beta-endorphin binding and activity: possible implications for opiate addiction. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:9608–13.
- [303] Oertel BG, Kettner M, Scholich K, Renne C, Roskam B, Geisslinger G, et al. A common human micro-opioid receptor genetic variant diminishes the receptor signaling efficacy in brain regions processing the sensory information of pain. *Journal of Biological Chemistry* 2009;284:6530–5.
- [304] Klepstad P, Rakvag TT, Kaasa S, Holthe M, Dale O, Borchgrevink PC, et al. The 118 A > G polymorphism in the human mu-opioid receptor gene may increase morphine requirements in patients with pain caused by malignant disease. *Acta Anaesthesiologica Scandinavica* 2004;48:1232–9.
- [305] Chou WY, Yang LC, Lu HF, Ko JY, Wang CH, Lin SH, et al. Association of mu-opioid receptor gene polymorphism (A118G) with variations in morphine consumption for analgesia after total knee arthroplasty. *Acta Anaesthesiologica Scandinavica* 2006;50:787–92.
- [306] Hirota T, Ieiri I, Takane H, Sano H, Kawamoto K, Aono H, et al. Sequence variability and candidate gene analysis in two cancer patients with complex clinical outcomes during morphine therapy. *Drug Metabolism and Disposition: The Biological Fate of Chemicals* 2003;31:677–80.
- [307] Oliveira A, Dinis-Oliveira RJ, Nogueira A, Azevedo AS, Goncalves F, Silva P, et al. Genetic profile and cancer-related pain: a tale from two outlier cases with bone metastatic disease. *Pain Medicine* 2014;15:710–2.
- [308] De Capraris A, Cinnella G, Marolla A, Salatto P, Da Lima S, Vetusch P, et al. Micro opioid receptor A118G polymorphism and post-operative pain: opioids' effects on heterozygous patients. *International Journal of Immunopathology and Pharmacology* 2011;24:993–1004.
- [309] Hayashida M, Nagashima M, Satoh Y, Katoh R, Tagami M, Ide S, et al. Analgesic requirements after major abdominal surgery are associated with OPRM1 gene polymorphism genotype and haplotype. *Pharmacogenomics* 2008;9:1605–16.
- [310] Gong XD, Wang JY, Liu F, Yuan HH, Zhang WY, Guo YH, et al. Gene polymorphisms of OPRM1 A118G and ABCB1 C3435T may influence opioid requirements in Chinese patients with cancer pain. *Asian Pacific Journal of Cancer Prevention* 2013;14:2937–43.
- [311] Reyes-Gibby CC, Shete S, Rakvag T, Bhat SV, Skorpen F, Bruera E, et al. Exploring joint effects of genes and the clinical efficacy of morphine for cancer pain: OPRM1 and COMT gene. *Pain* 2007;130:25–30.
- [312] Chidambaran V, Mavi J, Esslinger H, Pilipenko V, Martin LJ, Zhang K, et al. Association of OPRM1 A118G variant with risk of morphine-induced respiratory depression following spine fusion in adolescents. *The Pharmacogenomics Journal* 2015;15:255–62.
- [313] Lotsch J, Skarke C, Grosch S, Darimont J, Schmidt H, Geisslinger G. The polymorphism A118G of the human mu-opioid receptor gene decreases the pupil constrictory effect of morphine-6-glucuronide but not that of morphine. *Pharmacogenetics* 2002;12:3–9.
- [314] Olsen MB, Jacobsen LM, Schistad EI, Pedersen LM, Rygh LJ, Roe C, et al. Pain intensity the first year after lumbar disc herniation is associated with the A118G polymorphism in the opioid receptor mu 1 gene: evidence of a sex and genotype interaction. *Journal of Neuroscience* 2012;32:9831–4.
- [315] Sia AT, Lim Y, Lim EC, Ocampo CE, Lim WY, Cheong P, et al. Influence of mu-opioid receptor variant on morphine use and self-rated pain following abdominal hysterectomy. *The Journal of Pain* 2013;14:1045–52.
- [316] Zhang W, Chang YZ, Kan QC, Zhang LR, Lu H, Chu QJ, et al. Association of human micro-opioid receptor gene polymorphism A118G with fentanyl analgesia consumption in Chinese gynaecological patients. *Anaesthesia* 2010;65:130–5.
- [317] Fukuda K, Hayashida M, Ide S, Saita N, Kokita Y, Kasai S, et al. Association between OPRM1 gene polymorphisms and fentanyl sensitivity in patients undergoing painful cosmetic surgery. *Pain* 2009;147:194–201.
- [318] Ginosar Y, Davidson EM, Meroz Y, Blotnick S, Shacham M, Caraco Y. Mu-opioid receptor (A118G) single-nucleotide polymorphism affects alfentanil

- requirements for extracorporeal shock wave lithotripsy: a pharmacokinetic-pharmacodynamic study. *British Journal of Anaesthesia* 2009;103:420–7.
- [319] Landau R, Liu SK, Blouin JL, Carvalho B. The effect of OPRM1 and COMT genotypes on the analgesic response to intravenous fentanyl labor analgesia. *Anesthesia and Analgesia* 2013;116:386–91.
- [320] Liggett SB, Cresci S, Kelly RJ, Syed FM, Matkovich SJ, Hahn HS, et al. A GRK5 polymorphism that inhibits beta-adrenergic receptor signaling is protective in heart failure. *Nature Medicine* 2008;14:510–7.
- [321] Cresci S, Kelly RJ, Cappola TP, Diwan A, Dries D, Kardia SL, et al. Clinical and genetic modifiers of long-term survival in heart failure. *Journal of the American College of Cardiology* 2009;54:432–44.
- [322] Levitt P, Ebert P, Mirnics K, Nimgaonkar VL, Lewis DA. Making the case for a candidate vulnerability gene in schizophrenia: convergent evidence for regulator of G-protein signaling 4 (RGS4). *Biological Psychiatry* 2006;60:534–7.
- [323] Talkowski ME, Seltman H, Bassett AS, Brzustowicz LM, Chen X, Chowdari KV, et al. Evaluation of a susceptibility gene for schizophrenia: genotype based meta-analysis of RGS4 polymorphisms from thirteen independent samples. *Biological Psychiatry* 2006;60:152–62.
- [324] Campbell DB, Ebert PJ, Skelly T, Stroup TS, Lieberman J, Levitt P, et al. Ethnic stratification of the association of RGS4 variants with antipsychotic treatment response in schizophrenia. *Biological Psychiatry* 2008;63:32–41.
- [325] Lane HY, Liu YC, Huang CL, Chang YC, Wu PL, Huang CH, et al. RGS4 polymorphisms predict clinical manifestations and responses to risperidone treatment in patients with schizophrenia. *Journal of Clinical Psychopharmacology* 2008;28:64–8.
- [326] Lane HY, Chang YC, Chiu CC, Chen ML, Hsieh MH, Chang WH. Association of risperidone treatment response with a polymorphism in the 5-HT(2A) receptor gene. *American Journal of Psychiatry* 2002;159:1593–5.
- [327] Lane HY, Hsu SK, Liu YC, Chang YC, Huang CH, Chang WH. Dopamine D3 receptor Ser9Gly polymorphism and risperidone response. *Journal of Clinical Psychopharmacology* 2005;25:6–11.
- [328] Lane HY, Lee CC, Chang YC, Lu CT, Huang CH, Chang WH. Effects of dopamine D2 receptor Ser311Cys polymorphism and clinical factors on risperidone efficacy for positive and negative symptoms and social function. *The International Journal of Neuropsychopharmacology* 2004;7:461–70.
- [329] Manunta P, Lavery G, Lanzani C, Braund PS, Simonini M, Bodycote C, et al. Physiological interaction between alpha-adducin and WNK1-NEDD4L pathways on sodium-related blood pressure regulation. *Hypertension* 2008;52:366–72.
- [330] Cusi D, Barlassina C, Azzani T, Casari G, Citterio L, Devoto M, et al. Polymorphisms of alpha-adducin and salt sensitivity in patients with essential hypertension. *Lancet* 1997;349:1353–7.
- [331] Davis BR, Arnett DK, Boerwinkle E, Ford CE, Leidecker-Foster C, Miller MB, et al. Antihypertensive therapy, the alpha-adducin polymorphism, and cardiovascular disease in high-risk hypertensive persons: the Genetics of Hypertension-Associated Treatment Study. *The Pharmacogenomics Journal* 2007;7:112–22.
- [332] Wang D, Chen H, Momary KM, Cavallari LH, Johnson JA, Sadee W. Regulatory polymorphism in vitamin K epoxide reductase complex subunit 1 (VKORC1) affects gene expression and warfarin dose requirement. *Blood* 2008;112:1013–21.
- [333] Limdi NA, Wadelius M, Cavallari L, Eriksson N, Crawford DC, Lee MT, et al. Warfarin pharmacogenetics: a single VKORC1 polymorphism is predictive of dose across 3 racial groups. *Blood* 2010;115:3827–34.
- [334] Epstein RS, Moyer TP, Aubert RE, DJ OK XF, Verbrugge RR, et al. Warfarin genotyping reduces hospitalization rates results from the MM-WES (Medco-Mayo Warfarin Effectiveness study). *Journal of the American College of Cardiology* 2010;55:2804–12.
- [335] Anderson JL, Horne BD, Stevens SM, Woller SC, Samuelson KM, Mansfield JW, et al. Randomized and clinical effectiveness trial comparing two pharmacogenetic algorithms and standard care for individualizing warfarin dosing: CoumaGen-II. *Circulation* 2012;125:1997–2005.
- [336] Scharplatz M, Puhan MA, Steurer J, Bachmann LM. What is the impact of the ACE gene insertion/deletion (I/D) polymorphism on the clinical effectiveness and adverse events of ACE inhibitors?—Protocol of a systematic review. *BMC Medical Genetics* 2004;5:23.
- [337] Arnett DK, Davis BR, Ford CE, Boerwinkle E, Leidecker-Foster C, Miller MB, et al. Pharmacogenetic association of the angiotensin-converting enzyme insertion/deletion polymorphism on blood pressure and cardiovascular risk in relation to antihypertensive treatment: the Genetics of Hypertension-Associated Treatment (GenHAT) study. *Circulation* 2005;111:3374–83.
- [338] Tiret L, Rigat B, Visvikis S, Breda C, Corvol P, Cambien F, et al. Evidence, from combined segregation and linkage analysis, that a variant of the angiotensin I-converting enzyme (ACE) gene controls plasma ACE levels. *The American Journal of Human Genetics* 1992;51:197–205.
- [339] Danser AH, Derckx FH, Hense HW, Jeunemaitre X, Riegger GA, Schunkert H. Angiotensinogen (M235T) and angiotensin-converting enzyme (I/D) polymorphisms in association with plasma renin and prorenin levels. *Journal of Hypertension* 1998;16:1879–83.

- [340] Winkelmann BR, Nauck M, Klein B, Russ AP, Bohm BO, Siekmeier R, et al. Deletion polymorphism of the angiotensin I-converting enzyme gene is associated with increased plasma angiotensin-converting enzyme activity but not with increased risk for myocardial infarction and coronary artery disease. *Annals of Internal Medicine* 1996;125:19–25.
- [341] Andersson B, Sylven C. The DD genotype of the angiotensin-converting enzyme gene is associated with increased mortality in idiopathic heart failure. *Journal of the American College of Cardiology* 1996;28:162–7.
- [342] McNamara DM, Holubkov R, Postava L, Janosko K, MacGowan GA, Mathier M, et al. Pharmacogenetic interactions between angiotensin-converting enzyme inhibitor therapy and the angiotensin-converting enzyme deletion polymorphism in patients with congestive heart failure. *Journal of the American College of Cardiology* 2004;44:2019–26.
- [343] Wu CK, Luo JL, Tsai CT, Huang YT, Cheng CL, Lee JK, et al. Demonstrating the pharmacogenetic effects of angiotensin-converting enzyme inhibitors on long-term prognosis of diastolic heart failure. *The Pharmacogenomics Journal* 2010;10:46–53.
- [344] Zolk O. Disposition of metformin: variability due to polymorphisms of organic cation transporters. *Annals of Medicine* 2011;44:119–29.
- [345] Zhou K, Bellenguez C, Spencer CC, Bennett AJ, Coleman RL, Tavendale R, et al. Common variants near ATM are associated with glycemic response to metformin in type 2 diabetes. *Nature Genetics* 2011;43:117–20.
- [346] Tammimaki A, Mannisto PT. Catechol-O-methyltransferase gene polymorphism and chronic human pain: a systematic review and meta-analysis. *Pharmacogenetics and Genomics* 2012;22:673–91.
- [347] Berthele A, Platzer S, Jochim B, Boecker H, Buettner A, Conrad B, et al. COMT Val108/158Met genotype affects the mu-opioid receptor system in the human brain: evidence from ligand-binding, G-protein activation and preproenkephalin mRNA expression. *NeuroImage* 2005;28:185–93.
- [348] Zubieta JK, Heitzeg MM, Smith YR, Bueller JA, Xu K, Xu Y, et al. COMT val158met genotype affects mu-opioid neurotransmitter responses to a pain stressor. *Science* 2003;299:1240–3.
- [349] Rakvag TT, Klepstad P, Baar C, Kvam TM, Dale O, Kaasa S, et al. The Val158Met polymorphism of the human catechol-O-methyltransferase (COMT) gene may influence morphine requirements in cancer pain patients. *Pain* 2005;116:73–8.
- [350] Rakvag TT, Ross JR, Sato H, Skorpen F, Kaasa S, Klepstad P. Genetic variation in the catechol-O-methyltransferase (COMT) gene and morphine requirements in cancer patients with pain. *Molecular Pain* 2008;4:64.
- [351] Kannankeril P, Roden DM, Darbar D. Drug-induced long QT syndrome. *Pharmacological Reviews* 2010;62:760–81.
- [352] Roden DM. Personalized medicine and the genotype-phenotype dilemma. *Journal of Interventional Cardiac Electrophysiology* 2011;31:17–23.
- [353] Fernandez-Fernandez JM, Tomas M, Vazquez E, Orio P, Latorre R, Senti M, et al. Gain-of-function mutation in the KCNMB1 potassium channel subunit is associated with low prevalence of diastolic hypertension. *Journal of Clinical Investigation* 2004;113:1032–9.
- [354] Beitelshes AL, Gong Y, Wang D, Schork NJ, Cooper-Dehoff RM, Langae TY, et al. KCNMB1 genotype influences response to verapamil SR and adverse outcomes in the International Verapamil SR/Trandolapril Study (INVEST). *Pharmacogenetics and Genomics* 2007;17:719–29.
- [355] Gloyn AL, Pearson ER, Antcliff JF, Proks P, Bruining GJ, Slingerland AS, et al. Activating mutations in the gene encoding the ATP-sensitive potassium-channel subunit Kir6.2 and permanent neonatal diabetes. *New England Journal of Medicine* 2004;350:1838–49.
- [356] Babenko AP, Polak M, Cave H, Busiah K, Czernichow P, Scharfmann R, et al. Activating mutations in the ABCC8 gene in neonatal diabetes mellitus. *New England Journal of Medicine* 2006;355:456–66.
- [357] Pearson ER, Flechtner I, Njolstad PR, Malecki MT, Flanagan SE, Larkin B, et al. Switching from insulin to oral sulfonylureas in patients with diabetes due to Kir6.2 mutations. *New England Journal of Medicine* 2006;355:467–77.