

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Continuous-continuous and regressions

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

So for in part 2:

- ▶ continuous outcomes by categories (ie continuous outcome, categorical predictor)

Next up:

- ▶ continuous outcomes with continuous predictors
- ▶ a brief touch on multiple predictor variables with one continuous outcome

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Reminder of what we've done with continuous vs continuous variables in Part I of the course:

- ▶ Graph the data: scatter plot of the relationship between X and Y
 - ▶ Does the relationship look linear? If so, what is the correlation coefficient, \hat{r} ?
 - ▶ If not, can we transform X, Y, or both to have a linear relationship on the transformed scale?
- ▶ Fit the line of best fit using `lm()`
- ▶ Using `glance()` and `tidy()` from the library `broom` to summarize the linear model findings
- ▶ Interpret the slope (\hat{b}) and intercept (\hat{a}) parameters
- ▶ Interpret the \hat{r}^2 value

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Recap: Visualizing continuous-continuous relationships

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

- ▶ Scatterplots are a good way to visualize a relationship between two continuous variables
- ▶ When we look at a scatterplot we want to evaluate:
 - ▶ The overall Pattern of the dots
 - ▶ Any notable exceptions to the pattern
 - ▶ Direction (positive or negative)
 - ▶ Form (straight line or curved)
 - ▶ Strength (how closely the points follow a line)
 - ▶ Are there any obvious outliers

Scatterplot Syntax in R

```
name of plot <- ggplot(data = dataset, aes(x = xvariable, y = yvariable)) +  
geom_point(na.rm=TRUE) + theme_minimal(base_size = 15)+  
labs(x = "xlabel", y = "ylabel", title = "Title")
```

Remember the Manatees?


Manatee data set from your textbook:

```
mana_data <- read_csv("Ch03_Manatee-deaths.csv")  
head(mana_data)
```

```
## # A tibble: 6 x 3  
##   year powerboats deaths  
##   <dbl>      <dbl>   <dbl>  
## 1  1977         447     13  
## 2  1987         645     39  
## 3  1997         755     54  
## 4  2007        1027     73  
## 5  1978         460     21  
## 6  1988         675     43
```

```
mana_scatter <- ggplot(data = mana_data, aes(x = powerboats, y = deaths)) +  
  geom_point() + theme_minimal(base_size = 15)
```


Remember the Manatees?



Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Recap: Pearson's

- ▶ Pearson's correlation coefficient measures **linear association** between two continuous variables
- ▶ It characterizes the extent to which the points cluster around a straight line
- ▶ the correlation coefficient can take on any value between -1 to 1 (inclusive)
 - ▶ -1: A perfect, negative linear association
 - ▶ 1: A perfect, positive linear association
 - ▶ 0: No linear association
- ▶ usually we use ρ when referring to the correlation in a **population** and r when referring to the correlation observed in a **sample**

Recap: Pearson's

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

```
mana_cor <- mana_data %>%  
  summarize(corr_mana = cor(powerboats, deaths))  
mana_cor
```

```
## # A tibble: 1 x 1  
##   corr_mana  
##   <dbl>  
## 1      0.945
```

lm() of manatee deaths and powerboat purchases

Continuous-
continuous and
regressions

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Calculate the line of best fit:

```
mana_lm <- lm(deaths ~ powerboats, mana_data)
# we use the package broom to look at the output of the linear model
tidy(mana_lm)
```

```
## # A tibble: 2 x 5
##   term          estimate std.error statistic  p.value
##   <chr>          <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)  -46.8      6.03     -7.75  2.43e- 9
## 2 powerboats    0.136    0.00764    17.8  5.21e-20
```

Interpreting the intercept and slope

```
## # A tibble: 2 x 5
##   term          estimate std.error statistic  p.value
##   <chr>         <dbl>     <dbl>     <dbl>    <dbl>
## 1 (Intercept)  -46.8       6.03      -7.75 2.43e- 9
## 2 powerboats    0.136     0.00764    17.8 5.21e-20
```

- ▶ Intercept: The predicted number of deaths if there were no powerboats.
- ▶ Slope: A one unit change in the number of powerboats registered (X 1,000) is associated with an increase of manatee deaths of 0.1358. That is, an increase in the number of powerboats registered by 1,000 is association with 0.1358 more manatee deaths.

Getting the R-squared from your model

When we run a linear model, the r-squared is also calculated. Here is how to see the r-squared for the manatee data:

```
glance(mana_lm)
```

```
## # A tibble: 1 x 12
##   r.squared adj.r.squared sigma statistic  p.value    df logLik   AIC    BIC
##   <dbl>      <dbl> <dbl>      <dbl>    <dbl> <dbl> <dbl> <dbl> <dbl>
## 1    0.893      0.890  8.82      316. 5.21e-20     1  -143.  292.  297.
## # ... with 3 more variables: deviance <dbl>, df.residual <int>, nobs <int>
```

Focus on:

- ▶ Column called `r.squared` values only.
- ▶ Interpretation of r-squared: The fraction of the variation in the values of y that is explained by the line of best fit.

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Correlation vs R Squared

```
mana_cor <- mana_data %>%  
  summarize(corr_mana = cor(powerboats, deaths))  
mana_cor
```

```
## # A tibble: 1 x 1  
##   corr_mana  
##   <dbl>  
## 1      0.945
```

```
glance(mana_lm)
```

```
## # A tibble: 1 x 12  
##   r.squared adj.r.squared sigma statistic p.value    df logLik   AIC    BIC  
##   <dbl>      <dbl> <dbl>    <dbl>   <dbl> <dbl> <dbl> <dbl> <dbl>  
## 1      0.893      0.890   8.82    316. 5.21e-20     1  -143.  292.  297.  
## # ... with 3 more variables: deviance <dbl>, df.residual <int>, nobs <int>
```

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Recap of part 1 (chapters
3,4, lectures 4,5,6)

**Regression and assumptions
needed for inference**

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Regression and assumptions needed for inference

What are the regression “statistics?”

When we are estimating values from a sample, we often put a “hat” on them.

- ▶ \hat{e} , \hat{r}^2 , \hat{a} , and \hat{b} are all statistics based on the sample we chose. That is, if we chose a different SRS and re-plotted the data and re-run the regression, we would expect their values to change somewhat.
- ▶ When we are specifically interested in the **effect** of some explanatory variable x on y , then our main interest is often in the underlying parameter b , the slope coefficient for x .
- ▶ For now, we interpret b as an **association** rather than a causal effect because we have not learned in this class how to build causal models.
- ▶ Today we revisit the output from regression models and apply the inference techniques from Part III of the course to regression.

Assumptions that require checking for regression inference

Continuous-
continuous and
regressions

Recap of part 1 (chapters
3,4, lectures 4,5,6)

**Regression and assumptions
needed for inference**

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

- ▶ The way we state the assumptions is different from the text book
- ▶ Focus on the four assumptions stated on the next slide, **not** the textbook's version

Assumptions that require checking for regression inference

1. The relationship between x and y is linear in the population
2. y varies Normally about the line of best fit. That is, the residuals vary Normally around the line of best fit.
3. Observations are independent. Often we can't check this using a plot, it is based on what we know about the study design.
4. The standard deviation of the responses is the same for all values of x

Except for #3, these assumptions can be investigated by examining the **estimated residuals**

We also use these plots to keep an eye out for **outliers**, which can sometimes have a larger effect on \hat{a} and \hat{b}

Terminology needed to understand the assumptions

Observed value: y


Fitted value: $\hat{y} = \hat{a} + \hat{b}x$

Estimated residuals:

$\hat{e} = \text{observed value} - \text{fitted value}$

$$\hat{e} = y - (\hat{a} + \hat{b}x)$$

Terminology needed to understand the assumptions, visualized



Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Example 1: Investigating the assumptions


Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction



A good fit to the data

Some information about each of the four plots

Plot (a) shows a fitted regression line and the data. The estimated residuals are shown by the dashed lines. We want to see that the residuals are sometimes positive and sometimes negative with no trend in their location

Plot (b) shows a QQ plot of the residuals (to check if they're Normally distributed)

Plot (c) shows a plot of the fitted values vs. the residuals. We want this to look like a random scatter. If there is a pattern then an assumption has been violated. We will show examples of this.





Plot (d) shows a boxplot of the distribution of y vs. the distribution of the residuals. If x does a good job describing y , then the box plot for the residuals will be much shorter because the model fit is good

Example 1: Investigating the assumptions

- ▶ Plot (a): The residuals are sometimes positive and sometimes negative and their magnitude varies randomly as x increases
- ▶ Plot (b): The residuals appear to be Normally distributed
- ▶ Plot (c): A random scatter - good
- ▶ Plot (d): The model fits the data well because the variation in the residuals is much smaller than the variation in the y variable to begin with.

Example 2: Investigating the assumptions

```
## `geom_smooth()` using formula 'y ~ x'
```



The linear systematic assumption does not hold

Example 2: Investigating the assumptions

- ▶ Plot (a): While the residuals are small there is a pattern: they start positive, then turn negative and become positive again (as x increases).
- ▶ Plot (b): The QQ plot does not support Normality because it is much different from a line
- ▶ Plot (c): There is a trend in the residuals vs. fitted. This accentuates the pattern observed in plot (a)
- ▶ Plots (a)-(c) all provide evidence against the assumption that a linear fit is the most appropriate one. Because the fit is actually curved, this relationship would require a x^2 term in the model, i.e., $\hat{y} = \hat{a} + \hat{b}x + \hat{c}x^2$
- ▶ Plot (d): However, even though the linearity assumption is violated, the linear model still explains a lot of the variation so it still offers insight into explaining y , even if it isn't the best model

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference





Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Example 3: Investigating the assumptions

```
## `geom_smooth()` using formula 'y ~ x'
```



The constant variance assumption does not hold

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Example 3: Investigating the assumptions

- ▶ Plot (a): This might look okay at first glance, but notice that the magnitude of the residuals is very small for x -values < 2.5 , and then it increases
- ▶ Plot (b): Also shows some issues in the upper tail
- ▶ Plot (c): There is a definite pattern in this plot known as “fanning out”. Here, we see that as the fitted value increases, the residuals become further from 0.

A note on these diagnostic plots

- ▶ If you chose a different sample, the diagnostic plots would change
- ▶ Be careful not to over interpret them
- ▶ Our goal is to learn about the population, but we only have our one sample

A note on these diagnostic plots

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

- ▶ Regression procedures are not too sensitive to lack of Normality
- ▶ Outliers are important though because they have the potential to have a large effect on the intercept and/or slope terms.

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

**Hypothesis testing for
regression**

Confidence intervals for
regression coefficient

Inference for prediction

Hypothesis testing for regression

Hypothesis testing for regression

Continuous-
continuous and
regressions

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

What are the null and alternative hypotheses?

Hypothesis testing for regression


$H_0 : b = 0$ (i.e., There is no association between temperature and the frequency of mating calls)

$H_a : b \neq 0$ (i.e., There is an association between temperature and the frequency of mating calls)

side note: your book has a section on “Testing lack of correlation” please ignore this section

Frog data showing the estimates slope vs. null hypothesis slope

```
## `geom_smooth()` using formula 'y ~ x'
```



Hypothesis testing for regression

- ▶ The regression standard error is used as part of the test statistic for the slope coefficient

To test the null hypothesis, the t-test statistic is:

$$t = \frac{\hat{b}}{SE_b}$$

where $SE_b = \frac{s}{\sqrt{\sum (x - \bar{x})^2}}$ and $s = \sqrt{\frac{1}{n-2} \sum_{i=1}^n (y - \hat{y})^2}$

We will use R to compute the test statistic, SE_b and s . Be sure you know where all of these values come from and which functions we use to run a linear model and print these values.

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Two-sided hypothesis testing for regression using tidy()

```
tidy(frog_lm)
```

```
## # A tibble: 2 x 5
##   term          estimate std.error statistic    p.value
##   <chr>          <dbl>    <dbl>    <dbl>    <dbl>
## 1 (Intercept)   -6.19     8.24    -0.751  0.462
## 2 temp          2.33     0.347     6.72  0.00000266
```

Focus on the row of data for temp:

- ▶ estimate is the estimated slope coefficient \hat{b} : 2.33
- ▶ std.error is the standard error, $SE_b = 0.347$
- ▶ statistic is the t-test statistic: $\frac{\hat{b}}{SE_b} = 2.330816/0.3467893 = 6.72$
- ▶ The test has $n - 2$ degrees of freedom, where n is the number of observations in the data frame.
- ▶ p-value is the p-value corresponding to the test

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

p value for the slope

Remember we can check this in R using our `pt()` function

- ▶ statistic is the t-test statistic: $\frac{\hat{b}}{SE_b} = 2.330816/0.3467893 = 6.72$
- ▶ The test has $n - 2$ degrees of freedom, where n is the number of observations (in our frog data $n=20$)

```
pt(q = 6.7211302, df = 18, lower.tail = F)*2
```

```
## [1] 2.663401e-06
```

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

**Confidence intervals for
regression coefficient**

Inference for prediction

Confidence intervals for regression coefficient

Confidence intervals for the regression coefficient

We can also use the output from `tidy(your_lm)` to create a 95% confidence interval for the slope coefficient.

estimate \pm margin of error

$$\hat{b} \pm t^* SE_b$$

Where t^* is the critical value for the t distribution with $n - 2$ degrees of freedom with area C (e.g., 95%) between $-t^*$ and t^* .

Confidence intervals for the regression coefficient

First, find the critical value t^* , such that 95% of the area is between t^* and $-t^*$:
notice the p value I am entering - why is this not .95?

```
t_star<-qt(p = 0.975, df = 18)
t_star
```

```
## [1] 2.100922
```

95% CI:

$2.330816 \pm t^*0.3467893$ or $2.330816 \pm 2.100922 \times 0.3467893$

95% CI: 1.60 to 3.06

Interpretation: The estimate for the slope coefficient is 2.33 (95% CI: 1.60-3.06). We found this interval using a method that gives an interval that captures the true population slope parameter (b) 95% of the time.

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

Recap of part 1 (chapters
3,4, lectures 4,5,6)

Regression and assumptions
needed for inference

Hypothesis testing for
regression

Confidence intervals for
regression coefficient

Inference for prediction

Inference for prediction

Inference for prediction

- ▶ So far we've learned only about inference for the regression coefficient
- ▶ But what if you wanted to use the model to make a prediction?
- ▶ We already know how to predict the **average** number of mating calls corresponding to a specific x value, say of 21 degrees Celsius:

$$\hat{y} = -6.190332 + 2.330816x$$

$$\hat{y} = -6.190332 + 2.330816(21) = 42.8$$


We expect 42.8 mating calls, so 43 mating calls (rounding because the outcome is a discrete variable) when the temperature is 21 degrees Celsius.

Inference for prediction

How do we make a confidence interval for this prediction?

- It depends on whether you want to make a CI for the **average response** or for an **individual's response**

```
## `geom_smooth()` using formula 'y ~ x'
```



Inference for prediction

If you want to make inference for the **mean response** μ_y when x takes the value x^* ($x^*=21$ in our example):

$$\hat{y} \pm t * SE_{\hat{\mu}}, \text{ where } SE_{\hat{\mu}} = s \sqrt{\frac{1}{n} + \frac{(x^* - \bar{x})^2}{\sum (x - \bar{x})^2}}$$

If you want to make inference for a **single observation** y when x takes the value x^* ($x^*=21$ in our example):

$$\hat{y} \pm t * SE_{\hat{y}}, \text{ where } SE_{\hat{y}} = s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{\sum (x - \bar{x})^2}}$$

Corresponding R code for prediction and confidence interval:

```
# specify the value of the explanatory variable for which you want the prediction  
newdata = data.frame(temp = 21)
```

```
# use `predict()` to make prediction and confidence intervals  
prediction_interval <- predict(frog_lm, newdata, interval = "predict")  
prediction_interval
```

```
##          fit          lwr          upr  
## 1 42.7568 36.37187 49.14173
```

```
confidence_interval <- predict(frog_lm, newdata, interval = "confidence")  
confidence_interval
```


```
##          fit          lwr          upr  
## 1 42.7568 40.38472 45.12887
```

Inference for prediction, visualized


```
## `geom_smooth()` using formula 'y ~ x'
```

```
## `geom_smooth()` using formula 'y ~ x'
```

95% Confidence Interval



95% Prediction Interval



Recap on notation

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

| Term | Population | Sample |
|-----------|-----------------|-----------|
| Intercept | a or α | \hat{a} |
| Slope | b or β | \hat{b} |
| Residual | e | \hat{e} |

Note: Although many sources will use r to indicate residuals, we will try to be consistent and use e , because we use r and r^2 to represent the correlation coefficient and r-squared respectively and this is confusing.

Recap: Use `lm()` + broom functions to look at your linear model

- ▶ `tidy(your_lm)`: Presents the output of the linear model in a tidy way
- ▶ `glance(your_lm)`: Takes a quick (one line) look at the fit statistics.
- ▶ `augment(your_lm)`: Creates an augmented data frame that contains a column for the fitted y-values (\hat{y}) and the residuals ($\hat{e} = y - \hat{y}$) among other columns

Know these functions, what they do, and how to use them.

Parting humor

Continuous-continuous and regressions

Recap of part 1 (chapters 3,4, lectures 4,5,6)

Regression and assumptions needed for inference

Hypothesis testing for regression

Confidence intervals for regression coefficient

Inference for prediction

CURVE-FITTING METHODS AND THE MESSAGES THEY SEND

