

Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

## L26: More flavors of T

April 8, 2019

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

# Flavors of T

Last lecture we introduced the T test for two **independent** samples

In this lecture we will extend our t-testing framework to consider what happens when those samples are NOT independent.

###Example: Weight by gender Imagine for example that we want to show that weight is different among males and females in the United States. Imagine we have data from 100 randomly sampled males and 100 randomly sampled females in the United States.

We would test the null hypothesis that there is no difference between the mean weight of men and women in the united states

$$\bar{X}_{(group_a)} = \bar{X}_{(group_b)}$$

###Example: Weight by gender Would we consider these samples independent?

#standard two-sample t-test

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

## Paired t-test

# Paired t-test

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

In this case we are now testing the null hypothesis that the difference is 0

This is called a paired t-test.

$$t = \frac{\bar{d}_{(weightpost - weightpre)}}{\frac{S_d}{\sqrt{n}}}$$

# Paired t-test

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

```
weights %>% summarise(dif_mean = mean(dif2), dif_sd = sd(dif2),  
                      wt1_mean=mean(weight1),wt1_sd=sd(weight1),  
                      wt3_mean=mean(weight3),wt3_sd=sd(weight3))
```

```
##      dif_mean    dif_sd wt1_mean  wt1_sd wt3_mean  wt3_sd  
## 1 -5.008323  4.854787 180.9544 30.77072 175.946 31.59696
```

## Paired t-test

Notice the syntax here:

```
t.test(weights$weight1, weights$weight3, data=weights, paired=TRUE)
```

```
##
## Paired t-test
##
## data:  weights$weight1 and weights$weight3
## t = 14.589, df = 199, p-value < 2.2e-16
## alternative hypothesis: true mean difference is not equal to 0
## 95 percent confidence interval:
##  4.331380 5.685267
## sample estimates:
## mean difference
##      5.008323
```

### Paired test results Here we see that the estimate of difference is unchanged, but the t test is now using the standard deviation of the difference

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

t-test: More juice per squeeze?



## t-test: More juice per squeeze?

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

When we have

- ▶ The standard error was much lower using the paired test. Why?
- ▶ Only variation within a subject was used to calculate the SE of the mean difference
- ▶ there was much less variation within a subject than between subjects

# The Statistical Method

Problem

Plan

Data

Analysis

Conclusion

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

# Plan, a.k.a. experimental design

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ Once the **problem** has been stated, the next step is to determine a **plan** to best answer the question. One of the tenets of design is to maximize **efficiency**.
- ▶ When data are paired a paired test greatly maximizes the efficiency by removing the noise introduced by between-subject variability.

# When is a paired design the appropriate design?

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ Studies with multiple measures on the same units of observation
- ▶ Studies with inherently related observations
- ▶ Studies that match units of observation to reduce variability

# Studies with multiple measures on the same units

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

Cross-over or before and after studies - in our weight-loss example we were looking at measures before and after participation. . .

- ▶ When “the treatment alleviates a condition rather than affects a cure.” (Hills and Armitage, 1979)
- ▶ The effect of treatment is short-term. After  $x$  amount of time, participants return to baseline.
- ▶ The  $x$  above refers to the **wash-out** period. Before applying the second treatment, participants should have enough time to reach their baseline level. Otherwise there may be a **carry over** effect.

# Studies with multiple measures on the same units

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

Considerations for before/after or cross-over studies - The time between the alternative treatments isn't so long as to introduce confounding by other factors.

- For example, if you waited a year between applying treatments, other things may have changed in the world or in a person's life that affects the outcome.
- Thus, there is a balance between waiting too long or not waiting long enough.

If we wanted to look at changes in individual related to a treatment what other type of design might we consider?

### Inherently related observations - Matched body parts - Studies in identical twins - Studies of diet or health behaviors in couples or family members

# Studies that match to reduce variability

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ Matched communities

### Paired t-test

t-test: More juice per  
squeeze?

**Example small study of diet**

Examples - which flavor of  
T?


## Example small study of diet



# Cholesterol measurements following two alternative diets -

L26: More flavors  
of T

Suppose you received the following graphic illustrating cholesterol measurements following two alternate diets. What do you think about these data?



Paired t-test


t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

# Cholesterol measurements following two alternative diets -

L26: More flavors  
of T



## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ What do you notice about the variability between participants under each diet?
- ▶ What is the mean difference?

# Cholesterol measurements following two alternative diets -

L26: More flavors  
of T

An independent t-test reveals no evidence against the null hypothesis of no difference between the diets:

```
##  
## Welch Two Sample t-test  
##  
## data: chol_dat %>% pull(A) and chol_dat %>% pull(B)  
## t = -0.78557, df = 19.976, p-value = 0.4413  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## -28.20808 12.77511  
## sample estimates:  
## mean of x mean of y  
## 201.6364 209.3529
```

## Paired t-test


t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

# Better visualization for a very small study

Now, what do you notice about the paired data?



## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

## apply a paired t-test

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ The observed value of the test statistic is:  $t = \frac{\bar{x}_d - 0}{s_d / \sqrt{n}}$
- ▶ It can be compared to a critical value from the t distribution with  $n - 1$  degrees of freedom

# Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

► First let's have a look at the dataset as is:

##		A	B	id
##	1	155	159.7581	1
##	2	180	186.0793	2
##	3	190	203.2348	3
##	4	192	199.2820	4
##	5	203	210.5172	5
##	6	201	214.8603	6

## Calculate the test statistic, p-value, and 95% confidence interval

- ▶ We can use functions from the library `dplyr` to calculate the test statistic
- ▶ Use `mutate` to calculate each participant's difference:

```
chol_dat <- chol_dat %>%mutate(diff = B - A)
head(chol_dat)
```

##		A	B	id	diff
##	1	155	159.7581	1	4.758097
##	2	180	186.0793	2	6.079290
##	3	190	203.2348	3	13.234833
##	4	192	199.2820	4	7.282034
##	5	203	210.5172	5	7.517151
##	6	201	214.8603	6	13.860260

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

## Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

- Then use `summarize` to calculate the mean difference ( $\hat{\mu}_d$ ), its standard error ( $\hat{s}_d/\sqrt{n}$ ), and the observed t-statistic:

```
summary_stats <- chol_dat %>%  
  summarize(mean_diff = mean(diff), # mean difference  
            std_err_diff = sd(diff)/sqrt(n()), # SE of the mean  
            t_stat = mean_diff/std_err_diff) # test statistic  
summary_stats
```

```
##   mean_diff std_err_diff  t_stat  
## 1  7.716487   1.168587  6.603262
```

Paired t-test

t-test: More juice per  
squeeze?


Example small study of diet

Examples - which flavor of  
T?



# Calculate the test statistic, p-value, and 95% confidence interval

What is the probability of observing a t-stat  $\geq 6.6$  or  $\leq -6.6$  using the pt command.



## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

# Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- ▶ To calculate the 95% confidence interval, we need to know the quantile of the t distribution such that 2.5% of the data lies above or below it.
- ▶ Ask R: What is the quantile such that 97.5% of the t-distribution is below it on 10 degrees of freedom using the `qt` command.

## Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

```
q <- qt(p = 0.975, lower.tail = T, df = 10)
```

```
q
```

```
## [1] 2.228139
```

```
ucl <- summary_stats %>% pull(mean_diff) + (q * summary_stats %>% pull(std_err) *  
lcl <- summary_stats %>% pull(mean_diff) - (q * summary_stats %>% pull(std_err) *  
c(lcl, ucl)
```

```
## [1] 5.112712 10.320261
```

The confidence interval is (5.1127122, 10.3202611).

## Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- Or, have R do the work for you! Just be sure to specify that `paired = T`.

# Calculate the test statistic, p-value, and 95% confidence interval

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

```
##  
## Paired t-test  
##  
## data: chol_dat %>% pull(B) and chol_dat %>% pull(A)  
## t = 6.6033, df = 10, p-value = 6.053e-05  
## alternative hypothesis: true mean difference is not equal to 0  
## 95 percent confidence interval:  
## 5.112712 10.320261  
## sample estimates:  
## mean difference  
## 7.716487
```

## Compare the outputs from the independent and paired tests

L26: More flavors  
of T

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

	Independent	Paired
T statistic	-0.78557	6.6033
df	19.976	10
pvalue	0.4413	6.053e-05
mean	201.67 vs 209.35	7.72
95% CI	-28.21 to 12.78	5.11 to 10.32
SE	9.823	1.169

- ▶ What is the same?
- ▶ What is different?

### Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

**Examples - which flavor of  
T?**

Examples - which flavor of T?

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

## The “sleepy” mocktail

You may have seen “sleepy girl mocktail” recipes making the rounds on social media, promising deep and restful sleep. Usually, these non-alcoholic beverages include a combination of tart cherry juice with other ingredients, like a magnesium supplement powder.

We spoke with wellness dietitian Lindsey Wohlford to understand more about how these ingredients impact sleep and what cancer patients should know.



Image from an MD Anderson health website article



# RCT of magnesium

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

Abbasi B, Kimiagar M, Sadeghniiat K, Shirazi MM, Hedayati M, Rashidkhani B. The effect of magnesium supplementation on primary insomnia in elderly: A double-blind placebo-controlled clinical trial. J Res Med Sci. 2012 Dec;17(12):1161-9. PMID: 23853635; PMCID: PMC3703169.

Recruited elderly individuals with diagnosed insomnia, screened out other medical conditions including sleep apnea. Intervention was 8 weeks.

# RCT of magnesium

L26: More flavors of T

## Paired t-test

t-test: More juice per squeeze?

Example small study of diet

Examples - which flavor of T?

**Table 2.**

Comparison of sleep indices in magnesium supplementation and placebo groups before and after intervention

Variable	Magnesium supplementation (n=21)				Placebo (n=22)				P2†	P3‡
	Before intervention	After intervention	Difference (CI=95%)	P1*	Before intervention	After intervention	Difference (CI=95%)			
Insomnia severity index	16.52±2.01	14.14±2.68	-2.38±2.24	<0.001	16.27±1.69	15.77±1.92	-0.5±1.71	0.2	0.006	
Total sleep time (h)	7.8±1.1	7.9±0.6	0.1±0.7	0.4	7.6±0.9	7.6±0.8	-0.03±0.3	0.6	0.3	
Sleep time (h)	5.1±0.8	5.7±0.9	0.6±0.7	0.002	5.0±0.5	5.0±0.6	-0.02±0.3	0.7	0.002	
Sleep onset latency (h)	1.3±0.2	1.1±0.4	-0.2±0.4	0.04	1.4±0.2	1.4±0.2	0.04±0.1	0.1	0.02	
Early morning awakening (h)	1.04±0.02	1.01±0.05	-0.03±0.05	0.05	1.03±0.02	1.03±0.02	-0.01±0.01	0.09	0.08	
Sleep efficiency (h)	0.67±0.07	0.73±0.1	0.06±0.1	0.02	0.66±0.04	0.66±0.07	0.00±0.05	0.2	0.006	

P1\* P value of differences in magnesium group compared via paired t-test; P2† P value of differences in placebo group compared via paired t-test; P3‡ P value of differences between magnesium and placebo groups compared via independent samples t-test

# Which flavor of T?

L26: More flavors  
of T

## Paired t-test

t-test: More juice per  
squeeze?

Example small study of diet

Examples - which flavor of  
T?

- 1) You want to see if there is a difference in blood pressure among men and women. You randomly sample 10 households from each census tract in a city and measure blood pressure of a man and woman living in each household.
- 2) You are interested in the efficacy of a medication for rheumatoid arthritis. You measure severity of symptoms among individuals randomized to treatment or control.
- 3) You are interested in family size and hyperactivity. You measure hyperactive behavior among only children vs children with siblings.

## Which flavor of T?

- 4) You are testing a new medication for glaucoma. You randomize individuals with glaucoma in both eyes to put active medication in their right or left eye.
- 5) You are interested in educational attainment in charter schools. You measure scores on a standardized test among students and a charter school and compare the scores to the state average for public schools.

###coding notes

A one sample t- test will take the form:

```
t.test(x = x variable, alternative = greater, less or two.sided, mu = null hypothesis value)
```

A two sample t-test will take the form:

```
t.test(first sample data, second sample data, alternative = greater, less or two.sided)
```

A paired t-test will take the form:

### Paired t-test

t-test: More juice per squeeze?

Example small study of diet

Examples - which flavor of T?

# Parting Humor (from XKCD.com)

L26: More flavors of T

## Paired t-test

t-test: More juice per squeeze?

Example small study of diet

Examples - which flavor of T?

