

ENTYVIO infusions are about 30 minutes each, with an annual infusion time of approximately 4 hours for the first year.

After the first year, you will receive 6 infusions per year, which is approximately 3 hours.

ENTYVIO is administered by a healthcare provider, through a needle placed in a vein (intravenously) in your arm, either at the healthcare provider's office or at an alternate site of care. Your healthcare provider will monitor you during and after the ENTYVIO infusion for side effects to see if you have a reaction to the treatment.

Here is how it works

STARTING DOSES

You will receive ENTYVIO by infusion at Weeks 0, 2, and 6.

MAINTENANCE DOSES

Starting at Week 14, you'll receive 1 infusion every 8 weeks.

- In the first year, you'll receive 8 doses of ENTYVIO
- After the first year, you'll receive 6 doses each year

Infusion locations

Talk with your doctor and insurance provider to determine which locations work best for you.



Doctor's Office




Hospital



Infusion Center



At Home



Call *EntyvioConnect* at **1-844-ENTYVIO (1-844-368-9846)** to learn more about **Nurse Support** and **Financial Support** for eligible patients.

Uses of ENTYVIO[®] (vedolizumab)

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis
- moderately to severely active Crohn's disease

Important Safety Information for ENTYVIO


- Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients.

Please see additional Important Safety Information on the next page.


A few tips to prepare you for your infusion

**Enroll in *EntyvioConnect*.**


Once you have been prescribed ENTYVIO, you can enroll in *EntyvioConnect* to get access to support. Call **1-844-ENTYVIO (1-844-368-9846)** or visit www.ENTYVIO.com/Connect to sign up today.



Try to drink a lot of water. Staying hydrated is important.



Get a good night's sleep. It helps to be well rested.



Wear loose-fitting clothes. Make your infusion as comfortable as possible.

Talk to your doctor if you have any questions or concerns, or visit ENTYVIO.com.

Important Safety Information for ENTYVIO (continued)

- ENTYVIO may cause serious side effects, including:
 - Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. You may need treatment if you have an allergic reaction. Tell your healthcare provider or get immediate medical help if you get any of these symptoms during or after an infusion of ENTYVIO: rash, itching, swelling of your lips, tongue, throat or face, shortness of breath or trouble breathing, wheezing, dizziness, feeling hot, or palpitations (feel like your heart is racing).
 - ENTYVIO may increase your risk of getting a serious infection. Before receiving and during treatment with ENTYVIO, tell your healthcare provider if you think you have an infection or symptoms of an infection, such as fever, chills, muscle aches, cough, shortness of breath, runny nose, sore throat, red or painful skin or sores on your body, tiredness, or pain during urination.
 - People with weakened immune systems can get progressive multifocal leukoencephalopathy (PML) (a rare, serious brain infection caused by a virus). Although unlikely while receiving ENTYVIO, a risk of PML cannot be ruled out. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML. Tell your healthcare provider right away if you have any of the following symptoms: confusion or problems thinking, loss of balance, change in the way you walk or talk, decreased strength or weakness on one side of the body, blurred vision, or loss of vision.
 - Liver problems can happen in people who receive ENTYVIO. Tell your healthcare provider right away if you have any of the following symptoms: tiredness, loss of appetite, pain on the right side of your abdomen, dark urine, or yellowing of the skin and eyes (jaundice).
- The most common side effects of ENTYVIO include common cold, headache, joint pain, nausea, fever, infections of the nose and throat, tiredness, cough, bronchitis, flu, back pain, rash, itching, sinus infection, throat pain, and pain in extremities. These are not all the possible side effects of ENTYVIO. Call your healthcare provider for medical advice about side effects.
- Before receiving ENTYVIO, tell your healthcare provider about all of your medical conditions, including if you: have or think you may have an infection or have infections that keep coming back; have liver problems; have tuberculosis (TB) or have been in close contact with someone with TB; have recently received or are scheduled to receive a vaccine; or if you are pregnant, breastfeeding, plan to become pregnant, or plan to breastfeed.

Please see the [full Prescribing Information](#), including the [Medication Guide](#), for ENTYVIO and talk with your healthcare provider.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.