

Unit 3: Reflection, Refraction and Dispersion of Light

1. Reflection of Light. Images in Plane Mirrors.
2. Refraction of Light. Snell's Law.
3. Total Internal Reflection.
4. Dispersion of Light.

Reading: Ch.2 of the textbook and material posted on Blackboard.

What is the difference between these reflections?



Non-polished surfaces diffusely reflect (scatter) light in all directions.

1. Reflection of Light on Boundaries.

Specular Reflection: $i = r$




Fig.2.1 $i = \text{angle of incidence}$
 $r = \text{angle of reflection}$

Copyright © 2008 by Elsevier Inc. All rights reserved.

Terminology:

- Incident ray of light
- Reflected ray of light
- Angle of incidence (i) and angle of reflection (r) are measured in respect to the normal.

The law of specular (mirror) reflection:
the angle of incidence is equal to the angle of reflection,
 $i = r$

Plane Mirrors

The image is **virtual**, not real




Fig.2.2


The diverging reflected rays of object O behave as if originating from the image I.

- The image I is virtual, at a point behind the plane of the mirror.
- The image I is at the same distance behind the mirror as the object is from the mirror.
- No real light rays originate from the image.

How a plane mirror forms an image:

<http://www.physicsclassroom.com/mmedia/optics/ifpm.cfm>

The reflectivity curves determine the reflected colour:



Copyright © 2008 by Elsevier Inc. All rights reserved.


Fig.2.3

Gold absorbs blue – appears yellow;

Copper absorbs blue and green – looks orange-red;

Silver and Aluminum have flat reflectivity curves across the visible spectrum – they appear gray; they are the best materials for mirrors.

2. Refraction of Light.



Observation: when light travels from a less dense to a denser medium it bends toward the normal, and it slows down.

The speed of light in vacuum is $c = 3 \times 10^8$ m/s.


The index of refraction of a medium determines the speed of light in it:

$$v = \frac{c}{n}$$

The index of refraction depends on the density of the medium.


Material	n
<i>Vacuum</i>	<i>1.00000</i>
<i>Air at 0°C</i>	<i>1.00029</i>
<i>Water at 20°C</i>	<i>1.33</i>
<i>Plexiglass</i>	<i>1.50</i>
<i>Glass (crown)</i>	<i>1.52</i>
<i>Cubic zirconia</i>	<i>2.16</i>
<i>Diamond</i>	<i>2.42</i>

$$\text{Snell's Law: } n_a \sin\theta_a = n_m \sin\theta_m$$



Copyright © 2008 by Elsevier Inc. All rights reserved.


Refraction: Simplified Snell's Law



Copyright © 2008 by Elsevier Inc. All rights reserved.

Rays are reversible – they can be traced backward (from water to air in the figures).

3. Total Internal Reflection.



Copyright © 2008 by Elsevier Inc. All rights reserved.

Fig.2.7

Observed when

- the incident ray travels from a denser to less dense medium, and
- the angle of incidence is greater than the critical angle

To find the critical angle substitute $n_a = 1$ and $\sin 90^\circ = 1$ in Snell's law.

$$\sin \theta_c = \frac{1}{n_m}$$

Example: reflection and refraction in diamonds.

- Find the critical angle of total internal reflection in diamonds.






Fig.2.12-2.14 Rays 1 and 2 will undergo total internal reflection if they reach the inner surface at angles greater than the critical angle.

Fig.2.15 If the diamond is cut too shallow, or too deep, some light will be lost in refraction to air and there will be less sparkle.




Optical illusions as result of refraction from denser to less dense medium



A vertical stick under water appears to be shorter, due to refraction. If the stick is not vertical, it will appear bent at the air-water interface.




How deep is the fish?



Copyright © 2011 by Elsevier Inc. All rights reserved.

If you look from above the fish appears closer to the surface, but its size is not changed.



Copyright © 2011 by Elsevier Inc. All rights reserved.

If you look at an angle the fish appears closer to the surface, and vertically compressed.

4. Dispersion of Light.

The index of refraction depends slightly on wavelength, which causes dispersion of natural light.

Example: typical glass has $n_{\text{violet}} = 1.53$, but $n_{\text{red}} = 1.51$
(Violet has $\lambda = 400 \text{ nm}$; red has $\lambda = 700 \text{ nm}$).

In dispersion violet deflects more than red,
which is the opposite of
the case of diffraction.

Can you explain this?






Fig.6.1


This file is licensed under the [Creative Commons Attribution-Share Alike 3.0 Unported](#) license.
Author: Cmglee

Under the rainbow ...



Copyright © 2008 by Elsevier Inc. All rights reserved.

Fig.2.19



Copyright © 2008 by Elsevier Inc. All rights reserved.

Fig.2.21