

25 YEARS ANNIVERSARY  
SOICT

ĐẠI HỌC BÁCH KHOA HÀ NỘI  
VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG



ĐẠI HỌC BÁCH KHOA HÀ NỘI  
VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

# Chapter 5

## Approximation of Derivative and Integral

Vũ Văn Thiệu, Đinh Viết Sang, Nguyễn Khánh Phương

SCIENTIFIC COMPUTING

# Content

1. Approximation of Derivative
2. Approximation of Integral


# APPROXIMATION OF DERIVATIVE

# Derivative Problem

- The first order Derivative :

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- Geometric meaning:
  - $f'(x)$  is the slope of tangent at point  $x$  (solid – line)
- Approximation:
  - $h \neq 0$
  - $f'(x)$  is the slope of secant (dashed – line)



# Forward Difference Method (FD)

- Formulate the method: Consider the Taylor expansion of the function  $f$  at the neighborhood of  $x$ :

$$f(x+h) = f(x) + f'(x)h + f''(\xi)\frac{h^2}{2!} \quad (1)$$

where  $\xi$  belong to  $[x, x+h]$ .

From (1)  $\Rightarrow$ :

$$f'(x) = \frac{f(x+h) - f(x)}{h} + f''(\xi)\frac{h}{2!} \quad (2)$$

Given that  $f''(\xi) h/2$  is the truncation error, from (2)  $\Rightarrow$ :

$$f'(x) \approx \frac{f(x+h) - f(x)}{h} \quad (3)$$

(3) is the Forward Difference (FD) formula to approximate the derivative

# FD: Error analysis

- Truncation error :  $f''(\xi) h/2 = O(h)$

⇒ This method has accuracy of first order

- Rounding error: When calculating  $f(x)$  and  $f(x+h)$ , if there is a rounding error, the formula for  $f'$ :

$$\frac{f(x+h)(1+\delta_1) - f(x)(1+\delta_2)}{h} = \frac{f(x+h) - f(x)}{h} + \frac{\delta_1 f(x+h) - \delta_2 f(x)}{h}$$

Because  $|\delta_i|$  is smaller than the accuracy of the computer  $\varepsilon$ , so the rounding error when calculating  $f'$  is:

$$\frac{\varepsilon(|f(x+h)| + |f(x)|)}{h}$$

- The total error is minimal when:

$$h \approx \sqrt{\varepsilon}$$

# FD: Example

- Consider the function:  $f(x) = \sin(x)$ . Use the forward difference method to approximate  $f'(\pi/3)$ .
- Error analysis
  - Calculate with  $h=10^{-k}$ ,  $k = 1, \dots, 16$
  - Find  $h$  for minimum error



# Result

<b>h</b>	<b>Derivative</b>	<b>Error</b>
$10^{-1}$	0.455901885410761	-0.044098114589239
$10^{-2}$	0.495661575773687	-0.004338424226313
$10^{-3}$	0.499566904000770	-0.000433095999230
$10^{-4}$	0.499956697895820	-0.000043302104180
$10^{-5}$	0.499995669867026	-0.000004330132974
$10^{-6}$	0.499999566971887	-0.000000433028113
$10^{-7}$	0.499999956993236	-0.000000043006764
$10^{-8}$	0.499999996961265	-0.000000003038736
$10^{-9}$	0.5000000041370186	0.0000000041370185

# Backward Difference Method (BD)

- Formulate the method: Similar to the forward difference method, in Taylor expansion we use  $x-h$  instead of  $x+h$ , we have :

$$f'(x) \approx \frac{f(x) - f(x-h)}{h} \quad (1)$$

- Error: As same as in forward difference, backward difference has the first order accuracy

- Minimum error when:

$$h \approx \sqrt{\varepsilon}$$

- Exercise: Use the BD to approximate  $f'(\pi/3)$ , knowing that  $f(x) = \sin(x)$

# Central Difference Method (CD)

- Formulate the method: Consider the Taylor expansion of the function  $f$  at the neighborhood  $x$ :

$$f(x-h) = f(x) - f'(x)h + f''(x)\frac{h^2}{2!} - f'''(\xi^-)\frac{h^3}{3!} \quad (1)$$

$$f(x+h) = f(x) + f'(x)h + f''(x)\frac{h^2}{2!} + f'''(\xi^+)\frac{h^3}{3!} \quad (2)$$

where  $\xi^+$  belongs to  $[x, x+h]$ ,  $\xi^-$  belongs to  $[x-h, x]$ . From (1) and (2), we have:

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} \quad (3)$$

(3) is the Central Difference (CD) method

# CD: Error Analysis

- Truncation error:

$$-\frac{1}{6} f'''(\zeta) h^2, \quad \zeta \in [x - h, x + h]$$

- CD method has accuracy order of 2;
- Error minimal when  $h = \varepsilon^{1/3}$
- Exercise: Use the CD method to approximate  $f'(\pi/3)$ , knowing that  $f(x) = \sin(x)$ . Compare with FD and BD method

# Compare error of: FD, BD, CD

h	FD	BD	CC
$10^{-1}$	$\sim 10^{-2}$	$\sim 10^{-2}$	$\sim 10^{-4}$
$10^{-2}$	$\sim 10^{-3}$	$\sim 10^{-3}$	$\sim 10^{-6}$
$10^{-3}$	$\sim 10^{-4}$	$\sim 10^{-4}$	$\sim 10^{-8}$
$10^{-4}$	$\sim 10^{-5}$	$\sim 10^{-5}$	$\sim 10^{-10}$
$10^{-5}$	$\sim 10^{-6}$	$\sim 10^{-6}$	$\sim 10^{-12}$
$10^{-6}$	$\sim 10^{-7}$	$\sim 10^{-7}$	$\sim 10^{-11}$
$10^{-7}$	$\sim 10^{-8}$	$\sim 10^{-8}$	$\sim 10^{-10}$
$10^{-8}$	$\sim 10^{-9}$	$\sim 10^{-9}$	$\sim 10^{-9}$
$10^{-9}$	$\sim 10^{-8}$	$\sim 10^{-8}$	$\sim 10^{-8}$

# Approximation of second order derivative

- Consider the Taylor expansion of the function  $f$  at the neighborhood  $x$ :

$$f(x-h) = f(x) - f'(x)h + f''(x)\frac{h^2}{2!} - f'''(x)\frac{h^3}{3!} + f^{(4)}(x)\frac{h^4}{4!} - f^{(5)}(x)\frac{h^5}{5!} + \dots(1)$$

$$f(x+h) = f(x) + f'(x)h + f''(x)\frac{h^2}{2!} + f'''(x)\frac{h^3}{3!} + f^{(4)}(x)\frac{h^4}{4!} + f^{(5)}(x)\frac{h^5}{5!} + \dots(2)$$

- From (1) and (2), we have the approximate formula for 2nd order derivative

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2} \quad (3)$$

- Truncation error:

- Minimum when  $h = \varepsilon^{1/4}$

$$-\frac{1}{12} f'''(\zeta) h^2, \quad \zeta \in [x-h, x+h]$$

# Approximation of partial derivative

- Similarly, we can formulate the approximate formula for partial derivative, for example, central difference for partial derivatives of function  $f(x,y)$  as follows:

$$\frac{\partial f(x, y)}{\partial x} = \frac{f(x + h, y) - f(x - h, y)}{2h}$$

$$\frac{\partial f(x, y)}{\partial y} = \frac{f(x, y + h) - f(x, y - h)}{2h}$$

# APPROXIMATION OF INTEGRAL




# Integral Problem

- Integral formula:

$$I = \int_a^b f(x)dx,$$

where  $f(x)$  is an integrable function on the interval  $[a,b]$

- Geometrical meaning:



# Riemann Sum

- Suppose the function  $f$  defined on  $[a,b]$  and  $\Delta$  is the division of the interval  $[a,b]$  into  $n$  closed sub-intervals  $I_k=[x_{k-1},x_k]$ ,  $k=1,\dots,n$ , where  $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ . Choose  $n$  points  $\{c_k: k=1,\dots,n\}$ , each of which belongs to a sub-interval, that is:  $c_k$  belongs to  $I_k$  for all  $k$ . The sum:

$$\sum_{k=1}^n f(c_k)\Delta x_k = f(c_1)\Delta x_1 + f(c_2)\Delta x_2 + \dots + f(c_n)\Delta x_n$$

is called the Riemann sum of the function  $f(x)$  corresponding to the division  $\Delta$  and the selection points  $\{c_k: k=1,\dots,n\}$ .

# Approximating Integral

- The definite integral of the function  $f(x)$  wrt  $x$  from  $a$  to  $b$  is the limit of the Riemann sum

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta x_k,$$

- It is assumed that this limit exists.
  - The function  $f(x)$  is called the function to be integrated
  - $a, b$  are the integral limitations
  - $[a, b]$  is the integral interval

# Properties of definite integrals

$$\int_a^a f(x) dx = 0$$

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\int_a^b C \cdot f(x) dx = C \cdot \int_a^b f(x) dx$$

$$\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx, \quad c \in [a, b]$$

# Theorems

- Theorem 1: If  $f$  is continuous on  $[a,b]$  and  $F$  is a primitive of the function  $f$  ( $F' = f$ ), then:

$$\int_a^b f(x) dx = F(b) - F(a)$$

- Theorem 2 (Theorem about mean value): If  $f$  is continuous on  $[a,b]$ , there exists a number  $c$  in the interval  $[a,b]$  such that :

$$f(c) = \frac{1}{b-a} \int_a^b f(x) dx$$

# Newton-Cotes formula

- The first approach to formulate an integral approximation is to approximate the function  $f(x)$  over the integral interval  $[a,b]$  by a polynomial. In each subinterval we approximate the function  $f(x)$  by a polynomial:

$$p_m(x) = a_0 + a_1x + a_2x^2 + \dots + a_mx^m \quad (1)$$

- We can replace the function  $f(x)$  by an interpolated polynomial (1).
- And then easily calculate the exact integral of interpolated polynomial

# Newton-Cotes formula

- Replace  $f(x)$  with Lagrange interpolated polynomial, we have:

$$\begin{aligned}\int_a^b f(x) dx &= \int_a^b \left( \sum_{i=0}^m \prod_{\substack{j=0 \\ j \neq i}}^m \frac{x - x_j}{x_i - x_j} f(x_i) \right) dx \\ &= \sum_{i=0}^m f(x_i) \int_a^b \prod_{\substack{j=0 \\ j \neq i}}^m \frac{x - x_j}{x_i - x_j} dx\end{aligned}\quad (1)$$

# Newton-Cotes formula

- Error of the method is evaluated by:

$$\int_a^b f(x)dx - \int_a^b p_m(x)dx = \frac{1}{(m+1)!} \int_a^b f^{(m+1)}(\zeta_x) \left( \prod_{i=0}^m (x - x_i) \right) dx$$
$$\zeta_x \in [a, b] \quad (2)$$



# Newton-Cotes formula

- The integral approximation formulas are obtained with this approach which uses an equal grid in the integral interval, i.e.:

$$x_i = a + i \cdot h; i=0,1,\dots,m; h = (b-a)/m,$$

is called the Newton-Cotes formula.

- For different  $m$ , we have different Newton-Cotes formulas

m	Order	Formula	Error
1	1	Trapezoidal rule	$O(h^2)$
2	2	Simpson 1/3	$O(h^4)$
3	3	Simpson 3/8	$O(h^4)$

# Trapezoidal rule

- For  $n=1$ , the interpolated polynomial has the form:

$$p_1(x) = f(a) + \frac{f(b) - f(a)}{b - a}(x - a)$$

$$\Rightarrow I = \int_a^b f(x)dx \approx \int_a^b p_1(x)dx = \int_a^b \left( f(a) + \frac{f(b) - f(a)}{b - a}(x - a) \right) dx$$

$$\Rightarrow I = \frac{(f(a) + f(b))}{2}(b - a) \quad (1)$$


- (1) is called the Trapezoidal formula for approximating the integral

# Trapezoidal rule


- Error of trapezoidal rule:

$$-\frac{b-a}{12} f''(\zeta) h^2, \quad h = b - a, \quad \zeta \in [a, b]$$

- Geometrical meaning:



# Trapezoidal rule: Extension



- The idea of the extended trapezoidal formula: Divide the interval  $[a, b]$  into subintervals in order to reduce the error

# Trapezoidal rule: Extension

- Divide  $[a,b]$  into  $n$  equal intervals using  $n+1$  points:

$$x_0 = a, x_1 = a + h, x_{n-1} = a + (n-1)h, x_n = a + nh$$

where  $h = (b-a)/n$ , we have:

$$I = \int_a^b f(x)dx = \int_a^{a+h} f(x)dx + \int_{a+h}^{a+2h} f(x)dx + \dots + \int_{a+(n-1)h}^{a+nh} f(x)dx \quad (1)$$

- Applying the trapezoidal formula for each subinterval we have:

$$I = \frac{h}{2} \left[ f(a) + 2 \sum_{i=1}^{n-1} f(a + ih) + f(b) \right] \quad (2)$$

- (2) is called the expanded trapezoid formula

# Simpson 1/3 formula

- Substituting  $n=2$  into the Newton-Cotes formula and integrating, we get:

$$I = \int_a^b f(x)dx = \frac{h}{3} [f(x_0) + 4f(x_1) + f(x_2)]$$

$$x_0 = a, \quad x_1 = a + h, \quad x_2 = a + 2h = b, \quad (1)$$

- (1) is called Simpson's formula 1/3

# Simpson 1/3 formula: Extension

- Like the extended trapezoidal rule, we divide the interval  $[a,b]$  into several subintervals and apply the Simpson 1/3 to each subinterval, we get the extended Simpson 1/3:

$$I = \int_a^b f(x)dx = (b-a) \frac{f(x_0) + 4 \sum_{i=1,3,5,\dots}^{n-1} f(x_i) + 2 \sum_{j=2,4,6,\dots}^{n-2} f(x_j) + f(x_n)}{3n}$$

$$x_0 = a, \quad x_i = a + ih, \quad i = 1, \dots, n, \quad (1)$$

- Note: We need an even number of subintervals, or an odd number of points.

# Simpson 3/8 formula

- Substituting  $n=3$  into the Newton-Cotes formula and integrating, we get:

$$I = \int_a^b f(x)dx = \frac{3h}{8} [f(x_0) + 3f(x_1) + 3f(x_2) + f(x_3)]$$

$$x_0 = a, \quad x_1 = a + h, \quad x_2 = a + 2h, \quad x_3 = a + 3h \quad (1)$$

- (1) is called Simpson formula 3/8





25  
SOICT

VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG  
SCHOOL OF INFORMATION AND COMMUNICATION TECHNOLOGY

**Thank you  
for your  
attentions!**

 [soict.hust.edu.vn/](http://soict.hust.edu.vn/)  [fb.com/groups/soict](https://fb.com/groups/soict)

