



Sri Chaitanya
Educational Institutions



Sri Chaitanya IIT Academy., India.

✦ A.P ✦ T.S ✦ KARNATAKA ✦ TAMILNADU ✦ MAHARASTRA ✦ DELHI ✦ RANCHI

A right Choice for the Real Aspirant

ICON Central Office - Madhapur - Hyderabad

SEC: Sr.S60_Elite, Target & LIIT-BTs

Time: **09.00Am to 12.00Pm**

JEE-MAIN

GTM-11/06

Date: 29-12-2024

Max. Marks: 300

IMPORTANT INSTRUCTION:

1. Immediately fill in the Admission number on this page of the Test Booklet with **Blue/Black Ball Point Pen** only.
2. The candidates should not write their Admission Number anywhere (except in the specified space) on the Test Booklet/ Answer Sheet.
3. The test is of **3 hours** duration.
4. The Test Booklet consists of **75 Questions**. The maximum marks are **300**.
5. There are **three** parts in the question paper 1,2,3 consisting of **Mathematics, Physics and Chemistry** having **25 Questions** in each subject and subject having **two sections**.
(I) **Section –I** contains **20 Multiple Choice Questions** with only one correct option.
Marking scheme: +4 for correct answer, **0** if not attempt and **-1** in all other cases.
(II) **Section-II** contains **05 Numerical Value Type Questions**.
■ The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **Nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).
To cancel any attempted question bubble on the question number box.
For example: To cancel attempted Question 21. Bubble on 21 as shown below

For More Material Join: @JEEAdvanced_2025



Question Answered for Marking

Question Cancelled for Marking

Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases.

6. Use Blue / Black Point Pen only for writing particulars / marking responses on the Answer Sheet. Use of pencil is strictly prohibited.
7. No candidate is allowed to carry any textual material, printed or written, bits of papers, mobile phone any electron device etc, except the Identity Card inside the examination hall.
8. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
9. On completion of the test, the candidate must hand over the Answer Sheet to the invigilator on duty in the Hall. However, the candidate are allowed to take away this Test Booklet with them.
10. Do not fold of make any stray marks on the Answer Sheet

Name of the Candidate (in Capital): _____

Admission Number:

--	--	--	--	--	--	--	--	--	--

Candidate's Signature: _____

Invigilator's Signature: _____

29-12-2024_Sr.S60_Elite, Target & LIIT-BTs_Jee-Main-GTM-11/06_Test Syllabus

MATHEMATICS : TOTAL SYLLABUS

PHYSICS : TOTAL SYLLABUS

CHEMISTRY : TOTAL SYLLABUS

Sec: Sr.S60_Elite, Target & LIIT-BTs

Page 2

Sri Chaitanya
Educational Institutions

THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN
2023SINGARAJU
VEERAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class300
300
300RANK
1JEE Advanced
2023VAILALA
CHIVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class341
360
360RANK
1NEET
2023BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class720
720
720RANK
1

For More Material Join: @JEEAdvanced_2025


**MATHEMATICS****Max Marks: 100****SECTION-I (SINGLE CORRECT ANSWER TYPE)**

This section contains **20 Multiple Choice Questions**. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

- If $\int \frac{2 \sec x \tan x}{(\sec x - \tan x)^{10}} dx = (\sec x + \tan x)^{11} \left[p(\sec x - \tan x)^2 + q \right] + c$ then $\left(\frac{1}{p} + \frac{1}{q} \right)$ is equal to (where 'c' is constant of integration)

1) 1 2) -1 3) 2 4) -2
- Three identical circles, each of radius x , are drawn as shown in figure and tangent to a semi circle of radius a , then x in terms of a ?



1) $\frac{a}{4}$ 2) $\frac{a}{2}$ 3) $\frac{a}{3}$ 4) $\frac{2a}{3}$
- The length of the latus rectum of the parabola $169 \left[(x-1)^2 + (y-3)^2 \right] = (5x-12y+17)^2$ is

1) $\frac{14}{13}$ 2) $\frac{28}{13}$ 3) $\frac{12}{13}$ 4) $\frac{16}{13}$
- A chord of the circle $x^2 + y^2 - 4x - 6y = 0$ passing through origin subtends an angle $\tan^{-1}(7/4)$ at the point where the circle meets positive y-axis, then equation of the chord is

1) $2x + 3y = 0$ 2) $x + 2y = 0$ 3) $x - 2y = 0$ 4) $2x - 3y = 0$
- If the system of equations $2x - y + z = 0, x - 2y + z = 0$ and $ax - y + 2z = 0$ has infinitely many solutions and $f(x)$ be a continuous function such that $f(5+x) + f(x) = 2 \forall x \in R$, then $\int_0^{-2a} f(x) dx$ is equal to:

1) -10 2) -4a
3) $\int_0^a [x] dx$ [where $[x]$ denotes G.I.F of x] 4) 2a

Sec: Sr.S60_Elite, Target & LIIT-BTs

Page 3

**Sri Chaitanya**
Educational Institutions**Infinity**
Learn BY SRI CHAITANYA**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023****JEE MAIN
2023**SINGARAJU
VERKAT KOUNDINYA
AIR NO. 330102023
SRI CHAITANYA
6th-12th Class**300
300****RANK
1****JEE Advanced
2023**JAVILALA
CHIRUVILAS REDDY
AIR NO. 330102023
SRI CHAITANYA
6th-12th Class**341
360****RANK
1****NEET
2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 330102023
SRI CHAITANYA
6th-12th Class**720
720****RANK
1****For More Material Join: @JEEAdvanced_2025**



6. For $x \geq 0$, the least value of a , for which $5^{1+x} + 5^{1-x}, \frac{a}{2}, 25^x + 25^{-x}$ are three consecutive terms of an A.P., is equal to:
- 1) 8 2) 12 3) 10 4) 16
7. If $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2 \sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$, then $\lim_{x \rightarrow 0} \frac{f'(x)}{x}$ is equal to
- 1) 1 2) -1 3) 2 4) -2
8. If a, b, c are real numbers satisfying
- $$\begin{vmatrix} (a^2+1)^2 & (ab+1)^2 & (ac+1)^2 \\ (ab+1)^2 & (b^2+1)^2 & (bc+1)^2 \\ (ac+1)^2 & (bc+1)^2 & (c^2+1)^2 \end{vmatrix} = k(a-b)^2(b-c)^2(c-a)^2$$
- then $k =$
- 1) 0 2) 1 3) 2 4) 4
9. If $g(x) = \begin{cases} \frac{x(3e^{1/x} + 4)}{2 - e^{1/x}}, & x \neq 0 \\ 0 & x = 0 \end{cases}$, then $f(x)$ is
- 1) Continuous as well as differentiable at $x=0$
 2) Continuous but not differentiable at $x=0$
 3) Differentiable but not continuous at $x=0$
 4) Discontinuous every where
10. The equations of the sides AB, BC and CA of a triangle ABC are:
- $$2x + y = 0, x + py = 21a, (a \neq 0) \text{ and } x - y = 3$$
- respectively. Let $P(2, a)$ be the centroid of ΔABC . Then $(BC)^2$ is equal to
- 1) 25 2) 36 3) 121 4) 122

**JEE MAIN
2023**SINGARAJU
VEERAT KOUNDINYA
AIR NO. 30012023
Sri Chaitanya
6th-12th Class**300
300**
RANK**RANK
1****JEE Advanced
2023**VAMALA
CHIVILAS REDDY
AIR NO. 3411023
Sri Chaitanya
6th-12th Class**341
360**
RANK**RANK
1****NEET
2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 7201023
Sri Chaitanya
6th-12th Class**720
720**
RANK**RANK
1**



11. Statement-1: The sum of rational terms in $(\sqrt{2} + \sqrt[3]{3} + \sqrt[6]{5})^{10}$ is 1262
Statement-2: The number of irrational terms in the expansion of $(5^{1/6} + 2^{1/8})^{100}$ is 97
The correct statements among the following are
1) Statement-1 is true, statement-2 is true
2) Statement-1 is true, statement 2 is false
3) Statement-1 is false, statement 2 is true
4) Statement-1 is false, statement-2 is false
12. The length of perpendicular drawn from P(1,2,3) to the line $\frac{x-6}{3} = \frac{y-7}{2} = \frac{z-7}{-2}$ is
1) 9 2) 14 3) 7 4) 21
13. Nine balls of the same size and colour, numbered 1,2,...,9, were put into an Urn. Now A draws a ball from Urn, noted that it is of number x, and puts it back. Then B also drawn a ball from the Urn and noted that it is of number y. Then probability that the inequality $x-2y+10 > 0$ to hold is
1) $\frac{52}{18}$ 2) $\frac{59}{81}$ 3) $\frac{60}{81}$ 4) $\frac{61}{81}$
14. The Solution of the differential equation $(1+x^3)\frac{dy}{dx} + 3x^2y + \cos^2 x - 1 = 0$
1) $y(1+x^3) = \frac{x}{2} - \frac{1}{4} \sin 2x + C$ 2) $y(1+x^3) = \frac{x}{2} + \frac{1}{4} \sin 2x + C$
3) $y(1+x^3) = \frac{x}{4} - \frac{1}{2} \sin 2x + C$ 4) $y(1-x^3) = \frac{x}{2} - \frac{1}{4} \sin 2x + C$
15. Let $R = \{(a,b) | a-b \text{ is irrational} : a,b \text{ are real number}\}$, then relation R is
1) reflexive and symmetric relation 2) transitive and symmetric
3) symmetric relation 4) equivalence relation
16. The range of λ for which the line $z(1-\lambda i) - \bar{z}(1+\lambda i) = 0$ divides two chords drawn from point $z_1 = \sqrt{3} + i$ to the curve $|z| = 2$ in the ratio 1:2 is
1) $(0, \sqrt{3})$ 2) (1,2) 3) $(\frac{1}{\sqrt{3}}, 1)$ 4) $(0, \frac{1}{\sqrt{3}})$

**JEE MAIN
2023**SINGARAJU
VEERKAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class
300
RANK**1****JEE Advanced
2023**VAMILALA
CHIVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class
341
RANK**1****NEET
2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class
720
RANK**1**



17. The number of real solutions of the equation $\sqrt{1 + \cos 2x} = \sqrt{2} \cos^{-1}(\cos x)$ in $\left[\frac{\pi}{2}, \pi\right]$ is
- 1) 0 2) 1 3) 2 4) 4
18. Let the director circle of the hyperbola $H: \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ ($a > b$) be denoted by C_1 and let director circle of ellipse $E: \frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{1}{3}$ ($a > b$) be denoted by C_2 and C_2 is also the director circle of C_1 and let e_1 and e_2 be the eccentricities of the ellipse E and hyperbola H respectively. Then $\frac{e_2^2}{e_1^2}$ is equal to
- 1) 18 2) 12 3) 6 4) 24
19. $\sum_{i=1}^{10} (x_i - 5) = 5$ and $\sum_{i=1}^{10} (x_i - 5)^2 = 125$, variance of x_1, x_2, \dots, x_{10} will be
- 1) $\frac{9}{4}$ 2) $\frac{25}{4}$ 3) $\frac{49}{4}$ 4) $\frac{81}{4}$
20. Given that $x + y + z + t = 30$. Match the number of solutions to this equation in List-2 with constraints placed on x, y, z, t in List-1

List-I		List-II	
P)	$x, y, z, t \in I; x \geq -1, y \geq 3, z \geq 5, t \geq 0$	I)	585
Q)	$x, y, z, t \in I; x \geq 0, 2 \leq y \leq 7, 4 \leq z \leq 8, t \geq 1$	II)	680
R)	$x, y, z, t \in W$ and x, y are odd while z, t are even	III)	816
S)	$x, y, z, t \in N$ and t is a multiple of 5	IV)	580
		V)	2600

The correct option is:

- 1) $P \rightarrow (V), Q \rightarrow (I), R \rightarrow (II), S \rightarrow (IV)$ 2) $P \rightarrow (I), Q \rightarrow (V), R \rightarrow (III), S \rightarrow (II)$
- 3) $P \rightarrow (II), Q \rightarrow (V), R \rightarrow (III), S \rightarrow (I)$ 4) $P \rightarrow (III), Q \rightarrow (V), R \rightarrow (IV), S \rightarrow (IV)$

JEE MAIN
2023SINGARAJU
VEERKAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class300
300
RANKRANK
1JEE Advanced
2023VANILALA
CHIVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class341
360
RANKRANK
1NEET
2023BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class720
720
RANKRANK
1

**SECTION-II (NUMERICAL VALUE TYPE)**

This section contains **5 Numerical Value Type Questions**. The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **Nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).

Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases

21. Let $f(x) = \text{Max}\{x^2, (1-x)^2, 2x(1-x)\}$ where $0 \leq x \leq 1$, if the area of the region bounded by the curves $y = f(x)$, x -axis, $x = 0$ and $x = 1$ is $\frac{p}{q}$ then $p+q$ ____ (H.C.F of p,q is 1)
22. Let N denotes the sum of the numbers obtained when two dice are rolled. If the probability that $2^N < N!$ is $\frac{m}{n}$, where m and n are coprime, then $4m-3n$ is equal to
23. If $2 \sin \alpha \sin \beta + 3 \cos \beta + 5 \cos \alpha \sin \beta = \sqrt{38} \forall \alpha, \beta \in R$, then $\det(\text{adj}(\text{adj}A))$ is equal to (where $A = \begin{bmatrix} \tan \beta & -\frac{1}{3} \\ 1 & \sin \alpha \end{bmatrix}$ is equal to ____
24. Let $f: R \rightarrow R$ be defined as $f(x) = (2x - 3\pi)^3 + \frac{4x}{3} + \cos x$ and $g = f^{-1}$ then the value of $7g'(2\pi) + 3g''(2\pi)$ (where g' and g'' represent the 1st and 2nd order derivatives of g) is ____
25. Let $\vec{a} = -\hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} + \hat{k}$ and vector \vec{c} satisfying conditions (i) $[\vec{a} \ \vec{b} \ \vec{c}] = 0$ (ii) $\vec{b} \cdot \vec{c} = 0$
 $\vec{a} \cdot \vec{c} = 7$. Then the value of $\frac{2}{7} |\vec{c}|^2$ is equal to

**JEE MAIN**
2023SINGARAJU
VEERAT KOUNDINYA
AIR NO. 300102023
Sri Chaitanya
6th-12th Class**300**
300
RANK**RANK**
1**JEE Advanced**
2023VAILALA
CHIVILAS REDDY
AIR NO. 341102023
Sri Chaitanya
6th-12th Class**341**
360
RANK**RANK**
1**NEET**
2023BORA VARUN
CHAKRAVARTHI
AIR NO. 720102023
Sri Chaitanya
6th-12th Class**720**
720
RANK**RANK**
1

**PHYSICS****Max Marks: 100****SECTION-I (SINGLE CORRECT ANSWER TYPE)**

This section contains **20 Multiple Choice Questions**. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

26. **STATEMENT-1:** A block of mass m starts moving on a rough horizontal surface with a velocity v . It stops due to friction between the block and the surface after moving through a certain distance. The surface is now tilted to an angle of 30° with the horizontal and the same block is made to go up on the surface with the same initial velocity v . The decrease in the mechanical energy in the second situation is smaller than that in the first situation.
STATEMENT-2: The coefficient of friction between the block and the surface decreases with the increase in the angle of inclination.
- 1) Statement-1 is True, statement-2 is True; statement-2 is a correct explanation for statement-1
 - 2) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for statement-1
 - 3) Statement-1 is True, statement-2 is False
 - 4) Statement-1 is False, statement-2 is True
27. Given below are two statements one is labeled as Assertion (A) and other is labeled as Reason (R)
Assertion (A) : When a block is placed in a lift which is accelerating upwards then the body experiences the following three forces
 (A) Weight (Mg) (B) Normal reaction (N) and
 (C) Pseudo force (Ma) if 'a' is acceleration of lift
Reason (R) : Pseudo force is applied on a body only when the body is seen from an accelerated observer.
- In the light of the above statements, choose the most appropriate answer from the options given below :
- 1) Both (A) and (R) are correct and (R) is the correct explanation of (A)
 - 2) Both (A) and (R) are correct and (R) is not the correct explanation of (A)
 - 3) (A) is correct but (R) is not correct
 - 4) (A) is not correct but (R) is correct

Sec: Sr.S60_Elite, Target & LIIT-BTs

Page 8

**Sri Chaitanya**
Educational Institutions

THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

**JEE MAIN
2023**SINGARAJU
VERKAT KOUNDINNYA
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class**300
300**
SCORE**RANK
1****JEE Advanced
2023**JAVILALA
CHIVILAS REDDY
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class**341
360**
SCORE**RANK
1****NEET
2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class**720
720**
SCORE**RANK
1**

For More Material Join: @JEEAdvanced_2025



28. **STATEMENT-1:** The stream of water flowing at high speed from a garden hose pipe tends to spread like a fountain when held vertically up, but tends to narrow down when held vertically down.

STATEMENT-2: In any steady flow of an incompressible fluid, the volume flow rate of the fluid remains constant.

1) Statement-1 is True, statement-2 is True; statement-2 is a correct explanation for statement-1

2) Statement-1 is True, statement-2 is True; Statement-2 is NOT a correct explanation for statement-1

3) Statement-1 is True, statement-2 is False

4) Statement-1 is False, statement-2 is True

29. Match the List-I with List-II and select the correct answer using the codes given below the list

List-I

A. Boltzmann constant

B. Coefficient of viscosity

C. Planck constant

D. Thermal conductivity

List-II

I. $[ML^2T^{-1}]$

II. $[ML^{-1}T^{-1}]$

III. $[MLT^{-3}K^{-1}]$

IV. $[ML^2T^{-2}K^{-1}]$

Choose the correct answer from the options given below


1) A-III,B-I,C-II,D-IV

2) A-III,B-II,C-I,D-IV

3) A-IV,B-II,C-I,D-III

4) A-IV,B-I,C-II,D-III

30. Molten-wax of mass m drops on a block of mass M , which is oscillating on a frictionless table as shown. Select the **CORRECT** statements.



JEE MAIN
2023

SINGARAJU
VEERAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class

300
300
RANK



RANK
1

JEE Advanced
2023

VAHILA
CHIVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class

341
360
RANK



RANK
1

NEET
2023

BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class

720
720
RANK




RANK
1



34. Two identical spherically symmetric planets, each of mass M , are somehow held at rest with respect to each other. Each planet has radius R , and the distance between the centers of the planets is $4R$. If a rocket is launched from the surface of one planet with speed v , what is the minimum speed v so that the rocket can reach the other planet?


1) $\sqrt{\frac{2GM}{R}}$ 2) $\sqrt{\frac{GM}{R}}$ 3) $\sqrt{\frac{3GM}{4R}}$ 4) $\sqrt{\frac{2GM}{3R}}$

35. The equivalent capacitance between points A and B in below shown figure will be



1) $6\mu F$ 2) $\frac{32}{3}\mu F$ 3) $32\mu F$ 4) $2\mu F$

36. A container is partially filled with a liquid of density ρ_2 . A capillary tube of radius r is vertically inserted in this liquid. Now another liquid of density ρ_1 ($\rho_1 < \rho_2$) is slowly poured in the container to a height h as shown. There is only denser liquid in the capillary tube. The rise of denser liquid in the capillary tube is also h . Assuming zero contact angle, the surface tension of heavier liquid is




1) $r\rho_2gh$ 2) $2\pi r\rho_2gh$ 3) $\frac{r}{2}(\rho_2 - \rho_1)gh$ 4) $2\pi r(\rho_2 - \rho_1)gh$

JEE MAIN
2023SINGARAJU
VENKAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class300
300RANK
1JEE Advanced
2023JAVILALA
CHIRUVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class341
360RANK
1NEET
2023BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class720
720RANK
1



37. A hydrogen atom in an excited state emits a photon which has the longest wavelength of the Paschen series. Further emissions from the atom cannot include the
- 1) Longest wavelength of the Lyman series
 - 2) Second longest wavelength of the Lyman series
 - 3) Longest wavelength of the Balmer series
 - 4) Second longest wavelength of the Balmer series
38. For the circuit shown below, calculate the value of I_z :



- 1) 25 mA
 - 2) 15 mA
 - 3) 10 mA
 - 4) 50 mA
39. In Young's double slit experiment, if we increase the separation between the plane of slits and the screen by keeping everything constant, then
- 1) Intensity of central maxima will increase but fringe width will decrease
 - 2) Intensity of central maxima will decrease but fringe width will increase
 - 3) Both intensity of central maxima and fringe width will increase
 - 4) Both intensity of central maxima and fringe width will decrease
40. **STATEMENT-1:** Two cylinders, one hollow (metal) and the other solid (wood) with the same mass and identical dimensions are simultaneously allowed to roll without slipping down an inclined plane from the same height. The hollow cylinder will reach the bottom of the inclined plane first.
- and
- STATEMENT-2:** By the principle of conservation of energy, the total kinetic energies of both the cylinders are identical when they reach the bottom of the incline.






- 1) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1
 2) Statement-1 is True, Statement-2 is True, Statement-2 is NOT a correct explanation for Statement-2
 3) Statement-1 is True, Statement-2 is False
 4) Statement-1 is False, Statement-2 is True

41. Two particles of equal mass m have respective initial velocities $u \hat{i}$ and $u \left(\frac{\hat{i} + \hat{j}}{2} \right)$. They collide completely inelastically. The energy lost in the process is

- 1) $\frac{3}{4} mu^2$ 2) $\sqrt{\frac{2}{3}} mu^2$ 3) $\frac{1}{3} mu^2$ 4) $\frac{1}{8} mu^2$

42. A ray of light entering from air into a denser medium of refractive index $\frac{4}{3}$, as shown in figure. The light ray suffers total internal reflection at the adjacent surface as shown. The maximum value of angle θ should be equal to :



- 1) $\sin^{-1} \frac{\sqrt{7}}{3}$ 2) $\sin^{-1} \frac{\sqrt{5}}{4}$ 3) $\sin^{-1} \frac{\sqrt{7}}{4}$ 4) $\sin^{-1} \frac{\sqrt{5}}{3}$

43. A nucleus with mass number 184 initially at rest emits an α -particle. If the Q value of the reaction is 5.5 MeV, calculate the kinetic energy of the α -particle.

- 1) 5 MeV 2) 5.5 MeV 3) 0.12 MeV 4) 5.38 MeV

44. The magnetic flux through a coil perpendicular to its plane is varying according to the relation $\phi = (5t^3 + 4t^2 + 2t - 5)$ Weber. If the resistance of the coil is 5 ohm, then the induced current through the coil at $t = 2$ sec will be :

- 1) 15.6 A 2) 16.6 A 3) 17.6 A 4) 18.6 A





45. An object of mass m is projected with a momentum p at an angle θ with the horizontal such that its maximum height (H) is half of its Range (R). Minimum kinetic energy of the particle in its path will be

1) $\frac{p^2}{8m}$

2) $\frac{3p^2}{4m}$

3) $\frac{p^2}{10m}$

4) $\frac{p^2}{5m}$

SECTION-II (NUMERICAL VALUE TYPE)

This section contains **5 Numerical Value Type Questions**. The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **Nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).

Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases

46. A plane electromagnetic wave travelling in non-magnetic medium is given by

$$E = (9 \times 10^8 \text{ NC}^{-1}) \sin \left[(9 \times 10^8 \text{ rads}^{-1})t - (6m^{-1})x \right]$$

where x is in metre and t is in seconds. The dielectric constant of the medium is _____

47. In an LCR series circuit, an inductor 30 mH and a resistor 1Ω are connected to an AC source of angular frequency 300 rad/s. The value of capacitance for which, the current leads the voltage by 45° is $\frac{1}{x} \times 10^{-3} \text{ F}$. Then the value of x is ____.
48. A certain metallic surface is illuminated by monochromatic radiation of wavelength λ . The stopping potential for photoelectric current for this radiation is $3V_0$. If the same surface is illuminated with a radiation of wavelength 2λ , the stopping potential is V_0 . The threshold wavelength of this surface for photoelectric effect is _____ λ .
49. A system consists of two types of gas molecules A and B having same number density $2 \times 10^{25} / \text{m}^3$. The diameter of A and B are 10 \AA and 5 \AA respectively. They suffer collision at room temperature. The ratio of average distance covered by the molecule A to that of B between two successive collision is _____ $\times 10^{-2}$
50. An organ pipe 40 cm long is open at both ends. The speed of sound in air is 360 ms^{-1} . The frequency of the second harmonic is _____ Hz.


JEE MAIN
2023
SINGARAJU
VERKAT KOUNDINYA
AIR NO. 2020123000
SRI CHAITANYA
6th-12th Class
300
300
RANK

RANK
1
JEE Advanced
2023
VAHILALA
CHIVILAS REDDY
AIR NO. 202010000
SRI CHAITANYA
6th-12th Class
341
360
RANK

RANK
1
NEET
2023
BORA VARUN
CHAKRAVARTHI
AIR NO. 120201776
SRI CHAITANYA
6th-12th Class
720
720
RANK

RANK
1

**CHEMISTRY****Max Marks: 100****SECTION-I (SINGLE CORRECT ANSWER TYPE)**

This section contains **20 Multiple Choice Questions**. Each question has 4 options (1), (2), (3) and (4) for its answer, out of which **ONLY ONE** option can be correct.

Marking scheme: +4 for correct answer, 0 if not attempted and -1 in all other cases.

51. A gas mixture of 3 litres of propane (C_3H_8) and butane (C_4H_{10}) on complete combustion at $25^\circ C$ produced 10 litre CO_2 . Find out the composition of gas mixture (propane:Butane)
- 1) 2:1 2) 1:2 3) 1.5:1.5 4) 0.5:2.5
52. If λ_0 and λ be threshold wavelength and wavelength of incident light, the velocity of photoelectrons ejected from the metal surface is :
- 1) $\sqrt{\frac{2h}{m}(\lambda_0 - \lambda)}$ 2) $\sqrt{\frac{2hc}{m}(\lambda_0 - \lambda)}$
- 3) $\sqrt{\frac{2hc}{m}\left(\frac{\lambda_0 - \lambda}{\lambda\lambda_0}\right)}$ 4) $\sqrt{\frac{2h}{m}\left(\frac{1}{\lambda_0} - \frac{1}{\lambda}\right)}$
53. The formation of the oxide ion $O^{2-}(g)$, from oxygen atom requires first an exothermic and then an endothermic step as shown below:
- $$O(g) + e^- \rightarrow O^-(g); \Delta_1 H^\circ = -141 kJ mol^{-1}$$
- $$O^-(g) + e^- \rightarrow O^{2-}(g); \Delta_2 H^\circ = +780 kJ mol^{-1}$$
- Thus process of formation of O^{2-} in gas phase is unfavourable even though O^{2-} is isoelectronic with neon. It is due to the fact that
- 1) electron repulsion out weighs the stability gained by achieving noble gas configuration
- 2) O^- ion has comparatively smaller size than oxygen atom
- 3) Oxygen is more electronegative
- 4) Addition of electron in oxygen results in larger size of the ion.
54. Among the following ions, the $p\pi - d\pi$ overlap could be possible in
- 1) NO_3^- 2) PO_4^{3-} 3) CO_3^{2-} 4) NO_2^-




THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN 2023SINGARAJU
VEERAT KOUNDINYA
AIR NO. 300 (2023)
SRI CHAITANYA
6th-12th Class**300**
300
RANK**RANK**
1**JEE Advanced 2023**VAMILALA
CHIVILAS REDDY
AIR NO. 341 (2023)
SRI CHAITANYA
6th-12th Class**341**
360
RANK**RANK**
1**NEET 2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 720 (2023)
SRI CHAITANYA
6th-12th Class**720**
720
RANK**RANK**
1




55. Bond enthalpies of H_2 , X_2 and HX are in the ratio 2:1:2, if enthalpy of formation of HX is -50 kJ mol^{-1} , the bond enthalpy of X_2 is ?
 1) 100 kJ mol^{-1} 2) 300 kJ mol^{-1} 3) 200 kJ mol^{-1} 4) 400 kJ mol^{-1}
56. Concentration of Ag^+ ions in a saturated solution of Ag_2CrO_4 is $2.2 \times 10^{-4} \text{ mol L}^{-1}$. Solubility product of Ag_2CrO_4 is : –
 1) 2.66×10^{-12} 2) 4.5×10^{-11} 3) 5.3×10^{-12} 4) 2.42×10^{-8}
57. 25.3g of sodium carbonate, Na_2CO_3 is dissolved in enough water to make 250 mL of solution. If sodium carbonate dissociates completely, molar concentration of sodium ions, (Na^+) and carbonate ions, (CO_3^{2-}) respectively are (Molar mass of $Na_2CO_3 = 106 \text{ g mol}^{-1}$)
 1) 0.955 M and 1.910 M 2) 1.910 M and 0.955 M
 3) 1.90 M and 1.910 M 4) 0.477 M and 0.477 M
58. Benzene reacts with n-propyl chloride in the presence of anhydrous $AlCl_3$ to give
 1) 3 – Propyl-1- chlorobenzene 2) n-Propylbenzene
 3) No reaction 4) Isopropylbenzene
59. The correct order of basicity of the following compounds







- 1) $B > A > C$ 2) $A > B > C$ 3) $C > A > B$ 4) $C > B > A$


**JEE MAIN
2023**SINGARAJU
VEERAT KOUNDINYA
AIR NO. 20012023
SRI CHAITANYA
6th-12th Class**300
300****RANK
1****JEE Advanced
2023**VAMILALA
CHIVILAS REDDY
AIR NO. 20010000
SRI CHAITANYA
6th-12th Class**341
360****RANK
1****NEET
2023**BORA VARUN
CHAKRAVARTHI
AIR NO. 120010176
SRI CHAITANYA
6th-12th Class**720
720****RANK
1**







60.



61. The major product of the following reaction

**JEE MAIN**
2023SINGARAJU
VEERAT KOUNDINYA
APPL NO 230510125005
SRI CHAITANYA
6th-12th Class**300**
300
RANK**RANK**
1**JEE Advanced**
2023VAMILALA
CHIVILAS REDDY
APPL NO 230510125005
SRI CHAITANYA
6th-12th Class**341**
360
RANK**RANK**
1**NEET**
2023BORA VARUN
CHAKRAVARTHI
APPL NO 230510125005
SRI CHAITANYA
6th-12th Class**720**
720
RANK**RANK**
1



62. Match the compounds given List-I with List II and select the suitable option given below:

	List-I		List-II
A)	Para-cresol	i)	Phenolphthalein
B)	Phthalic anhydride	ii)	Gives +ve test with neutral $FeCl_3$
C)	Aniline	iii)	Oil of wintergreen
D)	Methyl salicylate	iv)	Carbyl amine test

1) A-iv; B-i; C- ii ; D- ii

2) A-iv; B-ii; C- iii; D- i

3) A-ii; B- iii; C- iv; D- i

4) A-ii; B-i; C- iv ; D- iii

63. If one stand DNA has the sequence ATGCTTGA, the sequence in the complimentary stand would be

1) TACGAAC 2) TCCGAAC 3) TACGTACT 4) TACGTAGT

JEE MAIN
2023SINGARAJU
VENKAT KOUNDINYA
AIR NO. 2020120001
SRI CHAITANYA
6th-12th Class300
300
300RANK
1JEE Advanced
2023VANILALA
CHIVILAS REDDY
AIR NO. 20210001
SRI CHAITANYA
6th-12th Class341
360
360RANK
1NEET
2023BORA VARUN
CHAKRAVARTHI
AIR NO. 12020174
SRI CHAITANYA
6th-12th Class720
720
720RANK
1



64. Number of SP-carbons in Benzyne





1) 2

2) 0

3) 6

4) 4

65. Which of the following ions are optically active?



1) I only

2) II only

3) II and III

4) IV only

66. Assertion: Boiling point of $PH_3 < AsH_3 < SbH_3 < NH_3$

Reason: NH_3 possess inter molecular H-Bonding

1) A is True, R is true R is correct explanation of A

2) A is True, R is True R is not correct explanation of A

3) A is True, R is false

4) A is false, R is True

67. Assertion: $H_3PO_4 + 2NaOH \rightarrow Na_2HPO_4 + 2H_2O$

Eq. wt of H_3PO_4 in above reaction is 49

Reason: In all reactions H_3PO_4 is a dibasic acid



JEE MAIN
2023

SINGARAJU
VERKAT KOUNDINYA
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

300
300
RANK



RANK
1

JEE Advanced
2023

VANILALA
CHIVILAS REDDY
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

341
360
RANK



RANK
1

NEET
2023

BORA VARUN
CHAKRAVARTHI
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

720
720
RANK



RANK
1



- 1) A is True, R is true R is correct explanation of A
 2) A is True, R is True R is not correct explanation of A
 3) A is True, R is false
 4) A is false, R is True
68. The $I.E_1$ among the group 13 member follows as
- 1) $B > Al > Ga < Tl < In$ 2) $B > Ga > Al > Tl > In$
 3) $B > Tl > Ga > Al > In$ 4) $Al > Ga > Tl > B > In$
69. Consider the following statement:
 (i) Atomic radii decreases across a row of the periodic table when we move from left to right
 (ii) Atomic radii increases down the column as we move from top to bottom
 (iii) screening effect order $s > p > d > f$
- Correct statements are
- 1) (i) and (ii) only 2) (i) and (iii) only
 3) (ii) and (iii) only 4) (i),(ii) and (iii)
70. Which of the following statements are correct?
- (i) In Sandmeyer reaction nucleophiles like Cl^- , Br^- and CN^- are introduced in benzene ring in the presence of Cu^+ ion
 (ii) In Gattermann reaction nucleophiles are introduced in benzene ring in the presence of copper powder and HCl.
 (iii) The yield in Gattermann reaction is found to be better than Sandmayer reaction.
- 1) (i) and (ii) only 2) (i), (ii) and (iii)
 3) (ii) and (iii) only 4) (i) and (iii) only

SECTION-II (NUMERICAL VALUE TYPE)

This section contains **5 Numerical Value Type Questions**. The Answer should be within **0 to 9999**. If the Answer is in **Decimal** then round off to the **Nearest Integer** value (Example i.e. If answer is above **10** and less than **10.5** round off is **10** and If answer is from **10.5** and less than **11** round off is **11**).

Marking scheme: +4 for correct answer, 0 if not attempt and -1 in all other cases

Sec: Sr.S60_Elite, Target & LIIT-BTs

Page 20



Sri Chaitanya
Educational Institutions



**THE PERFECT HAT-TRICK WITH ALL-INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023**

**JEE MAIN
2023**

SINGARAJU
VERKAT KOUNDINYA
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

**300
300
300**



**RANK
1**

**JEE Advanced
2023**

JAVILALA
CHIVILAS REDDY
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

**341
360
360**



**RANK
1**

**NEET
2023**

BORA VARUN
CHAKRAVARTHI
AIR NO. 1 (2023)
SRI CHAITANYA
6th-12th Class

**720
720
720**




**RANK
1**

For More Material Join: @JEEAdvanced_2025



71. In a fuel cell methanol is used as fuel and oxygen gas is used as an oxidizer. The reaction is
- $$CH_3OH(l) + \frac{3}{2}O_2(g) \rightarrow CO_2(g) + 2H_2O(l)$$
- At 298 K standard Gibb's energies of formation for $CH_3OH(l)$, $H_2O(l)$ and $CO_2(g)$ are -166.2 , -237.2 and $-394.4 kJ mol^{-1}$ respectively. If standard enthalpy of combustion of methanol is $-726 kJ mol^{-1}$, find the efficiency (in%) of the fuel cell:
72. The half life period for catalytic decomposition of AB_3 at initial pressure 50mm Hg is 4 hrs and at initial pressure 100 mm Hg it is 2hrs. Find the order of reaction.
73. Number of ambidentate nucleophiles among the following CN^- , SCN^- , OH^- , CNO^- , SO_4^{2-} , HCO_2^- , NO_2^- , NO_3^-
74. Number of compounds which forms aldehyde and ketone both in the same molecules on ozonolysis among the given compounds are



75. Number of lonepairs in I_3^-





Sri Chaitanya
Educational Institutions & Techno Schools

**Infinity
Learn**

39
Years of Excellence
1986-2024
★★★★★

300
300
MARKS



1
ALL INDIA
RANK

**EMPOWERING
EVERY STUDENT
TO BECOME EXTRAORDINARY**

**PROUDLY ACHIEVED
222 RANKS IN TOP 1000**

K C BASAVA REDDY
APPL.No. 240310618179*

SEIZES 4 RANKS IN TOP 10 IN ALL-INDIA RANKS

300
300
MARKS



ALL INDIA RANK

3
RANK

THOTAMSETTY NIKILESH
APPL.No. 240310813888*

300
300
MARKS



ALL INDIA RANK

6
RANK

HIMANSHU THAKOR
APPL.No. 240310580429*

300
300
MARKS



ALL INDIA RANK

9
RANK

REDDI ANIL
APPL.No. 240310238514

**SECURED 25 RANKS IN TOP 100
ALL INDIA OPEN CATEGORY**

**Sri Chaitanya - Nagpur
DLP Student**

1
RANK



G N NIRMALKUMAR
Appl.No. 240310385062*

9
RANK



REDDI ANIL
Appl.No. 240310238514*

14
RANK



K C BASAVA REDDY
Appl.No. 240310618179*

20
RANK



THOTAMSETTY NIKILESH
Appl.No. 240310813888*

21
RANK



A V TANISH REDDY
Appl.No. 240310807613

22
RANK



HIMANSHU THAKOR
Appl.No. 240310580429*

26
RANK



VEDANT SAINI
Appl.No. 240310182830

28
RANK



P MEET VIKRAMBHAI
Appl.No. 240310197524*

34
RANK



SANVI JAIN
Appl.No. 240310150036*

40
RANK



VISHARAD SRIVASTAVA
Appl.No. 240310046262*

43
RANK



T JAYADEV REDDY
Appl.No. 240310167365

46
RANK



ISHAAN GUPTA
Appl.No. 240310100229*

49
RANK



MAVURU JASWATHI
Appl.No. 240310542275*

52
RANK



DORISALA SRINIVASA REDDY
Appl.No. 240310682440*

53
RANK



ARCHIT RAHUL PATIL
Appl.No. 240310512311*

57
RANK



KRISHNA AGRAWAL
Appl.No. 240310285850*

60
RANK



AYUSH GANGAL
Appl.No. 240310270709

68
RANK



PALAGIRI SATHISH REDDY
Appl.No. 240310905497

70
RANK



MD K GHOUSE MOHIUDDIN
Appl.No. 240310176352

76
RANK



T V S SAI NAGA BHUSHAN
Appl.No. 240310889568

92
RANK



M M PRUTHVI RAJ
Appl.No. 24031084545

93
RANK



M SAI SIVA LOCHAN
Appl.No. 240310866829*

95
RANK



RAJDEEP MISHRA
Appl.No. 240310265621*

96
RANK



MANOJ SOHAN GAJULA
Appl.No. 240310529661

98
RANK



KRISHNA NARSARIA
Appl.No. 240310028285*



Below
100
All-India Open
Category Ranks

25

Below
500
All-India Open
Category Ranks

108

Below
1000
All-India Open
Category Ranks

222

Below
100
All-India All
Category Ranks

97

Below
1000
All Category
Ranks

888

**TOTAL QUALIFIED RANKS FOR
JEE ADVANCED-2024**

21,987

**FOR OFFER ON JEE MAIN &
JEE ADVANCED COURSES**



SCAN THE QR CODE

www.srichaitanya.net | Ph: 040 660 60606

* DLP | @ Gen-EWS | @ OBC-NCL | % SC | # ST | & GEN

For More Material Join: @JEEAdvanced_2025