



IIT-JEE
Batch – Growth (May) | Minor Test-01

Time: 3 Hours**Test Date: 9th June 2024****Maximum Marks: 300**

Name of the Candidate: _____ Roll No. _____

Centre of Examination (in Capitals): _____

Candidate's Signature: _____ Invigilator's Signature: _____

READ THE INSTRUCTIONS CAREFULLY

1. The candidates should not write their Roll Number anywhere else (except in the specified space) on the Test Booklet/Answer Sheet.
2. This Test Booklet consists of 90 questions.
3. This question paper is divided into three parts **PART A - MATHEMATICS, PART B - PHYSICS** and **PART C - CHEMISTRY** having 30 questions each and every **PART** has two sections.
 - (i) **Section-I** contains 20 multiple choice questions with only one correct option. Marking scheme: +4 for correct answer, 0 if not attempted and –1 in all other cases.
 - (ii) **Section-II** contains 10 questions the answer to only 5 questions, is an INTEGRAL VALUE.
Marking scheme: +4 for correct answer, 0 if not attempted and –1 in all other cases.
4. No candidate is allowed to carry any textual material, printed or written, bits of papers, mobile phone any electronic device etc., except the Identity Card inside the examination hall/room.
5. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
6. On completion of the test, the candidate must hand over the Answer Sheet to the invigilator on duty in the Room/Hall. However, the candidate is allowed to take away this Test Booklet with them.
7. **For the numerical based questions in Section-II of Mathematics, Physics, Chemistry, the answer should be in whole number only.**

TEST SYLLABUS

Batch – Growth (May) | Minor Test-01 9th June 2024

Mathematics:	"FOM-1 (Real Numbers Complex Numbers Even Numbers Odd Numbers, Prime Numbers, Composite Numbers, Co-Prime Numbers/ Relatively Prime Numbers, Twin Prime Numbers, LCM and HCF, Indices, Polynomial in One Variable, Degree of Polynomials, Some Special Types of Polynomials, Value and Zeros of a Polynomial, Roots of a Polynomial Equation, Remainder Theorem, Factor Theorem Factorization, Sets, Types of Sets, Laws of Algebra of Sets (Properties of Sets), Intervals as a subset of R Venn Diagram)"
Physics:	Basic Mathematics (Except Vector)
Chemistry:	Mole Concept & Concentration Terms-1 (Importance of Chemistry, Nature of matter, Sig.figure, Laws of chemical combination, Avogadro law, Dalton's atomic theory, Atomic and molecular masses, Till Average/Mean Atomic Mass) - Chemistry

Useful Data Chemistry:

Gas Constant	R	$= 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ $= 0.0821 \text{ Lit atm K}^{-1} \text{ mol}^{-1}$ $= 1.987 \approx 2 \text{ Cal K}^{-1} \text{ mol}^{-1}$
Avogadro's Number	N_a	$= 6.023 \times 10^{23}$
Planck's Constant	h	$= 6.626 \times 10^{-34} \text{ Js}$ $= 6.25 \times 10^{-27} \text{ erg.s}$
1 Faraday		$= 96500 \text{ Coulomb}$
1 calorie		$= 4.2 \text{ Joule}$
1 amu		$= 1.66 \times 10^{-27} \text{ kg}$
1 eV		$= 1.6 \times 10^{-19} \text{ J}$

Atomic No:

H = 1, D = 1, Li = 3, Na = 11, K = 19, Rb = 37, Cs = 55, F = 9, Ca = 20, He = 2, O = 8, Au = 79.

Atomic Masses:

He = 4, Mg = 24, C = 12, O = 16, N = 14, P = 31, Br = 80, Cu = 63.5, Fe = 56, Mn = 55, Pb = 207, Au = 197, Ag = 108, F = 19, H = 2, Cl = 35.5, Sn = 118.6

Useful Data Physics:

Acceleration due to gravity $g = 10 \text{ m / s}^2$

PART-A: MATHEMATICS

SECTION-I

1. Roster Form: $\{1, 2, 3, 4, 6, 8, 12, 24\}$, its set builder form is

- (A) $\{x : x \text{ is an odd natural number between } 20 \text{ and } 35\}$
 (B) $\{x : x \text{ is a natural number which divides } 24 \text{ completely}\}$
 (C) $\{x : x \text{ is an even natural number less than } 25\}$
 (D) $\{x : x \text{ is a letter used in the word 'MASSACHUSETTS'}\}$

Ans. (B)

Sol. Conceptual


2. Which of the following is a singleton set?

- (A) $\{x : -1 < x < 1, x \in \mathbb{Z}\}$
 (B) $\{x : -1 < x < 5, x \in \mathbb{Z}\}$
 (C) $\{x : x^2 = 1, x \in \mathbb{Z}\}$
 (D) None of these

Ans. (A)

Sol. $A = \{x : -1 < x < 1, x \in \mathbb{Z}\} = \{0\}$ i.e. a single for set: $-1 < o < 1$.

3. The shaded region in the given figure is



- (A) $A \cap (B \cup C)$
 (B) $A \cup (B \cap C)$
 (C) $A \cap (B - C)$
 (D) $A - (B \cup C)$


Ans. (D)

Sol. Conceptual

4. Which of the following are true?

- (A) $[3, 7] \subset (2, 10)$
 (B) $(0, \infty) \subset (4, \infty)$
 (C) $(5, 7] \subset [5, 7)$
 (D) $[2, 7] \subset (2.9, 8)$

Ans. (A)



5. The number of subsets of the power set of set $A = \{7, 10, 11\}$ is

- (A) 32
- (B) 16
- (C) 64
- (D) 256

Ans. (D)

Sol. Let $n(A) = m \Rightarrow n[P(A)] = 2^m$

$$n(P(P(A))) = 2^{2^n}$$

6. If $A = \{2, 3, 4, 8, 10\}$, $B = \{3, 4, 5, 10, 12\}$, $C = \{4, 5, 6, 12, 14\}$ then $(A \cap B) \cup (A \cap C)$ is equal to

- (A) $\{3, 4, 10\}$
- (B) $\{2, 8, 10\}$
- (C) $\{4, 5, 6\}$
- (D) $\{3, 5, 14\}$

Ans. (A)

Sol. $A \cap B = \{3, 4, 10\}$

$$A \cap C = \{4\}$$

$$(A \cap B) \cup (A \cap C) = \{3, 4, 10\}$$

7. Let $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$, $A = \{1, 2, 5\}$, $B = \{6, 7\}$, then $A \cap B'$ is

- (A) B'
- (B) A
- (C) A'
- (D) B

Ans. (B)

Sol. $B' = \{1, 2, 3, 4, 5, 8, 9, 10\}$ $A \cap B' = \{1, 2, 5\} = A$

8. $A \cup B = A \cap B$ iff:

- (A) $A \subset B$
- (B) $A = B$
- (C) $A \supset B$
- (D) $A \subset B$

Ans. (B)

Sol. Conceptual

9. The set $A = \{x : x \in \mathbb{R}, x^2 = 16 \text{ and } x^3 - 16x = 0\}$ is

- (A) Finite Set
- (B) Singleton set
- (C) Infinite set
- (D) not a well-defined collection

Ans. (A)

Sol. $x = \pm 4$ and $x = -4, 4, 0 \Rightarrow A = \{-4, 4\}$

10. The set of all type of triangles based on sides that may exist is

- (A) Null set
- (B) Not a well-defined collection
- (C) Universal set
- (D) Singleton set

Ans. (C)

Sol. Only three types of triangles are possible on based on side viz, isosceles, equilateral, scalene.

11. If $A = \{x : x \text{ is a multiple of } 4\}$ and $B = \{x : x \text{ is a multiple of } 6\}$, then $A \cap B$ consists of all multiple of

- (A) 16
- (B) 12
- (C) 8
- (D) 4

Ans. (B)

Sol. $\text{LCM}(4,6) = 12$

12. Let A and B be two sets, $(A \cup B)' \cup (A' \cap B)$ is equal to

- (A) A'
- (B) A
- (C) B'
- (D) None of these

Ans. (A)

Sol. $(A \cup B)' \cup (A' \cap B) = (A' \cap B') \cup (A' \cap B) = A' \cap (B' \cup B) = A'$

13. If $2 \leq 2x-4 < 6$ then $x \in$

- (A) $[3,5)$
- (B) $(3,5)$
- (C) $(3,5]$
- (D) None of these

Ans. (A)

Sol. $2 \leq 2x - 4 < 6$

$$6 \leq 2x < 10 \Rightarrow 3 \leq x < 5$$

14. If $5 < 5x - 10 < 15$, then

- (A) $-1 < x < 1$
- (B) $x > -1$
- (C) $-1 < x < 3$
- (D) $-x < 1$

Ans. (A)

Sol. $\Rightarrow 5 < 5x + 10 < 15$ (1)

Subtract "10" from eq.(1)

$$\Rightarrow -5 < 5x < 5 \quad \text{.....(2)}$$

divide by "5" to eq. (2)

$$\Rightarrow -1 < x < 1$$

15. If $-20 \leq 4x - 4 \leq 4$ then x is

- (A) $(-3, -1] \cup [1, 3)$
 (B) $[-4, 2]$
 (C) ϕ
 (D) None of these

Ans. (B)

Sol. $\Rightarrow -20 \leq 4x - 4 \leq 4$ (1)

Add "4" to eq.(1)

$$\Rightarrow -16 \leq 4x \leq 8 \quad \dots(2)$$

divide by "4" to eq. (2)

$$\Rightarrow -4 \leq x \leq 2$$

16. If $-9 < 3x - 6 < -3$, then

- (A) $x < 1$
 (B) $x > -1$
 (C) $-1 < x < 1$
 (D) $-x < 1$

Ans. (C)

Sol. $\Rightarrow -9 < 3x - 6 < -3$ (1)

Add "6" to eq.(1)

$$\Rightarrow -3 < 3x < 3 \quad \dots(2)$$

divide by "3" to eq.(2)

$$\Rightarrow -1 < x < 1$$

17. If $0 < 3x < 6$, then x belongs to

- (A) $(3, 4)$
 (B) $(0, 2)$
 (C) $\left(\frac{3}{2}, 4\right)$
 (D) None of these

Ans. (B)

Sol. $\Rightarrow 0 < 3x < 6$... (1)

Divide by "3" to eq.(1)

$$\Rightarrow 0 < x < 2$$

18. $(n^5 - n)$ is always divisible by $(n \in \mathbb{N})$

- (A) 30
 (B) 36
 (C) 6
 (D) 40

Ans. (A)

Sol. $n^5 - n = n(n^{5-1} - 1)$

Since 5 is a prime number, therefore, by Fermat's theorem $n^5 - n = n(n^{5-1} - 1)$ is divisible by 5.

$$\text{Also } n^5 - n = n(n-1)(n+1)(n^2+1)$$

But $n(n-1)(n+1)$ is divisible by $3!$. Therefore, $n^5 - n$ is divisible by both 5 and 6 i.e., 30. Hence correct answer is (A).

19. Simplify: $3^{-2} \times 81^{\frac{3}{4}} \div (729)^{\frac{-1}{3}}$

- (A) 9
(B) 27
(C) 81
(D) None of the Above

Ans. (B)

Sol. $3^{-2} \times 81^{\frac{3}{4}} \div (729)^{-1/3}$

$$\Rightarrow \frac{1}{9} \times (3^4)^{\frac{3}{4}} \div [(9)^3]^{-1/3}$$

$$\Rightarrow \frac{1}{9} \times 27 \div (9)^{-1}$$

$$\Rightarrow \frac{1}{9} \times 27 \times 9 \Rightarrow 27$$

20. If $i = \sqrt{-1}$, then $i^{324770} =$

- (A) -1
(B) 1
(C) 0
(D) None of these

Ans. (A)

Sol. $i^{324770} = i^2 = -1$


SECTION-II

21. If $U = \{1, 2, 3, 4, 5, 6\}$ be the universal set and $A = \{1, 3, 5\}$ and $B = \{2, 4, 5\}$ then $(A \cup B)' =$

Ans. (6)

Sol. $A \cup B = \{1, 2, 3, 4, 5\}$

$(A \cup B)' = U - (A \cup B) = \{6\}$



22. Number of integral values of x satisfying $-4 < 2x - 4 < 6$

Ans. (4)

Sol. $\Rightarrow -4 < 2x - 4 < 6 \quad \dots(1)$

Add "4" to eq.(1)

$\Rightarrow 0 < 2x < 10 \quad \dots(2)$

Divide by "2" to eq.(2)

$\Rightarrow 0 < x < 5$

as x is integer.

$$x = \{1, 2, 3, 4\}$$

23. If $7 + 4\sqrt{3} = (a + \sqrt{3})^2$, then $a =$

Ans. (2)

Sol. $7 + 4\sqrt{3} = (2 + \sqrt{3})^2$

24. If $f(x) \equiv 3x^3 - 2x^2 - 12x + 8$, then find the remainder when $f(x)$ is divided by $(x - 4)$.

Ans. (120)

Sol. $f(4) = 120$

25. The remainder when $12 \times 13 \times 14 + 13$ is divided by 13 is

Ans. (0)

Sol. The given number is $(12 \times 14 + 1) \times 13$ which is composite number having factor 13. Hence remainder is 0.

26. Number of prime numbers between 1 to 100 is

Ans. (25)

Sol. Conceptual

27. Number of integral values of x satisfying $-4 < 2x - 4 \leq 6$

Ans. (5)

Sol. $\Rightarrow -4 < 2x - 4 \leq 6 \dots(1)$

Add "4" to eq.(1)

$$\Rightarrow 0 < 2x \leq 10 \dots(2)$$

Divide by "2" to eq.(2)

$$\Rightarrow 0 < x \leq 5 \dots(3)$$

As x is integer then

$$x = \{1, 2, 3, 4, 5\}$$

28. If $3\frac{1}{2}$ m of wire is cut from a piece of 10 m long wire, length of remaining wire is $\frac{a}{2}$ m then $a =$

Ans. (13)

Sol. $10 - 3\frac{1}{2} = \frac{13}{2}$

29. If $x + y + z = 0$ are real then find the value of $\frac{x^2}{yz} + \frac{y^2}{zx} + \frac{z^2}{xy}$.

Ans. (3)

Sol. $\frac{x^2}{yz} + \frac{y^2}{zx} + \frac{z^2}{xy} = \frac{x^3 + y^3 + z^3}{xyz} = \frac{3xyz}{xyz}$

30. Sum of real roots of $x^3 - 7x - 6 = 0$ is

Ans. (0)

Sol. Roots are $x = -2, -1, 3$.

PART-B: PHYSICS

SECTION-I

31. What does the derivative represent in calculus?

- (A) Slope of the tangent to the curve
- (B) Area under the curve
- (C) Accumulated sum
- (D) Average value

Ans. (A)

Sol. Theory

32. The roots of quadratic equation: $6x^2 - x - 2 = 0$ are

- (A) $-\frac{1}{3}, \frac{1}{2}$
- (B) $-\frac{1}{3}, -\frac{1}{2}$
- (C) $\frac{1}{3}, -\frac{1}{2}$
- (D) $\frac{2}{3}, -\frac{1}{2}$

Ans. (D)

Sol. $x = \frac{1 \pm \sqrt{1+48}}{12} = \frac{2}{3}, -\frac{1}{2}$

33. The value of $(126)^{1/3}$ up to three decimal places is

- (A) 5.011
- (B) 5.012
- (C) 5.013
- (D) 5.014

Ans. (C)

Sol. $(126)^{1/3}$ can also be written as the cube root of 126.

$$\begin{aligned}
 & (126)^{1/3} \\
 &= (1+125)^{1/3} \\
 &= (1+5^3)^{1/3} \\
 &= 5 \left(\frac{1}{5^3} + 1 \right)^{1/3} \\
 &\approx 5 \left(1 + \frac{1}{3} \times \frac{1}{5^3} \right) \\
 &= 5.013
 \end{aligned}$$

34. Which of the following statements is not correct?

- (A) $\log_{10} 10 = 1$
 (B) $\log (2 + 3) = \log (2 \times 3)$
 (C) $\log_{10} 1 = 0$
 (D) $\log (1 + 2 + 3) = \log 1 + \log 2 + \log 3$

Ans. (B)

Sol. Conceptual

35. $\frac{m+n}{m-n} = \frac{7}{3}$, then $\frac{m^2 - n^2}{m^2 + n^2} =$

- (A) $\frac{29}{21}$
 (B) $\frac{21}{11}$
 (C) $\frac{21}{29}$
 (D) $\frac{11}{21}$

Ans. (C)

Sol. $\frac{m+n}{m-n} = \frac{7}{3} \Rightarrow \frac{m}{n} = \frac{7+3}{7-3} = \frac{5}{2} \Rightarrow \frac{m^2 - n^2}{m^2 + n^2} = \frac{21}{29}$

36. $\int (x^e + e^x + e^e) dx =$

- (A) $x^e + e^x + c$
 (B) $\frac{x^{e+1}}{e+1} + e^x + \frac{e^{e+1}}{e+1} + c$
 (C) $\frac{x^{e+1}}{e+1} + \frac{e^{x+1}}{x+1} + \frac{e^{e+1}}{e+1} + c$
 (D) $\frac{x^{e+1}}{e+1} + e^x + xe^e + c$

Ans. (D)

Sol. $\int (x^e + e^x + e^e) dx = \int x^e dx + \int e^x dx + \int e^e dx$
 $= \frac{x^{e+1}}{e+1} + e^x + e^e \int dx + c$
 $= \frac{x^{e+1}}{e+1} + e^x + xe^e + c$


37. Radius of circle $x^2 + y^2 = 400$ is

- (A) 400
 (B) 40
 (C) 20
 (D) 10

Ans. (C)

Sol. Equation of circle centered at origin: $x^2 + y^2 = R^2$.

38. In the given triangle, find the perimeter is _____ cm.



- (A) $15(1 + \sqrt{2} + \sqrt{3})$ cm
 (B) $15(3 + \sqrt{3} + \sqrt{2})$ cm
 (C) $15(2 + \sqrt{3} + \sqrt{5})$ cm
 (D) $15(3 - \sqrt{3} + \sqrt{5})$ cm

Ans. (B)


Sol. $AD = 30 \sin 30^\circ = 15$ cm

$$AB = AD \operatorname{cosec} 45^\circ = 15\sqrt{2} \text{ cm}$$

$$BD = AB \cos 45^\circ = 15 \text{ cm}$$

$$DC = 30 \cos 30^\circ = 15\sqrt{3} \text{ cm}$$

$$\text{Perimeter} = AB + BD + DC + CA = 15\sqrt{2} + 15 + 15\sqrt{3} + 30 = 15(3 + \sqrt{3} + \sqrt{2}) \text{ cm}$$



39. Evaluate the integral: $\int_0^2 (3x^2 + 4x - 5) dx$.

- (A) 3
 (B) 4
 (C) 5
 (D) 6

Ans. (D)

Sol. $\int_0^2 (3x^2 + 4x - 5) dx = \left[x^3 + 2x^2 - 5x \right]_0^2 = 8 + 8 - 10 = 6$

40. A ship is sailing along a straight line: $y = x$, while another ship sails along the line: $y + 2x = 150$. The coordinates of the point where the two ships may possibly collide are:

- (A) (50, 100)
 (B) (100, 100)
 (C) (150, 100)
 (D) (50, 50)

Ans. (D)

Sol. They may collide at the intersection points of the two lines.

$$\text{Solving } y = x \quad \dots(1)$$

$$y + 2x = 150 \quad \dots(2)$$

$$\text{We get } x = 50, y = 50$$

41. Which of the following is not a trigonometric identity?

(A) $1 - \sec^2 x = -\tan^2 x$

(B) $\sin(-x) = \sin x$

(C) $\cos 2x = \frac{1 - \tan^2 x}{1 + \tan^2 x}$

(D) $\sin 2x = \frac{2 \tan x}{1 + \tan^2 x}$

Ans. (B)

Sol. Trigonometric Identities.

42. A particle is moving along the x-axis, and its position (x-coordinate) varies with time t as $x = 7t^2 - 8t + 9$, where x is in meters and t is in time. Then the velocity of the particle at t = 0 is

$$\left(v = \frac{dx}{dt} \right)$$

(A) 9 m/s

(B) 8 m/s

(C) 6 m/s

(D) -8 m/s

Ans. (D)

Sol. $v = \frac{dx}{dt} = \frac{d(7t^2 - 8t + 9)}{dt} = 14t - 8$

v at t = 0 will be v = -8 m/s

43. Find $\frac{dy}{dx}$, if $y = (x + e^x) \sin x$.

(A) $e^x (\cos x + \sin x) + x \cos x + \sin x$

(B) $e^x (\cos x + \sin x) + x \sin x + \cos x$

(C) $(1 + e^x)(\cos x + \sin x)$

(D) $(1 + e^x)(\sin x - \cos x)$

Ans. (A)

Sol. $\frac{d(y)}{dx} = \frac{d((x + e^x) \sin x)}{dx}$

Using the product rule: $\frac{d(y)}{dx} = (x + e^x) \frac{d(\sin x)}{dx} + (\sin x) \frac{d(x + e^x)}{dx}$

$$= (x + e^x) \cos x + (\sin x)(1 + e^x) = e^x (\cos x + \sin x) + x \cos x + \sin x$$

44. $\sin 75^\circ =$

(A) $\frac{2 - \sqrt{3}}{2\sqrt{2}}$

(B) $\frac{\sqrt{3} - 1}{2\sqrt{2}}$

(C) $\frac{1 + \sqrt{3}}{2\sqrt{2}}$

(D) $\frac{1 + \sqrt{3}}{\sqrt{2}}$

Ans. (C)

Sol. $\sin 75^\circ = \sin (30^\circ + 45^\circ)$

$$= \sin 30^\circ \cos 45^\circ + \cos 30^\circ \sin 45^\circ$$

$$= \frac{1}{2} \times \frac{1}{\sqrt{2}} + \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} = \frac{1 + \sqrt{3}}{2\sqrt{2}}$$

45. If $f(x) = x^4 + 4e^x$, then $f(0) =$

(A) 0

(B) 1

(C) 2

(D) 4

Ans. (D)

Sol. $f(0) = 0^4 + 4e^0 = 0 + 4 \times 1 = 4$

46. $\int_0^{\pi/2} (\sin \theta + \cos \theta) d\theta$

(A) 1

(B) 0

(C) 2


(D) -1

Ans. (C)

Sol. $\int_0^{\pi/2} (\sin \theta + \cos \theta) d\theta = (-\cos \theta + \sin \theta)_0^{\pi/2}$

$$= [-0 + 1 - (-1) - 0] = 2$$

47. Equation of the straight line shown in the graph will be



(A) $\frac{y}{9} - \frac{x}{4} = 1$

(B) $\frac{y}{9} + \frac{x}{4} = 1$

(C) $\frac{y}{4} + \frac{x}{9} = 1$

(D) $9y = 4x - 36$

Ans. (D)**Sol.** Slope of line $= \frac{\Delta y}{\Delta x} = \frac{0 - 4}{9 - 0} = -\frac{4}{9}$; y intercept = -4

Equation of line $= y = mx + c \Rightarrow y = -\frac{4}{9}x - 4 \Rightarrow 9y = -4x - 36$

48. A stone is dropped into a quiet lake and waves move in circles at the speed of 5 cm/s. At the instant when the radius of the circular wave is 8 cm, how fast is the enclosed area increasing?

(A) $80 \pi \text{ cm}^2 / \text{s}$

(B) $90 \pi \text{ cm}^2 / \text{s}$

(C) $85 \pi \text{ cm}^2 / \text{s}$

(D) $89 \pi \text{ cm}^2 / \text{s}$

Ans. (A)**Sol.** Enclosed area: $A = \pi r^2$

So, $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$

Here $r = 8 \text{ cm}$, $\frac{dr}{dt} = 5 \text{ cm/s}$

$$\Rightarrow \frac{dA}{dt} = (2\pi)(8)(5) = 80 \pi \text{ cm}^2 / \text{s}$$

49. Which of the following expression is equal to $\cos 2\theta$?

(A) $\sin^2 \theta - 1$

(B) $2\sin^2 \theta - 1$

(C) $2\cos^2 \theta - 1$

(D) $2\tan^2 \theta - 1$

Ans. (C)**Sol.** Identity**50.** If $f(x) = (\sin x)^2$, then the value of $f(30^\circ)$ is

(A) 0.75

(B) 0.5

(C) 0.25

(D) $\sin^2(90^\circ)$ **Ans.** (C)

Sol. $f(30^\circ) = \sin^2(30^\circ) = \left(\frac{1}{2}\right)^2 = 0.25$

SECTION-II**51.** Distance between two points (5, 4) and (10, -8) is ____.**Ans.** (13)

Sol. Distance = $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(10 - 5)^2 + (-8 - 4)^2} = \sqrt{5^2 + 12^2} = 13$

52. Find the slope of a line passing through (8, 8) and (4, -4).**Ans.** (3)

Sol. Slope = $\frac{\Delta y}{\Delta x} = \frac{8 - (-4)}{8 - 4} = \frac{12}{4} = 3$

53. Distance of point (13, 4) from the line $y = -13$ is ____.**Ans.** (17)**Sol.** Conceptual**54.** If $f(1) = 2$, $f(2) = 4$ and $f(3) = 12$. Then $f(1) + f(2) =$ ____.**Ans.** (6)**Sol.** Conceptual

55. A particle is moving with initial velocity $u = 3$ m/s on x-axis. It's acceleration a varies with time t (in seconds) as $a = 3t - 1$ m/s², then the velocity of the particle at $t = 4$ seconds will be ____ m/s
 $\left(\int dv = \int a dt\right)$.

Ans. (23)

Sol. $\frac{dv}{dt} = a \Rightarrow dv = (3t - 1) dt$. Integrating both sides, we get

$$\int_3^v dv = \int_0^4 (3t - 1) dt \Rightarrow [v]_3^v = \left[3 \frac{t^2}{2} - t\right]_0^4 \Rightarrow v - 3 = 24 - 4 \Rightarrow v = 23$$

56. If $\cos(300^\circ) = \sqrt{\frac{\alpha}{\beta}}$, where α and β are co-prime numbers, then the value of $\alpha + \beta$ is ____.

Ans. (5)

Sol. $\cos(300^\circ) = \frac{1}{2} = \sqrt{\frac{1}{4}}$

57. If the position x of a particle varies with time t as $x = \frac{9}{\pi} \sin\left(2\pi t + \frac{\pi}{3}\right)$, where x is in meters and t is in seconds, then the velocity v of the particle at time $t = 2$ s is ____ m/s $\left(v = \frac{dx}{dt}\right)$.

Ans. (9)

$$\text{Sol. } v = \frac{dx}{dt} = \frac{d\left(\frac{9}{\pi} \sin\left(2\pi t + \frac{\pi}{3}\right)\right)}{dt} = \frac{9}{\pi} \frac{d\left(\sin\left(2\pi t + \frac{\pi}{3}\right)\right)}{dt} = \frac{9}{\pi} \times 2\pi \cos\left(2\pi t + \frac{\pi}{3}\right) = 18 \cos\left(2\pi t + \frac{\pi}{3}\right)$$


$$\text{time } t = 2 \text{ s, } v = 18 \cos\left(2\pi \times 2 + \frac{\pi}{3}\right) = 18 \cos \frac{\pi}{3} = 9$$

58. If $\int_0^{\pi} 40 \sin^2\left(\frac{\theta}{2}\right) d\theta = n\pi$. Then the value of n is ____.

Ans. (20)**Sol.** Using double angle formula

$$\begin{aligned} \int_0^{\pi} 40 \sin^2\left(\frac{\theta}{2}\right) d\theta &= \int_0^{\pi} 20(1 - \cos \theta) d\theta = 20 \int_0^{\pi} d\theta - 20 \int_0^{\pi} \cos \theta d\theta \\ &= 20[\theta]_0^{\pi} - 20[\sin \theta]_0^{\pi} = 20\pi \end{aligned}$$

59. Find the value of $20 \sin \theta$, where θ is the angle represented in the right angled triangle shown

**Ans.** (16)

$$\text{Sol. } \sin \theta = \frac{\sqrt{25^2 - 15^2}}{25} = \frac{4}{5}; \quad 20 \sin \theta = 16$$

60. If the velocity v of a particle varies with time t as $v = 4t - 2$ m/s, where t is in seconds. Then the displacement of the particle from $t = 0$ to $t = 4$ s is ____ m ($\int dx = \int v dt$).

Ans. (24)

$$\text{Sol. } \frac{dx}{dt} = v \Rightarrow dx = v dt \Rightarrow dx = (4t - 2) dt$$

Integrating both sides, we get:

$$\Rightarrow \int_{x_i}^{x_f} dx = \int_0^4 (4t - 2) dt \Rightarrow [x]_{x_i}^{x_f} = [2t^2 - 2t]_0^4 = 24$$

PART-C: CHEMISTRY SECTION-I

61. Which of the following is a mixture?

- (A) Oxygen
- (B) Sand
- (C) Pure Water
- (D) Mercury

Ans. (B)

Sol. Conceptual

62. How many basic laws are required to govern the combination of elements to form compounds?

- (A) 6
- (B) 5
- (C) 4
- (D) 1

Ans. (B)

Sol. Five basic laws are required to govern the combination of elements to form compounds. They are Law of Conservation of Mass, Law of Definite Proportions, Law of Multiple Proportions, Gay Lussac's Law of Gaseous Volumes, and Avogadro's Law.

63. 10 g of CaCO_3 contains

- (A) 10 mol of CaCO_3
- (B) 1 mol of Ca
- (C) 6×10^{22} atoms of Ca
- (D) None of these

Ans. (C)

Sol. Mole of $\text{CaCO}_3 = \frac{10}{100} = 0.1$

So, mole of Ca = 0.1

Number of atom of Ca = $0.1 \times 6 \times 10^{23} = 6 \times 10^{22}$

64. Who proposed Law of Conservation of Mass?

- (A) Antoine Lavoisier
- (B) Joseph Proust
- (C) Lorenzo Romano
- (D) Joseph Louis

Ans. (A)

Sol. Antoine Lavoisier conducted many experiments regarding combustion and noticed various physical and chemical changes and there is no change in overall mass. Hence he came to a conclusion that mass can neither be created nor destroyed i.e. Law of Conservation of Mass.

65. How many atoms of hydrogen are present in 40 molecules of $(\text{NH}_4)_2\text{SO}_4$.

- (A) 40
- (B) 80
- (C) 160
- (D) 320

Ans. (D)

Sol. 1 molecule of $(\text{NH}_4)_2\text{SO}_4$ has 8 atoms of Hydrogen.

66. A gaseous mixture contains oxygen and nitrogen in the ratio of 1:4 by mass. The ratio of their numbers of molecules is

- (A) 1:4
- (B) 1:8
- (C) 7:32
- (D) 3:16

Ans. (C)

Sol. $\frac{W}{32} : \frac{4W}{28}$

67. The largest number of molecules is in

- (A) 36 g of water
- (B) 28 g of carbon monoxide
- (C) 46 g of ethyl alcohol [C₂H₅OH]
- (D) 54 g of nitrogen pentoxide [N₂O₅]

Ans. (A)

Sol. $n_{\text{H}_2\text{O}} = \frac{36}{18} = 2$

$$N_{\text{H}_2\text{O}} = 2 \times N_A$$

68. 3.42 g of sucrose are dissolved in 18 g of water in a beaker. The number of oxygen atoms in the solution are

- (A) 6.68×10^{23}
- (B) 6.09×10^{22}
- (C) 6.022×10^{23}
- (D) 6.022×10^{21}

Ans. (A)

Sol. 0.01 mol of sucrose $0.01 \times 11 \times N_A$ atoms of O

$$\Rightarrow 0.11 \times N_A \text{ atoms of O}$$

$$n_{\text{H}_2\text{O}} = m/M$$

$$n_{\text{H}_2\text{O}}$$

$$= 18 \text{ g} / 18 \text{ g}$$

$$= 1 \text{ mole}$$

1 mol of water has N_A atoms of O

$$\text{Total number of O atoms} = (0.11 + 1) \times N_A$$

$$= 1.11 \times N_A$$

$$= 1.11 \times 6.022 \times 10^{23}$$

$$= 6.68 \times 10^{23}$$

69. How many molecules of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ will have 954 atoms of oxygen?

- (A) 954
- (B) 238
- (C) 191
- (D) 106

Ans. (D)

Sol. 1 molecule of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ has 9 atoms of Oxygen.

70. Law of Definite Composition is also known as

- (A) Law of Multiple Proportions
- (B) Avogadro's Law
- (C) Law of Definite Proportion
- (D) Law of Conservation of Mass

Ans. (C)

Sol. Joseph Proust worked about the composition of elements present in a compound experimentally, he concluded that from any source, a particular compound always contains the same elements in the same proportion by mass/weight. Hence it can also be known as the Law of Definite Proportion.

71. Which of the following correctly represents 360 g of water?

- (i) 2 moles of H_2O
- (ii) 20 moles of water
- (iii) 6.022×10^{23} molecules of water
- (iv) 1.2044×10^{25} molecules of water
- (A) (i)
- (B) (i) and (iv)
- (C) (ii) and (iii)
- (D) (ii) and (iv)

Ans. (D)

Sol. (i) Mass of 2 mol of water = $2 \times 18 = 36$ g

(ii) Mass of 20 mol of water = $20 \times 18 = 360$ g

(iii) Mass of 6.022×10^{23} molecules of water = 1 mol = 18 g

(iv) Mass of 1.2044×10^{25} molecules of water = $\frac{1.2044 \times 10^{25}}{6.022 \times 10^{23}}$ mol = 20 mol = 20×18 g = 360 g

72. Which of the following is not a law of chemical combination?

- (A) Law of Multiple Proportions
- (B) Avogadro's Law
- (C) Law of Definite Proportion
- (D) Law of Conservation of volume

Ans. (D)

Sol. Five basic laws are required to govern the combination of elements to form compounds. They are Law of Conservation of Mass, Law of Definite Proportions, Law of Multiple Proportions, Gay Lussac's Law of Gaseous Volumes, and Avogadro's Law.

73. Which sample contains the largest number of atoms?

- (A) 1 mg of C_4H_{10}
- (B) 1 mg of N_2
- (C) 1 mg of Na
- (D) 1 ml of H_2O

Ans. (D)

Sol. 1 mg of C_4H_{10} (14 atoms in 1 molecule)

$$= (14 \times N_A \times 10^{-3}) / 58 = 2.41 \times 10^{-4} \times N_A \text{ atoms}$$

1 mg of N_2 (2 atoms in 1 molecule)

$$= (2N_A \times 10^{-3}) / 28 = 7.14 \times 10^{-5} \times N_A \text{ atoms}$$

1 mg of Na (1 atom in 1 molecule)

$$= (N_A \times 10^{-3}) / 23 = 4.34 \times 10^{-5} \times N_A \text{ atoms}$$

1 ml of H_2O (3 atoms in 1 molecule)

$$\text{Weight of } H_2O = 1 \text{ ml} \times (1 \text{ gm/ml}) = 1 \text{ g}$$

$$= 1 \text{ g of } H_2O = 3N_A / 18 = 2.41 \times 10^{-4} \times N_A \text{ atoms}$$

Thus, 1 ml of H_2O has largest number of atoms

74. Who proposed the Law of Definite Composition?

- (A) Joseph Proust
- (B) Lorenzo Romano
- (C) Joseph Louis
- (D) Antoine Lavoisier

Ans. (A)

Sol. Joseph Proust worked about the composition of elements present in a compound experimentally, he concluded that from any source, a particular compound always contains the same elements in the same proportion by mass/weight.

75. Which has the maximum number of molecules?

- (A) 7 g N_2
- (B) 2 g H_2
- (C) 16 g NO_2
- (D) 16 g O_2

Ans. (B)

Sol. Number of moles = $\frac{\text{Given mass of substance}}{\text{Molecular mass of substance}}$

$$n = \frac{m}{M}$$

$$7\text{g } N_2 \rightarrow \frac{7}{28} = 0.25 \text{ moles of molecules}$$

$$2\text{ g } H_2 \rightarrow \frac{2}{2} = 1 \text{ mole of molecules}$$

$$16\text{ g } NO_2 \rightarrow \frac{16}{46} = 0.347 \text{ moles of molecules}$$

$$16\text{ g } O_2 \rightarrow \frac{16}{32} = 0.5 \text{ moles of molecules}$$

Thus, 2 g H_2 has the most number of molecules.

76. A sample of $Mg_3(PO_4)_2$ contains 72 g Mg in the sample. Calculate the number of g molecule of oxygen in the sample.

- (A) 2
- (B) 3
- (C) 3.5
- (D) 4

Ans. (D)

Sol. 1 mole $Mg_3(PO_4)_2$
8 moles of O atoms
4 moles of O_2 molecules
4 gm molecules of O_2 are present.

77. What mass (in gram) of Na_2SO_4 would contain the same number of total ions as present in 930 g of $Ca_3(PO_4)_2$?

- (A) 710
(B) 800
(C) 200
(D) 500

Ans. (A)

Sol. 930 gm of $\text{Ca}_3(\text{PO}_4)_2 = 3$ mole of $\text{Ca}_3(\text{PO}_4)_2$

$\left[1\text{Ca}_3(\text{PO}_4)_2 \text{ contains } 3\text{Ca}^{+2}, 2\text{PO}_4^{-3} \right]$

= 15 moles of ions

= 5 moles of Na_2SO_4

= 15 mole of ions weight of $\text{Na}_2\text{SO}_4 = 5 \times 142$

= 710 g

78. A sample of Cl contains 80% Cl^{35} and 20% Cl^{37} isotopes. The average number of neutrons per atom is

- (A) 18.4
(B) 19.4
(C) 35.4
(D) 20.4

Ans. (A)

Sol. Average atomic weight = $\sum X_i M_i = 35 \times 0.8 + 37 \times 0.2 = 35.4$

Cl^{35} contains neutron = $35 - 17 = 18$

Cl^{37} contains neutron = $37 - 17 = 20$

Average number of neutrons = $\frac{18 \times 80 + 20 \times 20}{100} = 18.4$

79. A sample of $\text{K}_2\text{SO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 24\text{H}_2\text{O}$ contains 6.4 kg oxygen. Calculate mass of sulphur (in gram) in same sample.

- (A) 1280
(B) 1000
(C) 128
(D) 12.8

Ans. (A)

Sol. $\text{K}_2\text{SO}_4 \cdot \text{Al}_2(\text{SO}_4)_3 \cdot 24 \text{H}_2\text{O}$

6.4 kg oxygen

$[\text{moles of substance} \times 40] = \text{number of mole of oxygen } 40 \times x \times 16 = 6400 \text{ g wt of oxygen}$

$x = 10$ moles

number of moles of S = 4×10 moles = 40 moles

weight of sulphur = $40 \times 32 = 1280$ g

80. A 448 L vessel contains $\text{O}_2(\text{g})$ and $\text{CO}_2(\text{g})$ in 2: 3 ratio at 1 atm and 0°C . Calculate the number of moles of $\text{CO}_2(\text{g})$ present in the vessel.

- (A) 12
(B) 1.1
(C) 1.2
(D) 120

Ans. (A)

Sol. 448ℓ O_2 and $CO_2(g)$ 2 : 3

$$\text{Volume of } CO_2 = \frac{3}{5} \times 448 = 268.8\ell$$


$$\text{Number of moles of } CO_2 = \frac{268.8L}{22.4L} = 12 \text{ mole}$$

SECTION-II

81. Find the value of 'x', if hydrated salt $A_2SO_4 \cdot xH_2O$ undergoes 45% loss in mass on heating and becomes anhydrous (where atomic weight of 'A' is 7).

Ans. (5)

Sol. $A_2SO_4 \cdot H_2O$



Molar mass of $A_2SO_4 = 110$

$$\% \text{ of } H_2O = \frac{x \times 18}{110 + 18x} \times 100 = 45$$

$$1800x = 4950 + 810x$$

$$1800x = 810x = 4950$$

$$990x = 4950$$

$$x = 5$$

82. Find number of moles of H_2 in 10 g of H_2 .

Ans. (5)

Sol. Given mass of $H_2 = m = 10 \text{ g}$

Molar Mass of $H_2 = M = 2 \text{ g}$

We know that,

$$\text{Number of moles of a substance} = \frac{\text{Given mass of the substance}}{\text{Molar mass of the substance}}$$

$$n = \frac{m}{M} = \frac{10 \text{ g}}{2 \text{ g}} = 5 \text{ moles}$$

83. How many moles is 12.044×10^{23} atoms of He?

Ans. (2)

Number of atoms of the He = $N = 12.044 \times 10^{23}$ Avogadro's Constant = $N_A = 6.022 \times 10^{23}$

We know that,

$$\text{Number of moles of a substance} = \frac{\text{Number of particles of the substance}}{\text{Avogadro's Constant}}$$

$$= N/N_A = 12.044 \times 10^{23} / 6.022 \times 10^{23} = 2 \text{ moles of He atoms}$$

84. How many moles are there in 52 g of Helium?

Ans. (13)

Sol. Atomic Mass of Helium = 4u

So, Molar Mass of Helium = 4 gram

It means that Mass of 1 mole of Helium = 4 grams

We can also write as

$$4 \text{ grams} = 1 \text{ mole}$$

1 gram = $\frac{1}{4}$ mole 52 grams = $(\frac{1}{4}) \times 52$ mole

52 grams = 13 moles

So, $n = \text{Given mass} / \text{Molar mass} = 52/4 = 13$ moles

85. Calculate the molar mass of Calcium Carbonate (CaCO_3) in g/mol units.

Ans. (100)

Sol. M of $\text{CaCO}_3 = (1 \times \text{mass of Ca}) + (1 \times \text{mass of C}) + (3 \times \text{mass of O})$

M of $\text{CaCO}_3 = (1 \times 40) + (1 \times 12) + (3 \times 16)$

M of $\text{CaCO}_3 = (1 \times 40) + (1 \times 12) + (3 \times 16)$

M of $\text{CaCO}_3 = 100$ g/mol

86. From 160 g of SO_2 (g) sample, 1.2046×10^{24} molecules of SO_2 are removed then find out the volume (in L) of left over SO_2 (g) at 0°C and 1 atm. Report your answer to nearest integer.

Ans. (11)

Sol. $n_{\text{removed}} \text{ of } \text{SO}_2 = \frac{1.2046 \times 10^{24}}{6.022 \times 10^{23}} = 2$

$n_{\text{initial}} = \frac{160}{64} = 2.5$

$n_{\text{remaining}} = 2.5 - 2 = 0.5$

Volume of $\text{SO}_2 = 22.4 \times 0.5 = 11.2$ L

87. 14 g of Nitrogen gas and 22 g of CO_2 gas are mixed together. Find the volume (in L) of gaseous mixture at 0°C and 1 atm. Report your answer to nearest integer.

Ans. (22)

Sol. Mole of $\text{N}_2 = \frac{14}{28} = \frac{1}{2}$

Mole of $\text{CO}_2 = \frac{22}{44} = \frac{1}{2}$

Total mole = $\frac{1}{2} + \frac{1}{2} = 1$

Total volume = $22.4 \times 1 = 22.4$


88. Calculate the volume in litres of 20 g hydrogen gas at 0°C and 1 atm.

Ans. (224)

Sol. No. of moles of hydrogen gas = mass + atomic weight

$= \frac{20 \text{ gm}}{2 \text{ gm}} = 10 \text{ mol}$

Volume of hydrogen gas = 10×22.4 lt



89. Calculate the volume in litres of 142 g chlorine gas at 0°C and 1 atm. Report your answer to nearest integer.

Ans. (45)

Sol. Volume of $\text{CO}_2 = \left(\frac{142}{71}\right) \times 22.4 = 44.8$

90. Find the volume occupied by 16 g of ozone at 0°C and 1 atm. If your answer is x L. Then $\frac{x}{7.5}$ is

Ans. (1)

Sol. Volume of Ozone = $\frac{16}{48} \times (22.4) = \text{Approx } 7.5$

Unacademy Centres across India

