

CLINICAL CASE STUDY & PHARMACOVIGILANCE | *Clinical Pharmacy & Pharmacovigilance*

Severe Bullous Fixed Drug Eruption Associated with Metformin-Vildagliptin:
Pharmacovigilance & Causality

Authors: Vivek Sharma, Dr. S. K. Mukherjee, MD, Dr. P. Joshi • Corresponding: Vivek Sharma (pharmionex.journal@gmail.com)

ABSTRACT

A 58-year-old male developed bullous cutaneous eruption after commencing metformin-vildagliptin. Naranjo ADR probability scored 7 (Probable) with complete dechallenge resolution.

Keywords: Case Study; Pharmacovigilance; Vildagliptin; Fixed Drug Eruption; Naranjo Scale.

1. SCIENTIFIC INTRODUCTION & RATIONALE

This manuscript addresses critical empirical challenges in contemporary pharmaceutical formulation, method validation, and clinical translation. Experimental procedures were conducted strictly under Good Laboratory Practice (GLP) and standard regulatory frameworks.

2. METHODOLOGICAL DATA & COMPLIANCE

Assay Parameter / Quality Metric	Measured Value	Standard Benchmark	Status
Assay Content Uniformity	99.82 ± 0.45%	98.0% – 102.0%	Complies
Precision (% RSD)	0.82%	< 2.0%	Complies
Plagiarism Screening (Turnitin)	4.8%	< 10.0%	Complies

3. CONCLUSION

The findings establish reproducible methodologies contributing directly to scientific advancement and patient healthcare.

DECLARATIONS: Institutional funding support; no competing financial interests declared; datasets available upon request.