
Phase Theory: A Unified and Deployable Framework for Matter, Geometry, and Emergence

Marlon Hanks

Independent Researcher, Dust LLC, United States

Submitted: January 9, 2026 • Revised: May 2026

Subject Areas: Foundations of Physics / Unified Theory • **MSC:** 81T13, 83C45, 57R57

Abstract

Phase Theory proposes that phase is the sole fundamental constituent of physical reality. Space, time, matter, light, and geometry are not primitive entities but emergent structures arising from the global organization, topology, and consistency of a single phase-bearing substrate. The present work revises the original Phase Theory preprint into a fully deployable framework: the central dynamical object is identified as a gauged nonlinear sigma model (NLSM) with target manifold $T = U(1) \times SU(2) \times SU(3)$, coupled to induced gravity via the Sakharov mechanism, with matter identified as topological solitons — skyrmions, vortices, and holonomy defects. Three core ontological commitments are preserved: (i) all structure emerges from a single phase-valued field, (ii) particles are topological defects, and (iii) spacetime geometry is emergent. The framework is fully specified, admits a lattice formulation suitable for numerical simulation with existing codes, and yields six falsifiable predictions with quantitative numbers. Two predictions from the original preprint are explicitly retracted with corrected parametric estimates. The theoretical status of all open problems is assessed

honestly.

Keywords: phase ontology, nonlinear sigma model, topological solitons, induced gravity, emergent spacetime, quantum foundations, Skyrme model, lattice QCD

I. Introduction

The two pillars of twentieth-century physics — quantum mechanics and general relativity — have resisted unification for nearly a century. Quantum field theory treats spacetime as a fixed background and populates it with fields whose quanta are particles. General relativity treats spacetime as dynamical, encoding gravity through curvature, but remains fundamentally classical. Direct quantization of gravity encounters perturbative non-renormalizability [1, 2], while the Standard Model depends on approximately twenty-four free parameters whose values must be inserted by hand [3].

Beyond technical difficulties, a deeper conceptual tension persists. Quantum mechanics assigns a privileged role to measurement; general relativity describes a universe independent of observation. The ontological status of spacetime is contested. The proliferation of unification proposals — string theory [4, 5], loop quantum gravity [6], causal set theory [7], causal dynamical triangulations [8] — reflects the absence of experimental guidance at the Planck scale.

Phase Theory takes a different starting point: rather than quantizing geometry or extending the field content of the Standard Model, it asks whether all physical structure — particles, geometry, and dynamics — might emerge from a single topological substrate. The key insight, formalized in this revision, is that this substrate is precisely described by a gauged nonlinear sigma model (NLSM) with the Standard Model gauge group as target manifold, coupled to emergent gravity through the Sakharov induced-gravity mechanism [9].

This paper proceeds as follows. Section II establishes the translation between original Phase Theory language and standard physics. Section III presents the

complete action. Section IV derives the equations of motion. Section V classifies the topological defect spectrum. Section VI develops the perturbative expansion and Feynman rules. Section VII addresses renormalization and UV completion. Section VIII presents the complete lattice formulation. Section IX derives the emergent metric and gravitational dynamics. Section X discusses the particle spectrum and generation mass ratios. Section XI analyzes the dark sector. Section XII addresses causality. Section XIII enumerates falsifiable predictions. Section XIV provides an honest assessment of open problems. Appendices collect parameter tables, computation priorities, and comparison with the original preprint claims.

II. Translation Key and Core Identification

The original Phase Theory preprint introduces terminology that maps precisely onto standard mathematical physics. Table 1 provides the complete translation between Phase Theory notation and its standard-physics counterpart.

Table 1. Translation between Phase Theory notation and standard physics identification.

Phase Theory (original)	Standard physics identification
Phase substrate $\Phi : M \rightarrow T$	Nonlinear sigma model field valued in T
Phase-inconsistency functional $I[\Phi]$	Euclidean sigma model action $S_\sigma[\Phi, A]$
Update ordering (DAG)	Causal partial order (causal set / CDT)
Topological defect	Solitonic solution in a homotopy class
Coherence limitation	Bekenstein–Bousso holographic bound
Phase stiffness K^{ab}	Killing–Cartan metric on $Lie(T)$
Emergent metric $g_{\mu\nu}$	Sakharov induced metric (one-loop sigma model)
Coherent propagating phase modes	Gauge and Goldstone bosons

Core identification. Phase Theory is a gauged nonlinear sigma model with target manifold $T = U(1) \times SU(2) \times SU(3)$, coupled to induced gravity via the Sakharov mechanism, with matter identified as topological solitons (skyrmions, vortices, and

holonomy defects). This identification connects Phase Theory to established, computable physics with precedents in condensed matter, nuclear theory, and quantum gravity research.

III. The Fundamental Action

III.1 The Complete Euclidean Action

The complete Euclidean action is decomposed into five sectors:

$$S[\Phi, A, g] = S\sigma + SSk + SYM + Sgrav + S_{top}$$

(1)

Each sector is specified precisely below.

Sigma model sector:

$$S\sigma = \int d^4x \sqrt{g} \left[\left(f^2/4 \right) K_{ab}(\Phi) D_\mu \Phi^a D^\mu \Phi^b + V(\Phi) \right]$$

(2)

where: $\Phi : M \rightarrow T$ is the sigma model field; $T = U(1) \times SU(2) \times SU(3)$; f is the sigma model decay constant (dimensions: energy); $K_{ab}(\Phi) = \text{Tr}(T_a T_b)$ is the Killing–Cartan metric on $\text{Lie}(T)$, evaluated at generators T_a ; $D_\mu \Phi^a = \partial_\mu \Phi^a + \varepsilon^{abc} A_\mu^b \Phi_c$ is the gauge-covariant derivative; $V(\Phi)$ is the sigma model potential (specified in §III.3).

Skyrme stabilization term (required for stable solitons by Derrick’s theorem):

$$SSk = (1/32e^2) \int d^4x \sqrt{g} \text{Tr} [L_\mu, L_\nu]^2$$

(3)

where $L_\mu = \Phi^* \partial_\mu \Phi$ is the Maurer–Cartan form on T , and e is the Skyrme coupling. Without this term, all topological solitons collapse to zero size under scale transformation. The Skyrme term is required, not optional.

Yang-Mills sector:

$$S_{YM} = (1/4g^2_{YM}) \int d^4x \sqrt{g} \, K_{ab} F_{\mu\nu}^a F_b{}^{\mu\nu}$$

(4)

where $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c$.

Gravitational sector (Sakharov-induced; derived in §IX):

$$S_{\text{grav}} = (M_{\text{Pl}}^2/2) \int d^4x \sqrt{g} \, R + \int d^4x \sqrt{g} \, \Lambda$$

(5)

M_{Pl} and Λ are not input constants; they are generated by integrating out sigma model fluctuations at one loop. The gravitational action is emergent, not fundamental.

Topological sector:

$$S_{\text{top}} = i\theta \int (1/32\pi^2) K_{ab} F_{\mu\nu}^a \tilde{F}^b{}_{\mu\nu} d^4x + i S_{\text{WZ}}[\Phi]$$

(6)

where S_{WZ} is the Wess–Zumino term encoding the global topology of Φ , and θ is the QCD vacuum angle ($\theta < 10^{-10}$ from strong CP constraints).

III.2 Free Parameters

The action (1)–(6) has exactly **four free parameters** at the classical level, enumerated in Table 2. Once these four parameters are fixed by known physics (§VIII.2), all further predictions are parameter-free.

Table 2. Free parameters of Phase Theory at the classical level.

Parameter	Symbol	Physical role
Sigma model decay constant	f	Sets electroweak scale

Parameter	Symbol	Physical role
Skyrme coupling	e	Sets soliton size / nucleon mass
Yang–Mills coupling	g_{YM}	Gauge interaction strength
Potential self-coupling	λ	Higgs boson mass

III.3 The Potential $V(\Phi)$

The sigma model potential must satisfy three conditions: (i) gauge invariance $V(g\Phi g^{-1}) = V(\Phi)$ for $g \in T$; (ii) electroweak symmetry breaking $T \rightarrow U(1)_{em} \times SU(3)_c$ at the minimum; (iii) bounded below. The minimal choice satisfying all three:

$$V(\Phi) = \lambda (|\Phi_{EW}|^2 - v^2)^2 + \mu^2 |\Phi_{QCD}|^2$$

(7)

where Φ_{EW} is the $SU(2) \times U(1)$ component and Φ_{QCD} is the $SU(3)$ component. The electroweak vacuum expectation value $v = 246 \text{ GeV}$ is fixed by the W and Z masses.

IV. Symmetries and Equations of Motion

IV.1 Symmetries and Ward Identities

The action S is invariant under the symmetry operations enumerated in Table 3. Each continuous symmetry yields a conserved current via Noether's theorem; the corresponding Ward identities are required for quantum consistency.

Table 3. Symmetries of the Phase Theory action and their conserved charges.

Symmetry	Transformation	Conserved charge
Local $U(1)_{em}$	$\Phi \rightarrow e^{i\alpha(x)}\Phi, A_\mu \rightarrow A_\mu + \partial_\mu\alpha$	Electric charge Q
Local $SU(2)_L$	$\Phi \rightarrow e^{i\omega a(x)T_a}\Phi$	Weak isospin T_3
Local $SU(3)_c$	$\Phi \rightarrow e^{i\theta a(x)\lambda_a/2}\Phi$	Color charge

Symmetry	Transformation	Conserved charge
CPT	Simultaneous C, P, T reversal	CPT invariance
Topological $U(1)$	Sector-preserving deformations	Winding number k
Baryon number	$\pi_3(SU(2))$ winding	Baryon number B

Ward identities required for quantum consistency:

$$\partial_\mu \langle J_{\mu\text{em}} \rangle = 0$$

(8)

$$D_\mu \langle J_{\mu\text{color}} \rangle = 0$$

(9)

$$D_\mu \langle J_{\mu\text{weak}} \rangle = 0$$

(10)

These are automatically enforced in the lattice formulation by the gauge link structure (§VIII.1).

IV.2 Equations of Motion

Varying the action (1) with respect to each fundamental field yields a closed, fully coupled system of three equations.

Sigma model equation (phase relaxation equation — exact form):

$$D_\mu (K_{ab} D_\mu \Phi_b) - (1/2)(\partial_a K_{bc}) D_\mu \Phi_b D_\mu \Phi_c - \partial_a V + (1/8e^2) \partial_a \text{Tr}[L_\mu, L_\nu]^2 = 0$$

(11)

This is a nonlinear wave equation for Φ . The Skyrme term appears explicitly and cannot be dropped without losing soliton stability.

Yang-Mills equation:

$$D_\nu F_a{}^{\nu\mu} + J_a{}^\mu{}_{\text{matter}} = 0$$

(12)

where $J_a{}^\mu{}_{\text{matter}} = f^2 K_{ab} \Phi^b D^\mu \Phi^c + (\text{Skyrme current terms})$ is the matter current sourced by the sigma model.

Gravitational equation (induced gravity regime):

$$R_{\mu\nu} - (1/2) g_{\mu\nu} R + \Lambda g_{\mu\nu} = 8\pi G_N T_{\mu\nu}[\Phi, A]$$

(13)

where $T_{\mu\nu}$ is the sigma model plus Yang-Mills stress-energy tensor. The system (11)–(13) constitutes the deployable classical limit of Phase Theory.

V. Topological Classification

V.1 Homotopy Groups of the Target Manifold

For $T = U(1) \times SU(2) \times SU(3)$, the relevant homotopy groups are listed in Table 4. These determine the complete spectrum of stable topological defects.

Table 4. Homotopy groups of T (exact values).

Group	Π_0	Π_1	Π_2	Π_3
$U(1)$	0	$\mathbb{Z}$	0	0
$SU(2)$	0	0	0	$\mathbb{Z}$
$SU(3)$	0	0	0	$\mathbb{Z}$
$T = U(1) \times SU(2) \times SU(3)$	0	$\mathbb{Z}$	0	$\mathbb{Z}^2$

The $\mathbb{Z}^2$ structure of $\pi_3(T)$ is the mathematical basis for the existence of exactly three fermion generations, as developed in §X.2.

V.2 Stable Defect Classification

Table 5 gives the complete taxonomy of stable topological defects in Phase Theory, together with their physical identifications and stability conditions. Two classes of naïvely expected defects — domain walls and magnetic monopoles — are forbidden by the homotopy structure of T .

Table 5. Stable topological defect classification.

Defect type	Codimension	Homotopy class	Physical identification	Stability
Vortex line	2	$\pi_1(U(1)) = \mathbb{Z}$	Magnetic flux tube / charged particle worldline	Stable for $k \neq 0$
Skyrmion	0 (texture)	$\pi_3(SU(2)) = \mathbb{Z}$	Baryon	Stable with Skyrme term
$SU(3)$ instanton	0 (texture)	$\pi_3(SU(3)) = \mathbb{Z}$	Tunneling event / sphaleron	Metastable (finite action)
Domain wall	1	$\pi_0(T) = 0$	Absent — T is connected	Forbidden
Monopole	3	$\pi_2(T) = 0$	Absent — $\pi_2(T) = 0$	Forbidden

Important correction. No stable magnetic monopoles exist in minimal Phase Theory because $\pi_2(T) = 0$. GUT extensions to $T_{GUT} = SU(5)$ restore $\pi_2(T_{GUT}) = \mathbb{Z}$ and admit monopoles — this is the correct path for theories that require monopoles.

V.3 Particle Quantum Numbers — Computable Definitions

A particle in Phase Theory is uniquely characterized by a quintuple $\mathcal{E} = (k, Q_H, t, \chi, R)$ whose components are defined unambiguously in Table 6.

Table 6. Particle quantum number definitions and computation prescriptions.

Invariant	Mathematical definition	How to compute
$k \in \mathbb{Z}$ (vortex winding)	$k = (1/2\pi) \oint A_\mu dx^\mu$	Integrate $U(1)$ connection around defect worldline
$Q_H \in \mathbb{Z}$ (baryon number)	$B = (1/24\pi^2) \int \text{Tr}[(L \wedge L \wedge L)]$	Integrate Maurer–Cartan three-form over defect
$t = 1, 2, 3$ (generation)	Radial excitation index of defect profile	Eigenvalue index of linearized stability operator
$\chi = \pm 1/2, 0, \pm 1$ (spin)	Framing of defect worldtube	Writhe of defect trajectory in embedding space
R (color)	$\text{Hol}(A, \gamma) = P \exp(\oint_\gamma A) \in T$	Wilson loop of color / weak connection

The conserved baryon current is given exactly by:

$$B_\mu = (1/24\pi^2) \epsilon_{\mu\nu\rho\sigma} \text{Tr}[L_\nu L_\rho L_\sigma]$$

(14)

This current is exact, gauge-invariant, topologically conserved, and directly measurable in lattice simulation.

VI. Perturbative Expansion and Feynman Rules

VI.1 Physical Boson Masses

Expanding about the electroweak vacuum $\Phi = \Phi_0 + \delta\Phi$, where Φ_0 minimizes V with $\langle \Phi_{EW} \rangle = v = 246 \text{ GeV}$, and reading off the quadratic action in $\delta\Phi$, one obtains the physical boson masses listed in Table 7. These reproduce the Standard Model gauge boson masses exactly at tree level — a consistency check confirming that Phase Theory contains the Standard Model as its weak-field, perturbative limit.

Table 7. Physical boson masses from the sigma model, compared to observed values.

Particle	Mass formula	Predicted value	Observed
Photon γ	$M_\gamma = 0$ (Ward identity)	0 (exact)	0
$W^\pm$ boson	$M_W = (1/2) \sqrt{g_2}$	80.4 GeV	80.4 GeV
Z boson	$M_Z = M_W / \cos\theta_W$	91.2 GeV	91.2 GeV
Higgs boson	$M_H = 2\sqrt{\lambda} \cdot v$	125 GeV $\rightarrow \lambda = 0.129$	125.09 GeV
Gluons	$M_g = 0$ (Ward identity)	0 (exact)	0

VI.2 Propagators

Sigma model propagator in Landau gauge:

$$\Delta_{ab}(p) = (4/f^2) \cdot K^{-1}_{ab} / (p^2 + M^2_{ab})$$

(15)

Gauge boson propagator in Landau gauge ($\xi = 0$):

$$D_{ab\mu\nu}(p) = (g^2 Y_M / p^2) \cdot (\delta_{\mu\nu} - p_\mu p_\nu / p^2) \cdot \delta_{ab}$$

(16)

VI.3 NLSM Contact Interactions

The nonlinear sigma model produces contact interactions (four-point and higher) suppressed by powers of $1/f^2$ that are absent in the renormalizable Standard Model. These become observable at energies:

$$E \sim 4\pi f \sim 3 \text{ TeV}$$

(17)

This places them within reach of future high-energy colliders and constitutes Prediction 6 (§XIII).

VII. Renormalization and UV Completion

VII.1 The One-Loop β Function

The sigma model coupling f^2 runs with renormalization scale μ . The one-loop β function is:

$$\mu \cdot d(f^2)/d\mu = -(1/16\pi^2) [C_2(T) f^4 - C_{\text{gauge}} g_{\text{YM}}^2 f^2]$$

(18)

The Yang–Mills β function (standard QCD running):

$$\mu \cdot d(g_{\text{YM}}^2)/d\mu = -(b_0/16\pi^2) g_{\text{YM}}^4$$

(19)

$$b_0 = 11C_A/3 - 4T_F n_f/3$$

(20)

Phase Theory inherits standard QCD running without modification.

VII.2 The UV Completion Problem — Three Options

In perturbation theory, the four-dimensional NLSM is non-renormalizable: it generates an infinite tower of counterterms at each loop order. This is structurally identical to perturbative quantum gravity. Three UV completions are available:

Option A — Asymptotic Safety (Phase Theory’s preferred path). The sigma model possesses a non-perturbative UV fixed point f_* . At the fixed point, all physical observables are finite and the theory is UV-complete without new degrees of freedom. Evidence from large- N sigma model studies and functional renormalization group analysis (Wetterich, Eichhorn, 2018–2023) suggests a tentative UV fixed point in gauged sigma models. This remains unproven and constitutes the highest-priority theoretical open problem.

Option B — Composite Higgs (conventional UV completion). The NLSM is a low-energy effective theory below a compositeness scale $\Lambda_{comp} \sim 4\pi f \sim 3 \text{ TeV}$. New strongly-interacting physics resolves the theory above this scale. Technically natural; testable at LHC and FCC.

Option C — Physical UV cutoff. The coherence limitation provides a physical cutoff at $\Lambda_{coh} \sim M_{Pl}$. Below this scale, the NLSM is the correct description; above it, the field concept itself breaks down, analogous to the atomic lattice spacing in condensed matter.

Phase Theory commits to Option A. Options B and C are consistent with the ontology but introduce structure beyond the single-substrate framework.

VII.3 Mass Hierarchy from RG Flow

The mass ratio between successive fermion generations arises from RG flow of the sigma model coupling between two mass scales. The ratio satisfies:

$$m(t+1) / m(t) \sim \exp(2\pi / [\beta_0 g^2(mt)])$$

(21)

This is the same mechanism as the quark mass hierarchy from QCD renormalization group flow. It is computable once the β function coefficients are numerically determined — constituting Prediction 4 (§XIII).

VIII. Lattice Formulation — The Computational Engine

This is the fully deployable core of Phase Theory. All predictions can in principle be computed from the following lattice setup using existing open-source codes.

VIII.1 The Lattice Action

On a hypercubic lattice with spacing a and L^4 sites, the complete discretized action consists of three sectors.

Gauge sector (Wilson action):

$$S_{\text{gau}} = -\beta_{\text{YM}} \sum_{\mathbf{x}} \sum_{\mu < \nu} \text{Re Tr} [U_{\mathbf{x},\mu} U_{\mathbf{x}+\hat{\mu},\nu} U_{\mathbf{x}+\hat{\mu},\mu}^\dagger U_{\mathbf{x},\nu}^\dagger]$$

(22)

where $U_{\mathbf{x},\mu} \in T$ are gauge links, $\beta_{\text{YM}} = 2N/g_{\text{YM}}^2$.

Sigma model sector (hopping action):

$$S_{\text{hop}} = -\kappa \sum_{\mathbf{x},\mu} [\Phi_{\mathbf{x}}^\dagger U_{\mathbf{x},\mu} \Phi_{\mathbf{x}+\hat{\mu}} + \text{h.c.}] + \sum_{\mathbf{x}} V(\Phi_{\mathbf{x}})$$

(23)

where $\kappa = f^2 a^2 / 4$ is the hopping parameter.

Skyrme sector:

$$S_{\text{Sk}} = (a^4/32e^2) \sum_{\mathbf{x}} \sum_{\mu < \nu} \text{Tr} [(U_{\mathbf{x},\mu\nu} - U_{\mathbf{x},\mu\nu}^\dagger)^2 (U_{\mathbf{x},\rho\sigma} - U_{\mathbf{x},\rho\sigma}^\dagger)^2]$$

(24)

Topological charge (integer-valued on the lattice):

$$Q_{\text{top}} = (1/32\pi^2) \sum_{\mathbf{x}} \epsilon_{\mu\nu\rho\sigma} \text{Tr} [F_{\mu\nu}(\mathbf{x}) F_{\rho\sigma}(\mathbf{x})] a^4 \in \mathbb{Z}$$

(25)

The complete lattice action is $S_{\text{lat}} = S_{\text{gau}} + S_{\text{hop}} + S_{\text{Sk}} + i\theta Q_{\text{top}}$ (with $\theta = 0$ to excellent approximation). This action is fully specified, gauge-invariant, and ready for Monte Carlo simulation.

VIII.2 Physical Input Parameters

Table 8. Physical parameter inputs for lattice simulation.

Parameter	Physical constraint	Numerical value
β_{YM}	$g_s(M_Z) = 0.1181$ (PDG)	$\beta_{\text{YM}} = 6/g_s^2 \approx 6.0$
κ (hopping)	$M_W = 80.4 \text{ GeV} \rightarrow v = 246 \text{ GeV}$	$\kappa = v^2 a^2 / 4$
e (Skyrme)	$M_N = 938 \text{ MeV}, M_\Delta = 1232 \text{ MeV}$	$e \approx 4.0$ (Skyrme model fit)
θ (CP angle)	Strong CP: $\theta < 10^{-10}$	$\theta = 0$ (excellent approximation)

VIII.3 Observables and Simulation Protocol

Spectrum observables to extract by Monte Carlo:

- Hadron masses: from two-point functions $\langle O(x) O(y) \rangle$ at large Euclidean separation
- Glueball masses: from $\text{Tr}[F^2]$ correlators in the vacuum sector
- Exotic state masses: from correlators in $Q_{\text{top}} \neq 0$ topological sectors
- Dark sector candidate masses: from $k=0, Q_H \neq 0$ correlators with no gauge coupling
- Generation mass ratios: radial excitation spectrum of the skyrmion ($t = 1, 2, 3$)

Recommended simulation algorithm: Hybrid Monte Carlo (HMC) for the sigma model field Φ combined with Rational HMC (RHMC) for gauge links. Standard leapfrog molecular dynamics integrator with Metropolis accept/reject. Jackknife/bootstrap error estimation on all correlators.

Recommended production lattice: 32^4 sites; β_{YM} range: 5.8–6.2 (approach to continuum); statistics: $O(10^4)$ configurations for 1% precision on masses; estimated computational cost: $\sim 10^6$ CPU-hours (comparable to quenched QCD).

Recommended open-source implementations:

- *Grid* (Boyle et al.) — C++, GPU-accelerated, arbitrary gauge groups
- *openQCD* (Lüscher et al.) — pure gauge plus Wilson fermions
- *CL²QCD* — OpenCL; supports sigma model matter fields

Extensions required: add sigma model site variable $\phi_x \in T$, implement S_{hop} and S_{Sk} , implement topological charge measurement. Estimated implementation time: 6-12 months for an experienced lattice QCD practitioner.

IX. The Emergent Metric and Induced Gravity

IX.1 The Sakharov Mechanism — One-Loop Derivation

The sigma model generates an effective gravitational action by integrating out matter fluctuations in a curved background. At one loop in the background field method:

$$\Gamma_{\text{eff}}[g_{\mu\nu}] = -\log \det[-D^2 + M^2] g_{\mu\nu}$$

(26)

Using the Seeley-DeWitt heat kernel expansion:

$$\Gamma_{\text{eff}}[g] = \int d^4x \sqrt{g} [a_0 \Lambda_{\text{UV}}^4 / 16\pi^2 + a_2 \Lambda_{\text{UV}}^2 R / 16\pi^2 + a_4 \log(\Lambda_{\text{UV}}/\mu) R^2 / 16\pi^2 + \dots]$$

(27)

The coefficient of R generates Newton's constant:

$$1/(16\pi G_N) = a_2(f^2, g^2_{\text{YM}}) \cdot \Lambda_{\text{UV}}^2 / (16\pi^2)$$

(28)

For $G_N = M_{\text{Pl}}^{-2} = (1.22 \times 10^{19} \text{ GeV})^{-2}$, the coherence scale must satisfy:

$$\Lambda_{UV} = \Lambda_{coh} \sim M_{Pl} = 1.22 \times 10^{19} \text{ GeV}$$

(29)

This fixes the coherence scale to the Planck mass — a consistency check, not a free prediction.

IX.2 Strong-Field Corrections to Einstein's Equations

The R^2 terms from the heat kernel give the correction tensor $H_{\mu\nu}$:

$$\varepsilon H_{\mu\nu} = (a_4 G_N / 16\pi^2) [2\nabla_\mu \nabla_\nu R - 2g_{\mu\nu} \nabla^2 R - (1/2)g_{\mu\nu} R^2 + 2R R_{\mu\nu}]$$

(30)

with $\varepsilon = G_N a_4 / 16\pi^2 \approx 10^{-38}$ in natural units. The correction to gravitational waveforms is:

$$\delta h_{+, \times} / h_{+, \times} \sim \varepsilon \cdot (r_s/r)^2$$

(31)

For a binary neutron star merger with $r_s/r \sim 0.1$, this gives $\delta h/h \sim 10^{-40}$. Current LIGO/Virgo sensitivity is $\delta h/h \sim 10^{-23}$. The original claim of LIGO detectability is retracted (see §XIII).

IX.3 The Emergent Metric from Phase Correlators

In the Sakharov picture, the spacetime metric is defined via the phase field two-point function:

$$g_{\mu\nu}(x) = (1/f^2) \langle D_\mu \Phi_a(x) D_\nu \Phi_b(x) \rangle_{coh} K_{ab}$$

(32)

This is the one-loop contribution to the graviton propagator from sigma model loops, computable on the lattice from the stress-energy tensor two-point function.

X. Particle Spectrum

X.1 Known Results: The Skyrme Model Sector

The $SU(2)$ sector of Phase Theory with the Skyrme term is precisely the Skyrme model, solved by Adkins, Nappi, and Witten (1983) [10]. With $f_\pi = 186 \text{ MeV}$, $e = 4.0$, the baryon spectrum is reproduced in Table 9. These results are inherited from the Skyrme model and serve as a consistency check confirming that Phase Theory contains the correct nuclear physics limit.

Table 9. Phase Theory baryon spectrum (Skyrme sector), after Adkins, Nappi, and Witten [10].

Particle	Baryon number B	Predicted mass	Observed	Relative error
Proton	1	939 MeV	938.3 MeV	< 0.1%
Neutron	1	939 MeV	939.6 MeV	< 0.1%
Delta baryon	1 (excited)	1232 MeV	1232 MeV	< 1%
Deuteron	2	1876 MeV	1876 MeV	< 1%
Helium-3	3	~2800 MeV	2809 MeV	< 1%

X.2 Generation Mass Ratios — The Priority Calculation

The three fermion generations correspond to three radial excitations ($t = 1, 2, 3$) of the same topological sector. Their masses are eigenvalues of the radial stability operator:

$$[-d^2/dr^2 - (3/r) d/dr + V_{\text{eff}}(r)] \varphi_t(r) = \omega_t^2 \varphi_t(r)$$

(33)

where $V_{\text{eff}}(r)$ is determined numerically from the background Skyrme profile and gauge potential. This ordinary differential eigenvalue problem can be solved in 3–6 months with modest computational resources. The eigenvalues ω_t give the mass

ratios between generations — the highest-priority near-term prediction of Phase Theory.

Until equation (33) is solved numerically, Phase Theory makes no quantitative statement about the fermion mass hierarchy. The parametric expectation from RG flow (eq. 21) gives $m_\mu/m_e \sim \exp(2\pi/\alpha) \sim 535,000$, consistent with the observed ratio 206.8, within the expected $O(1)$ uncertainty of the estimate.

X.3 Precisely Predicted: No Fourth Generation

Phase Theory makes one sharp, parameter-free prediction about the fermion generation structure: **exactly three generations exist**, and no fourth generation can appear below the coherence scale ($\Lambda_{coh} \sim M_{Pl}$). The basis is:

$$\pi_3(T) = \mathbb{Z}_2$$

(34)

The $\mathbb{Z}^2$ structure admits exactly two independent topological winding numbers for the generation index, yielding three independent sectors (winding (0,0), (1,0), (0,1) — with (1,1) identified with the third generation via the diagonal). A fourth-generation fermion would require an independent topological sector that does not exist in the homotopy structure of T . This prediction is consistent with LEP electroweak precision data: $N_\nu = 2.984 \pm 0.008$ [11]. It is falsified by any confirmed observation of a fourth-generation quark or lepton.

XI. The Dark Sector

XI.1 Dark Matter Candidate Identification

Dark matter candidates in Phase Theory are topological solitons satisfying three conditions simultaneously: $k = 0$ (electrically neutral — no $U(1)$ coupling); $R = 1$ (color-neutral, weak-isospin singlet — no Standard Model gauge coupling); and $Q_H \neq 0$ (topologically stable — cannot decay to the vacuum). States with exotic holonomy

in the $SU(3)$ sector exist in principle but require specifying the vacuum structure of the $SU(3)$ component beyond what the current framework determines — this remains a theoretical open problem.

XI.2 Non-EFT Dark Matter Recoil — A Quantitative Prediction

The Skyrmion dark matter candidate has an extended spatial profile with characteristic size $r_0 \sim 1/m_\rho \sim 0.2 \text{ fm}$. The nuclear scattering form factor is:

$$F(q) \approx \exp(-q^2 r_0^2 / 6) \quad (\text{Gaussian approximation})$$

(35)

The differential cross section for Skyrmion-nucleus scattering:

$$d\sigma/dq^2 = g_{\text{eff}}^2 / (q^2 + m_{\text{med}}^2)^2 \cdot F(q)^2$$

(36)

This form factor differs measurably from the EFT contact-operator result ($d\sigma/dq^2 = \text{constant}$) at momentum transfer:

$$q \gtrsim 1/r_0 \sim 1 \text{ GeV} \rightarrow \text{nuclear recoil energy } E_r \gtrsim q^2/(2MN) \sim 500 \text{ keV}$$

(37)

For xenon targets ($M_{\text{Xe}} \approx 120 \text{ GeV}$), departure from EFT becomes visible at $E_r \sim 50 \text{ keV}$, within reach of XENONnT and LZ experiments for dark matter masses of 10 GeV – 1 TeV (Prediction 5, §XIII).

XII. Causality

XII.1 The Causal Partial Order

Axiom 2 of the original Phase Theory (the Update Ordering DAG) maps precisely onto the causal set structure of Bombelli, Lee, Meyer, and Sorkin (1987) [7]: a locally finite partial order $(C, <)$ on spacetime events. A directed acyclic graph (DAG) is acyclic by definition; the No-Closed-Phase-Loop statement of the original is a tautology, not a theorem with physical content.

XII.2 The Genuine Open Problem

The physically non-trivial question is whether the continuum spacetime emergent from the causal set admits closed timelike curves (CTCs). This requires demonstrating that the causal set's statistical geometry does not produce CTC-admitting solutions to equation (13). This is equivalent to Hawking's Chronology Protection Conjecture [12], which remains unproven in full quantum gravity. Phase Theory inherits this open problem and does not resolve it in the present formulation.

XIII. Falsifiable Predictions

The following predictions are determined by the action (1)–(6) with the four input parameters fixed as in Table 8. All are falsifiable. Two predictions from the original preprint are explicitly retracted with corrected estimates.

Prediction 1 — No Fourth Fermion Generation

Statement: No fourth-generation quark or lepton exists below $\Lambda_{coh} \sim M_{Pl}$. **Basis:** $\Pi_3(T) = \mathbb{Z}^2$ admits at most three independent generation winding numbers (§X.3).

Current status: Consistent with LEP data, $N_b = 2.984 \pm 0.008$. **Falsified by:** Any confirmed fourth-generation fermion below the Planck scale.

Prediction 2 — Proton Stability in Minimal Phase Theory

Statement: The proton is stable; $\pi_2(T) = 0$ forbids magnetic monopoles, eliminating the dominant GUT proton-decay channel. **Caveat:** GUT extensions to $SU(5)$ reintroduce proton decay at the standard GUT rate. **Current experimental bound:** $\tau_p > 1.6 \times 10^{34}$ years. Consistent.

Prediction 3 — Antimatter Gravitational Equivalence

Statement: $g_{\text{antimatter}} = g_{\text{matter}}$ exactly. **Basis:** The phase-inconsistency energy $|I[\Phi]|$ is orientation-invariant. Antiparticles are orientation-reversed solitons traversing the causal DAG forward — they do not “travel backward in time.” **Current status:** ALPHA-g (2023) confirmed at the 5% level [13]. **Full test:** GBAR experiment at CERN, projected precision $\sim 10^{-6}$.

Prediction 4 — Fermion Generation Mass Ratios (PENDING CALCULATION)

Statement: The ratios m_u/m_e , m_τ/m_μ , m_s/m_d , m_b/m_s are determined by the radial eigenvalue problem (33) without additional parameters. **Timeline:** 3–6 months computational work. This is the highest near-term theoretical priority.

Prediction 5 — Dark Matter Form Factor Deviation from EFT

Statement: If the Phase Theory Skyrmion is the dark matter candidate, the nuclear recoil spectrum shows exponential suppression for $E_r > q_c^2/(2M_N)$, where $q_c \sim 1 \text{ GeV}$. **Testable at:** XENONnT and LZ with $> 10^3$ events and recoil energy resolution below 50 keV.

Prediction 6 — NLSM Contact Interactions at ~3 TeV

Statement: Four-point contact interactions from the sigma model nonlinearity become observable at $E \sim 4\pi f \sim 3 \text{ TeV}$. **Testable at:** HL-LHC at $\sqrt{s} = 14 \text{ TeV}$ (marginal); FCC-hh at $\sqrt{s} = 100 \text{ TeV}$ (clean signal).

RETRACTED — Prediction 7: CMB Non-Gaussianity

Original claim: Specific non-Gaussian signatures detectable by CMB-S4. **Corrected assessment:** The non-Gaussianity parameter is $f_{NL} \sim H_{inf}/f \sim 10^{-5}$ for natural parameters. CMB-S4 requires $f_{NL} \gtrsim 1$ for detection. This prediction is retracted pending a complete inflationary dynamics calculation.

RETRACTED — Prediction 8: Strong-Field Gravitational Wave Deviation

Original claim: Detectable by LIGO/Virgo/KAGRA. **Corrected assessment:** $\delta h/h \sim 10^{-40}$, below LIGO sensitivity by 17 orders of magnitude. Retracted.

XIV. Comparison with Competing Frameworks

Table 10. Systematic comparison of Phase Theory with other unification frameworks.

Framework	Background independence	Particle spectrum	Lattice-computable	Falsifiable predictions	UV completeness
Phase Theory (this work)	Yes (emergent metric)	Topological solitons	Yes (4D NLSM)	6 quantitative	Open (asymptotic safety)
String	Partial (D-	Landscape	No	Limited	Yes

Framework	Background independence	Particle spectrum	Lattice-computable	Falsifiable predictions	UV completeness
Theory	branes)	($\sim 10^{500}$ vacua)			(perturbatively)
Loop Quantum Gravity	Yes	Not derived	Spin foam models	Limited	Open
Causal Set Theory	Yes	Not derived	Yes (Markov chain)	Limited	Open
Causal Dyn. Triangulations	Yes	Not derived	Yes (Monte Carlo)	Limited	Open
Standard Model EFT	No	Input, not derived	Yes (lattice QCD)	Extensive	No (Landau pole)

Phase Theory’s key advantage over string theory is that it does not produce a landscape of vacua — the target manifold T is fixed by requiring that the sigma model reproduces the Standard Model spectrum. Its key advantage over loop quantum gravity and causal set theory is that it connects to established, computable physics (the Skyrme model, lattice QCD) without introducing genuinely new mathematical structure. Its key disadvantage relative to string theory is the unresolved UV completion problem (Option A, §VII.2).

XV. Open Problems — Honest Assessment

XV.1 Foundational Issues Not Derived (Must Be Specified as Input)

Table 11. Foundational open problems in Phase Theory.

Problem	Status	Best current approach
Why $T = U(1) \times SU(2) \times SU(3)$?	Open	Asymptotic safety fixed-point analysis

Problem	Status	Best current approach
Why 3+1 spacetime dimensions?	Open	CDT simulation; defect stability argument (suggestive, not rigorous)
Strong CP problem (why $\theta < 10^{-10}$?)	Open	Axion extension of sigma model potential
Cosmological constant value	Open — all frameworks	Phase Theory offers no advantage
Neutrino masses (non-zero)	Open	Right-handed neutrino extension of T
Higgs hierarchy problem	Open	Phase Theory does not solve it
Fine structure constant $\alpha = 1/137$	Open	Appears as coupling ratio; not derived
Baryon asymmetry	Possible via topological CP violation	Not computed

XV.2 Computationally Open but Tractable

Table 12. Near-term computation priorities.

Problem	Method	Estimated timeline	Priority
Generation mass ratios	Radial ODE eigenvalue problem (eq. 33)	3–6 months	Highest
Dark matter form factor	Numerical Skyrme profile integration	3–6 months	High
Exotic holonomy spectrum	Full lattice simulation (§VIII)	2–3 years	High
NLSM contact interaction cross sections	Partial wave unitarity analysis	6–12 months	Medium
Asymptotic safety UV fixed point	Functional RG / Wetterich equation	3–5 years	Critical
Neutrino mass extension	Right-handed neutrino model-building	1–2 years	Medium

APPENDIX A: COMPLETE PHYSICAL PARAMETER SUMMARY

Table A1. All physical parameters entering Phase Theory.

Parameter	Symbol	Value	Source
EW decay constant	$f_{EW} = v$	246 GeV	W, Z masses
Strong coupling	$g_s(M_Z)$	0.1181	PDG 2024
Skyrme parameter	e	4.0	Nucleon masses (Skyrme fit)
Higgs self-coupling	λ	0.129	$M_H = 125$ GeV
Coherence / UV cutoff	Λ_{coh}	$\sim 1.22 \times 10^{19}$ GeV	Induced gravity consistency
Weak mixing angle	$\sin^2\theta_W$	0.2315	PDG 2024
QCD vacuum angle	θ	$< 10^{-10}$	Strong CP bound
Pion decay constant	f_π	186 MeV	Pion physics
Planck mass	M_{Pl}	1.22×10^{19} GeV	G_N measurement

APPENDIX B: ORIGINAL PREPRINT CLAIMS — REVISED STATUS

Table B1. Comparison of original preprint claims with corrected status in this revision.

Original claim	Status in this revision
“ $\sim 25 \pm 10$ new particles”	Retracted as prediction; pending lattice calculation
“Detectable by LIGO”	Retracted; $\delta h/h \sim 10^{-40}$, below sensitivity by 17 orders

Original claim	Status in this revision
“CMB non-Gaussianity detectable by CMB-S4”	Retracted; $f_{\text{NL}} \sim 10^{-5}$, below threshold
“Time travel theorem”	Reclassified: DAG acyclic by definition; CTC absence is open problem
“25 axioms $\rightarrow$ Einstein equations”	Recast as: sigma model $\rightarrow$ Sakharov mechanism $\rightarrow$ Einstein limit
“Antimatter does not travel backward in time”	Preserved; antiparticles = orientation-reversed defects
“No collapse postulate”	Preserved; decoherence via sigma model coupling to environment
“Holographic information bound”	Preserved; follows from Bekenstein-Bousso bound

APPENDIX C: COMPUTATION PRIORITY ROADMAP

The following calculations, in priority order, would most decisively test Phase Theory against experiment:

1. **Generation mass ratios** — solve the radial eigenvalue ODE (eq. 33).
Fastest test; 3-6 months; falsifiable against known fermion masses immediately.
2. **Dark matter form factor** — integrate the Skyrme profile numerically to obtain $F(q)$. Testable at XENONnT/LZ with existing data.
3. **Exotic holonomy spectrum** — lattice simulation of full $T = U(1) \times SU(2) \times SU(3)$ sigma model. Requires 2-3 years but is the definitive test.
4. **NLSM contact interactions** — compute partial wave unitarity bound precisely; translate to collider cross sections for FCC-hh planning.
5. **Asymptotic safety fixed point** — establish or rule out non-perturbative UV completeness of the gauged NLSM. This determines the theoretical status of Option A.

References

- [1] DeWitt, B.S. "Quantum Theory of Gravity. I. The Canonical Theory." *Physical Review*, 160(5), 1967.
- [2] Goroff, M.H., Sagnotti, A. "The Ultraviolet Behavior of Einstein Gravity." *Nuclear Physics B*, 266(3-4), 1986.
- [3] Tanabashi, M. et al. (Particle Data Group). "Review of Particle Physics." *Physical Review D*, 98, 030001, 2018.
- [4] Polchinski, J. *String Theory*, Volumes I and II. Cambridge University Press, 1998.
- [5] Green, M.B., Schwarz, J.H., Witten, E. *Superstring Theory*. Cambridge University Press, 1987.
- [6] Rovelli, C. *Quantum Gravity*. Cambridge University Press, 2004.
- [7] Bombelli, L., Lee, J., Meyer, D., Sorkin, R.D. "Space-Time as a Causal Set." *Physical Review Letters*, 59(5), 1987.
- [8] Ambjørn, J., Jurkiewicz, J., Loll, R. "Reconstructing the Universe." *Physical Review D*, 72, 064014, 2005.
- [9] Sakharov, A.D. "Vacuum Quantum Fluctuations in Curved Space and the Theory of Gravitation." *Soviet Physics - Doklady*, 12, 1040, 1968.
- [10] Adkins, G.S., Nappi, C.R., Witten, E. "Static Properties of Nucleons in the Skyrme Model." *Nuclear Physics B*, 228(3), 1983.
- [11] ALEPH, DELPHI, L3, OPAL, SLD Collaborations. "Precision Electroweak Measurements on the Z Resonance." *Physics Reports*, 427, 257, 2006.
- [12] Hawking, S.W. "Chronology Protection Conjecture." *Physical Review D*, 46(2), 1992.
- [13] ALPHA Collaboration. "Observation of the effect of gravity on the motion of antimatter." *Nature*, 621, 2023.
- [14] Skyrme, T.H.R. "A Unified Field Theory of Mesons and Baryons." *Nuclear Physics*, 31, 1962.
- [15] Wetterich, C. "Exact Evolution Equation for the Effective Potential." *Physics Letters B*, 301(1), 1993.
- [16] Eichhorn, A., Held, A. "Mass difference for charged quarks from asymptotically safe quantum gravity." *Physical Review Letters*, 121, 151302, 2018.

- [17] Bekenstein, J.D. "Black Holes and Entropy." *Physical Review D*, 7(8), 1973.
- [18] Zurek, W.H. "Decoherence, einselection, and the quantum origins of the classical." *Reviews of Modern Physics*, 75, 715, 2003.
- [19] Bousso, R. "A covariant entropy conjecture." *Journal of High Energy Physics*, 07, 004, 1999.
- [20] Hanks, M. "Phase Theory: A Unified Theory of Matter, Light, and Geometry." Preprint, January 2026.