

Chapter 1

Derivation of the used inverse covariance matrix from the Aleph data

While performing a **Generalized least squares** (GLS) we estimate our regression coefficients $\hat{\beta}$ as follows:

$$\hat{\beta} = \underset{\mathbf{b}}{\operatorname{argmin}} (\mathbf{y} - \mathbf{X}\mathbf{b})^T \mathbf{\Omega}^{-1} (\mathbf{y} - \mathbf{X}\mathbf{b}), \quad (1.1)$$

with \mathbf{b} being an candidate estimate of β , \mathbf{X} being the design matrix, \mathbf{y} being the response values and $\mathbf{\Omega}^{-1}$ being the **inverse covariance matrix**.

The Aleph data includes the **standard error** (SE), which are equal to the **standard deviation** as per definition. Furthermore Aleph provides the **correlation coefficients** of the errors. We will use these two quantities in combination with **Gaussian error propagation** to derive an approximation of the covariance matrix.

1.1 Propagation of experimental errors and correlation

Let $\{f_k(x_1, x_2, \dots, x_n)\}$ be a set of m functions, which are linear combinations of n variables x_1, x_2, \dots, x_n with combination coefficients $A_{k1}, A_{k2}, \dots, A_{kn}$, where $k \in \{1, 2, \dots, m\}$. Let the covariance matrix of x_n be denoted by

$$\Sigma^x = \begin{pmatrix} \sigma_1^2 & \sigma_{12} & \sigma_{13} & \cdots \\ \sigma_{12} & \sigma_2^2 & \sigma_{23} & \cdots \\ \sigma_{13} & \sigma_{23} & \sigma_3^2 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \quad (1.2)$$

Then the covariance matrix of the functions Σ^f is given by

$$\Sigma_{ij}^f = \sum_k^n \sum_l^n A_{ik} \sum_{kl}^x A_{jl}, \quad \Sigma^f = A \Sigma^x A^T. \quad (1.3)$$

In our case we are dealing with non-linear functions, which we will linearized with the help of the **Taylor expansion**

$$f_k \approx f_k^0 + \sum_i^n \frac{\partial f_k}{\partial x_i} x_i, \quad f \approx f^0 + Jx. \quad (1.4)$$

Therefore, the propagation of error follows from the linear case, replacing the Jacobian matrix with the combination coefficients ($J = A$)

Chapter 2

Coefficients

2.1 β function

There are several conventions for defining the β coefficients, depending on a minus sign and/or a factor of two (if one substitutes $\mu \rightarrow \mu^2$) in the β -function ?? . We follow the convention from Pascual and Tarrach (except for the minus sign) and have taken the values from ??

$$\beta_1 = \frac{1}{6}(11N_c - 2N_f) \quad (2.1)$$

$$\beta_2 = \frac{1}{12}(17N_c^2 - 5N_cN_f - 3C_fN_f) \quad (2.2)$$

$$\beta_3 = \frac{1}{32} \left(\frac{2857}{54}N_c^3 - \frac{1415}{54}N_c^2N_f + \frac{79}{54}N_cN_f^2 - \frac{205}{18}N_cC_fN_f + \frac{11}{9}C_fN_f^2 + C_f^2N_f \right) \quad (2.3)$$

$$\beta_4 = \frac{140599}{2304} + \frac{445}{16}\zeta_3, \quad (2.4)$$

where we used $N_f = 6$ and $N_c = 3$ for β_4 .

2.2 Adler function