

Chapter 1

Introduction

In particle physics we are concerned about small objects and their interactions. Their dynamics are currently best described by the Standard Model (SM).

The SM contains two groups of fermionic, Spin 1/2 particles. The former group, the Leptons consist of: the electron (e), the muon (μ), the tau (τ) and their corresponding neutrinos ν_e , ν_μ and ν_τ . The latter group, the Quarks contain: u , d (up and down, the so called light quarks), s (strange), c (charm), b (beauty or beauty) and t (top or truth). The SM furthermore differentiates between three fundamental forces (and its carriers): the electromagnetic (γ photon), weak (Z- or W-Boson) and strong (g gluon) interactions. The before mentioned Leptons solely interact through the electromagnetic and the weak force (also referred to as electroweak interaction), whereas the quarks additionally interact through the strong force.

The strong force is denominated Quantumchromodynamics (QCD). As the name suggest¹ the force is characterized by the color charge. Every quark has next to its type one of the three colors blue, red or green. The color force is mediated through eight gluons, which each being bi-colored², interact with quarks and each other. The strength of the strong force is given by the coupling constant α_s . The coupling constants are a function of energy E and $\alpha_s(E)$ increases with energy³. This is exclusive for QCD and leads to *asymptotic freedom* and *confinement*. The former phenomenon describes the decreasing strong force between quarks and gluons, which become asymptotically free at large energies. The latter expresses the fact, that no isolated quark has been found until today. Quarks appear confined as *Hadrons*, the so called *Mesons*⁴ and *Baryons*⁵. As we measure *Hadrons* in our experiments but calculate with quarks within our theoretical QCD model we have to assume *Quark-Hadron Duality*, which states that QCD is still valid for Hadrons for energies suffi-

¹Chromo is the greek word for color.

²Each gluon carries a color and an anti-color.

³In contrast to the electromagnetic force, where $\alpha(E)$ decreases!

⁴Composite of a quark and an anti-quark.

⁵Composite of three quarks or three anti-quarks.

cently high energies. There exist *Duality Violations* (DV), which will be investigated within this work.

In the following (section 1.1) we will describe the τ -decays, which play an essential role in our QCD analysis. Then (section 1.2) we want to explain some more details of QCD, especially about the coupling constant $\alpha_s(s)$ (which is not constant at all) and the *QCD sum rules*.

1.1 τ -Decays

The τ -particle is an elementary particle with negative electric charge and a spin of $1/2$. Together with the lighter electron and muon it forms the *charged Leptons*⁶. Even though it is an elementary particle it decays via the *weak interaction* with a lifetime of $\tau_\tau = 2.9 \times 10^{-13} \text{ s}$ and a mass of $1776.86(12) \text{ MeV}$ [22]. It is the only lepton massive enough to decay into Hadrons. The final states of a decay are limited by *conservation laws*. In case of a τ -decay they must conserve the electric charge (-1) and *invariant mass* of the system. Thus, as we can see from the corresponding Feynman diagram (see section 1.1)⁷ the τ decays by the emission of a W boson and a tau-neutrino ν_τ into different pairs of $(e^-, \bar{\nu}_e)$, $(\mu^-, \bar{\nu}_\mu)$ or (q, \bar{q}) .




Figure 1.1: Feynman diagram of common decay of a τ -lepton into pairs of lepton-antineutrino or quark-antiquark by the emission of a W boson.

We are foremost interested into the hadronic decay channels, meaning τ -decays that have quarks in their final states. Unfortunately the quarks have never been measured isolated, but appear always in combination of *mesons* and *baryons*. Due to its mass of $m_\tau \approx 1.8 \text{ GeV}$ the τ -particle decays into light mesons (pions- π , kaons-K, and eta- η , see section 1.1), which can be experimentally detected.

The hadronic τ – decay provides one of the most precise ways to determine the strong coupling [18] and can be calculated to high precision within the framework of QCD.

⁶Leptons do not interact via the strong force.

⁷The τ -particle can also decay into strange quarks or charm quarks, but these decays are rather uncommon due to the heavy masses of s and c .

Name	Symbol	Quark content	Rest mass (MeV)
Pion	π^-	$\bar{u}d$	139.570 61(24) MeV
Pion	π^0	$(u\bar{u} - d\bar{d})/\sqrt{2}$	134.9770(5) MeV
Kaon	K^-	$\bar{u}s$	493.677(16) MeV
Kaon	K^0	$d\bar{s}$	497.611(13) MeV
Eta	η	$(u\bar{u} + d\bar{d} - 2s\bar{s})/\sqrt{6}$	547.862(17) MeV

Table 1.1: List of mesons produced by a τ -decay. Rare final states with branching Ratios smaller than 0.1 have been omitted. The list is taken from [9] with corresponding rest masses taken from [22].

Flavour	Mass	comment
u	$2.2^{+0.5}_{-0.4}$ MeV	\overline{MS}
d	$4.7^{+0.5}_{-0.3}$ MeV	
s	95^{+9}_{-3} MeV	
c	$1.275^{+0.025}_{-0.035}$ GeV	
b	$4.18^{+0.04}_{-0.03}$ GeV	
t	173.0(40) GeV	

Table 1.2: List of Quarks and their masses[22].

1.2 Quantumchromodynamics

QCD describes the strong interaction, which occur between *quarks* and are transmitted through *gluons*. A list of quarks can be found in 1.2.

The QCD Lagrange density is similar to that of QED[12],

$$\mathcal{L}_{\text{QCD}}(x) = -\frac{1}{4} G_{\mu\nu}^a(x) G^{\mu\nu a}(x) + \sum_A \left[\frac{i}{2} \bar{q}^A(x) \gamma^\mu \overleftrightarrow{D}_\mu q^A(x) - m_A \bar{q}^A(x) q^A(x) \right], \quad (1.1)$$

where $q^A(x)$ represents the quark fields and $G_{\mu\nu}^a$ being the *gluon field strength tensor* given by:

$$G_{\mu\nu}^a(x) \equiv \partial_\mu B_\nu^a(x) - \partial_\nu B_\mu^a(x) + g f^{abc} B_\mu^b(x) B_\nu^c(x), \quad (1.2)$$

where B_μ^a are the *gluon fields*, given in the *adjoint representation* of the $SU(3)$ gauge group with f^{abc} as *structure constants*. Furthermore we have used $A, B, \dots = 0, \dots, 5$ as flavour indices, $a, b, \dots = 0, \dots, 8$ as color indices and $\mu, \nu, \dots = 0, \dots, 3$ as lorentz indices.

1.2.1 Renormalisation Group

The perurbations of the QCD Lagrangian 1.1 lead to divergencies, which have to be *renormalized*. There are different approaches to ‘make’ these divergencies

finite. The most popular one is *dimensional regularisation*. In *Dimensional regularisation* we expand the four space-time dimensions to arbitrary dimensions. Consequently the in QCD calculations appearing *Feynman integrals* have to be continued to D-dimensions like

$$\mu^{2\epsilon} \int \frac{d^D p}{(2\pi)^D} \frac{1}{[p^2 - m^2 + i0][(q - p)^2 - m^2 + i0]}, \quad (1.3)$$

where we introduced the scale parameter μ to account for the extra dimensions and conserve the mass dimension of the non continued integral.

In addition *physical quantities*⁸ cannot depend on the renormalisation scale μ . Thus examining a *physical quantity* $R(q, a_s, m)$ that depends on the external momentum q , the renormalised coupling $a_s = \alpha_s/\pi$ and the renormalized quark mass m

$$\mu \frac{d}{d\mu} R(q, a_s, m) = \left[\mu \frac{\partial}{\partial \mu} + \mu \frac{da_s}{d\mu} \frac{\partial}{\partial a_s} + \mu \frac{dm}{d\mu} \frac{\partial}{\partial m} \right] R(q, a_s, m) = 0 \quad (1.4)$$

we can define the *renormalisation group functions*:

$$\beta(a_s) \equiv -\mu \frac{da_s}{d\mu} = \beta_1 a_s^2 + \beta_2 a_s^3 + \dots \quad \beta - \text{function} \quad (1.5)$$

$$\gamma(a_s) \equiv -\frac{\mu}{m} \frac{dm}{d\mu} = \gamma_1 a_s + \gamma_2 a_s^2 + \dots \quad \text{anomalous mass dimension.} \quad (1.6)$$

Running gauge coupling

Regarding the β -function we notice, that $a_s(\mu)$ is not a constant, but *runs* by varying the scale μ . Integrating the β -function yields

$$\int_{a_s(\mu_1)}^{a_s(\mu_2)} \frac{da_s}{\beta(a_s)} = - \int_{\mu_1}^{\mu_2} \frac{d\mu}{\mu} = \log \frac{\mu_1}{\mu_2}. \quad (1.7)$$

To analytically evaluate the above integral we can approximate the β -function to first order, with the known coefficient

$$\beta_1 = \frac{1}{6}(11N_c - 2N_f), \quad (1.8)$$

yielding

$$a_s(\mu_2) = \frac{a_s(\mu_1)}{\left(1 - a_s(\mu_1)\beta_1 \log \frac{\mu_1}{\mu_2}\right)}. \quad (1.9)$$

As we have three colours $N_c = 3$ and six flavours $N_f = 6$ the first β -function 1.5 is positive. Thus for $\mu_2 > \mu_1$ $a_s(\mu_2)$ decreases logarithmically and vanishes for $\mu_2 \rightarrow \infty$. This behaviour is known as *asymptotic freedom*. The coefficients of the β -function are currently known up to the 5th order, which are displayed in the appendix 4.1.

⁸Observables that can be measured.

Running quark mass

The properties of the running quark mass can be derived similar to the gauge coupling. Starting from integrating the *anomalous mass dimension* 1.6

$$\log \frac{m(\mu_2)}{m(\mu_1)} = \int_{a_s(\mu_1)}^{a_s(\mu_2)} da_s \frac{\gamma(a_s)}{\beta(a_s)} \quad (1.10)$$

we can approximate the *anomalous mass dimension* to first order and solve the integral analytically [19]

$$m(\mu_2) = m(\mu_1) \left(\frac{a(\mu_2)}{a(\mu_1)} \right)^{\frac{\gamma_1}{\beta_1}} (1 + \mathcal{O}(\beta_2, \gamma_2)). \quad (1.11)$$

As β_1 and γ_1 (see 4.2) are positive the quark mass decreases with increasing μ . The general relation between different scales is given by

$$m(\mu_2) = m(\mu_1) \exp \left(\int_{a_s(\mu_1)}^{a_s(\mu_2)} da_s \frac{\gamma(a_s)}{\beta(a_s)} \right) \quad (1.12)$$

and can be solved numerically to run the quark mass to the needed scale μ_2 .

1.2.2 Sum Rules

We need to relate the measurable hadronic final states of a QCD process (e.g. τ -decays into Hadrons) to a theoretical calculable value. Consequently we will employ **QCD Sum Rules**[20], which is a combination of the **operator product expansion** (OPE), the **optical theorem**, a **dispersion relation** the analyticity of the **two-point function** and the **quark hadron duality**.

Starting from the the vacuum expectation value of the product of the conserved noether current $J_\mu(x)$ at different space-times points x and y , which is known as the *two-point function* (or simply correlator)

$$\Pi(q^2) = \langle 0 | J_\mu(x) J_\nu(y) | 0 \rangle, \quad (1.13)$$

where the noether current is given by

$$J_\mu(x) = \Psi^\dagger(x) \gamma_\mu (\gamma_5) \Psi(x). \quad (1.14)$$

The two-point function, within the framework of QCD sum rules, is improved by the OPE expansion

$$\Pi_{\text{OPE}}(s) = \sum_n C_{2n}(s, \mu) \frac{\langle \hat{\mathcal{O}}(\mu) \rangle}{s^n}, \quad (1.15)$$

where we used $q^2 = s$. It is furthermore related to the hadronic **spectral function** $\rho(q^2)$ through the *Källén-Lehmann spectral representation* [13][16]

$$\Pi(q^2) = \int_0^\infty ds \frac{\rho(s)}{s - q^2 - i\epsilon}, \quad (1.16)$$

where the spectral function $\rho(s)$ is defined as

$$\rho(s) \equiv \frac{1}{\pi} \text{Im} \Pi(s). \quad (1.17)$$

Equation 1.16 is referred to as **dispersion relation** analogous to similar relations which arise for example in electrodynamics. The main contribution from the spectral function given in eq. (1.16) are the hadronic final states

$$2\pi\rho(m^2) = \sum_n \langle 0 | J_\mu(x) | n \rangle \langle n | J_\nu(y) \rangle (2\pi^2)^4 \delta^{(4)}(p - p_n), \quad (1.18)$$

which lead to a series of continuous poles on the positive real axis for the two-point function, see Fig. 1.2.2. As the experimental data, which solely




Figure 1.2: Analytic structure in the complex q^2 -plane of the Fourier transform of the two-point function. The hadronic final states are responsible for poles appearing on the real-axis. The one-particle states contribute as isolated pole and the multi-particle states contribute as bound-states poles or a continuous “discontinuity cut” (see [17]).

contributes to the spectral function $\rho(q^2)$, is only accessible on the positive real axis, we have to use Cauchy’s theorem to access the theoretical values of the two-point function close to the positive real axis (see section 1.2.2).

The final ingredient of the QCD sum rules is the *optical theorem*, relating experimental data with the imaginary part of the correlator. E.g. taking the total e^+e^- cross section scattering into hadrons

$$R_q(s) \equiv \frac{\sigma(e^+e^- \rightarrow \gamma^* \rightarrow q\bar{q})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} = 12\pi \text{Im} \Pi_{\text{Had}}(s). \quad (1.19)$$




Figure 1.3: Analytical structure of $\Pi(s)$ with the used contour \mathcal{C} for the final QCD Sum Rule expression [eq. \(1.20\)](#).

Due to asymptotic freedom⁹ experiments can only detect Hadrons (note the exp-index in $\text{Im}\Pi_{\text{exp}}(s)$), but on the theory side we are calculating with quarks as degrees of freedom. Consequently we assume that Π_{Had} can be set equal to Π_{OPE} , which is referred to as *quark hadron duality*.

In total, with the help Cauchy's theorem, the QCD sum rules can be summed up in the following expression

$$\frac{1}{\pi} \int_0^\infty \frac{\text{Im} \Pi_{\text{Had}}(t)}{t-s} dt = \frac{1}{\pi} \oint_{\mathcal{C}} \frac{\text{Im} \Pi_{\text{OPE}}(t)}{t-s} dt, \quad (1.20)$$

where the l.h.s. is given by the experiment and the r.h.s. can be theoretically evaluated with by applying the OPE of the correlator $\Pi_{\text{OPE}}(s)$.

⁹There are no free quarks. They are bound in pairs of two or three.

Chapter 2

τ decays into hadrons

The τ -lepton is the only lepton heavy enough to decay into Hadrons. It permits one of the most precise determinations of the strong coupling α_s . The inclusive τ -decay ratio

$$R_\tau = \frac{\Gamma(\tau \rightarrow \nu_\tau + \text{Hadrons})}{\Gamma(\tau \rightarrow \nu_\tau e^+ e^-)} \quad (2.1)$$

can be precisely calculated and is sensitive to α_s . Due to the mass of the τ -lepton $m_\tau \approx 1.8 \text{ GeV}$ it is a good phenomenon to perform low-energy QCD analysis. The theoretical expression of the hadronic τ -decay ratio was first derived by [23] (using current algebra, a more recent derivation making use of the *optical theorem* can be taken from [19]):

$$R_\tau = 12\pi \int_0^{m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right) \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im} \Pi^{(T)}(s) + \text{Im} \Pi^{(L)}(s) \right]. \quad (2.2)$$

The τ -decay ratio depends on the two-point function

$$\Pi_{\mu\nu,ij}^{V/A}(s) \equiv i \int dx e^{ipx} \langle \Omega | T \{ J_{\mu,ij}^{V/A}(x) J_{\nu,ij}^{V/A}(0)^\dagger \} | \Omega \rangle, \quad (2.3)$$

with $|\Omega\rangle$ being the physical vacuum. The vectorial and axial-vector currents are given by

$$J_{\mu,ij}^V(x) = \bar{q}_j(x) \gamma_\mu q_i(x) \quad \text{and} \quad J_{\mu,ij}^A(x) = \bar{q}_j(x) \gamma_\mu \gamma_5 q_i(x) \quad (2.4)$$

where i, j stand for the light quark flavours u, d and s .

The general correlator $\Pi^{\mu\nu}(q^2)$ can be decomposed into a vector/ axial-vector (V/A) and scalar/ pseudo-scalar (S/P) part containing a correction [6]

$$\begin{aligned} \Pi^{\mu\nu}(q^2) = & (q^\mu q^\nu - q^2 g^{\mu\nu}) \Pi^{V,A}(q^2) + \frac{g^{\mu\nu}}{q^2} (m_i \mp m_j) \Pi^{S,P}(q^2) \\ & + g^{\mu\nu} \frac{(m_i \mp m_j)}{q^2} [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle], \end{aligned} \quad (2.5)$$

which is composed of a vector $\Pi^{V,A}$ and scalar $\Pi^{S,P}$ part. The third term are corrections arising due to the physical vacuum $|\Omega\rangle$.

The general correlator $\Pi^{\mu\nu}$ can also be decomposed into transversal and longitudinal components:

$$\Pi^{\mu\nu}(q^2) = (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T)}(q^2) + q^\mu q^\nu \Pi^{(L)}(q^2). \quad (2.6)$$

To relate the two different contributions we note, that only the scalar components of [eq. \(2.5\)](#) carry a mass term. Using the *Ward identity*

$$q_\mu \Pi^{\mu\nu}(q^2) = 0 \quad (2.7)$$

we can introduce two four-momenta into [eq. \(2.5\)](#)

$$q_\mu q_\nu \Pi^{\mu\nu}(q^2) = (m_i \mp m_j)^2 \Pi^{S,P}(q^2) + (m_i \mp m_j) [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle] \quad (2.8)$$

to relate the longitudinal of [eq. \(2.6\)](#)

$$q_\mu q_\nu \Pi^{\mu\nu}(q^2) = q^4 \Pi^{(L)}(q^2) = s^2 \Pi^{(L)}(s), \quad (2.9)$$

where we defined $s \equiv q^2$. Thus

$$s^2 \Pi^{(L)}(s) = (m_i \mp m_j)^2 \Pi^{(S,P)}(s) + (m_i \mp m_j) [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle]. \quad (2.10)$$

Furthermore we can relate the transversal and vectorial components via

$$\begin{aligned} \Pi^{\mu\nu}(s) &= \underbrace{(q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T)}(s) + (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(L)}(s)}_{=(q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T+L)}(s)} + \frac{g^{\mu\nu} s^2}{q^2} \Pi^{(L)}(s) \end{aligned} \quad (2.11)$$

where $\Pi^{(T+L)}(s) \equiv \Pi^{(T)}(s) + \Pi^{(L)}(s)$, such that

$$\Pi^{(V,A)}(s) = \Pi^{(T)}(s) + \Pi^{(L)}(s) = \Pi^{(T+L)}(s). \quad (2.12)$$

Inspecting the inclusive ratio R_τ in [eq. \(2.1\)](#) introduces a problematic integral over the real axis of $\Pi(s)$ from 0 up to m_τ . The integral is problematic for two reasons:

- The *perturbative Quantum Chromodynamics (pQCD)* and the OPE breaks down for low energies (over which we have to integrate).
- The positive euclidean axis of $\Pi(s)$ has a discontinuity cut and can theoretically not be evaluated.

To literally circumvent these issues we make use of *Cauchy's Theorem*

$$\int_{\mathcal{C}} f(z) dz = 0, \quad (2.13)$$

where $f(z)$ is an analytic function on a closed contour \mathcal{C} .

In our case we have to deal with the two-point correlator $\Pi(s)$, which is analytic except for the positive real axis (with which we will deal with to a later point¹) Consequently, to rewrite we can rewrite the definite integral of eq. (2.2) into a contour integral over a closed circle with radius m_τ^2 . The closed contour consists of four line integrals, which have been visualized in fig. 2.1. Summing over the four line integrals, performing a *analytic continuation* of the two-point correlator $\Pi(s) \rightarrow \Pi(s + i\epsilon)$ and finally taking the limit of $\epsilon \rightarrow 0$ gives us the needed relation between eq. (2.2) and the closed contour:

$$\begin{aligned}
\oint_{s=m_\tau} \Pi(s) &= \int_0^{m_\tau} \Pi(s + i\epsilon) ds + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{m_\tau}^0 \Pi(s - i\epsilon) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\
&= \int_0^{m_\tau} \Pi(s + i\epsilon) - \Pi(s - i\epsilon) ds + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\
&= \int_0^{m_\tau} \Pi(s + i\epsilon) - \overline{\Pi(s + i\epsilon)} + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\
&\stackrel{\lim \epsilon \rightarrow 0}{=} 2i \int_0^{m_\tau} \text{Im} \Pi(s) ds + \oint_{s=m_\tau} \Pi(s) ds
\end{aligned} \tag{2.14}$$

where we made use of $\Pi(z) = \overline{\Pi(\bar{z})}$ (due to $\Pi(s)$ is analytic) and $\Pi(z) - \overline{\Pi(\bar{z})} = 2i \text{Im} \Pi(z)$. The result can be rewritten in a more intuitive form, which we also visualized in fig. 2.1

$$\int_0^{m_\tau} \Pi(s) ds = \frac{i}{2} \oint_{s=m_\tau} \Pi(s) ds \tag{2.15}$$

Finally combining eq. (2.15) with eq. (2.2) we get

$$R_\tau = 6\pi i \oint_{s=m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right) \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(T)}(s) + \Pi^{(L)} \right] \tag{2.16}$$

for the hadronic τ -decay ratio.

The contour integral obtained is an import result as we can now theoretically evaluate the hadronic τ -decay ratio sufficiently large energy scales ($m_\tau \approx 1.78 \text{ MeV}$) at which $\alpha_s(m_\tau) \approx 0.33$ [18] is tolerable heigh for applying perturbation theory and the OPE. Obviously we would benefit from a contour integral over a bigger circumference, but τ -decays are limited by the m_τ . Nevertheless there are promising e^+e^- annihilation data, which yields valuable R-ratio values up to 2 GeV [4][14].

It is convenient to rewrite the

$$\Pi^{(L+T)} = \Pi^{(L)} + \Pi^{(T)} \tag{2.17}$$

¹To not evaluate $\Pi(s)$ at the positive real axis we have to introduce *pinched weights*. The *pinched weights* vanish for $s \rightarrow m_\tau$.




Figure 2.1: Visualization of the usage of Cauchy's theorem to transform eq. (2.2) into a closed contour integral over a circle of radius m_τ^2 .

$$R_\tau = 6\pi i \oint_{|s|=m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(L+T)}(s) - \left(\frac{2s}{m_\tau^2}\right) \Pi^{(L)}(s) \right] \quad (2.18)$$

$$D^{(L+T)}(s) \equiv -s \frac{d}{ds} \Pi^{(L+T)}(s), \quad D^{(L)}(s) \equiv \frac{s}{m_\tau^2} \frac{d}{ds} (s \Pi^{(L)}(s)) \quad (2.19)$$

Integration by parts

$$\int_a^b u(x) V(x) dx = [u(x) V(x)]_a^b - \int_a^b u(x) v(x) dx \quad (2.20)$$

$$\begin{aligned} R_\tau^{(1)} &= \frac{6\pi i}{m_\tau^2} \oint_{|s|=m_\tau^2} \underbrace{\left(1 - \frac{s}{m_\tau^2}\right)^2}_{=u(x)} \underbrace{\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(L+T)}(s)}_{=V(x)} \\ &= \frac{6\pi i}{m_\tau^2} \left\{ \left[-\frac{m_\tau^2}{2} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left(1 + \frac{s}{m_\tau^2}\right) \Pi^{(L+T)}(s) \right]_{|s|=m_\tau^2} \right. \\ &\quad \left. + \oint_{|s|=m_\tau^2} \underbrace{-\frac{m_\tau^2}{2} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left(1 + \frac{s}{m_\tau^2}\right)}_{=u(x)} \underbrace{\frac{d}{ds} \Pi^{(L+T)}(s)}_{=v(x)} \right\} \\ &= -3\pi i \oint_{|s|=m_\tau^2} \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left(1 + \frac{s}{m_\tau^2}\right) \frac{d}{ds} D^{(L+T)} \end{aligned} \quad (2.21)$$

where we fixed the integration constant to $C = -\frac{m_\tau^2}{2}$ in the second line and left

the antiderivatives contained in the squared brackets untouched. Parametrizing the expression in the squared brackets

$$\left[-\frac{m_\tau^2}{2} (1 - e^{-i\phi})^3 (1 + e^{-i\phi}) \Pi^{(L+T)}(m_\tau^2 e^{-i\phi}) \right]_0^{2\pi} = 0 \quad (2.22)$$

where $s \rightarrow m_\tau^2 e^{-i\phi}$ and $(1 - e^{-i \cdot 0}) = (1 - e^{-i \cdot 2\pi}) = 0$.

$$\begin{aligned} R_\tau^{(2)} &= \oint_{|s|=m_\tau^2} ds \left(1 - \frac{s}{m_\tau^2} \right)^2 \left(-\frac{2s}{m_\tau^2} \right) \Pi^{(L)}(s) \\ &= -4\pi i \oint \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2} \right)^3 D^{(L)}(s) \end{aligned} \quad (2.23)$$

$$R_\tau = -\pi i \oint_{|s|=m_\tau^2} \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2} \right)^3 \left[3 \left(1 + \frac{s}{m_\tau^2} D^{(L+T)}(s) + 4 D^{(L)}(s) \right) \right] \quad (2.24)$$

$$R_\tau = -\pi i \oint_{|s|=m_\tau^2} \frac{dx}{x} (1-x)^3 \left[3(1+x) D^{(L+T)}(m_\tau^2 x) + 4 D^{(L)}(m_\tau^2 x) \right], \quad (2.25)$$

where $x = s/m_\tau^2$.

$$R_{\tau,V/A}^\omega = \frac{N_c}{2} S_{EW} |V_{ud}|^2 \left(1 + \delta_\omega^{(0)} + \delta_\omega^{EW} + \delta_\omega^{DV} s + \sum_{D \leq 2} \delta_{ud,\omega}^{(D)} \right) \quad (2.26)$$


2.1 The perturbative expansion

We will treat the correlator in the chiral limit for which the longitudinal components $\Pi^L(s)$ vanish (see eq. (2.11)) and the axial and vectorial contributions are equal. Consequently [2] we can write the vector correlation function $\Pi(s)$ as:

$$\Pi_V^{T+L}(s) = -\frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=0}^{n+1} c_{n,k} L^k \quad \text{with} \quad L \equiv \ln \frac{-s}{\mu^2}. \quad (2.27)$$

The coefficient $c_{n,k}$ up to two-loop order can be obtained by Feynman-diagram calculations. **add complete calculation** E.g. we can compare the zero-loop result of the correlator [12]

$$\Pi_{\mu\nu}^B(q^2) \Big|^{1\text{-loop}} = \frac{N_c}{12\pi^2} \left(\frac{1}{\hat{\epsilon}} - \log \frac{(-q^2 - i0)}{\mu^2} + \frac{5}{3} + \mathcal{O}(\epsilon) \right) \quad (2.28)$$



with eq. (2.27) and extract the first two coefficients

$$c_{00} = -\frac{5}{3} \quad \text{and} \quad c_{01} = 1, \quad (2.29)$$

where $\Pi_{\mu\nu}^B(q^2)$ is not renormalized²

The second loop can also be calculated by diagram techniques resulting in [3]

$$\Pi_V^{(1+0)}(s) \Big|^{2\text{-loop}} = -\frac{N_c}{12\pi^2} a_\mu \log\left(\frac{-s}{\mu^2}\right) + \dots \quad (2.30)$$

yielding $c_{11} = 1$.

Beginning from three loop diagrams the algebra becomes exhausting and one has to use dedicated algorithms to compute the heigher loops. The third loop calculations have been done in the late seventies by [8, 10, 7]. The four loop evaluation have been completed a little more than ten years later by [11, 21]. The heighest loop published, that amounts to α_s^4 , was published in 2008 [1] almost 20 years later.

Fixing the number of colors to $N_c = 3$ the missing coefficients up to order four in α_s read:

$$\begin{aligned} c_{2,1} &= \frac{365}{24} - 11\zeta_3 - \left(\frac{11}{12} - \frac{2}{3}\zeta_3\right) N_f \\ c_{3,1} &= \frac{87029}{288} - \frac{1103}{4}\zeta_3 + \frac{275}{6}\zeta_5 \\ &\quad - \left(\frac{7847}{216} - \frac{262}{9}\zeta_3 + \frac{25}{9}\zeta_5\right) N_f + \left(\frac{151}{162} - \frac{19}{27}\zeta_3\right) N_f^2 \\ c_{4,1} &= \frac{78631453}{20736} - \frac{1704247}{432}\zeta_3 + \frac{4185}{8}\zeta_3^2 + \frac{34165}{96}\zeta_5 - \frac{1995}{16}\zeta_7, \end{aligned} \quad (2.31)$$

where used the flavour number $N_f = 3$ for the last line.

The 6-loop calculation has until today not been achieved, but Beneke und Jamin [2] used and educated guess to estimate the coefficient

$$c_{5,1} \approx 283 \pm 283. \quad (2.32)$$

Until know we have mentioned the coefficients $c_{n,k}$ with a fixed $k = 1$. This is due to the RGE, which relates coefficients with a different k to the coefficients mentioned above. To make usage of the RGE $\Pi_V^{T+L}(s)$ needs to be

²The term $1/\hat{\epsilon}$, which is of order 0 in α_s , will be cancelled by renormalization.

a physical quantity, which can be achieved by rewriting eq. (2.19) to:

$$D_V^{(T+L)} = -s \frac{d\Pi_V^{(T+L)}(s)}{ds} = \frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=1}^{n+1} k c_{n,k} L^{k-1}, \quad (2.33)$$

where we used $dL^k/ds = k \ln(-s/\mu^2)^{k-1} (-1/\mu^2)$. D_V^{1+0} being a physical quantity has to fulfill the RGE ??

$$-\mu \frac{d}{d\mu} D_V^{(T+L)} = -\mu \frac{d}{d\mu} \left(\frac{\partial}{\partial L} L + \frac{\partial}{\partial a_s} a_s \right) D_V^{T+L} = \left(2 \frac{\partial}{\partial L} + \beta \frac{\partial}{\partial a_s} \right) D_V^{T+L} = 0, \quad (2.34)$$

where we defined the β -function in eq. (1.5) and used $dL/d\mu = -2/\mu$. The RGE puts constraints on the $c_{n,k}$ -coefficients, ... not independent

$$D(s) = \frac{N_c}{12\pi^2} \left[c_{01} + a_\mu (c_{11} + 2c_{12}L) + a_\mu^2 (c_{21} + 2c_{22}L + 3c_{23}L^2) \right] \quad (2.35)$$

inserting into RGE

$$4a_\mu c_{12} + 2a_\mu^2 (2c_{22} + 6c_{23}L) + \beta_1 a_\mu^2 (c_{11} + 2c_{12}L) + \mathcal{O}(a_\mu^3) = 0 \quad (2.36)$$

Thus

$$c_{12} = 0 \quad c_{22} = \frac{\beta_1 c_{11}}{4} \quad c_{23} = 0 \quad (2.37)$$

or $D(s)$ to the first order in α_s

$$D(s) = \frac{N_c}{12\pi^2} \left[c_{01} + c_{11} a_\mu \left(c_{21} - \frac{1}{2} \beta_1 c_{11} L \right) a_\mu^2 \right] + \mathcal{O}(a_\mu^3) \quad (2.38)$$

2.1.1 Renormalisation group summation

We can express the perturbative contribution $\delta^{(0)}$ to R_τ (see eq. (2.26)) as

$$\delta^{(0)} = \sum_{n=1}^{\infty} a_\mu^n \sum_{k=1}^n k c_{n,k} \frac{1}{2\pi i} \oint_{|x|=1} \frac{dx}{x} (1-x)^3 (1+x) \log \left(\frac{-M_\tau^2 x}{\mu^2} \right)^{k-1}, \quad (2.39)$$

where we inserted the expansion of $D_V^{(T+L)}$ eq. (2.19) into R_τ eq. (2.25). Keep in mind that we are working in the chiral limit, such that $D^L = 0$ vanishes and the contributions from the vector and axialvector correlator are identical

$$D^{(T+L)} = D_V^{(T+L)} + D_A^{(T+L)} = 2D_V^{(T+L)}. \quad (2.40)$$

The perturbative contribution $\delta^{(0)}$ is a physical quantity and satisfies the homogeneous RGE, thus is independent on the scale μ . Consequently we have the freedom to choose μ , which leads to two main descriptions **fixed-order perturbation theory** (FOPT) and **contour-improved perturbation theory** (CIPT). The two resulting series should converge to equal values, but differ notably.

By using the FOPT prescription we fix $\mu^2 = m_\tau^2$ leading to

$$\delta_{\text{FO}}^{(0)} = \sum_{n=1}^{\infty} a(m_\tau^2)^n \sum_{k=1}^n k c_{n,k} J_{k-1} \quad (2.41)$$

where the contour integrals J_l are defined by

$$J_l \equiv \frac{1}{2\pi i} \oint_{|x|=1} \frac{dx}{x} (1-x)^3 (1+x) \log^l(-x). \quad (2.42)$$

The integrals J_l up to order α_s^4 are given by [2]:

$$J_0 = 1, \quad J_1 = -\frac{19}{12}, \quad J_2 = \frac{265}{72} - \frac{1}{3}\pi^2, \quad J_3 = -\frac{3355}{288} + \frac{19}{12}\pi^2. \quad (2.43)$$

Using FOPT the strong coupling $a(\mu)$, which runs with the scale μ , is fixed at $a(m_\tau^2)$ and can be taken out of the closed-contour integral. We still have to integrate over the logarithms $\log(-s/m_\tau^2)$.

Using CIPT we can sum the logarithms by setting the scale to $\mu^2 = -m_\tau^2 x$ in eq. (2.39), resulting in:

$$\delta_{\text{CI}}^{(0)} = \sum_{n=1}^{\infty} c_{n,1} J_n^a(m_\tau^2), \quad (2.44)$$

where the contour integrals J_l are defined by

$$J_n^a(m_\tau^2) \equiv \frac{1}{2\pi i} \oint_{|x|=1} \frac{dx}{x} (1-x)^3 (1+x) a^n(-m_\tau^2 x). \quad (2.45)$$

All logarithms vanish except the ones for $k = 1$:

$$\log(1)^{k-1} = \begin{cases} 1 & \text{if } k = 1, \\ 0 & \text{if } k \neq 1 \end{cases} \quad (2.46)$$

which selects adler function coefficients $c_{n,1}$ with a fixed $k = 1$. Handling the logarithms left us with the integration of $a_s(-m_\tau^2 x)$ over the closed-contour $\oint_{|x|=1}$, which now depends on the integration variable x .

Calculating the perturbative contribution $\delta^{(0)}$ to R_τ for the two different prescriptions yields [2]

$$\begin{array}{cccccc} \alpha_s^2 & \alpha_s^2 & \alpha_s^3 & \alpha_s^4 & \alpha_s^5 & \\ \delta_{\text{FO}}^{(0)} = 0.1082 + 0.0609 + 0.0334 + 0.0174(+0.0088) = 0.2200(0.2288) & (2.47) \end{array}$$

$$\delta_{\text{CI}}^{(0)} = 0.1479 + 0.0297 + 0.0122 + 0.0086(+0.0038) = 0.1984(0.2021). \quad (2.48)$$

The series indicate, that CIPT converges faster and that both series approach a different value. This discrepancy represents currently the biggest theoretical uncertainty while extracting the strong coupling α_s .

As today we do not know which if FOPT or CIPT is the correct approach of measuring α_s . Therefore there are currently three ways of stating result:

- Quoting the average of both results.
- Quoting the CIPT result.
- Quoting the FOPT result.

We follow the approach of Beneke and Jamin [Benke2008] who prefer FOPT.

2.2 Non-Perturbative OPE Contribution

The perturbative contribution to the Sum-Rule, that we have seen so far, is the dominant one. With

$$\begin{aligned} R_\tau^{\text{FOPT}} &= \\ R_\tau^{\text{CIPT}} &= \end{aligned} \quad (2.49)$$

The NP vs perturbative contributions can be varied by chosen different weights than ω_τ .

2.2.1 Dimension four

For the OPE contributions of dimension four we have to take into account the terms with masses to the fourth power m^4 , the quark condensate multiplied by a mass $m\langle\bar{q}q\rangle$ and the gluon condensate $\langle GG\rangle$. The resulting expression can be taken from the appendix of [Pich1999], yielding:

$$D_{ij}^{(L+T)}(s)\Big|_{D=4} = \frac{1}{s^2} \sum_n \Omega^{(1+0)}(s/\mu^2) a^n, \quad (2.50)$$

where

$$\begin{aligned} \Omega_n^{(1+0)}(s/\mu^2) &= \frac{1}{6} \langle aGG \rangle p_n^{(L+T)}(s/\mu^2) + \sum_k m_k \langle \bar{q}_k q_k \rangle r_n^{(L+T)}(s/\mu^2) \\ &\quad + 2 \langle m_i \bar{q}_i q_i + m_j \bar{q}_j q_j \rangle q_n^{(L+T)}(s/\mu^2) \pm \frac{8}{3} \langle m_j \bar{q}_i q_i + m_i \bar{q}_j q_j \rangle t_n^{(L+T)} \\ &\quad - \frac{3}{\pi^2} (m_i^4 + m_j^4) h_n^{(L+T)}(s/\mu^2) \mp \frac{5}{\pi^2} m_i m_j (m_i^2 + m_j^2) k_n^{(L+T)}(s/\mu^2) \\ &\quad + \frac{3}{\pi^2} m_i^2 m_j^2 g_n^{(L+T)}(s/\mu^2) + \sum_k m_k^4 j_n^{(L+T)}(s/\mu^2) + 2 \sum_{k \neq l} m_k^2 m_l^2 u_n^{(L+T)}(s/\mu^2) \end{aligned} \quad (2.51)$$

The perturbative expansion coefficients are known to $\mathcal{O}(a^2)$ for the condensate contributions,

$$\begin{aligned} p_0^{(L+T)} &= 0, & p_1^{(L+T)} &= 1, & p_2^{(L+T)} &= \frac{7}{6}, \\ r_0^{(L+T)} &= 0, & r_1^{(L+T)} &= 0, & r_2^{(L+T)} &= -\frac{5}{3} + \frac{8}{3} \zeta_3 - \frac{2}{3} \log(s/\mu^2), \\ q_0^{(L+T)} &= 1, & q_1^{(L+T)} &= -1, & q_2^{(L+T)} &= -\frac{131}{24} + \frac{9}{4} \log(s/\mu^2), \\ t_0^{(L+T)} &= 0, & t_1^{(L+T)} &= 1, & t_2^{(L+T)} &= \frac{17}{2} + \frac{9}{2} \log(s/\mu^2). \end{aligned} \quad (2.52)$$

while the m^4 terms have been only computed to $\mathcal{O}(a)$

$$\begin{aligned}
h_0^{(L+T)} &= 1 - 1/2 \log(s/\mu^2), & h_1^{(L+T)} &= \frac{25}{4} - 2\zeta_3 - \frac{25}{6} \log(s/\mu^2) - 2 \log(s/\mu^2)^2, \\
k_0^{(L+T)} &= 0, & k_1^{(L+T)} &= 1 - \frac{2}{5} \log(s/\mu^2), \\
g_0^{(L+T)} &= 1, & g_1^{(L+T)} &= \frac{94}{9} - \frac{4}{3} \zeta_3 - 4 \log(s/\mu^2), \\
j_0^{(L+T)} &= 0, & j_1^{(L+T)} &= 0, \\
u_0^{(L+T)} &= 0, & u_2^{(L+T)} &= 0.
\end{aligned} \tag{2.53}$$

2.2.2 Dimension six and eight

Our application of dimension six contributions is founded in [5] and has previously been calculated beyond leading order by [15]. The operators appearing are the masses to the power six m^6 , the four-quark condensates $\langle \bar{q}q\bar{q}q \rangle$, the three-gluon condensates $\langle g^3 G^3 \rangle$ and lower dimensional condensates multiplies by the corresponding masses, such that in total the mass dimension of the operator will be six. As there are too many parameters to be fitted with experimental data we have to omit some of them, starting with the three-gluon condensate, which does not contribute at leading order. The four-quark condensates known up to $\mathcal{O}(a^2)$, but we will make use of the *vacuum saturation approach* [2, 5, 20] to express them in quark, anti-quark condensates $\langle q\bar{q} \rangle$. In our work we take the simplest approach possible: Introducing an effective dimension six coefficient $\rho_{V/A}^{(6)}$ divided by the appropriate power in s

$$D_{ij,V/A}^{(1+0)} \Big|_{D=6} = 0.03 \frac{\rho_{V/A}^{(6)}}{s^3} \tag{2.54}$$

As for the dimension eighth contribution the situation is not better than the dimension six one we keep the simplest approach, leading to

$$D_{ij,V/A}^{(1+0)} \Big|_{D=8} = 0.04 \frac{\rho_{V/A}^{(8)}}{s^4}. \tag{2.55}$$

Chapter 3

Derivation of the used inverse covariance matrix from the Aleph data

While performing a **Generalized least squares** (GLS) we estimate our regression coefficients $\hat{\beta}$ as follows:

$$\hat{\beta} = \underset{\mathbf{b}}{\operatorname{argmin}} (\mathbf{y} - \mathbf{X}\mathbf{b})^T \mathbf{\Omega}^{-1} (\mathbf{y} - \mathbf{X}\mathbf{b}), \quad (3.1)$$

with \mathbf{b} being an candidate estimate of β , \mathbf{X} being the design matrix, \mathbf{y} being the response values and $\mathbf{\Omega}^{-1}$ being the **inverse covariance matrix**.

The Aleph data includes the **standard error** (SE), which are equal to the **standard deviation** as per definition. Furthermore Aleph provides the **correlation coefficients** of the errors. We will use these two quantities in combination with **Gaussian error propagation** to derive an approximation of the covariance matrix.

3.1 Propagation of experimental errors and correlation

Let $\{f_k(x_1, x_2, \dots, x_n)\}$ be a set of m functions, which are linear combinations of n variables x_1, x_2, \dots, x_n with combination coefficients $A_{k1}, A_{k2}, \dots, A_{kn}$, where $k \in \{1, 2, \dots, m\}$. Let the covariance matrix of x_n be denoted by

$$\Sigma^x = \begin{pmatrix} \sigma_1^2 & \sigma_{12} & \sigma_{13} & \cdots \\ \sigma_{12} & \sigma_2^2 & \sigma_{23} & \cdots \\ \sigma_{13} & \sigma_{23} & \sigma_3^2 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \quad (3.2)$$

Then the covariance matrix of the functions Σ^f is given by

$$\Sigma_{ij}^f = \sum_k^n \sum_l^n A_{ik} \sum_{kl}^x A_{jl}, \quad \Sigma^f = A \Sigma^x A^T. \quad (3.3)$$

In our case we are dealing with non-linear functions, which we will linearized with the help of the **Taylor expansion**

$$f_k \approx f_k^0 + \sum_i^n \frac{\partial f_k}{\partial x_i} x_i, \quad f \approx f^0 + Jx. \quad (3.4)$$

Therefore, the propagation of error follows from the linear case, replacing the Jacobian matrix with the combination coefficients ($J = A$)

Chapter 4

Coefficients

4.1 β function

There are several conventions for defining the β coefficients, depending on a minus sign and/or a factor of two (if one substitutes $\mu \rightarrow \mu^2$) in the β -function [1.5](#). We follow the convention from Pascual and Tarrach (except for the minus sign) and have taken the values from [\[3\]](#)

$$\beta_1 = \frac{1}{6}(11N_c - 2N_f) \quad (4.1)$$

$$\beta_2 = \frac{1}{12}(17N_c^2 - 5N_cN_f - 3C_fN_f) \quad (4.2)$$

$$\beta_3 = \frac{1}{32} \left(\frac{2857}{54}N_c^3 - \frac{1415}{54}N_c^2N_f + \frac{79}{54}N_cN_f^2 - \frac{205}{18}N_cC_fN_f + \frac{11}{9}C_fN_f^2 + C_f^2N_f \right) \quad (4.3)$$

$$\beta_4 = \frac{140599}{2304} + \frac{445}{16}\zeta_3, \quad (4.4)$$

where we used $N_f = 6$ and $N_c = 3$ for β_4 .

4.2 Anomalous mass dimension

4.3 Adler function

Bibliography

- [1] P. A. Baikov, K. G. Chetyrkin, and Johann H. Kuhn. “Order α_s^4 QCD Corrections to Z and tau Decays”. In: *Phys. Rev. Lett.* 101 (2008), p. 012002. DOI: [10.1103/PhysRevLett.101.012002](https://doi.org/10.1103/PhysRevLett.101.012002). arXiv: [0801.1821](https://arxiv.org/abs/0801.1821) [hep-ph].
- [2] Martin Beneke and Matthias Jamin. “ α_s and the tau hadronic width: fixed-order, contour-improved and higher-order perturbation theory”. In: *JHEP* 09 (2008), p. 044. DOI: [10.1088/1126-6708/2008/09/044](https://doi.org/10.1088/1126-6708/2008/09/044). arXiv: [0806.3156](https://arxiv.org/abs/0806.3156) [hep-ph].
- [3] Diogo Boito. “QCD phenomenology with τ and charm decays”. PhD thesis. Universitat Autònoma de Barcelona, Sept. 2011.
- [4] Diogo Boito et al. “Strong coupling from $e^+e^- \rightarrow$ hadrons below charm”. In: *Phys. Rev. D* 98.7 (2018), p. 074030. DOI: [10.1103/PhysRevD.98.074030](https://doi.org/10.1103/PhysRevD.98.074030). arXiv: [1805.08176](https://arxiv.org/abs/1805.08176) [hep-ph].
- [5] E. Braaten, Stephan Narison, and A. Pich. “QCD analysis of the tau hadronic width”. In: *Nucl. Phys. B* 373 (1992), pp. 581–612. DOI: [10.1016/0550-3213\(92\)90267-F](https://doi.org/10.1016/0550-3213(92)90267-F).
- [6] D. J. Broadhurst. “A strong constraint on chiral symmetry breaking at short distances”. In: *Nuclear Physics B* 85 (Jan. 1975), pp. 189–207. DOI: [10.1016/0550-3213\(75\)90564-7](https://doi.org/10.1016/0550-3213(75)90564-7).
- [7] William Celmaster and Richard J. Gonsalves. “An Analytic Calculation of Higher Order Quantum Chromodynamic Corrections in e^+e^- Annihilation”. In: *Phys. Rev. Lett.* 44 (1980), p. 560. DOI: [10.1103/PhysRevLett.44.560](https://doi.org/10.1103/PhysRevLett.44.560).
- [8] K. G. Chetyrkin, A. L. Kataev, and F. V. Tkachov. “Higher Order Corrections to $\Sigma(e^+e^- \rightarrow \text{Hadrons})$ in Quantum Chromodynamics”. In: *Phys. Lett.* 85B (1979), pp. 277–279. DOI: [10.1016/0370-2693\(79\)90596-3](https://doi.org/10.1016/0370-2693(79)90596-3).
- [9] Michel Davier, Andreas Höcker, and Zhiqing Zhang. “The physics of hadronic tau decays”. In: *Rev. Mod. Phys.* 78 (4 Oct. 2006), pp. 1043–1109. DOI: [10.1103/RevModPhys.78.1043](https://doi.org/10.1103/RevModPhys.78.1043). URL: <https://link.aps.org/doi/10.1103/RevModPhys.78.1043>.

- [10] Michael Dine and J. R. Sapiirstein. “Higher Order QCD Corrections in $e^+ e^-$ Annihilation”. In: *Phys. Rev. Lett.* 43 (1979), p. 668. DOI: [10.1103/PhysRevLett.43.668](https://doi.org/10.1103/PhysRevLett.43.668).
- [11] S. G. Gorishnii, A. L. Kataev, and S. A. Larin. “The $O(\alpha_s^3)$ -corrections to $\sigma_{\text{tot}}(e^+ e^- \rightarrow \text{hadrons})$ and $\Gamma(\tau^- \rightarrow \nu_\tau + \text{hadrons})$ in QCD”. In: *Phys. Lett.* B259 (1991), pp. 144–150. DOI: [10.1016/0370-2693\(91\)90149-K](https://doi.org/10.1016/0370-2693(91)90149-K).
- [12] Matthias Jamin. *QCD and Renormalisation Group Methods*. Lecture presented at Herbstschule für Hochenergiephysik Maria Laach. Sept. 2006.
- [13] Gunnar Kallen. “On the definition of the Renormalization Constants in Quantum Electrodynamics”. In: *Helv. Phys. Acta* 25 (417). [509(1952)]. DOI: [10.1007/978-3-319-00627-7_90](https://doi.org/10.1007/978-3-319-00627-7_90).
- [14] Alexander Keshavarzi, Daisuke Nomura, and Thomas Teubner. “ μ_{g-2} and $\alpha(M_Z^2)$: a new data-based analysis”. In: *Phys. Rev.* D97.11 (2018), p. 114025. DOI: [10.1103/PhysRevD.97.114025](https://doi.org/10.1103/PhysRevD.97.114025). arXiv: [1802.02995](https://arxiv.org/abs/1802.02995) [hep-ph].
- [15] L. V. Lanin, V. P. Spiridonov, and K. G. Chetyrkin. “Contribution of Four Quark Condensates to Sum Rules for ρ and A_1 Mesons. (In Russian)”. In: *Yad. Fiz.* 44 (1986), pp. 1372–1374.
- [16] H. Lehmann. “On the Properties of propagation functions and renormalization constants of quantized fields”. In: *Nuovo Cim.* 11 (1954), pp. 342–357. DOI: [10.1007/BF02783624](https://doi.org/10.1007/BF02783624).
- [17] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to quantum field theory*. Reading, USA: Addison-Wesley, 1995. URL: <http://www.slac.stanford.edu/~mpeskin/QFT.html>.
- [18] Antonio Pich and Antonio Rodríguez-Sánchez. “Determination of the QCD coupling from ALEPH τ decay data”. In: *Phys. Rev. D* 94 (3 Aug. 2016), p. 034027. DOI: [10.1103/PhysRevD.94.034027](https://doi.org/10.1103/PhysRevD.94.034027). URL: <https://link.aps.org/doi/10.1103/PhysRevD.94.034027>.
- [19] Felix Schwab. “Strange Quark Mass Determination From Sum Rules For Hadronic τ -Decays”. German. MA thesis. somewhere, 2002.
- [20] Mikhail A. Shifman, A. I. Vainshtein, and Valentin I. Zakharov. “QCD and Resonance Physics. Theoretical Foundations”. In: *Nucl. Phys.* B147 (1979), pp. 385–447. DOI: [10.1016/0550-3213\(79\)90022-1](https://doi.org/10.1016/0550-3213(79)90022-1).
- [21] Levan R. Surguladze and Mark A. Samuel. “Total hadronic cross-section in $e^+ e^-$ annihilation at the four loop level of perturbative QCD”. In: *Phys. Rev. Lett.* 66 (1991). [Erratum: *Phys. Rev. Lett.* 66,2416(1991)], pp. 560–563.
- [22] M. Tanabashi et al. “Review of Particle Physics”. In: *Phys. Rev.* D98.3 (2018), p. 030001. DOI: [10.1103/PhysRevD.98.030001](https://doi.org/10.1103/PhysRevD.98.030001).

- [23] Yung-Su Tsai. “Decay Correlations of Heavy Leptons in $e^+ + e^- \rightarrow l^+ + l^-$ ”. In: *Phys. Rev. D* 4 (9 Nov. 1971), pp. 2821–2837. DOI: [10 . 1103/PhysRevD.4.2821](https://link.aps.org/doi/10.1103/PhysRevD.4.2821). URL: <https://link.aps.org/doi/10.1103/PhysRevD.4.2821>.