

Chapter 1

τ decays into hadrons

$$R_\tau = \frac{\Gamma(\tau \rightarrow \nu_\tau + \text{Hadrons})}{\Gamma(\tau \rightarrow \nu_\tau e^+ e^-)} \quad (1.1)$$

in terms of V/A, S/P [Broadhurst1975]

$$\begin{aligned} \Pi^{\mu\nu}(q^2) &= (q^\mu q^\nu - q^2 g^{\mu\nu}) \Pi^{V,A}(q^2) + \frac{g^{\mu\nu}}{q^2} (m_i \mp m_j) \Pi^{S,P}(q^2) \\ &+ g^{\mu\nu} \frac{(m_i \mp m_j)}{q^2} [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle] \end{aligned} \quad (1.2)$$

$$q_\mu q_\nu \Pi^{\mu\nu}(q^2) = (m_i \mp m_j)^2 \Pi^{S,P}(q^2) + (m_i \mp m_j) [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle] \quad (1.3)$$

in terms of T and L

$$\Pi^{\mu\nu}(q^2) = (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T)}(q^2) + q^\mu q^\nu \Pi^{(L)}(q^2) \quad (1.4)$$

$$q_\mu q_\nu \Pi^{\mu\nu}(q^2) = q^4 \Pi^{(L)}(q^2) = s^2 \Pi^{(L)}(s), \quad (1.5)$$

where $s \equiv q^2$

relation L and S,P

$$s^2 \Pi^{(L)}(s) = (m_i \mp m_j)^2 \Pi^{(S,P)}(s) + (m_i \mp m_j) [\langle \bar{q}_i q_i \rangle \mp \langle \bar{q}_j q_j \rangle] \quad (1.6)$$

need relation T and V,A

$$\begin{aligned} \Pi^{\mu\nu}(s) &= \underbrace{(q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T)}(s) + (q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(L)}(s)}_{=(q^\mu q^\nu - g^{\mu\nu} q^2) \Pi^{(T+L)}(s)} + \frac{g^{\mu\nu} s^2}{q^2} \Pi^{(L)}(s) \end{aligned} \quad (1.7)$$

where $\Pi^{(T+L)}(s) \equiv \Pi^{(T)}(s) + \Pi^{(L)}(s)$

$$\Pi^{(V,A)}(s) = \Pi^{(T)}(s) + \Pi^{(L)}(s) = \Pi^{(T+L)}(s) \quad (1.8)$$

$$q_\mu q_n u \quad (1.9)$$

The theoretical expression of the hadronic τ -decay ratio was first derived by [Tsai1971] (using current algebra, a more recent derivation making use of the *optical theorem* can be taken from [Schwab2002]):

$$R_\tau = 12\pi \int_0^{m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right) \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im } \Pi^{(T)}(s) + \text{Im } \Pi^{(L)} \right]. \quad (1.10)$$

R_τ introduces a problematic integral over the real axis of $\Pi(s)$ from 0 up to m_τ . The integral is problematic for two reasons:

- The *perturbative Quantum Chromodynamics* (**pQCD**) and the OPE breaks down for low energies (over which we have to integrate).
- The positive euclidean axis of $\Pi(s)$ has a discontinuity cut and can theoretically not be evaluated.

To literally circumvent these issues we make use of *Cauchy's Theorem*

$$\int_{\mathcal{C}} f(z) dz = 0, \quad (1.11)$$

where $f(z)$ is an analytic function on a closed contour \mathcal{C} .

In our case we have to deal with the two-point correlator $\Pi(s)$, which is analytic except for the positive real axis (with which we will deal with to a later point¹) Consequently, to rewrite we can rewrite the definite integral of eq. (1.10) into a contour integral over a closed circle with radius m_τ^2 . The closed contour consists of four line integrals, which have been visualized in fig. 1.1. Summing over the four line integrals, performing a *analytic continuation* of the two-point correlator $\Pi(s) \rightarrow \Pi(s + i\epsilon)$ and finally taking the limit of $\epsilon \rightarrow 0$ gives us the needed relation between eq. (1.10) and the closed contour:

$$\begin{aligned} \oint_{s=m_\tau} \Pi(s) &= \int_0^{m_\tau} \Pi(s + i\epsilon) ds + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{m_\tau}^0 \Pi(s - i\epsilon) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\ &= \int_0^{m_\tau} \Pi(s + i\epsilon) - \Pi(s - i\epsilon) ds + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\ &= \int_0^{m_\tau} \Pi(s + i\epsilon) - \overline{\Pi(s + i\epsilon)} + \int_{\mathcal{C}_2} \Pi(s) ds + \int_{\mathcal{C}_4} \Pi(s) ds \\ &\stackrel{\lim \epsilon \rightarrow 0}{=} 2i \int_0^{m_\tau} \text{Im } \Pi(s) ds + \oint_{s=m_\tau} \Pi(s) ds \end{aligned} \quad (1.12)$$

where we made use of $\Pi(z) = \overline{\Pi(\bar{z})}$ (due to $\Pi(s)$ is analytic) and $\Pi(z) - \overline{\Pi(\bar{z})} = 2i \text{Im } \Pi(z)$. The result can be rewritten in a more intuitive form, which we also

¹To not evaluate $\Pi(s)$ at the positive real axis we have to introduce *pinched weights*. The *pinched weights* vanish for $s \rightarrow m_\tau$.




Figure 1.1: Visualization of the usage of Cauchy's theorem to transform eq. (1.10) into a closed contour integral over a circle of radius m_τ .

visualized in fig. 1.1

$$\int_0^{m_\tau} \Pi(s) ds = \frac{i}{2} \oint_{|s|=m_\tau} \Pi(s) ds \quad (1.13)$$

Finally combining eq. (1.13) with eq. (1.10) we get

$$R_\tau = 6\pi i \oint_{|s|=m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right) \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(T)}(s) + \Pi^{(L)} \right] \quad (1.14)$$

for the hadronic τ -decay ratio.

The contour integral obtained is an import result as we can now theoretically evaluate the hadronic τ -decay ratio sufficiently large energy scales ($m_\tau \approx 1.78 \text{ MeV}$) at which $\alpha_s(m_\tau) \approx 0.33$ [Pich2016] is tolerable heigh for applying perturbation theory and the OPE. Obviously we would benefit from a contour integral over a bigger circumference, but τ -decays are limited by the m_τ . Nevertheless there are promising e^+e^- annihilation data, which yields valuable R-ratio values up to 2 GeV [Boito2018][Keshavarzi2018].

It is convenient to rewrite the

$$\Pi^{(L+T)} = \Pi^{(L)} + \Pi^{(T)} \quad (1.15)$$

$$R_\tau = 6\pi i \oint_{|s|=m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(L+T)}(s) - \left(\frac{2s}{m_\tau^2}\right) \Pi^{(L)}(s) \right] \quad (1.16)$$

$$D^{(L+T)}(s) \equiv -s \frac{d}{ds} \Pi^{(L+T)}(s), \quad D^{(L)}(s) \equiv \frac{s}{m_\tau^2} \frac{d}{ds} (s \Pi^{(L)}(s)) \quad (1.17)$$

Integration by parts

$$\int_a^b u(x)V(x) dx = [u(x)V(x)]_a^b - \int_a^b u(x)v(x) dx \quad (1.18)$$

$$\begin{aligned} R_\tau^{(1)} &= \frac{6\pi i}{m_\tau^2} \oint_{|s|=m_\tau^2} \underbrace{\left(1 - \frac{s}{m_\tau^2}\right)^2}_{=u(x)} \underbrace{\left(1 + 2\frac{s}{m_\tau^2}\right)}_{=V(x)} \Pi^{(L+T)}(s) \\ &= \frac{6\pi i}{m_\tau^2} \left\{ \left[-\frac{m_\tau^2}{2} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left(1 + \frac{s}{m_\tau^2}\right) \Pi^{(L+T)}(s) \right]_{|s|=m_\tau^2} \right. \\ &\quad \left. + \oint_{|s|=m_\tau^2} \underbrace{-\frac{m_\tau^2}{2} \left(1 - \frac{s}{m_\tau^2}\right)^3}_{=U(x)} \underbrace{\left(1 + \frac{s}{m_\tau^2}\right) \frac{d}{ds} \Pi^{(L+T)}(s)}_{=v(x)} \right\} \\ &= -3\pi i \oint_{|s|=m_\tau^2} \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left(1 + \frac{s}{m_\tau^2}\right) \frac{d}{ds} D^{(L+T)} \end{aligned} \quad (1.19)$$

where we fixed the integration constant to $C = -\frac{m_\tau^2}{2}$ in the second line and left the antiderivatives contained in the squared brackets untouched. Parametrizing the expression in the squared brackets

$$\left[-\frac{m_\tau^2}{2} (1 - e^{-i\phi})^3 (1 + e^{-i\phi}) \Pi^{(L+T)}(m_\tau^2 e^{-i\phi}) \right]_0^{2\pi} = 0 \quad (1.20)$$

where $s \rightarrow m_\tau^2 e^{-i\phi}$ and $(1 - e^{-i \cdot 0}) = (1 - e^{-i \cdot 2\pi}) = 0$.

$$\begin{aligned} R_\tau^{(2)} &= \oint_{|s|=m_\tau^2} ds \left(1 - \frac{s}{m_\tau^2}\right)^2 \left(-\frac{2s}{m_\tau^2}\right) \Pi^{(L)}(s) \\ &= -4\pi i \oint \frac{ds}{s} \left(1 - \frac{s}{m_\tau^2}\right)^3 D^{(L)}(s) \end{aligned} \quad (1.21)$$

$$R_\tau = -\pi i \oint_{|s|=m_\tau^2} \frac{d}{s} \left(1 - \frac{s}{m_\tau^2}\right)^3 \left[3 \left(1 + \frac{s}{m_\tau^2}\right) D^{(L+T)}(s) + 4 D^{(L)}(s) \right] \quad (1.22)$$

$$R_\tau = -\pi i \oint_{|s|=m_\tau^2} \frac{d}{x} (1-x)^3 \left(3(1+x) D^{(L+T)}(s) + 4 D^{(L)}(s) \right), \quad (1.23)$$

where $x = s/m_\tau^2$.

1.1 The perturbative expansion

We will treat the correlator in the chiral limit for which the longitudinal components $\Pi^L(s)$ vanish (see eq. (1.7)) and the axial and vectorial contributions are equal. Consequently [Beneke2008] we can write the vector correlation function $\Pi(s)$ can be written as follows:

$$\Pi_V^{T+L} = -\frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=1}^{n+1} k c_{n,k} L^{k-1} \quad \text{with} \quad L \equiv \ln \frac{-s}{\mu^2}. \quad (1.24)$$

$\Pi_V^{T+L}(s)$ is not a physical quantity but we can use eq. (1.17) to rewrite it into one:

$$D_V^{T+L} = -s \frac{d\Pi_V^{1+0}(s)}{ds} = \frac{N_c}{12\pi^2} \sum_{n=0}^{\infty} a_\mu^n \sum_{k=1}^{n+1} k c_{n,k} L^{k-1}, \quad (1.25)$$

where we used $dL^k/ds = k \ln(-s/\mu^2)^{k-1} (-1/\mu^2)$. D_V^{1+0} being a physical quantity has to fulfill the RGE ??

$$-\mu \frac{d}{d\mu} D_V^{T+L} = -\mu \frac{d}{d\mu} \left(\frac{\partial}{\partial L} D_V^{T+L} + \frac{\partial}{\partial a_s} D_V^{T+L} \right) = \left(2 \frac{\partial}{\partial L} + \beta \frac{\partial}{\partial a_s} \right) D_V^{T+L} = 0, \quad (1.26)$$

where we defined the β -function in ?? and used $dL/d\mu = -2/\mu$. The RGE puts constraints on the $c_{n,k}$ -coefficients, ... not independent

$$D(s) = \frac{N_c}{12\pi^2} \left[c_{01} + a_\mu (c_{11} + 2c_{12}L) + a_\mu^2 (c_{21} + 2c_{22}L + 3c_{23}L^2) \right] \quad (1.27)$$

inserting into RGE

$$4a_\mu c_{12} + 2a_\mu^2 (2c_{22} + 6c_{23}L) + \beta_1 a_\mu^2 (c_{11} + 2c_{12}L) + \mathcal{O}(a_\mu^3) = 0 \quad (1.28)$$

Thus

$$c_{12} = 0 \quad c_{22} = \frac{\beta_1 c_{11}}{4} \quad c_{23} = 0 \quad (1.29)$$

or $D(s)$ to the first order in α_s

$$D(s) = \frac{N_c}{12\pi^2} \left[c_{01} + c_{11} a_\mu \left(c_{21} - \frac{1}{2} \beta_1 c_{11} L \right) a_\mu^2 \right] + \mathcal{O}(a_\mu^3) \quad (1.30)$$