Challenge Stigma / Find Peace

FrontlineFolks | Guide #32

Porn, Solo Sex, and Secret Coping

A Field Guide for Frontline Folks and Their Families

Field notes for secret coping and compulsive sexual habits.

By Michael Jenkins, MS, MSW, LCSW
Never Forgotten Services, PLLC

Disclaimer

This guide is for general education only.

It is not:

- Therapy
- Medical care
- A crisis service
- Individualized advice

Reading this guide does not create a therapist–client relationship with the author or with Never Forgotten Services, PLLC.

If you are in crisis or thinking about harming yourself or someone else, call 988 in the United States, go to your nearest emergency room, or contact your local emergency services.

If you want mental health treatment, please contact a licensed provider in your state or call the number on the back of your insurance card.

Table of Contents

1. What Is Happening
2. Why It Escalates
3. What To Do This Week
4. For Partners and Team
5. When To Get More Help
6. 7-Day Field Drill
7. Worksheets
    - Worksheet 1: Worksheet: Porn, Solo Sex, and Secret Coping
    - Worksheet 2: Worksheet: Porn, Solo Sex, and Secret Coping - Partner/Team Check-In

1. What Is Happening

Porn, masturbation, and sexual fantasy are tools you're using to regulate. They work fast: activation goes up, you dissociate into solo sex, dopamine floods, and for a moment you feel numb to the pain. The problem isn't masturbation (that's normal)—it's using it as your primary regulation tool. You're doing it multiple times daily, you're sneaking around about it, you're using more intense content to get the same effect, and your partner doesn't know the extent. This is sexual compulsion as a coping mechanism: using sexual stimulation to numb dysregulation, loneliness, or pain.

Signs sexual compulsion is becoming a problem

- Using porn/masturbation as your primary regulation tool: When you feel bad, you go straight to sexual content.
- Frequency escalating: What used to be weekly is now daily or multiple times daily.
- Tolerance building: You need more intense content to get the same effect.
- Time loss: You're spending 1-3 hours daily in sexual content/activity, losing time with partner or work.
- Shame and secrecy: You're hiding it from your partner, clearing browser history, using private browsers.
- Conflict with partner: Partner has found evidence or confronted you, creating rupture and blame.
- Impact on partner sex: You're less interested in partnered sex because solo sex is less demanding.
- Unable to reduce despite trying: You've promised to stop/reduce and can't maintain it.

> **This is not moral failure:**
> Compulsive sexual behavior is a self-regulation strategy for dysregulation, trauma, or loneliness. It's a symptom, not a character flaw.

2. Why It Escalates

Frontline work leaves you activated, numb, lonely. Solo sex is a regulation tool that works immediately: you're alone (safe), in control (not vulnerable), and you get dopamine without emotional risk. Your partner wants connection, but connection requires vulnerability and energy you don't have. Solo sex is easier. Add shame about it, and you hide it. The secrecy creates rupture, which creates more loneliness, which drives more compulsive use.

How compulsive sexual behavior escalates

- Using sex to regulate dysregulation: Bad shift? Solo sex. Argument? Solo sex. Lonely? Solo sex compounds the loneliness.
- Tolerance escalation: Vanilla content stops working, so you seek more intense, more niche, more extreme content.
- Preference shift: Increasingly, your arousal is tied to specific, intense content rather than your partner.
- Secrecy creating shame: You hide it, feel guilty, then use it again to numb the guilt.
- Partner rupture: Partner finds out, feels betrayed/rejected, conflict escalates, you use more to escape the conflict.
- Isolation deepening: Using more solo sex means less partnered sex, less vulnerability, less connection.
- Compulsive cycle: You can't control it despite wanting to, which creates more shame and more compulsion.

> *Why this works: reducing total load and increasing predictability lowers baseline arousal and improves regulation.*

3. What To Do This Week

You can reduce compulsive sexual behavior by finding alternative regulation tools and rebuilding intimacy with your partner.

1. Identify what you're avoiding with sexual compulsion: Loneliness? Dysregulation? Intimacy? Name it so you can address it directly.
2. Build a regulation toolkit outside of sex: Movement, breathing, time in nature, music—go-to tools that don't involve sexual content.
3. Increase partner time without sexual expectation: Talking, touching, being together without sex looming. Rebuild non-sexual intimacy.
4. Set a realistic limit, not zero: 'I'm reducing to X times per week' is more sustainable than 'I'll never do it again.'
5. Remove access when activated: Delete apps, use parental controls, do anything that puts a pause between impulse and action.
6. Rebuild transparency: Tell partner what you're working on. Regular check-ins about what's helping. Rebuild trust through honesty.
7. Address the root: If it's loneliness, build connection. If it's dysregulation, build regulation tools. If it's trauma, get trauma treatment.

8. Consider accountability support: 12-step groups, therapist specializing in sex addiction, structured treatment if it's severe.

> Field rule: set the bar low enough to do it even on your worst day.

4. For Partners and Team

Stress communication should be clear, brief, and repeatable. Predictable routines protect relationships.

- Use concrete language: 'I've been using porn/masturbation to regulate. I'm not doing it to hurt you or avoid sex with you. It's about my coping.'
- Agree on transparency: No more hiding. Regular check-ins. Partner has access to what you're working on.
- Debrief without blame: 'This is what I'm working on. This is what I need from you.' Not 'You're never interested in sex so I...'
- Consent and pace: Rebuilding partnered sex takes time. Don't expect immediate change.

---

5. When To Get More Help

Self-guided tools are a start. Escalate care when symptoms persist, spread, or begin harming safety and function.

- Compulsive use is continuing despite attempts to reduce.
- It's significantly impacting your relationship or work function.
- You've escalated to content that worries or shames you.
- Your partner has threatened to leave over the issue.
- Sex therapy or specialized addiction treatment can help identify roots and rebuild healthy sexual expression.

6. 7-Day Field Drill

Use the worksheet every day for one week. Track trigger, body signal, and reset action. Keep what works and remove what does not.

> **Daily drill:**
> 1) Name the trigger.
> 2) Name the first body signal.
> 3) Run one reset action.
> 4) Log response quality.

Worksheet 1: Worksheet: Porn, Solo Sex, and Secret Coping

Use this to track secret coping and compulsive sexual habits and build a practical response you can repeat under load.

Where did this pattern show up this week?

List specific moments at work, at home, and in transition.

What did it cost you when it took over?

Name impact on body, mood, decisions, and relationships.

What two actions will you run for the next 7 days?

Choose actions small enough to use on hard days.

1. Trigger:

What activated the pattern?

2. Early body signal:

What did your body do first?

3. Reset action used:

What fast action lowered intensity?

4. Better next response:

What exact behavior will you use next time?

> Track this daily for one week, then tighten the plan around what actually worked.

Worksheet 2: Worksheet: Porn, Solo Sex, and Secret Coping - Partner/Team Check-In

Use this page for one short practical check-in each week.

What do others notice first when this pattern is active?

Capture observable behavior, not labels.

What support helps most and least?

List what helps and what escalates the issue.

What agreement will you both practice this week?

One clear action from each side.

1. Signal phrase:

What phrase means pause and reset?

2. Time-to-return commitment:

When will re-engagement happen?

3. Repair step:

What specific repair happens after a rupture?

> Keep check-ins short, specific, and repeatable.

Closing

You've seen and carried more than most people will ever know.

Blow ups and shutdowns are not proof that you're broken or that your relationship is doomed.

They are signs that your nervous system has been running in survival mode for a long time.

The fact that you picked up this guide and read it matters.

You are choosing to be intentional about how you show up for the people who matter to you.

That choice is a lot more powerful than having flawless days.

About the Author

Michael Jenkins, MS, MSW, LCSW

Founder, Never Forgotten Services, PLLC

Michael is a U.S. Army combat veteran and licensed clinical social worker who specializes in working with military, veterans, and first responders—our frontline folks. After leaving the military, he spent years fighting his own demons and trying to figure out why he could hold it together on the job but feel completely wrecked at home. The turning point was realizing he wasn't the only one living that split life—and that nobody should have to feel that alone, confused, or broken by what the job asks of them.

Since then, his work has focused on trauma, the impact of high-stress jobs on home life, and helping families make sense of what the job brings home. He uses plain language, zero shame, and a "you're not crazy, this actually makes sense" approach to help frontline folks and their families understand what's happening in their nervous systems and learn how to move through the chaos together.