Challenge Stigma / Find Peace

FrontlineFolks | Guide #29

Low Desire and Low Drive

A Field Guide for Frontline Folks and Their Families

Libido loss is usually hormonal and neurological, not relationship failure.

By Michael Jenkins, MS, MSW, LCSW
Never Forgotten Services, PLLC

Disclaimer

This guide is for general education only.

It is not:

- Therapy
- Medical care
- A crisis service
- Individualized advice

Reading this guide does not create a therapist–client relationship with the author or with Never Forgotten Services, PLLC.

If you are in crisis or thinking about harming yourself or someone else, call 988 in the United States, go to your nearest emergency room, or contact your local emergency services.

If you want mental health treatment, please contact a licensed provider in your state or call the number on the back of your insurance card.

Table of Contents

1. What Is Happening
2. Why It Escalates
3. What To Do This Week
4. For Partners and Team
5. When To Get More Help
6. 7-Day Field Drill
7. Worksheets
    - Worksheet 1: Worksheet: Low Desire and Low Drive
    - Worksheet 2: Worksheet: Low Desire and Low Drive - Partner/Team Check-In

1. What Is Happening

Your libido is gone. You used to want sex, now you don't think about it. Your partner tries to initiate and you feel a mix of obligation, guilt, and lack of interest. This isn't about not loving your partner—it's about a neurobiological problem. Chronic stress, sleep debt, depression, and trauma all suppress sexual desire. High cortisol suppresses testosterone. Hypervigilance makes relaxation (required for arousal) impossible. Anxiety and hyperarousal activate the sympathetic nervous system, which blocks parasympathetic arousal.

Common signs of low desire and low drive

- No spontaneous sexual thoughts or desire: You don't think about it, and partner initiating doesn't spark anything.
- Going along without desire: You have sex because partner wants it, not because you do.
- Difficulty with foreplay: The stuff that used to turn you on now feels boring or uncomfortable.
- Long refractory period: Even if you do have sex, it takes forever to recover interest.
- Loss of morning erections or baseline genital engagement: You're not waking up aroused anymore.
- No fantasy life: You can't access sexual thoughts even in private.
- Attraction unaffected: You can find your partner attractive, but desire doesn't follow.
- Fatigue as the driver: You're too tired, too stressed, too depleted to want anything, including sex.

> **This is not broken sexuality:**
> This is a dysregulated nervous system. When the nervous system is in survival mode, reproduction is off the priority list.

2. Why It Escalates

Frontline work keeps cortisol elevated, which suppresses testosterone. Sleep deprivation (from shift work and hyperarousal) tanks libido. Depression and trauma are strongly linked to anhedonia (loss of pleasure, including sexual pleasure). Chronic activation of the sympathetic nervous system (fight-flight) blocks the parasympathetic activation needed for arousal. Your body is essentially saying 'we're not safe to relax and have sex.'

How desire disappears

- Elevated cortisol: Chronically high cortisol suppresses testosterone production and sexual interest.
- Poor sleep: Sleep debt directly lowers libido and makes arousal slower and harder.
- Anhedonia: Loss of pleasure response means sexual pleasure doesn't register like it used to.
- Performance pressure: Partner's frustration creates pressure, which further suppresses arousal.
- Hypervigilance: Brain scanning for threats can't be in relaxation mode simultaneously.
- Numbing protection: Emotional numbness extends to sexual desire.
- Avoidance cycle: You avoid sex → partner feels rejected → pressure increases → avoidance deepens.

3. What To Do This Week

You can rebuild desire by addressing the nervous system dysregulation first. Desire often returns when arousal capacity returns.

1. Get sleep: Prioritize sleep recovery above almost everything else. Low libido without sleep is mechanical—it's not a relationship issue.
2. Reduce stress and activation: Movement, breathing work, time outdoors all lower cortisol and create space for desire.
3. Address depression or anhedonia: Often desire returns when depression treatment starts working.
4. Rebuild parasympathetic activation: Activities that activate rest-and-digest (massage, slow breathing, warm baths, yoga) help.
5. Take sex off the agenda: Stop trying to have sex. This relieves performance pressure and creates space for desire to emerge naturally.
6. Build small moments of physical affection without expectations: Kissing, touching, being close—without it leading to sex.
7. Get hormone levels checked: If cortisol is very high or testosterone is very low, medical intervention can help.
8. Check medications: Some antidepressants and anxiety meds suppress libido; talking to your prescriber about alternatives might help.

> Field rule: set the bar low enough to do it even on your worst day.

4. For Partners and Team

Stress communication should be clear, brief, and repeatable. Predictable routines protect relationships.

- Use concrete language: 'My libido is low from stress and sleep debt, not because I'm not attracted to you.'
- Agree on a timeline: Libido recovery is slow. Ask for patience while you rebuild.
- Debrief without pressure: 'I'm working on this. I don't have a timeline.' No promises of when it will return.
- Consent, pacing, and emotional safety are part of intimacy, not separate from it: No coercion or guilt. No 'you should be able to by now.'

---

5. When To Get More Help

Self-guided tools are a start. Escalate care when symptoms persist, spread, or begin harming safety and function.

- Desire has been absent for 6+ months despite stress reduction and sleep recovery.
- Relationship conflict is centered on sex, and it's deteriorating.
- You're using substances or compulsive behaviors to stimulate desire (alcohol, porn, risky behavior).
- Medical causes have been ruled out, but desire still isn't returning.
- Sex therapy or trauma-informed couples care can identify blocks and rebuild desire gradually.

6. 7-Day Field Drill

Use the worksheet every day for one week. Track trigger, body signal, and reset action. Keep what works and remove what does not.

> **Daily drill:**
> 1) Name the trigger.
> 2) Name the first body signal.
> 3) Run one reset action.
> 4) Log response quality.

Worksheet 1: Worksheet: Low Desire and Low Drive

Use this to track reduced desire and low drive and build a practical response you can repeat under load.

Where did this pattern show up this week?

List specific moments at work, at home, and in transition.

What did it cost you when it took over?

Name impact on body, mood, decisions, and relationships.

What two actions will you run for the next 7 days?

Choose actions small enough to use on hard days.

1. Trigger:

What activated the pattern?

2. Early body signal:

What did your body do first?

3. Reset action used:

What fast action lowered intensity?

4. Better next response:

What exact behavior will you use next time?

> Track this daily for one week, then tighten the plan around what actually worked.

Worksheet 2: Worksheet: Low Desire and Low Drive - Partner/Team Check-In

Use this page for one short practical check-in each week.

What do others notice first when this pattern is active?

Capture observable behavior, not labels.

What support helps most and least?

List what helps and what escalates the issue.

What agreement will you both practice this week?

One clear action from each side.

1. Signal phrase:

What phrase means pause and reset?

2. Time-to-return commitment:

When will re-engagement happen?

3. Repair step:

What specific repair happens after a rupture?

> Keep check-ins short, specific, and repeatable.

Closing

You've seen and carried more than most people will ever know.

Blow ups and shutdowns are not proof that you're broken or that your relationship is doomed.

They are signs that your nervous system has been running in survival mode for a long time.

The fact that you picked up this guide and read it matters.

You are choosing to be intentional about how you show up for the people who matter to you.

That choice is a lot more powerful than having flawless days.

About the Author

Michael Jenkins, MS, MSW, LCSW

Founder, Never Forgotten Services, PLLC

Michael is a U.S. Army combat veteran and licensed clinical social worker who specializes in working with military, veterans, and first responders—our frontline folks. After leaving the military, he spent years fighting his own demons and trying to figure out why he could hold it together on the job but feel completely wrecked at home. The turning point was realizing he wasn't the only one living that split life—and that nobody should have to feel that alone, confused, or broken by what the job asks of them.

Since then, his work has focused on trauma, the impact of high-stress jobs on home life, and helping families make sense of what the job brings home. He uses plain language, zero shame, and a "you're not crazy, this actually makes sense" approach to help frontline folks and their families understand what's happening in their nervous systems and learn how to move through the chaos together.