Challenge Stigma / Find Peace

FrontlineFolks | Guide #33

Touch Without Pressure: Relearning Safe Closeness

A Field Guide for Frontline Folks and Their Families

Field notes for the pattern in this chapter.

By Michael Jenkins, MS, MSW, LCSW
Never Forgotten Services, PLLC

Disclaimer

This guide is for general education only.

It is not:

- Therapy
- Medical care
- A crisis service
- Individualized advice

Reading this guide does not create a therapist–client relationship with the author or with Never Forgotten Services, PLLC.

If you are in crisis or thinking about harming yourself or someone else, call 988 in the United States, go to your nearest emergency room, or contact your local emergency services.

If you want mental health treatment, please contact a licensed provider in your state or call the number on the back of your insurance card.

Table of Contents

1. What Is Happening
2. Why It Escalates
3. What To Do This Week
4. For Partners and Team
5. When To Get More Help
6. 7-Day Field Drill
7. Worksheets
    - Worksheet 1: Worksheet: Touch Without Pressure: Relearning Safe Closeness
    - Worksheet 2: Worksheet: Touch Without Pressure: Relearning Safe Closeness - Partner/Team Check-

1. What Is Happening

Your partner tries to touch you and your nervous system sends a threat signal. Maybe you flinch. Maybe you go rigid. Maybe you snap at them. Maybe you become numb and can't feel the touch. This is hypervigilance extending into your body: your nervous system is scanning everything (including safe touch) for danger. Safe, loving touch should feel comforting. Instead, it feels intrusive. This creates a painful paradox: you need physical closeness to feel connected, but your body won't allow it.

Common touch sensitivity patterns

- Flinching to touch: Even gentle, expected touch causes you to startle.
- Stiffening when touched: Your body goes rigid, partner feels the resistance, they withdraw.
- Irritability to touch: Partner touches you affectionately and you snap at them.
- Needing to control touch: You can touch them, but them touching you feels threatening.
- Numbness to touch: They're touching you but you can't feel it, or it feels like they're touching someone else.
- Certain areas untouchable: Specific body parts are off-limits because they trigger or feel violated.
- Hyperawareness of touch: You're hyper-attuned to exactly how they're touching, waiting for threat.
- Dissociation during touch: You check out, feel separated from your body, can't be present.

> **This is not rejection of your partner:**
> This is your nervous system in protection mode. Your body has learned that touch equals vulnerability, which equals danger. That's not about them, it's about your threat detection system.

2. Why It Escalates

Hypervigilance means you're scanning everything for threats. Touch includes vulnerability: letting someone close, relinquishing control, allowing someone to affect your body. Your nervous system says 'that's dangerous.' If you have trauma history (especially touch-based trauma), the association is even stronger. Your body has learned 'touch = violation', so safe touch gets treated like dangerous touch.

How touch sensitivity escalates

- Partner confusion: They don't understand why you flinch, they feel rejected, they stop trying.
- Isolation deepening: Less touch means less physical connection, which means less bonding.
- Your shame mounting: You feel broken because you can't accept touch, you snap at partner for trying.
- Partner resentment: They've been rejected so many times they're considering leaving.
- Hypervigilance intensifying: The less touch you receive, the more your nervous system assumes touch is dangerous.
- Intimacy erosion: Without physical affection, emotional intimacy deteriorates too.

> *Why this works: reducing total load and increasing predictability lowers baseline arousal and improves regulation.*

3. What To Do This Week

You can relearn safe touch, but it requires slow, gradual, consensual rebuilding. Your nervous system needs to learn 'this touch is safe' through repeated safe experiences.

1. Start with self-touch: Massage your own hands, arms, face. Practice receiving sensation from your own body in a safe way.
2. Agree on one safe touch: You and partner identify one way of being touched that feels (even slightly) safe. Start there only.
3. Practice micro-touches: 30 seconds of hand-holding, brief hug, short back rub. Short duration = lower threat.
4. Use grounding during touch: Name what you're sensing ('I feel their hand, it's warm, it's steady') so you stay present.
5. Permission to pause: If touch becomes uncomfortable, you can say 'pause' and stop—no explanation, no guilt. Partner honors it immediately.
6. Rebuild from non-sexual touch: Hugging, hand-holding, massaging get reestablished before any sexual touch.
7. Communicate need explicitly: 'I need slow touch' or 'I need you to ask before touching me' or 'I need to be the one to initiate.'
8. Use trauma treatment to lower baseline threat: EMDR, somatic therapy, or trauma work specifically helps if touch-based trauma is the root.
9. Partner participation in rebuilding: They need to be patient, predictable, and honoring your boundaries.

> Field rule: set the bar low enough to do it even on your worst day.

4. For Partners and Team

Stress communication should be clear, brief, and repeatable. Predictable routines protect relationships.

- Use concrete language: 'I'm hypervigilant to touch right now. This isn't about you or your love.'
- Agree on complete permission to pause: Touch can stop any time, no questions asked, no guilt.
- Debrief without blame: 'I flinched because my nervous system is protecting me, not because I don't love you.'
- Consent, pacing, and emotional safety are part of intimacy: Slow rebuilding is the only way this works.

---

5. When To Get More Help

Self-guided tools are a start. Escalate care when symptoms persist, spread, or begin harming safety and function.

- Touch sensitivity is preventing any physical affection, and it's been ongoing for 6+ months.
- Partner has withdrawn from all touch attempts and the relationship is suffering.
- Touch sensitivity is spreading to other contexts (avoiding handshakes, hugs with friends, etc).
- You can't identify any safe touch—everything feels threatening.
- Trauma-informed therapy (somatic work, EMDR, or body-based modalities) can help rebuild safety in your body and with touch.

6. 7-Day Field Drill

Use the worksheet every day for one week. Track trigger, body signal, and reset action. Keep what works and remove what does not.

> **Daily drill:**
> 1) Name the trigger.
> 2) Name the first body signal.
> 3) Run one reset action.
> 4) Log response quality.

Worksheet 1: Worksheet: Touch Without Pressure: Relearning Safe Closeness

Use this to track the pattern in this chapter and build a practical response you can repeat under load.

Where did this pattern show up this week?

List specific moments at work, at home, and in transition.

What did it cost you when it took over?

Name impact on body, mood, decisions, and relationships.

What two actions will you run for the next 7 days?

Choose actions small enough to use on hard days.

1. Trigger:
What activated the pattern?

2. Early body signal:
What did your body do first?

3. Reset action used:
What fast action lowered intensity?

4. Better next response:
What exact behavior will you use next time?

> Track this daily for one week, then tighten the plan around what actually worked.

Worksheet 2: Worksheet: Touch Without Pressure: Relearning Safe Closeness - Partner/Team Check-In

Use this page for one short practical check-in each week.

What do others notice first when this pattern is active?

Capture observable behavior, not labels.

What support helps most and least?

List what helps and what escalates the issue.

What agreement will you both practice this week?

One clear action from each side.

1. Signal phrase:

What phrase means pause and reset?

2. Time-to-return commitment:

When will re-engagement happen?

3. Repair step:

What specific repair happens after a rupture?

> Keep check-ins short, specific, and repeatable.

Closing

You've seen and carried more than most people will ever know.

Blow ups and shutdowns are not proof that you're broken or that your relationship is doomed.

They are signs that your nervous system has been running in survival mode for a long time.

The fact that you picked up this guide and read it matters.

You are choosing to be intentional about how you show up for the people who matter to you.

That choice is a lot more powerful than having flawless days.

About the Author

Michael Jenkins, MS, MSW, LCSW

Founder, Never Forgotten Services, PLLC

Michael is a U.S. Army combat veteran and licensed clinical social worker who specializes in working with military, veterans, and first responders—our frontline folks. After leaving the military, he spent years fighting his own demons and trying to figure out why he could hold it together on the job but feel completely wrecked at home. The turning point was realizing he wasn't the only one living that split life—and that nobody should have to feel that alone, confused, or broken by what the job asks of them.

Since then, his work has focused on trauma, the impact of high-stress jobs on home life, and helping families make sense of what the job brings home. He uses plain language, zero shame, and a "you're not crazy, this actually makes sense" approach to help frontline folks and their families understand what's happening in their nervous systems and learn how to move through the chaos together.