Challenge Stigma / Find Peace

FrontlineFolks | Guide #34

Body Image, Scars, and Sexual Confidence

A Field Guide for Frontline Folks and Their Families

Field notes for scars, visible trauma, and reclaiming sexual confidence.

By Michael Jenkins, MS, MSW, LCSW
Never Forgotten Services, PLLC

Disclaimer

This guide is for general education only.

It is not:

- Therapy
- Medical care
- A crisis service
- Individualized advice

Reading this guide does not create a therapist–client relationship with the author or with Never Forgotten Services, PLLC.

If you are in crisis or thinking about harming yourself or someone else, call 988 in the United States, go to your nearest emergency room, or contact your local emergency services.

If you want mental health treatment, please contact a licensed provider in your state or call the number on the back of your insurance card.

Table of Contents

1. What Is Happening
2. Why It Escalates
3. What To Do This Week
4. For Partners and Team
5. When To Get More Help
6. 7-Day Field Drill
7. Worksheets
    - Worksheet 1: Worksheet: Body Image, Scars, and Sexual Confidence
    - Worksheet 2: Worksheet: Body Image, Scars, and Sexual Confidence - Partner/Team Check-In

1. What Is Happening

You have a visible marker of what happened. A scar from an injury, a tattoo you regret, burns, amputations, or maybe just a body that looks different than before. When you're with your partner and they see your body, you wonder what they think. You feel watched, ashamed, or like you have to explain. Maybe you've stopped taking off your shirt during sex. Maybe you're having less sex because you feel less attractive. This isn't about vanity—it's about how trauma gets written on your body, and how that changes how you feel in your own skin and with your partner.

Common signs

- Avoiding undressing: You keep clothes on during sex, avoid showers/intimate moments, or request darkness.
- Self-consciousness about scars: You cover them, won't look at them, or constantly apologize for them.
- Reduced sexual desire: You don't want sex because you feel undesirable or ashamed of your body.
- Difficulty accepting compliments: Partner tells you they find you attractive and you dismiss it or feel guilty.
- Over-focusing on the scar/injury: You can't be present during sex because you're hyper-aware of that specific spot.
- Avoidance of mirrors: You don't look at your body directly, or you only look at certain parts.
- Partner rejection fears: You assume your partner finds you less attractive, even if they haven't said so.
- Sex performance pressure: You feel like you have to compensate for your appearance by being extra attentive or skilled.

> **This is not vanity or weakness:**
> Your body is a map of what happened. Feeling self-conscious about visible trauma is a normal grief response, not a character flaw.

2. Why It Escalates

Your nervous system learned that your body isn't fully safe. The scar or injury is a reminder of violation, pain, or helplessness. Your brain associates that part of your body with danger. Add hypervigilance (you're watching your partner's face, your own reflection, scanning for rejection) and the vulnerability required for sex (being seen, touched, exposed), and you get avoidance. If the injury was inflicted by someone (assault, abuse), the shame is even deeper.

Common amplifiers

- Shame spiraling: One moment of self-consciousness becomes a spiral of 'I'm permanently broken' and 'They don't want me.'
- Avoidance escalating: You skip sex more. Partner feels rejected. You feel guilty. You avoid again.
- Connection eroding: Less sex means less physical touch, which means less bonding and more isolation.
- Comparison and rumination: You compare your body to before the injury, or to other people, reinforcing the 'I'm less' narrative.
- Partner withdrawal: Your partner senses your discomfort and they back off to avoid conflict, which you interpret as rejection.
- Performance anxiety: Even if your partner is accepting, you create pressure to compensate (extra enthusiasm, intensity, etc).
- Deeper shame: The longer you avoid, the more shame builds, the harder it is to be seen.

> *Why this works: reducing total load and increasing predictability lowers baseline arousal and improves regulation.*

3. What To Do This Week

You can rebuild body confidence and sexual pleasure, but it requires gradual exposure and partner collaboration.

1. Start by looking: Spend 5-10 minutes looking at your body directly (whole body, then scars/marks specifically) without judgment. Name what you see without shame words: 'I see a scar. It's part of my history.'
2. Touch your own scars: Gentle self-massage or simply touching the scarred area helps your nervous system learn 'this is mine, it's not dangerous.'
3. Tell your partner what you need: 'I need you to be patient while I rebuild confidence in my body.' Give them specific instructions: 'Please don't comment on the scar' OR 'Please tell me you're attracted to me' (whatever helps).

4. Gradual exposure: Start with lights dimmed, then partial nudity, then full nudity in light. Move at your pace.
5. Separate scar from sexy: Understand that your scars don't define your sexual value. You can be scarred AND desirable.
6. Practice compliment receiving: When partner says 'I find you attractive,' practice saying 'thank you' instead of arguing.
7. Change the sexual context: Maybe sex in dim light feels easier. Maybe specific positions feel less vulnerable. Build from what works.
8. Address trauma if needed: If the scar came from assault/abuse, trauma processing (EMDR, CPT) helps reduce the shame association.

> Field rule: set the bar low enough to do it even on your worst day.

4. For Partners and Team

Stress communication should be clear, brief, and repeatable. Predictable routines protect relationships.

- Use concrete language: 'I feel self-conscious about my scar. This isn't about you or your attraction. It's about my healing.'
- Give specific requests: 'I need the light off' or 'I need you to touch this scar gently' or 'I need you to tell me you find me sexy.'
- Debrief without defensiveness: 'I had a moment of shame during sex. It's not your fault. I'm working on it.'
- Acceptance without fixing: They don't need to convince you that you're attractive. They just need to honor your pace and support your healing.

---

5. When To Get More Help

Self-guided tools are a start. Escalate care when symptoms persist, spread, or begin harming safety and function.

- Body image shame is preventing sexual intimacy and it's been ongoing for 6+ months.
- You're avoiding medical appointments or showers because of shame about your body.
- Partner is feeling rejected and your relationship is suffering because of this avoidance.
- You're unable to look at scars/marks without severe emotional distress.
- Sex therapy or trauma-informed therapy (especially if the injury was interpersonal) can help rebuild body confidence and sexual pleasure.

6. 7-Day Field Drill

Use the worksheet every day for one week. Track trigger, body signal, and reset action. Keep what works and remove what does not.

> **Daily drill:**
> 1) Name the trigger.
> 2) Name the first body signal.
> 3) Run one reset action.
> 4) Log response quality.

Worksheet 1: Worksheet: Body Image, Scars, and Sexual Confidence

Use this to track the pattern in this chapter and build a practical response you can repeat under load.

Where did this pattern show up this week?

List specific moments at work, at home, and in transition.

What did it cost you when it took over?

Name impact on body, mood, decisions, and relationships.

What two actions will you run for the next 7 days?

Choose actions small enough to use on hard days.

1. Trigger:

What activated the pattern?

2. Early body signal:

What did your body do first?

3. Reset action used:

What fast action lowered intensity?

4. Better next response:

What exact behavior will you use next time?

> Track this daily for one week, then tighten the plan around what actually worked.

Worksheet 2: Worksheet: Body Image, Scars, and Sexual Confidence - Partner/Team Check-In

Use this page for one short practical check-in each week.

What do others notice first when this pattern is active?

Capture observable behavior, not labels.

What support helps most and least?

List what helps and what escalates the issue.

What agreement will you both practice this week?

One clear action from each side.

1. Signal phrase:

What phrase means pause and reset?

2. Time-to-return commitment:

When will re-engagement happen?

3. Repair step:

What specific repair happens after a rupture?

> Keep check-ins short, specific, and repeatable.

Closing

You've seen and carried more than most people will ever know.

Blow ups and shutdowns are not proof that you're broken or that your relationship is doomed.

They are signs that your nervous system has been running in survival mode for a long time.

The fact that you picked up this guide and read it matters.

You are choosing to be intentional about how you show up for the people who matter to you.

That choice is a lot more powerful than having flawless days.

About the Author

Michael Jenkins, MS, MSW, LCSW

Founder, Never Forgotten Services, PLLC

Michael is a U.S. Army combat veteran and licensed clinical social worker who specializes in working with military, veterans, and first responders—our frontline folks. After leaving the military, he spent years fighting his own demons and trying to figure out why he could hold it together on the job but feel completely wrecked at home. The turning point was realizing he wasn't the only one living that split life—and that nobody should have to feel that alone, confused, or broken by what the job asks of them.

Since then, his work has focused on trauma, the impact of high-stress jobs on home life, and helping families make sense of what the job brings home. He uses plain language, zero shame, and a "you're not crazy, this actually makes sense" approach to help frontline folks and their families understand what's happening in their nervous systems and learn how to move through the chaos together.