Challenge Stigma / Find Peace

FrontlineFolks | Guide #12

Overreacting to Small Stuff

A Field Guide for Frontline Folks and Their Families

When minor irritations feel like major threats.

By Michael Jenkins, MS, MSW, LCSW
Never Forgotten Services, PLLC

Disclaimer

This guide is for general education only.

It is not:

- Therapy
- Medical care
- A crisis service
- Individualized advice

Reading this guide does not create a therapist–client relationship with the author or with Never Forgotten Services, PLLC.

If you are in crisis or thinking about harming yourself or someone else, call 988 in the United States, go to your nearest emergency room, or contact your local emergency services.

If you want mental health treatment, please contact a licensed provider in your state or call the number on the back of your insurance card.

Table of Contents

1. What Overreaction Looks Like Under Stress
2. Why Your Threat Threshold Drops
3. Field Tools: Catching It Earlier
4. For Partners and Family
5. When To Get More Help
6. 7-Day Overreaction Tracking Drill
7. Worksheets
    - Worksheet 1: Worksheet: Overreacting to Small Stuff
    - Worksheet 2: Worksheet: Overreacting to Small Stuff - Partner/Team Check-In

1. What Overreaction Looks Like Under Stress

Overreacting isn't about being dramatic—it's about a nervous system calibrated for high-stakes threats that's now responding to low-stakes irritations with the same intensity. A slow driver, a dropped plate, a forgotten errand—these trigger the same physiological response as genuine danger. Your body goes full-threat while your brain knows it's not warranted. The result: explosive reactions that feel disproportionate to the trigger but perfectly aligned with your nervous system's current threat threshold.

What it looks like in frontline folks

- Snapping at people over minor issues (forgot milk, asked a question at the wrong time).
- Road rage or intense irritation at slow traffic, bad drivers, parking issues.
- Explosive frustration over technology, broken items, or things not working.
- Disproportionate anger when plans change or things don't go as expected.
- Immediate 0-to-100 activation over small annoyances.
- Feeling guilty or confused after: 'Why did I lose it over that?'

> **Not a character flaw:**
> Overreaction is a symptom of a sensitized nervous system. Your threat threshold is set too low. The fix isn't willpower—it's recalibration.

2. Why Your Threat Threshold Drops

Frontline work keeps your nervous system in a constant state of readiness. When baseline arousal stays elevated (chronic hypervigilance), your brain starts treating everything as a potential threat. Low-level stressors that normally wouldn't register now trigger full activation. It's like an alarm system stuck on high sensitivity—every breeze sets it off. Add sleep debt, unprocessed stress, and no recovery time, and your threshold drops even lower.

What lowers your overreaction threshold

- Chronic hypervigilance: Always scanning for threats means everything looks threatening.
- Sleep debt: Emotional regulation fails first when tired.
- Stacked stress: Multiple small stressors combine into one big reaction.
- Unprocessed critical incidents: Past events prime you to react faster.
- Control loss: Frontline work demands control. Losing it anywhere else triggers disproportionate responses.

- Transition failures: Not downshifting from work mode to home mode.

3. Field Tools: Catching It Earlier

You can't always stop the initial surge, but you can catch it before it escalates. The goal is to name the activation early, reset fast, and repair quickly when you miss the window.

Immediate strategies

1. Name early body signals: Jaw tight, fists clenched, chest pressure. These show up 5-10 seconds before explosion.
2. Use a 10-second pause script: 'I'm activated. I need 60 seconds.' Then step away, breathe, reset.
3. Lower baseline arousal daily: 10 minutes of intentional downshift after work (breathing, movement, quiet).
4. Track your threshold: Log when overreactions happen. Pattern = tired, hungry, transition times, specific people/places.
5. Repair fast: When you overreact, own it within 30 minutes. 'I blew that out of proportion. I'm sorry.'
6. Protect sleep for one week: Overreaction threshold rises dramatically with even small sleep improvements.

> Field rule: The overreaction isn't the problem—it's a signal. Your nervous system is telling you it's overloaded. Listen.

4. For Partners and Family

When your partner explodes over something small, it's not about the dropped keys or forgotten item—it's about a nervous system at capacity. Pointing out the disproportion mid-reaction doesn't help. Wait, then address the pattern.

What helps

- Don't engage during activation: Let the wave pass. Arguing escalates.
- Name the pattern later (not during): 'You seemed really activated earlier. What was going on?'
- Agree on a pause signal: A word or gesture that means 'I need a reset before I explode.'
- Track patterns together: When do overreactions happen? Tired? Transition times? Specific triggers?

- Support recovery: Overreactions spike when sleep is bad and stress is high. Help protect rest.

What makes it worse

- 'You're overreacting' (mid-explosion): True, but doesn't de-escalate.
- 'It's just [small thing]': Minimizing the reaction increases defensiveness.
- Demanding explanation during activation: They can't think clearly. Wait.
- Walking on eggshells: Avoiding all triggers enables the pattern instead of addressing it.

---

5. When To Get More Help

Overreactions that escalate, persist despite recovery efforts, or harm relationships and work need professional assessment. Chronic irritability can indicate underlying conditions (depression, TBI, PTSD).

Escalate if:

- Overreactions are increasing in frequency or intensity.
- You're damaging relationships or facing consequences at work.
- Physical aggression shows up (throwing things, punching walls, physical intimidation).
- You can't recover after overreacting—shame and guilt spiral for hours or days.
- Sleep and stress reduction don't improve the threshold.

6. 7-Day Overreaction Tracking Drill

Use the worksheet to track overreactions for one week. Pattern recognition is your first tool. Identify when, where, and under what conditions your threshold drops.

> **Daily drill:**
> 1) Log one overreaction (what happened, what triggered it).
> 2) Identify your state before it happened (tired, hungry, just got home?).
> 3) Name the first body signal (jaw, fists, chest).
> 4) Practice one early-catch tool tomorrow (pause, name it, reset).

Worksheet 1: Worksheet: Overreacting to Small Stuff

Use this to track overreactions and identify patterns so you can catch them earlier.

When did you overreact to something small this week?

Examples: snapped at someone, road rage, explosive frustration over minor issue.

What was your state just before each overreaction?

Tired, hungry, just got home, stacked stress, transition time, specific person/place.

What two early-catch strategies will you practice this week?

Examples: name body signals early, use pause script, lower baseline arousal daily.

1. Trigger:
What activated the pattern?

2. Early body signal:
What did your body do first?

3. Reset action used:
What fast action lowered intensity?

4. Better next response:
What exact behavior will you use next time?

> Overreaction is a signal, not a character flaw. Track the pattern, then adjust.

Worksheet 2: Worksheet: Overreacting to Small Stuff - Partner/Team Check-In

Use this with a partner to build shared awareness and early-warning agreements.

What does my overreaction look like to you?

Observable signs: snapping, explosive tone, throwing things, storming out, silent withdrawal.

What helps most when I'm starting to overreact?

Examples: pause signal, space without pursuit, calm tone, waiting to discuss later.

What's one agreement we'll try this week?

Example: 'I'll say "I need a reset" when activated. You'll give me 5 minutes without following.'

1. Signal phrase:

What phrase means pause and reset?

2. Time-to-return commitment:

When will re-engagement happen?

3. Repair step:

What specific repair happens after a rupture?

> Keep check-ins short, specific, and repeatable.

Closing

You've seen and carried more than most people will ever know.

Blow ups and shutdowns are not proof that you're broken or that your relationship is doomed.

They are signs that your nervous system has been running in survival mode for a long time.

The fact that you picked up this guide and read it matters.

You are choosing to be intentional about how you show up for the people who matter to you.

That choice is a lot more powerful than having flawless days.

About the Author

Michael Jenkins, MS, MSW, LCSW

Founder, Never Forgotten Services, PLLC

Michael is a U.S. Army combat veteran and licensed clinical social worker who specializes in working with military, veterans, and first responders—our frontline folks. After leaving the military, he spent years fighting his own demons and trying to figure out why he could hold it together on the job but feel completely wrecked at home. The turning point was realizing he wasn't the only one living that split life—and that nobody should have to feel that alone, confused, or broken by what the job asks of them.

Since then, his work has focused on trauma, the impact of high-stress jobs on home life, and helping families make sense of what the job brings home. He uses plain language, zero shame, and a "you're not crazy, this actually makes sense" approach to help frontline folks and their families understand what's happening in their nervous systems and learn how to move through the chaos together.