

Careless/Insufficient Effort Responding (C/IER) Detection

Pablo Rogers

2026-09-25

This notebook presents a tutorial on the detection of careless or insufficient effort responding (C/IER) in self-report questionnaires. We use real database from WHOQOL-Bref scale as an example ($n = 1,546$), and we apply the ‘careless’ package and Laz.R (Biemann et al., 2025) to detect C/IER (post hoc indicators). We also discuss the importance of procedural C/IER detection in the quality control of data obtained through self-report questionnaires.

1 Contextualization

The detection of Careless or Insufficient Effort Responding (C/IER) is an essential step in the quality control of data obtained through self-report questionnaires. C/IER occurs when respondents complete the instrument without attention to the item content, compromising the validity of the psychometric and substantive inferences derived from the data (Curran, 2016; Huang et al., 2026).

The analytical flow of this document follows two sequential levels of detection:

1. **Questionnaire:** procedural exclusions (a priori), including dropout, response time, and quality control items.
2. **WHOQOL Scale:** diagnostic post-hoc indicators (Laz.R, Longstring, Mahalanobis Distance, and IRV) with cut-off points estimated via the Kneedle algorithm (Biemann et al., 2025).

2 Data reading and preparation

The `data.csv` file contains the raw survey responses (separator `;`). In this step, dropout records are removed, variables for the analytical flow are selected, and the total response time (`TIME`) is derived.

Table 1: Structure and variable dictionary of the prepared survey dataset (`imported_data`)

Group	Variables	Class / Type	Description
Identification	ID	Numeric (Double)	Unique respondent identifier
Timing	CREATED, MODIFIED, TIME	Date-time / Numeric (s)	Survey timestamps and total response duration in seconds
Attention Checks	CQ1, CQ2, CQ3	Binary (0/1)	Quality control check items (1 = passed, 0 = failed)
WHOQOL-Bref Scale	Q1 to Q26 (26 items)	Ordinal (1 to 5)	Self-reported quality of life items

Source: [Procedural exclusions](#)

3 Procedural exclusions

Level 1 applies three procedures prior to calculating post-hoc indicators: dropout removal, exclusion for response time below 208 seconds ($104 \text{ items} \times 2 \text{ s/item}$) (Huang et al., 2026), and exclusion for two or more failures on quality control items (CQ1–CQ3) (Curran, 2016). Every exclusion is recorded in `exclusion_log.csv` with its reason and criterion.

Table 2: Procedural exclusions summary

Step	N
Raw data	1546
Dropouts removed	220
Excluded by time ($< 208 \text{ s}$)	0
Excluded by QC (≥ 2 fails)	28
Excluded by missings (> 5 in Q1-Q26)	3
Eligible for Level 2	1295
Remaining NAs in Q1-Q26	90

Source: [Procedural exclusions](#)

4 Calculation of C/IER indicators

Each indicator captures distinct manifestations of C/IER; their concurrent presence offers stronger evidence than any single indicator alone (Curran, 2016; Hong et al., 2020; Huang et al., 2026).

4.1 Univariate visualization




Figure 1: Univariate distributions of C/IER indicators

4.2 Cut-off points via Kneedle algorithm

The Kneedle algorithm is applied to the sorted (descending) curve of each indicator. For **Laz.R**, **Longstring**, and **MD**, kneedle with **sign** = -1 identifies where extremely high values separate from the majority (right tail). For **IRV**, **sign** = -1 is used to detect excessively high IRV and **sign** = +1 for excessively low IRV, since both extremes can be considered suspicious.

Table 3: Cut-off points estimated with the Kneedle algorithm

Indicator	Critical value	% flagged
Laz.R	0.60	6.8
Longstring	7.00	14.8
MD	40.12	15.4
IRV (high end)	1.34	4.9
IRV (low end)	0.69	3.8

Source: [Post-hoc C/IER Indicators](#)

4.3 Plots with cut-off points




Figure 2: Cumulative frequency curves with Kneedle cut-off points

4.4 C/IER Flags

For each indicator, a `_FLAG` variable is created (1 = flagged, 0 = not flagged) based on the Kneedle cut-off points.

Note: Post-hoc indicators are **diagnostic**, they do not trigger automatic exclusion in isolation. The final decision below requires corroboration from at least two of four selected indicators.

Table 4: Summary of respondents flagged by each C/IER indicator

Indicator	N flagged	% flagged
LazR_FLAG	91	7.0
Longstr_FLAG	191	14.7
MD_FLAG	201	15.5
IRV_high_FLAG	66	5.1
IRV_low_FLAG	49	3.8

Source: [Post-hoc C/IER Indicators](#)

4.5 From diagnostic flags to final exclusions

The cut-offs above identify response patterns that warrant attention; they do not, by themselves, prove that a respondent answered carelessly. Detection methods are sensitive to different manifestations of C/IER and may also flag unusual but valid response profiles. Consequently, combining every available flag with an **OR** rule can increase sensitivity at the cost of specificity and sample size (Curran, 2016; Hong et al., 2020). Indiscriminate post-hoc removal may also distort statistical and psychometric estimates when valid atypical respondents are classified as inattentive (Wang et al., 2025).

4.5.1 Consequences of alternative decision rules

The scenarios below make the cost of each possible rule explicit. The first two use all five diagnostic flags. The next two remove Longstring from the decision set, while retaining it as a descriptive diagnostic. The final two show the Laz.R–MD screening alternative.

Table 5: Consequences of alternative post-hoc decision rules

Decision rule	Excluded	% excluded	Retained
Any of the five diagnostic flags	436	33.7	859

Table 5: Consequences of alternative post-hoc decision rules

Decision rule	Excluded	% excluded	Retained
At least two of the five diagnostic flags	139	10.7	1156
Any of the four decision flags (excluding Longstring)	344	26.6	951
At least two of the four decision flags (Plan A)	62	4.8	1233
Laz.R or MD (case-review screening)	290	22.4	1005
Laz.R and MD	2	0.2	1293

Source: [Post-hoc C/IER Indicators](#)

An OR rule across all five flags would remove 436 respondents (33.7%), leaving 859 observations. Even after removing Longstring from the decision set, an OR rule would remove 344 respondents (26.6%). These reductions are difficult to defend when a single post-hoc indicator can produce false positives. In particular, Mahalanobis distance may overflag respondents when ordinal item responses depart from the multivariate-normal assumptions used in conventional outlier detection (Hong et al., 2020).

4.5.2 What the intersections show

Longstring and low IRV are invariability-oriented diagnostics, high IRV targets excessive variation, and MD targets multivariate atypicality. Laz.R detects predictable transitions, including straightlining, alternating, and diagonal patterns, but it is not a general detector of random responding. Laz.R should therefore be combined with complementary approaches, and its Kneedle cut-off should support scrutiny and possible exclusion rather than be treated as proof of carelessness (Biemann et al., 2025).

Table 6: Pairwise overlap among diagnostic C/IER flags

Indicator 1	Indicator 2	N intersection	% of indicator 1	% of indicator 2
Laz.R	Longstring	76	83.5	39.8
Laz.R	MD	2	2.2	1.0
Laz.R	IRV high	2	2.2	3.0
Laz.R	IRV low	18	19.8	36.7
Longstring	MD	8	4.2	4.0
Longstring	IRV high	11	5.8	16.7
Longstring	IRV low	26	13.6	53.1
MD	IRV high	42	20.9	63.6
MD	IRV low	0	0.0	0.0
IRV high	IRV low	0	0.0	0.0

Source: [Post-hoc C/IER Indicators](#)

Of the 91 Laz.R flags, 76 (83.5%) are also Longstring flags. Counting both in the composite rule would allow one narrow manifestation - repetitive or highly predictable responding - to contribute two votes. Longstring remains useful for diagnosis and teaching, but its narrower scope and empirical redundancy with Laz.R justify excluding it from the final flag count. Conversely, 42 of the 66 IRV-high cases are also flagged by MD, providing convergent evidence for excessive variability or multivariate atypicality.

Table 7: Intersection of Laz.R and Mahalanobis distance flags

Classification	N	% of sample
Neither flag	1005	77.6
MD only	199	15.4
Laz.R only	89	6.9
Both Laz.R and MD	2	0.2

Source: [Post-hoc C/IER Indicators](#)

Only two respondents fail both Laz.R and MD, whereas 199 fail MD only and 89 fail Laz.R only. Requiring both would remove almost no cases, while using either would identify 290 cases (22.4%) for case-by-case review. This can be a useful screening pool, but it was not adopted as an automatic exclusion rule because the indicators are nearly disjoint in this sample and MD-only evidence is not sufficient to infer carelessness from ordinal responses (Hong et al., 2020; Ward & Meade, 2023).

4.5.3 Final rule: at least two of four decision flags

The final rule counts **Laz.R**, **MD**, **IRV high**, and **IRV low** and excludes a respondent only when at least two of these four flags are present. It requires corroboration, avoids deletion based on MD alone, and retains IRV as evidence for both insufficient and excessive within-person variability. This follows the recommendation to use multiple, context-appropriate indicators rather than rely blindly on one index (Curran, 2016; Ward & Meade, 2023).

Hong et al. (2020) best-subset simulations support selecting a smaller group of complementary indicators instead of a kitchen-sink combination, although their specific subset and cut-offs are design-dependent and should not be imported unchanged into the WHOQOL-Bref (Hong et al., 2020). Plan A is therefore a transparent teaching decision for this dataset, not a universal threshold.

Table 8: Evidence combinations among respondents excluded by Plan A

decision_evidence	Excluded	% of exclusions
MD + IRV high	41	66.1
Laz.R + IRV low	18	29.0
Laz.R + IRV high	1	1.6
Laz.R + MD	1	1.6
Laz.R + MD + IRV high	1	1.6

Source: [Post-hoc C/IER Indicators](#)

Plan A removes 62 of the 1,295 eligible respondents (4.8%). The full audit log preserves each excluded ID, indicator value, flag, composite count, and evidence combination in `posthoc-exclusion-log.csv`.

4.5.4 Final analysis dataset

Table 9: Derivation of the final analysis sample

Stage	N
Eligible after procedural exclusions	1295
Excluded by Plan A post-hoc rule	62
Final analysis dataset	1233

Source: [Post-hoc C/IER Indicators](#)

The final `whoqol_analysis.csv` contains 1,233 respondents, the ID column, and Q1-Q26. Items Q3, Q4, and Q26 are reverse-scored only after the C/IER indicators have been calculated on the response sequence as administered. For substantive analyses, the unfiltered 1,295-case dataset can be retained as a sensitivity condition. Large changes between filtered and unfiltered results should be reported rather than used to choose whichever cleaning rule produces the preferred result (Wang et al., 2025).

References

Biemann, T., Koch-Bayram, I. F., Meier-Barthold, M., & Aguinis, H. (2025). Using Markov Chains to Detect Careless Responding in Survey Research. *Organizational Research Methods*, 28(4), 543–568. <https://doi.org/10.1177/10944281251334778>

- Curran, P. G. (2016). Methods for the detection of carelessly invalid responses in survey data. *Journal of Experimental Social Psychology*, 66, 4–19. <https://doi.org/10.1016/j.jesp.2015.07.006>
- Hong, M., Steedle, J. T., & Cheng, Y. (2020). Methods of Detecting Insufficient Effort Responding: Comparisons and Practical Recommendations. *Educ Psychol Meas*, 80(2), 312–345. <https://doi.org/10.1177/0013164419865316>
- Huang, J. L., Wang, Z., Huang, R., Wu, D., & Shi, H. (2026). *Insufficient Effort Responding in Management Research: A Critical Review and Future Directions*. 52(1). <https://doi.org/10.1177/01492063251330268>
- Wang, M. D., Xiao, L., & Zang, X. (2025). When Cleaning Data Introduces Bias: A Critical Examination of Post-Hoc Methods in Detecting Insufficient Effort Responding. *Journal of Official Statistics*, 41(4). <https://doi.org/10.1177/0282423X251337843>
- Ward, M. K., & Meade, A. W. (2023). Dealing with Careless Responding in Survey Data: Prevention, Identification, and Recommended Best Practices. *Annu. Rev. Psychol.*, 74(Volume 74, 2023), 577–596. <https://doi.org/10.1146/annurev-psych-040422-045007>