

TT Projection Curved Space v2

1 Maximally Symmetric Space TT

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{g_{\mu\nu}}{D-1}(\nabla^\sigma W_\sigma - h) + \frac{2-D}{D-1} \left(\nabla_\mu \nabla_\nu - \frac{g_{\mu\nu} R}{D(D-1)} \right) \int D(x, x') \nabla^\sigma W_\sigma - \frac{1}{D-1} \left(\nabla_\mu \nabla_\nu - \frac{g_{\mu\nu} R}{D(D-1)} \right) \int D(x, x') h \quad (1.1)$$

$$\left(\nabla_\alpha \nabla^\alpha - \frac{R}{D-1} \right) D(x, x') = g^{-1/2} \delta^{(D)}(x - x') \quad (1.2)$$

$$\nabla^\mu h_{\mu\nu} = \left(\nabla_\alpha \nabla^\alpha - \frac{R}{D} \right) W_\nu \quad (1.3)$$

With the covariant operator $(\nabla^2 - R/D)$ mixing indices of W_ν , the particular integral solution for W_ν involves a bi-tensor Green's function $D_{\sigma\rho'}$ which obeys

$$\left(\nabla^\alpha \nabla_\alpha - \frac{R}{D} \right) D_{\sigma\rho'}(x, x') = g_{\sigma\rho'} g^{-1/2} \delta^4(x - x'). \quad (1.4)$$

Here $g_{\sigma\rho'}$ represents a parallel propagator, defined in terms of Vierbeins e_μ^a :

$$g^{\alpha'}_{\beta}(x, x') = e_a^{\alpha'}(x') e_\beta^a(x), \quad g_{\mu\nu} = \eta_{ab} e_\mu^a e_\nu^b. \quad (1.5)$$

In terms of (1.4), W_ν has particular solution

$$W_\nu = g^{1/2} \int D_{\nu}{}^{\rho'}(x, x') \nabla^{\sigma'} h_{\rho'\sigma'}. \quad (1.6)$$

2 Curved Space TT

To generalize to curved space, we assume $h_{\mu\nu}$ to be of the form:

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \underbrace{\left(\nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{2}{D} g_{\mu\nu} \nabla^\alpha W_\alpha \right)}_{W_{\mu\nu}} + \underbrace{\frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu)}_{S_{\mu\nu}} \Psi \quad (2.1)$$

Taking the trace of (2.1), we find the vector sector $W_{\mu\nu}$ is decoupled from the trace and Ψ can easily be inverted,

$$g^{\mu\nu} W_{\mu\nu} = 0 \quad (2.2)$$

$$g^{\mu\nu} S_{\mu\nu} = \nabla_\alpha \nabla^\alpha \Psi = h \quad \rightarrow \Psi = \int g^{1/2} D(x, x') h \quad (2.3)$$

Taking the divergence of (2.1), we have

$$\nabla^\mu h_{\mu\nu} = \nabla^\mu W_{\mu\nu} + \nabla^\mu S_{\mu\nu}(h) \quad (2.4)$$

By substituting (2.3), the above serves to define an equation for W_μ in terms of h and $h_{\mu\nu}$, namely

$$\nabla_\alpha \nabla^\alpha W_\nu + \nabla^\alpha \nabla_\nu W_\alpha - \frac{2}{D} \nabla_\nu \nabla^\alpha W_\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \int g^{1/2} D(x, x') h \quad (2.5)$$

Commuting derivatives, (2.5) can be expressed in the equivalent forms,

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \nabla_\alpha \nabla_\nu - \frac{2}{D} \nabla_\nu \nabla_\alpha \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \int g^{1/2} D(x, x') h, \quad (2.6)$$

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha - R_{\nu\alpha} \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} R_{\nu\alpha} \nabla^\alpha \int g^{1/2} D(x, x') h. \quad (2.7)$$

Similar to (1.4), the requisite Green's function that solves W_α is a bi-tensor defined as

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha - R_{\nu\alpha} \right] D^{\alpha\gamma'} = g^{\alpha\gamma'} g^{-1/2} \delta^{(D)}(x, x'). \quad (2.8)$$

Hence, W_μ takes the form

$$W_\mu = \int g^{1/2} D_\mu{}^{\sigma'} \left[\nabla^{\rho'} h_{\sigma'\rho'} - \frac{1}{D-1} R_{\sigma'\rho'} \nabla^{\rho'} \int g^{1/2} D(x', x'') h \right]. \quad (2.9)$$

To summarize, in curved space $h_{\mu\nu}$ may be decomposed according to

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \left(\nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{2}{D} g_{\mu\nu} \nabla^\alpha W_\alpha \right) + \frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu) \Psi \quad (2.10)$$

with Ψ and W_μ obeying

$$\Psi = \int g^{1/2} D(x, x') h \quad (2.11)$$

$$W_\mu = \int g^{1/2} D_\mu{}^{\sigma'} \left[\nabla^{\rho'} h_{\sigma'\rho'} - \frac{1}{D-1} R_{\sigma'\rho'} \nabla^{\rho'} \int g^{1/2} D(x', x'') h \right] \quad (2.12)$$

$$\nabla_\alpha \nabla^\alpha D(x, x') = g^{-1/2} \delta^{(D)}(x - x') \quad (2.13)$$

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha - R_{\nu\alpha} \right] D^{\alpha\gamma'} = g^{\alpha\gamma'} g^{-1/2} \delta^{(D)}(x, x'). \quad (2.14)$$

2.1 SVTD Decomposition

Starting with

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \left(\nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{2}{D} g_{\mu\nu} \nabla^\alpha W_\alpha \right) + \frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu) \Psi, \quad (2.15)$$

we decompose W_μ into transverse and longitudinal components viz.

$$W_\mu = \underbrace{W_\mu - \nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma}_{F_\mu} + \underbrace{\nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma}_H. \quad (2.16)$$

Setting $h_{\mu\nu}^{T\theta} = 2F_{\mu\nu}$, (2.28) becomes

$$h_{\mu\nu} = 2F_{\mu\nu} + \nabla_\mu F_\nu + \nabla_\nu F_\mu + 2\nabla_\mu \nabla_\nu H - \frac{2}{D} g_{\mu\nu} \nabla_\alpha \nabla^\alpha H + \frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu) \Psi. \quad (2.17)$$

Upon further defining

$$F = H - \frac{1}{2(D-1)} \Psi \quad (2.18)$$

$$\chi = \frac{1}{D} \nabla_\alpha \nabla^\alpha H - \frac{1}{2(D-1)} \nabla_\alpha \nabla^\alpha \Psi, \quad (2.19)$$

we may express (2.28) as the desired SVTD form:

$$h_{\mu\nu} = -2g_{\mu\nu} \chi + 2\nabla_\mu \nabla_\nu F + \nabla_\mu F_\nu + \nabla_\nu F_\mu + 2F_{\mu\nu}. \quad (2.20)$$

$$\chi = \frac{1}{D} \nabla^\sigma W_\sigma - \frac{1}{2(D-1)} h \quad (2.21)$$

$$F = \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma - \frac{1}{2(D-1)} \int g^{1/2} D(x, x') h \quad (2.22)$$

$$F_\mu = W_\mu - \nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma \quad (2.23)$$

$$2F_{\mu\nu} = 2g_{\mu\nu} \chi - 2\nabla_\mu \nabla_\nu F - \nabla_\mu F_\nu - \nabla_\nu F_\mu - h_{\mu\nu} \quad (2.24)$$

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha - R_{\nu\alpha} \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \Psi \quad (2.25)$$

$$\frac{2(D-1)}{D} \nabla_\alpha \nabla^\alpha \nabla^\sigma W_\sigma - \nabla^\alpha R W_\alpha - 2R^{\alpha\beta} \nabla_\alpha W_\beta = \nabla^\alpha \nabla^\beta h_{\alpha\beta} - \frac{1}{(D-1)} \left[\frac{1}{2} \nabla^\alpha R \nabla_\alpha + R^{\alpha\beta} \nabla_\alpha \nabla_\beta \right] \Psi \quad (2.26)$$

2.2 Curved TT in Max. Symmetric Space (Incomplete)

In a space of maximal symmetry defined by

$$\begin{aligned} R_{\lambda\mu\nu\kappa} &= k(g_{\mu\nu} g_{\lambda\kappa} - g_{\lambda\nu} g_{\mu\kappa}) \\ R_{\mu\kappa} &= k(1-D)g_{\mu\kappa} = \frac{R}{D} g_{\mu\kappa} \\ R &= kD(1-D), \end{aligned} \quad (2.27)$$

the defining equation for W_μ reduces to

$$\left[g_{\nu\alpha} \left(\nabla_\beta \nabla^\beta - \frac{R}{D} \right) + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{R}{D-1} \nabla_\nu \int g^{1/2} D(x, x') h \quad (2.28)$$

2.3 Curved TT in Minkowski

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \left(\nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{2}{D} g_{\mu\nu} \nabla^\alpha W_\alpha \right) + \frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu) \Psi, \quad (2.29)$$

In a Minkowski geometry, the defining equation for W_μ reduces to

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} \quad (2.30)$$

$$W_\mu = \int g^{1/2} D_\mu^{\sigma'} \nabla^{\rho'} h_{\sigma' \rho'} \quad (2.31)$$

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha \right] D^{\alpha\gamma'} = g^{\alpha\gamma'} \delta^{(D)}(x, x') \quad (2.32)$$

$$\nabla_\alpha \nabla^\alpha \Psi = h \quad (2.33)$$

$$\Psi = \int g^{1/2} D(x, x') h \quad (2.34)$$

$$\nabla_\alpha \nabla^\alpha D(x, x') = g^{-1/2} \delta^{(D)}(x - x') \quad (2.35)$$

Decompose W_μ into transverse and longitudinal components viz.

$$W_\mu = \underbrace{W_\mu - \nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma}_{F_\mu} + \underbrace{\nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma}_{H}. \quad (2.36)$$

$$h_{\mu\nu} = h_{\mu\nu}^{T\theta} + \left(\nabla_\mu W_\nu + \nabla_\nu W_\mu - \frac{2}{D} g_{\mu\nu} \nabla^\alpha W_\alpha \right) + \frac{1}{D-1} (g_{\mu\nu} \nabla_\alpha \nabla^\alpha - \nabla_\mu \nabla_\nu) \Psi \quad (2.37)$$

$$h_{\mu\nu} = -2g_{\mu\nu}\chi + 2\nabla_\mu \nabla_\nu F + \nabla_\mu F_\nu + \nabla_\nu F_\mu + 2F_{\mu\nu}. \quad (2.38)$$

$$\chi = \frac{1}{D} \nabla^\sigma W_\sigma - \frac{1}{2(D-1)} h \quad (2.39)$$

$$F = \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma - \frac{1}{2(D-1)} \int g^{1/2} D(x, x') h \quad (2.40)$$

$$F_\mu = W_\mu - \nabla_\mu \int g^{1/2} D(x, x') \nabla^\sigma W_\sigma \quad (2.41)$$

$$\left[g_{\nu\alpha} \nabla_\beta \nabla^\beta + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla_\alpha - R_{\nu\alpha} \right] W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \Psi \quad (2.42)$$

$$\nabla_\alpha \nabla^\alpha W_\nu + \left(\frac{D-2}{D} \right) \nabla_\nu \nabla^\alpha W_\alpha - R_{\nu\alpha} W^\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \Psi \quad (2.43)$$

$$\nabla_\alpha \nabla^\alpha W_\nu + \nabla^\alpha \nabla_\nu W_\alpha - \frac{2}{D} \nabla_\nu \nabla^\alpha W_\alpha = \nabla^\alpha h_{\alpha\nu} - \frac{1}{D-1} (\nabla_\nu \nabla_\alpha \nabla^\alpha - \nabla_\alpha \nabla^\alpha \nabla_\nu) \Psi \quad (2.44)$$

$$\frac{2(D-1)}{D} \nabla_\alpha \nabla^\alpha \nabla^\sigma W_\sigma - \nabla^\alpha R W_\alpha - 2R^{\alpha\beta} \nabla_\alpha W_\beta = \nabla^\alpha \nabla^\beta h_{\alpha\beta} - \frac{1}{(D-1)} \left[\frac{1}{2} \nabla^\alpha R \nabla_\alpha + R^{\alpha\beta} \nabla_\alpha \nabla_\beta \right] \Psi \quad (2.45)$$