

Some Title

And Maybe a Subtitle

Peter Illig Eli Orvis Yukihiro Segawa Nick Spinale

Carleton College

February 20th, 2018

Contents

- 1 Introduction
- 2 Simplifying the Problem

A Title

Contents of the slide

Roots of $f^n(x)$

$$f(x) = (x - \gamma)^2 + \gamma + m$$

- The roots of $f(x)$ are $\gamma \pm \sqrt{-m - \gamma}$
- If α is a root of $f^n(x)$, then $\gamma \pm \sqrt{\alpha - m - \gamma}$ are roots of $f^{n+1}(x)$

Observation

For $n > 0$, the roots of $f^n(x)$ are, with n radicals:

$$\gamma \pm \sqrt{-m \pm \sqrt{-m \pm \sqrt{-m \pm \dots \sqrt{-m - \gamma}}}}$$

Roots of $f^n(x)$

Observation

For $n > 0$, the roots of $f^n(x)$ are, with n radicals:

$$\gamma \pm \sqrt{-m \pm \sqrt{-m \pm \sqrt{-m \pm \dots \sqrt{-m - \gamma}}}}$$

For notational convenience, define $\beta : \Sigma^* \rightarrow \mathbb{C}$ where

$$\begin{aligned}\beta_\epsilon &= -\gamma \\ \beta_{0s} &= \sqrt{-m + \beta_s} \\ \beta_{1s} &= -\sqrt{-m + \beta_s}\end{aligned}$$

For $n > 0$, the roots of $f^n(x)$ are exactly $\{ \gamma + \beta_s \mid s \in \Sigma^n \}$.