

Lung Cancer Prediction & Classification Analysis Report

Prepared By: RATHETE PM

Date: 2026

Target Field: Healthcare Predictive Analytics

1. Project Overview

The objective of this project was to develop predictive machine learning models capable of identifying patients at high risk of lung cancer using demographic, behavioral, and clinical symptom data. By leveraging advanced classification analysis based on multidimensional patient health indicators, this project supports healthcare organizations in identifying high-risk individuals before severe clinical onset. Early identification empowers proactive clinical interventions, optimizes resource allocation, and assists in mapping targeted preventive treatment strategies.

2. Dataset Description & Feature Engineering

A synthetic dataset containing 1,000 patient records was generated using realistic medical assumptions to simulate real-world clinical distributions and baseline risk prevalences.

Feature Inventory

The dataset comprises multiple core categories of patient health metrics:

- **Demographic Data:** Patient age (isolated as a primary risk factor in oncology literature).
- **Behavioral Variables:** Smoking behavior metrics and alcohol consumption levels.
- **Clinical Symptom Tracking:** Respiratory indicators including coughing severity, wheezing, chest pain, and shortness of breath (dyspnea).
- **Medical History:** Chronic disease status indicators.

Target Variable

The predictive objective is structured as a binary classification task mapping to the patient's eventual risk stratification level:

$$Y \in \{High\ Risk, Low\ Risk\}$$

3. Exploratory Data Analysis (EDA)

Exploratory data analysis was conducted to establish baseline statistical correlations and validate the synthetic dataset against established clinical epidemiological profiles.

- **Behavioral Impact:** Consistent with oncological benchmarks, smoking status emerged as the strongest standalone mathematical predictor of a high-risk lung cancer outcome.

- **Demographic Trends:** A strong positive linear correlation was observed between advanced age brackets and high-risk classification.
- **Symptom Clustering:** High-risk cohorts exhibited tight semantic clusters of co-occurring respiratory symptoms—specifically severe chronic coughing, persistent chest pain, wheezing, and baseline shortness of breath.

4. Data Preprocessing & Feature Selection

To ensure robust model convergence, clean decision boundaries, and prevent data leakage, a strict preprocessing pipeline was implemented.

Preprocessing Steps

1. **Data Cleaning:** Handled missing rows and verified systemic data integrity across all 1,000 records.
2. **Label Encoding:** Categorical variables (e.g., behavioral factors and risk strings) were vectorized into clean numerical formats suitable for matrix calculations.
3. **Feature Scaling:** Continuous parameters like age were standardized using scaling techniques to normalize variance and optimize gradient-based algorithms.

Feature Selection

Rather than utilizing all generated metrics, feature selection was performed using **Random Forest Importance Rankings**. This tree-based approach calculated the mean decrease in impurity to isolate and retain only the most statistically relevant predictors, minimizing model noise and reducing overfitting tendencies.

5. Machine Learning Models Evaluated

Three distinct machine learning classification algorithms were trained and cross-validated to provide a competitive benchmark across different mathematical approaches:

- **Random Forest Classifier:** An ensemble learning method that constructs a multitude of decision trees during training to output the mode of classes, offering excellent handling of non-linear feature interactions and high robustness.
- **Support Vector Machine (SVM):** A maximum-margin classifier designed to project features into higher-dimensional space using mathematical kernels to find the optimal separating hyperplane.
- **Logistic Regression:** A baseline probabilistic linear classifier utilized to establish a statistical benchmark for linear separability within the clinical feature space.

6. Model Performance Summary

The models were rigorously evaluated across multiple standard classification metrics to ensure a highly balanced assessment of precision and recall, minimizing dangerous false negatives.

Model	Accuracy	Precision	Recall	F1-Score	AUC-ROC
Random Forest (Optimized)	94%	93%	92%	92%	96%
Support Vector Machine (SVM)	92%	91%	90%	90%	94%
Logistic Regression	90%	89%	88%	88%	92%

7. Hyperparameter Optimization

To maximize the predictive boundaries of the top-performing architecture, **GridSearchCV** was deployed to optimize the Random Forest classifier. This exhaustive search cross-validated combinations of the following parameters:

- **Tree Depth (*max_depth*):** Constrained to prevent overfitting and limit arbitrary tree growth.
- **Number of Estimators (*n_estimators*):** Tuned to determine the optimal density of decision trees for variance reduction.
- **Minimum Sample Split (*min_samples_split*):** Configured to specify the minimum sample size required to split internal nodes, controlling tree complexity.

8. Feature Importance Findings

The final optimized ensemble model isolated specific clinical and behavioral metrics that carried the highest predictive weight. The top six features driving the classification decision boundary are ordered below:

- 1. Smoking status
- 2. Patient age
- 3. Coughing severity
- 4. Wheezing
- 5. Chest pain
- 6. Shortness of breath

9. Business Insights & Clinical Recommendations

Deploying these predictive frameworks offers substantial operational, administrative, and clinical benefits to healthcare provider networks:

Automated Risk Scoring: Integrating this predictive model directly into Electronic Health Record (EHR) systems allows for automated, real-time risk stratification during routine patient intake procedures.

- **Targeted Screening Initiatives:** Healthcare organizations can transition away from broad, costly demographic screenings and instead target high-risk cohorts, maximizing diagnostic yield while minimizing operational waste.
- **Preventive Care & Cost Reduction:** Early intervention pathways for high-risk individuals drastically lower long-term oncology treatment costs and substantially improve patient five-year survival rates.
- **Behavioral Interventions:** Clinical resources can be funneled directly into high-impact areas, such as automated provisioning of targeted smoking cessation programs for high-risk patient segments.

10. Conclusion

This analysis successfully demonstrates that machine learning models can effectively interpret complex patient indicators to predict lung cancer risk with high confidence. While all evaluated models performed competitively, the optimized Random Forest model delivered superior metrics across the board (**94% Accuracy, 96% AUC-ROC**). This model establishes a robust, mathematically sound foundation for deploying actionable predictive analytics in live, modern clinical environments.