

# Index Reduction for Second Order Singular Systems of Difference Equations

Vu Hoang Linh<sup>a,\*</sup>, Ha Phi<sup>a</sup>

<sup>a</sup>*Faculty of Mathematics, Mechanics, and Informatics, Vietnam National University,  
334 Nguyen Trai, Thanh Xuan, Hanoi, Vietnam*

---

## Abstract

This paper is devoted to the analysis of linear second-order *discrete-time descriptor systems* (or singular difference equations (SiDEs) with control). Following the algebraic approach proposed by Kunkel and Mehrmann for pencils of matrix valued functions, first we present a theoretical framework based on a procedure of reduction to analyze the corresponding initial value problem for SiDEs, which is followed by the analysis of descriptor systems. We also describe methods to analyze structural properties related to the solvability analysis of these systems. Namely, two numerical algorithms for reduction to the so-called *strangeness-free forms* are presented. Two associated index notions are also introduced and discussed. This work extends and complement some recent results for high-order continuous-time descriptor systems and first-order discrete-time descriptor systems.

*Keywords:* Singular systems, Descriptor systems, Second-order difference equations, Index reduction, Regularization, Strangeness-index  
*2010 MSC:* 15A23, 39A05, 39A06, 93C05

---

## 1. Introduction

In this paper we study second-order discrete-time descriptor systems of the form

$$A_n x(n+2) + B_n x(n+1) + C_n x(n) + D_n u(n) = f(n), \quad \text{for all } n \geq n_0. \quad (1)$$

We will also discuss the initial value problem of the associated singular difference equation (SiDE)

$$A_n x(n+2) + B_n x(n+1) + C_n x(n) = f(n), \quad \text{for all } n \geq n_0, \quad (2)$$

together with some given initial conditions

$$x(n_0+1) = x_1, \quad x(n_0) = x_0. \quad (3)$$

Here the solution/state  $x = \{x(n)\}_{n \geq n_0}$ , the inhomogeneity  $f = \{f(n)\}_{n \geq n_0}$ , the input  $u = \{u(n)\}_{n \geq n_0}$ , where  $x(n) \in \mathbb{R}^d$ ,  $f(n) \in \mathbb{R}^m$  and  $u(n) \in \mathbb{R}^p$  for each

---

\*Corresponding author

Email addresses: [linhvh@vnu.edu.vn](mailto:linhvh@vnu.edu.vn) (Vu Hoang Linh), [haphi.hus@vnu.edu.vn](mailto:haphi.hus@vnu.edu.vn) (Ha Phi)  
Preprint submitted to Linear Algebra and its Applications November 7, 2019

13  $n \geq n_0$ . Three matrix sequences  $\{A_n\}_{n \geq n_0}$ ,  $\{B_n\}_{n \geq n_0}$ ,  $\{C_n\}_{n \geq n_0}$  take values in  
14  $\mathbb{R}^{m,d}$ , and  $\{D_n\}_{n \geq n_0}$  takes values in  $\mathbb{R}^{m,p}$ . We notice that all the results in this  
15 paper also can be carried over to the complex case and they can also be easily  
16 extended to systems of higher than second order. However, for ease of notation  
17 and because this is the most important case in practice, we restrict ourselves to  
18 the case of real and second order systems.

19 The SiDE (2), on one side, can be considered as a resulting equation, ob-  
20 tained by discretization (e.g. finite difference) of some differential-algebraic  
21 equations (DAEs) or constrained partial differential equations (PDEs). On the  
22 other side, there are also many models/applications in real-life, which lead to  
23 SiDEs, for example Leontief economic models, biological backward Leslie model,  
24 etc, see e.g. [1, 5, 9, 13].

25 While both DAEs and SiDEs of first order have been well-studied from both  
26 theoretical and numerical sides, the same maturity has not been reached for  
27 higher order systems. In classical literature [1, 5, 9], usually new variables are  
28 introduced to present some chosen derivatives of the state variable  $x$  such that  
29 a high order system can be reformulated as a first order one. This method,  
30 however, is not only non-unique but also has presented some substantial dis-  
31 advantages. As have been fully discussed in [12, 16] for continuous-time sys-  
32 tems, these disadvantages include: (1st) increase the index of the system, and  
33 therefore the complexity of a numerical method to solve it; (2nd) increase the  
34 computational effort, due to the bigger size of a new system; (3rd) affect the  
35 controllability/observability of the corresponding descriptor system, since there  
36 exist situations where a new system is uncontrollable while the original one is.  
37 Therefore, the *algebraic approach*, which treats the system directly without re-  
38 formulating it, has been presented in [12, 16, 21, 22] in order to overcome the  
39 disadvantages mentioned above. Nevertheless, even for second order SiDEs, this  
40 method has not yet been considered.

41 Another motivation of this work comes from recent research on the stability  
42 analysis of high order, discrete-time systems with time-dependent coefficients  
43 [11, 17]. In these works, systems are supposed to be given in either strangeness-  
44 free form or linear state-space form. Nevertheless, it is not always the case in  
45 applications, and hence, a reformulation procedure would be required.

46 Therefore, the main aim of this article is to set up a comparable framework  
47 for second order SiDEs and for discrete-time descriptor systems as well. It  
48 is worth marking that the algebraic method proposed in [12, 16] is applicable  
49 theoretically but not numerically, due to two reasons: (1st) The condensed forms  
50 of the matrix coefficients are really big and complicated; (2nd) The system's  
51 transformations are not orthogonal, and hence, not numerically stable. In this  
52 work, we will modify this method to make it more concise and also computable  
53 in a stable way.

54 The outline of this paper is as follows. After giving some auxiliary results in  
55 Section 2, in Sections 3 and 4 we consecutively introduce *index reduction proce-*  
56 *dures* for SiDEs and for descriptor systems. The main results of these sections  
57 are Theorem 3.10 and Algorithm 1 (Section 3) and Theorem 4.5 (Section 4).  
58 Resulting systems from these procedures allow us to determine structural prop-

erties such as existence and uniqueness of a solution, consistency and hidden constraints, etc. In order to get stable numerical solutions of these systems, in Section 5 we study the *difference array approach* in Algorithm 2 and Theorem 5.7 aiming at bringing out the strangeness-free form of a given system. Finally, we finish with some conclusions.

## 2. Preliminaries

In the following example we demonstrate some difficulties that may arise in the analysis of second order SiDEs.

**Example 2.1.** Consider the following second order descriptor system, motivated from Example 2, [16].

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} x(n+2) + \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} x(n+1) + \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} x(n) + \begin{bmatrix} -1 \\ -1 \end{bmatrix} u(n) = \begin{bmatrix} f_1(n) \\ f_2(n) \end{bmatrix}, \quad n \geq n_0. \quad (4)$$

Clearly, from the second equation  $\begin{bmatrix} 1 & 0 \end{bmatrix} x(n) = u(n) + f_2(n)$ , we can shift forward the time  $n$  by one to obtain

$$\begin{bmatrix} 1 & 0 \end{bmatrix} x(n+1) = u(n+1) + f_2(n+1) \quad \text{and} \quad \begin{bmatrix} 1 & 0 \end{bmatrix} x(n+2) = u(n+2) + f_2(n+2).$$

Inserting these into the first equation of (4), we find out the hidden constraint

$$f_2(n+2) + u(n+2) + f_2(n+1) + u(n+1) + \begin{bmatrix} 0 & 1 \end{bmatrix} x(n) = f_1(n).$$

Consequently, we deduce the following system, which possess a unique solution

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} x(n) = \begin{bmatrix} f_1(n) - f_2(n+2) - f_2(n+1) - u(n+2) - u(n+1) \\ u(n) + f_2(n) \end{bmatrix}, \quad n \geq n_0.$$

Let  $n = n_0$  in this new system, we obtain a constraint that  $x(n_0)$  must obey. This example showed us some important facts. Firstly, one can use some shift operators and row-manipulation (Gaussian eliminations) to derive hidden constraints. Secondly, a solution only exists if initial conditions and an input fulfill certain consistency conditions. Finally, in this example the solution depends on the future input. This property is called *non-causality* and cannot happen in the case of regular difference equations.

For matrices  $Q \in \mathbb{R}^{q,d}$ ,  $P \in \mathbb{R}^{p,d}$ , the pair  $(Q, P)$  is said to *have no hidden redundancy* if

$$\text{rank} \left( \begin{bmatrix} Q \\ P \end{bmatrix} \right) = \text{rank}(Q) + \text{rank}(P).$$

Otherwise,  $(Q, P)$  is said to *have hidden redundancy*. The geometrical meaning of this concept is that the intersection space  $\text{span}(P^T) \cap \text{span}(Q^T)$  contains only the zero-vector  $\vec{0}$ . Here by  $\text{span}(P^T)$  (resp.,  $\text{span}(Q^T)$ ) we denote the real

vector space spanned by the rows of  $P$  (resp., rows of  $Q$ ). We further notice that, if  $\begin{bmatrix} Q \\ P \end{bmatrix}$  is of full row rank then obviously, the pair  $(Q, P)$  has no hidden redundancy. However, the converse is not true as is obvious for  $Q = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ ,  
 $P = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ .

**Lemma 2.2.** ([7]) Suppose that for  $Q \in \mathbb{R}^{q,d}$ ,  $P \in \mathbb{R}^{p,d}$ , the pair  $(Q, P)$  has no hidden redundancy. Then, for any matrix  $U \in C^{q,q}$  and any  $V \in C^{p,p}$ , the pair  $(UQ, VP)$  has no hidden redundancy.

**Lemma 2.3.** ([7]) Consider  $k + 1$  full row rank matrices  $R_0 \in \mathbb{R}^{r_0,d}, \dots, R_k \in \mathbb{R}^{r_k,d}$ , and assume that for  $j = k, \dots, 1$  none of the matrix pairs  $\left( R_j, \begin{bmatrix} R_{j-1} \\ \vdots \\ R_0 \end{bmatrix} \right)$  has a hidden redundancy. Then,  $\begin{bmatrix} R_k \\ \vdots \\ R_0 \end{bmatrix}$  has full row rank.

Lemma 2.4 below will be very useful later for our analysis, in order to remove hidden redundancy in the coefficients of (2).

**Lemma 2.4.** Consider two matrix sequences  $\{P_n\}_{n \geq n_0}$ ,  $\{Q_n\}_{n \geq n_0}$  which take values in  $\mathbb{R}^{p,d}$  and  $\mathbb{R}^{q,d}$ , respectively. Furthermore, assume that they satisfy the constant rank assumptions

$$\text{rank}(Q_n) = r_Q, \quad \text{and} \quad \text{rank} \left( \begin{bmatrix} P_n \\ Q_n \end{bmatrix} \right) = r_{[P;Q]}, \quad \text{for all } n \geq n_0.$$

Then, there exists a matrix sequence  $\left\{ \begin{bmatrix} S_n & 0 \\ Z_n^{(1)} & Z_n^{(2)} \end{bmatrix} \right\}_{n \geq n_0}$  in  $\mathbb{R}^{p,p+q}$  such that the following conditions hold.

- i)  $S_n \in \mathbb{R}^{r_{[P;Q]} - r_Q, p}$ ,  $Z_n^{(1)} \in \mathbb{R}^{p - r_{[P;Q]} + r_Q, p}$ ,  $Z_n^{(2)} \in \mathbb{R}^{p - r_{[P;Q]} + r_Q, q}$ ,
- ii)  $\begin{bmatrix} S_n \\ Z_n^{(1)} \end{bmatrix} \in \mathbb{R}^{p,p}$  is orthogonal, and  $Z_n^{(1)} P_n + Z_n^{(2)} Q_n = 0$ ,
- iii)  $S_n P_n$  has full row rank, and the pair  $(S_n P_n, Q_n)$  has no hidden redundancy.

PROOF. First, using a Singular Value Decomposition (SVD) we factorize  $Q_n$  and then partition  $P_n$  conformably to get

$$U_1^T Q_n V_1 = \begin{bmatrix} \Sigma_n & 0 \\ 0 & 0 \end{bmatrix}, \quad \text{and} \quad P_n V_1 = \begin{bmatrix} P_{n,1} & P_{n,2} \end{bmatrix}, \quad (5)$$

where the matrices  $U_1 = \begin{bmatrix} U_{11} & U_{12} \end{bmatrix} \in \mathbb{R}^{q,q}$ ,  $V_1 = \begin{bmatrix} V_{11} & V_{12} \end{bmatrix} \in \mathbb{R}^{d,d}$  are orthogonal and  $\Sigma_n \in \mathbb{R}^{r_Q, r_Q}$  is nonsingular and diagonal. Now we use a second SVD

to factorize  $P_{n,2}$  and to find an orthogonal matrix  $U_2^T = \begin{bmatrix} S \\ Z_n^{(1)} \end{bmatrix} \in \mathbb{R}^{p,p}$  such that  $U_2^T P_{n,2} = \begin{bmatrix} P_{n,12} \\ 0 \end{bmatrix}$ , where  $P_{n,12}$  has full row rank. Thus, we obtain

$$\begin{bmatrix} S_n & 0 \\ Z_n^{(1)} & 0 \\ 0 & U_{11}^T \\ 0 & U_{12}^T \end{bmatrix} \begin{bmatrix} P_n \\ Q_n \end{bmatrix} \begin{bmatrix} V_{11} & V_{12} \end{bmatrix} = \begin{bmatrix} P_{n,11} & P_{n,12} \\ P_{n,21} & 0 \\ \Sigma_n & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} r_{[P;Q]} - r_Q \\ p - r_{[P;Q]} + r_Q \\ r_Q \\ q - r_Q \end{bmatrix}.$$

Since  $P_{n,12}$  has full row rank,  $S_n P_n = \begin{bmatrix} P_{n,11} & P_{n,12} \end{bmatrix} V_1^{-1}$  also has full row rank. Moreover, one sees that

$$\text{rank} \left( \begin{bmatrix} S_n P_n \\ Q_n \end{bmatrix} \right) = \text{rank} \left( \begin{bmatrix} 0 & P_{n,12} \end{bmatrix} \right) + \text{rank} \left( \begin{bmatrix} \Sigma_n & 0 \end{bmatrix} \right) = \text{rank}(S_n P_n) + \text{rank}(Q_n),$$

which follows that the pair  $(S_n P_n, Q_n)$  has no hidden redundancy.

Finally, setting  $Z_n^{(2)} := -P_{n,21} \Sigma_n^{-1} U_{11}^T$ , we obtain

$$Z_n^{(1)} P_n + Z_n^{(2)} Q_n = ([P_{n,21} \ 0] - P_{n,21} \Sigma_n^{-1} [\Sigma_n \ 0]) V_1^{-1} = 0,$$

96 which completes the proof.

97 **Remark 2.5.** i) In the special case, where  $P_n$  has full row rank and the pair  
 98  $(P_n, Q_n)$  has no hidden redundancy, we will adapt the notation of an empty  
 99 matrix and take  $S_n = I_p$ ,  $Z_n^{(1)} = [\ ]_{0,p}$ ,  $Z_n^{(2)} = [\ ]_{0,q}$ .  
 100 ii) Furthermore, we notice that the matrices  $U_1$ ,  $U_2$ ,  $V_1$  in the proof of Lemma  
 101 2.4 are orthogonal. Therefore, in case that the smallest singular value of  $Q_n$  and  
 102 the largest one do not differ very much in size, then  $\Sigma_n^{-1}$  is well-conditioned, and  
 103 hence we can stably compute the matrix  $Z_n^{(2)}$ . Both matrices  $Z_n^{(1)}$  and  $Z_n^{(2)}$  will  
 104 play the key role in our *index reduction procedure* presented in the next section.

105 For any given matrix  $M$ , by  $M^T$  we denote its transpose. By  $T_0(M)$  we  
 106 denote an orthogonal matrix whose columns span the left null space of  $M$ . By  
 107  $T_\perp(M)$  we denote an orthogonal matrix whose columns span the vector space  
 108  $\text{range}(M)$ . From basic linear algebra, we have the following three lemmas.

**Lemma 2.6.** *The matrix  $\begin{bmatrix} T_\perp^T(M) \\ T_0^T(M) \end{bmatrix}$  is nonsingular, the matrix  $T_\perp^T(M) M$  has full row rank, and the following identity holds*

$$\begin{bmatrix} T_\perp^T(M) \\ T_0^T(M) \end{bmatrix} M = \begin{bmatrix} T_\perp^T(M) & M \\ 0 & \end{bmatrix}.$$

109 **PROOF.** A simple proof can be found, for example, in [6].

**Lemma 2.7.** Given four matrices  $\check{A}, \check{B}, \check{C}$  in  $\mathbb{R}^{m,d}$  and  $\check{D}$  in  $\mathbb{R}^{m,p}$ . Let us consider the following matrices whose columns span orthogonal bases of the associated vector spaces

$$\begin{array}{llll} T_1 & \text{basis of } \ker(\check{A}^T), & \text{and} & T_{1,\perp} \text{ basis of } \text{range}(\check{A}), \\ W_1 & \text{basis of } \ker(T_1^T \check{D})^T, & \text{and} & W_{1,\perp} \text{ basis of } \text{range}(T_1^T \check{D}), \\ & & & J_D := W_{1,\perp}^T T_1^T \check{D}, \\ J_{B_1} & := W_1^T T_1^T \check{B}, & \text{and} & J_{B_2} := W_{1,\perp}^T T_{1,\perp}^T \check{B}, \\ J_{C_1} & := W_1^T T_1^T \check{C}, & \text{and} & J_{C_2} := W_{1,\perp}^T T_{1,\perp}^T \check{C}, \\ T_2 & \text{basis of } \ker(J_{B_1}^T), & \text{and} & T_{2,\perp} \text{ basis of } \text{range}(J_{B_1}), \\ T_3 & \text{basis of } \ker(J_{B_2}^T), & \text{and} & T_{3,\perp} \text{ basis of } \text{range}(J_{B_2}), \\ T_4 & \text{basis of } \ker(T_2^T J_{C_1})^T, & \text{and} & T_{4,\perp} \text{ basis of } \text{range}(T_2^T J_{C_1}). \end{array}$$

110 Then, the following assertions hold true.

- 111 i) The matrices  $\begin{bmatrix} T_{i,\perp} \\ T_i \end{bmatrix}$ ,  $i = 1, \dots, 4$ ,  $\begin{bmatrix} W_{1,\perp} \\ W_1 \end{bmatrix}$  are orthogonal.
- 112 ii) The matrices  $T_{1,\perp}^T \check{A}$ ,  $T_{2,\perp}^T J_{B_1}$ ,  $T_{3,\perp}^T J_{B_2}$ ,  $T_{4,\perp}^T T_2^T J_{C_1}$ , and  $J_D$  have full row
- 113 rank.
- iii) Moreover, there exists an orthogonal matrix  $\check{U}$  such that

$$\check{U} \begin{bmatrix} \check{A} & \check{B} & \check{C} & | & \check{D} \end{bmatrix} = \begin{bmatrix} \check{A}_1 & \check{B}_1 & \check{C}_1 & | & \check{D}_1 \\ 0 & \check{B}_2 & \check{C}_2 & | & 0 \\ 0 & 0 & \check{C}_3 & | & 0 \\ 0 & 0 & 0 & | & 0 \\ \hline 0 & \check{B}_4 & \check{C}_4 & | & \check{D}_4 \\ 0 & 0 & \check{C}_5 & | & \check{D}_5 \end{bmatrix}, \quad (6)$$

114 where the matrices  $\check{A}_1, \check{B}_2, \check{B}_4, \check{C}_3, \begin{bmatrix} \check{D}_4 \\ \check{D}_5 \end{bmatrix}$  have full row rank.

PROOF. The first two claims followed directly from Lemma 2.6. To prove the third claim, we construct the desired matrix  $\check{U}$  as follows

$$\check{U} := \begin{bmatrix} I & & & \\ & I & & \\ & & T_{4,\perp}^T & \\ & & T_4^T & \\ & & & I \end{bmatrix} \cdot \begin{bmatrix} I & & \\ & T_{2,\perp}^T & \\ & T_2^T & \\ & & T_{3,\perp}^T \\ & & T_3^T \end{bmatrix} \cdot \begin{bmatrix} I & \\ & W_{1,\perp}^T \end{bmatrix} \cdot \begin{bmatrix} T_{1,\perp}^T \\ T_1^T \end{bmatrix}.$$

Thus, we have that

$$\check{U} \begin{bmatrix} \check{A} & \check{B} & \check{C} & | & \check{D} \end{bmatrix} = \begin{bmatrix} T_{1,\perp}^T \check{A} & T_{1,\perp}^T \check{B} & T_{1,\perp}^T \check{C} & | & T_{1,\perp}^T \check{D} \\ 0 & T_{2,\perp}^T J_{B_1} & T_{2,\perp}^T J_{C_1} & | & 0 \\ 0 & 0 & T_{4,\perp}^T T_2^T J_{C_1} & | & 0 \\ 0 & 0 & 0 & | & 0 \\ \hline 0 & T_{3,\perp}^T J_{B_2} & T_{3,\perp}^T J_{C_2} & | & T_{3,\perp}^T J_D \\ 0 & 0 & T_3^T J_{C_2} & | & T_3^T J_D \end{bmatrix}.$$

Due to the parts i) and ii), we see that this is exactly the desired form (6).

**Lemma 2.8.** *Let  $P \in \mathbb{R}^{p,d}$ ,  $Q \in \mathbb{R}^{q,d}$  be two full row rank matrices, where  $p + q \leq d$ . Then, the following assertions hold true.*

- i) *There exists a matrix  $F \in \mathbb{R}^{d,d}$  such that  $H := \begin{bmatrix} P \\ QF \end{bmatrix}$  has full row rank.*
- ii) *For any  $G \in \mathbb{R}^{q,d}$ , there exists a matrix  $F \in \mathbb{R}^{d,d}$  such that  $\begin{bmatrix} P \\ G + QF \end{bmatrix}$  has full row rank.*

PROOF. i) First we consider the SVDs of  $P$  and  $Q$  that reads

$$U_P P V_P = [\Sigma_P \quad 0_{p,d-p}], \quad U_Q Q V_Q = [\Sigma_Q \quad 0_{q,d-q}],$$

where  $\Sigma_P, \Sigma_Q$  are nonsingular, diagonal matrices, and  $0_{p,d-p}$  (resp.  $0_{q,d-q}$ ) are the zero matrix of size  $p$  by  $d-p$  (resp.  $q$  by  $d-q$ ).

By choosing  $F := V_Q \begin{bmatrix} 0 & I_q \\ I_{d-q} & 0 \end{bmatrix} V_P^T$  we see that

$$\begin{bmatrix} U_P & 0 \\ 0 & U_Q \end{bmatrix} \begin{bmatrix} P \\ QF \end{bmatrix} V_P = \begin{bmatrix} U_P P V_P \\ U_Q Q F V_P \end{bmatrix} = \begin{bmatrix} \Sigma_P & 0_{p,d-p-q} & 0_{p,q} \\ 0_{q,p} & 0_{p,d-p-q} & \Sigma_Q \end{bmatrix},$$

and hence, the claim i) is proven.

ii) Clearly, in case that the matrix  $F$  is very big, then  $G$  is only a small perturbation, and hence for sufficiently large  $\eta$ , by choosing

$$F := \eta V_Q \begin{bmatrix} 0 & I_q \\ I_{d-q} & 0 \end{bmatrix} V_P^T,$$

we obtain the full row rank property of  $\begin{bmatrix} P \\ G + QF \end{bmatrix}$ .

**Remark 2.9.** It should be noted that, the proof of Lemmas 2.7 and 2.8 are constructive, and all the matrices  $T_{i,\perp}$ ,  $T_i$ ,  $i = 1, \dots, 4$ ,  $W_{1,\perp}$ ,  $W_1$  and  $F$  can be stably computed.

### 3. Strangeness-index of second-order SiDEs

In this section, we study the solvability analysis of the second-order SiDE (2) and of its corresponding IVP (2)–(3). Many regularization procedures and their associated index concepts have been proposed for first order systems, see the survey [15] and the references therein. Nevertheless, for second order systems, only the strangeness-index has been proposed for only continuous but not discrete-time systems in [16, 22]. Thus, it is our purpose to construct a comparable regularization and index concept for system (2).

Let

$$M_n := [A_n \quad B_n \quad C_n], \quad X(n) := \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix},$$

we call  $\{M_n\}_{n \geq n_0}$  the *behavior matrix sequence* of system (2). Thus, (2) can be rewritten as

$$M_n X(n) = f(n), \text{ for all } n \geq n_0. \quad (7)$$

Clearly, by scaling (2) with a pointwise nonsingular matrix sequence  $\{P_n\}_{n \geq n_0}$  in  $\mathbb{R}^{m,m}$ , we obtain a new system

$$\begin{bmatrix} P_n A_n & P_n B_n & P_n C_n \end{bmatrix} X(n) = P_n f(n), \text{ for all } n \geq n_0, \quad (8)$$

without changing the solution space. This motivates the following definition.

**Definition 3.1.** Two behavior matrix sequences  $\{M_n = \begin{bmatrix} A_n & B_n & C_n \end{bmatrix}\}_{n \geq n_0}$  and  $\{\tilde{M}_n = \begin{bmatrix} \tilde{A}_n & \tilde{B}_n & \tilde{C}_n \end{bmatrix}\}_{n \geq n_0}$  are called (*strongly*) *left equivalent* if there exists a pointwise nonsingular matrix sequence  $\{P_n\}_{n \geq n_0}$  such that  $\tilde{M}_n = P_n M_n$  for all  $n \geq n_0$ . We denote this equivalence by  $\{M_n\}_{n \geq n_0} \stackrel{\ell}{\sim} \{\tilde{M}_n\}_{n \geq n_0}$ . If this is the case, we also say that two SiDEs (2), (8) are left equivalent.

**Lemma 3.2.** Consider the behavior matrix sequence  $\{M_n\}_{n \geq n_0}$  of system (2). Then, for all  $n \geq n_0$ , we have that

$$\{M_n\}_{n \geq n_0} \stackrel{\ell}{\sim} \left\{ \begin{bmatrix} A_{n,1} & B_{n,1} & C_{n,1} \\ 0 & B_{n,2} & C_{n,2} \\ 0 & 0 & C_{n,3} \\ 0 & 0 & 0 \end{bmatrix} \right\}_{n \geq n_0}, \quad \begin{matrix} r_{2,n} \\ r_{1,n} \\ r_{0,n} \\ v_n \end{matrix} \quad (9)$$

where the matrices  $A_{n,1}$ ,  $B_{n,2}$ ,  $C_{n,3}$  on the main diagonal have full row rank. Here the numbers  $r_{2,n}$ ,  $r_{1,n}$ ,  $r_{0,n}$ ,  $v_n$  are row-sizes of the block rows of  $M_n$ . Furthermore, these numbers are invariant under left equivalent transformations. Thus, we can call them the local characteristic invariants of the SiDE (2).

*Proof.* The block diagonal form (9) is obtained directly by consecutively compressing the block columns  $A_n$ ,  $B_n$ ,  $C_n$  of  $M_n$  via Lemma 2.6. In details, we have that

$$\begin{aligned} \text{rows of } A_{n,1} & \text{ form the basis of the space } \text{range}(A_n^T), \\ \text{rows of } B_{n,2} & \text{ form the basis of the space } \text{range}(T_0^T(A_n) B_n)^T, \\ \text{rows of } C_{n,3} & \text{ form the basis of the space } \text{range}\left(T_0^T\left(\begin{bmatrix} A_n \\ B_n \end{bmatrix}\right) C_n\right)^T. \end{aligned}$$

Moreover, from (9), we obtain the following identities

$$\begin{aligned} r_{2,n} &= \text{rank}(A_n), \\ r_{1,n} &= \text{rank}\left(\begin{bmatrix} A_n & B_n \end{bmatrix}\right) - \text{rank}(A_n), \\ r_{0,n} &= \text{rank}\left(\begin{bmatrix} A_n & B_n & C_n \end{bmatrix}\right) - \text{rank}\left(\begin{bmatrix} A_n & B_n \end{bmatrix}\right), \\ v_n &= m - r_{2,n} - r_{1,n} - r_{0,n}, \end{aligned}$$

which proves the second claim.  $\square$



145 Analogous to the continuous-time case, we will apply an *algebraic approach*  
 146 (see [2, 16]), which aims to reformulate (2) into a so-called *strangeness-free* form,  
 147 as stated in the following definition.

**Definition 3.3.** ([11]) System (2) is called *strangeness-free* if there exists a pointwise nonsingular matrix sequence  $\{P_n\}_{n \geq n_0}$  such that by scaling the SiDE (2) at each point  $n$  with  $P_n$ , we obtain a new system of the form

$$\begin{matrix} \hat{r}_2 \\ \hat{r}_1 \\ \hat{r}_0 \\ \hat{v} \end{matrix} \begin{bmatrix} \hat{A}_{n,1} \\ 0 \\ 0 \\ 0 \end{bmatrix} x(n+2) + \begin{bmatrix} \hat{B}_{n,1} \\ \hat{B}_{n,2} \\ 0 \\ 0 \end{bmatrix} x(n+1) + \begin{bmatrix} \hat{C}_{n,1} \\ \hat{C}_{n,2} \\ \hat{C}_{n,3} \\ 0 \end{bmatrix} x(n) = \begin{bmatrix} \hat{f}_1(n) \\ \hat{f}_2(n) \\ \hat{f}_3(n) \\ \hat{f}_4(n) \end{bmatrix}, \text{ for all } n \geq n_0, \quad (10)$$

148 where the matrix  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}$  has full row rank for all  $n \geq n_0$ .

149 In order to perform an algebraic approach, an additional assumption below  
 150 is usually needed.

151 **Assumption 3.4.** Assume that the local characteristic invariants  $r_{2,n}$ ,  $r_{1,n}$ ,  
 152  $r_{0,n}$  become global, i.e., they are constant for all  $n \geq n_0$ . Furthermore, assume  
 153 that two matrix sequences  $\left\{ \begin{bmatrix} A_{n,1} \\ B_{n,2} \\ C_{n,3} \end{bmatrix} \right\}_{n \geq n_0}$  and  $\left\{ \begin{bmatrix} B_{n,2} \\ C_{n,3} \end{bmatrix} \right\}_{n \geq n_0}$  have constant  
 154 rank for all  $n \geq n_0$ .

**Remark 3.5.** Following directly from the proof of Lemma 3.2, we see that Assumption 3.4 is satisfied if and only if five following constant rank conditions are satisfied

$$\begin{aligned} \text{rank}(A_n) &\equiv \text{const.}, \quad \text{rank}\begin{bmatrix} A_n & B_n \end{bmatrix} \equiv \text{const.}, \quad \text{rank}\begin{bmatrix} A_n & B_n & C_n \end{bmatrix} \equiv \text{const.}, \\ \text{rank}(T_0^T(A_n) \ B_n) &\equiv \text{const.}, \quad \text{rank}\left(T_0^T\left(\begin{bmatrix} A_n \\ B_n \end{bmatrix}\right) C_n\right) \equiv \text{const.} \end{aligned} \quad (11)$$

155 **Remark 3.6.** In system (10), the quantities  $r_2$ ,  $r_1$ , and  $r_0$  are the dimensions of  
 156 the second-order dynamics part, the first-order dynamics part, and the algebraic  
 157 (zero-order) part, respectively. Furthermore,  $r_2 + r_1$  is exactly the degree of  
 158 freedoms.

Let us call the number

$$r_u := 3r_2 + 2r_1 + r_0$$

the *upper rank* of system (2). Clearly,  $r_u$  is invariant under left equivalence transformations. Rewrite (7) block row-wise, we obtain the following system for

all  $n \geq n_0$ .

$$A_{n,1}x(n+2) + B_{n,1}x(n+1) + C_{n,1}x(n) = f_1(n), \quad r_2 \text{ equations}, \quad (12a)$$

$$B_{n,2}x(n+1) + C_{n,2}x(n) = f_2(n), \quad r_1 \text{ equations}, \quad (12b)$$

$$C_{n,3}x(n) = f_3(n), \quad r_0 \text{ equations}, \quad (12c)$$

$$0 = f_4(n), \quad v \text{ equations}. \quad (12d)$$

Since the matrices  $A_{n,1}$ ,  $B_{n,2}$ ,  $C_{n,3}$  have full row rank, the number of scalar difference equations of order 2 (resp. 1, and 0) in (2) is exactly  $r_2$  (resp.  $r_1$  and  $r_0$ ), while  $v$  is the number of redundant equations. Now we are able to define the shift-forward operator  $\Delta$ , which acts on some or whole equations of system (12). This operator maps each equation of system (12) at the time instant  $n$  to the equation itself at the time  $n+1$ , for example

$$\Delta : C_{n,3}x(n) = f_3(n) \mapsto C_{n+1,3}x(n+1) = f_3(n+1). \quad (13)$$

Clearly, under Assumption 3.4, this shift operator can be applied to equations of system (12). In order to reveal all hidden constraints of (12) we propose the idea, that for each  $j = 1, 2$ , we use equations of order less than  $j$  to reduce the number of scalar equations of order  $j$ . This task will be performed in Lemmas 3.8 and 3.9 below. In details, if the matrix pair  $(B_{n,2}, C_{n+1,3})$  has hidden redundancy then we will make use of the shifted equation (13). Analogously, if the pair  $\left(A_{n,1}, \begin{bmatrix} B_{n+1,2} \\ C_{n+2,3} \end{bmatrix}\right)$  has hidden redundancy then we will make use of the shifted equation

$$B_{n+1,2}x(n+2) + C_{n+1,2}x(n+1) = f_2(n+1), \quad (14)$$

and may be also the double shifted equation

$$C_{n+2,3}x(n+2) = f_3(n+2). \quad (15)$$

**Lemma 3.7.** *Consider the SiDE (2) and the equivalent system (12). Then, (2) has the same solution set as that of the following extended system*

$$\begin{array}{c|ccc} r_2 & A_{n,1} & B_{n,1} & C_{n,1} \\ r_1 & 0 & B_{n,2} & C_{n,2} \\ r_0 & 0 & 0 & C_{n,3} \\ v & 0 & 0 & 0 \\ \hline r_0 & 0 & C_{n+1,3} & 0 \\ r_1 & B_{n+1,2} & C_{n+1,2} & 0 \\ r_0 & C_{n+2,3} & 0 & 0 \end{array} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} = \begin{bmatrix} f_1(n) \\ f_2(n) \\ f_3(n) \\ f_4(n) \\ f_3(n+1) \\ f_2(n+1) \\ f_3(n+2) \end{bmatrix}, \quad (16)$$

for all  $n \geq n_0$ .

*Proof.* Since all equations in the lower part of (16) at any time point  $n$  is the consequence of the upper part (which is exactly (12)) at the time instants  $n+1$  and  $n+2$ , the proof is directly followed.  $\square$



- 180 i) scaling a block row equation with a nonsingular matrix,  
 181 ii) adding to one row a linear combination of some other rows.

Let the matrix sequences  $\{S_n^{(i)}\}_{n \geq n_0}$ ,  $i = 1, 2$ , and  $\{Z_n^{(j)}\}_{n \geq n_0}$ ,  $j = 1, \dots, 5$  be defined as in Lemma 3.8. Firstly, by scaling the first (resp., second) block row equation of (16) with an orthogonal matrix  $\begin{bmatrix} S_n^{(2)} \\ Z_n^{(2)} \end{bmatrix}$  (resp.,  $\begin{bmatrix} S_n^{(1)} \\ Z_n^{(1)} \end{bmatrix}$ ), we obtain an equivalent system to (12), as follows

$$\begin{array}{c|ccc} d_2 & S_n^{(2)} A_{n,1} & S_n^{(2)} B_{n,1} & S_n^{(2)} C_{n,1} \\ s_2 & Z_n^{(2)} A_{n,1} & Z_n^{(2)} B_{n,1} & Z_n^{(2)} C_{n,1} \\ \hline d_1 & 0 & S_n^{(1)} B_{n,2} & S_n^{(1)} C_{n,2} \\ s_1 & 0 & Z_n^{(1)} B_{n,2} & Z_n^{(1)} C_{n,2} \\ \hline r_0 & 0 & 0 & C_{n,3} \\ v & 0 & 0 & 0 \\ \hline r_0 & 0 & C_{n+1,3} & 0 \\ r_1 & B_{n+1,2} & C_{n+1,2} & 0 \\ r_0 & C_{n+2,3} & 0 & 0 \end{array} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} = \begin{array}{c} \begin{bmatrix} S_n^{(2)} f_1(n) \\ Z_n^{(2)} f_1(n) \end{bmatrix} \\ \begin{bmatrix} S_n^{(1)} f_2(n) \\ Z_n^{(1)} f_2(n) \end{bmatrix} \\ \begin{bmatrix} f_3(n) \\ f_4(n) \end{bmatrix} \\ \begin{bmatrix} f_3(n+1) \\ f_2(n+1) \\ f_3(n+2) \end{bmatrix} \end{array}. \quad (19)$$

By adding the seventh row scaled with  $Z_n^{(3)}$  to the fourth row of (19) and making use of (17a) we obtain the first hidden constraint

$$Z_n^{(1)} C_{n,2} x(n) = Z_n^{(1)} f_2(n) + Z_n^{(3)} f_3(n+1),$$

182 which is exactly the fourth row of (18).

We continue by adding the seventh row scaled with  $Z_n^{(4)}$  and the eighth row scaled with  $Z_n^{(5)}$  to the second row of (19) and making use of (17b) to obtain

$$\begin{aligned} & \left( Z_n^{(2)} B_{n,1} + Z_n^{(4)} C_{n+1,2} \right) x(n+1) + Z_n^{(2)} C_{n,1} x(n) \\ & = Z_n^{(2)} f_1(n) + Z_n^{(4)} f_2(n+1) + Z_n^{(5)} f_3(n+2). \end{aligned}$$

This is exactly the second row of (18). Therefore, any solution to (12) is also a solution to (18).

**Sufficiency:** Let  $x$  be an arbitrary solution to (18). Thus,  $x$  is also a solution to the shifted system

$$\begin{array}{c|ccc} d_2 & S_n^{(2)} A_{n,1} & S_n^{(2)} B_{n,1} & S_n^{(2)} C_{n,1} \\ s_2 & 0 & Z_n^{(2)} B_{n,1} + Z_n^{(4)} C_{n+1,2} & Z_n^{(2)} C_{n,1} \\ \hline d_1 & 0 & S_n^{(1)} B_{n,2} & S_n^{(1)} C_{n,2} \\ s_1 & 0 & 0 & Z_n^{(1)} C_{n,2} \\ \hline r_0 & 0 & 0 & C_{n,3} \\ v & 0 & 0 & 0 \\ \hline r_0 & 0 & C_{n+1,3} & 0 \\ r_0 & C_{n+2,3} & 0 & 0 \end{array} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} =$$

$$= \begin{bmatrix} S_n^{(2)} f_1(n) \\ Z_n^{(2)} f_1(n) + Z_n^{(4)} f_2(n+1) + Z_n^{(5)} f_3(n+2) \\ S_n^{(1)} f_2(n) \\ Z_n^{(1)} f_2(n) + Z_n^{(3)} f_3(n+1) \\ f_3(n) \\ f_4(n) \\ f_3(n+1) \\ f_3(n+2) \end{bmatrix}, \text{ for all } n \geq n_0. \quad (20)$$

183 Since elementary matrix row operations are reversible, we can reverse the trans-  
 184 formations performed in the necessity part. Consequently, we see that any so-  
 185 lution to (20) is also a solution to (19), and hence, this completes the proof.  $\square$

186 Consider system (18), we see that the upper rank of the behavior matrix is

$$\begin{aligned} r_u^{new} &\leq 3d_2 + 2(s_2 + d_1) + (s_1 + r_0) \\ &= 3(r_2 - s_2) + 2(s_2 + r_1 - s_1) + (s_1 + r_0) \\ &= r - (s_2 + s_1) \leq r. \end{aligned}$$

187 In conclusion, after performing a so-called *index reduction step*, which passes  
 188 from (12) to (18), we have reduced the upper rank  $r_u$  at least by  $s_2 + s_1$ .  
 189 Continue in this fashion until  $s_1 = s_2 = 0$ , we obtain the following algorithm.

---

**Algorithm 1** Index reduction steps for SiDEs at the time point  $n$

---

**Input:** The SiDE (2) and its behavior form (7).

**Output:** A strangeness-free SiDE of the form (10) and the strangeness-index  $\mu$ .

- 1: Set  $i = 0$ .
- 2: Transform the behavior matrix  $[A_n \ B_n \ C_n]$  to the block upper triangular form

$$\tilde{M}_n := \begin{bmatrix} A_{n,1} & B_{n,1} & C_{n,1} \\ 0 & B_{n,2} & C_{n,2} \\ 0 & 0 & C_{n,3} \\ 0 & 0 & 0 \end{bmatrix},$$

where all the matrices  $A_{n,1}$ ,  $B_{n,2}$ ,  $C_{n,3}$  on the main diagonal have full row rank. The system now takes the form (12).

- 3: **if** both matrix pairs  $\left(A_{n,1}, \begin{bmatrix} B_{n+1,2} \\ C_{n+2,3} \end{bmatrix}\right)$  and  $(B_{n,2}, C_{n+1,3})$  have no hidden redundancy **then** set  $\mu = i$  and **STOP**.
  - 4: **else** set  $i := i + 1$
  - 5: Find the matrices  $S_n^{(j)}$ ,  $j = 1, 2$ , and  $Z_n^{(j)}$ ,  $j = 1, \dots, 5$  as in Lemma 3.8.
  - 6: Transform the system to the new form (18) as in Lemma 3.9.
  - 7: **end if**
  - 8: Go back to Step 2 with the updated behavior matrix.
-

190 After each index reduction step the upper rank  $r_u^i$  has been decreased at  
 191 least by  $s_2^i + s_1^i$ , so Algorithm 1 terminates after a finite number  $\mu$  of iterations,  
 192 which will be called the *strangeness-index* of the SiDE (2).

**Theorem 3.10.** *Consider the SiDE (2) and assume that Assumption 3.4 is satisfied for any  $n$  and any  $i$  considered within the loop, such that the strangeness-index  $\mu$  is well-defined by Algorithm 1. Then, the SiDE (2) has the same solution set as the strangeness-free SiDE*

$$\begin{matrix} r_2^\mu \\ r_1^\mu \\ r_0^\mu \\ v^\mu \end{matrix} \begin{bmatrix} \hat{A}_{n,1} & \hat{B}_{n,1} & \hat{C}_{n,1} \\ 0 & \hat{B}_{n,2} & \hat{C}_{n,2} \\ 0 & 0 & \hat{C}_{n,3} \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} = \begin{bmatrix} \hat{g}_1(n) \\ \hat{g}_2(n) \\ \hat{g}_3(n) \\ \hat{g}_4(n) \end{bmatrix}, \text{ for all } n \geq n_0, \quad (21)$$

193 where the matrix  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}$  has full row rank for all  $n \geq n_0$ . Here  $\hat{g}_2$  and  $\hat{g}_3$   
 194 consist of the components of  $f(n), f(n+1), \dots, f(n+2\mu)$  (at most).

195 *Proof.* The proof is a direct consequence of Algorithm 1, where the matrix  
 196  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}$  has full row rank due to Lemma 2.3. □

197 To illustrate Algorithm 1, we consider the following example.

**Example 3.11.** Given a parameter  $\alpha \in \mathbb{R}$ , we consider the second order SiDE

$$\begin{bmatrix} 1 & n+1 & n+4 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} x(n+2) + \begin{bmatrix} 0 & \alpha & 2n+3 \\ 1 & n & 1 \\ 0 & 0 & 0 \end{bmatrix} x(n+1) + \begin{bmatrix} 0 & n+1 & 0 \\ 0 & 0 & n \\ 0 & 0 & n+1 \end{bmatrix} x(n) = \begin{bmatrix} f_1(n) \\ f_2(n) \\ f_3(n) \end{bmatrix}, \quad (22)$$

for all  $n \geq 0$ . Fortunately, the behavior matrix

$$M = \left[ \begin{array}{ccc|ccc|ccc} 1 & n+1 & n+4 & 0 & \alpha & 2n+3 & 0 & n+1 & 0 \\ 0 & 0 & 0 & 1 & n & 1 & 0 & 0 & n \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & n+1 \end{array} \right] = \begin{bmatrix} A_{n,1} & B_{n,1} & C_{n,1} \\ 0 & B_{n,2} & C_{n,2} \\ 0 & 0 & C_{n,3} \end{bmatrix}$$

is already in the block diagonal form, so we do not need to perform Step 2 in Algorithm 1. Furthermore, all constant rank conditions required in Assumption 3.4 are satisfied. We observe that

$$\begin{aligned} B_{n+1,2} &= \begin{bmatrix} 1 & n+1 & 1 \end{bmatrix}, & C_{n+1,2} &= \begin{bmatrix} 0 & 0 & n+1 \end{bmatrix}, \\ C_{n+1,3} &= \begin{bmatrix} 0 & 0 & n+2 \end{bmatrix}, & C_{n+2,3} &= \begin{bmatrix} 0 & 0 & n+3 \end{bmatrix}. \end{aligned}$$

By directly verifying, we see that the matrix pair  $\left( A_{n,1}, \begin{bmatrix} B_{n+1,2} \\ C_{n+2,3} \end{bmatrix} \right)$  has hidden redundancy, while the pair  $(B_{n,2}, C_{n+1,3})$  does not. Due to Lemma 3.8 we

choose  $S_n^{(2)} = \begin{bmatrix} & & \end{bmatrix}$ ,  $Z_n^{(2)} = 1$ ,  $Z_n^{(4)} = -1$ ,  $Z_n^{(5)} = -1$ . Notice that the fact  $Z_n^{(5)}$  is non-empty leads to the appearance of  $f_3(n+2)$ . Furthermore, the resulting system (18) reads

$$\begin{bmatrix} 0 & \alpha & n+2 \\ 1 & n & 1 \\ 0 & 0 & 0 \end{bmatrix} x(n+1) + \begin{bmatrix} 0 & n+1 & 0 \\ 0 & 0 & n \\ 0 & 0 & n+1 \end{bmatrix} x(n) = \begin{bmatrix} f_1(n) - f_2(n+1) - f_3(n+2) \\ f_2(n) \\ f_3(n) \end{bmatrix}. \quad (23)$$

Here the leading coefficient matrix associated with  $x(n+2)$  becomes zero, so for notational convenience we do not write this term. Go back to Step 3, we see that two following cases may happen.

i) If  $\alpha \neq 0$ , then Algorithm 1 terminates here, and the strangeness-index is  $\mu = 1$ . Here the number of time-shift appear in the inhomogeneity  $f$  in the strangeness-free formulation (23) is 2.

ii) If  $\alpha = 0$ , then the matrix pair  $\left( \begin{bmatrix} 0 & \alpha & n+2 \\ 1 & n & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & n+2 \end{bmatrix} \right)$  have hidden redundancy. Due to Lemma 3.8 we choose  $S_n^{(1)} = \begin{bmatrix} 1 & 0 \end{bmatrix}$ ,  $Z_n^{(1)} = \begin{bmatrix} 0 & 1 \end{bmatrix}$ ,  $Z_n^{(3)} = -\begin{bmatrix} 0 & 1 \end{bmatrix}$ . The resulting system (18) now reads

$$\begin{bmatrix} 1 & n & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} x(n+1) + \begin{bmatrix} 0 & 0 & n \\ 0 & n+1 & 0 \\ 0 & 0 & n+1 \end{bmatrix} x(n) = \begin{bmatrix} f_2(n) \\ f_1(n) - f_2(n+1) - f_3(n+2) - f_3(n+1) \\ f_3(n) \end{bmatrix}. \quad (24)$$

Algorithm 1 terminates here, and the strangeness-index is  $\mu = 2$ . However, the number of time-shifts appearing in the inhomogeneity  $f$  in the strangeness-free formulation (24) remains 2.

As a direct consequence of Theorem 3.10, we obtain the solvability for (2) as follows.

**Corollary 3.12.** *Under the assumption of Theorem 3.10, the following statements hold true.*

- i) *The corresponding IVP for the SiDE (2) is solvable if and only if either  $v^\mu = 0$  or  $\hat{g}_4(n) = 0$  for all  $n \geq n_0$ . Furthermore, it is uniquely solvable if, in addition, we have  $d = m - v^\mu$ .*
- ii) *The initial condition (3) is consistent if and only if the following equalities hold.*

$$\begin{aligned} \hat{B}_{n_0,2}x_1 + \hat{C}_{n_0,2}x_0 &= \hat{g}_2(n_0), \\ \hat{C}_{n_0,3}x_0 &= \hat{g}_3(n_0). \end{aligned}$$

Another direct consequence of Theorem 5.3 is that we can obtain an inherent regular difference equation as follows.

**Corollary 3.13.** *Assume that the IVP (2)-(3) is uniquely solvable for any consistent initial condition. Under the assumption of Theorem 3.10, the solution  $x$  to this IVP is also a solution to the (implicit) inherent regular difference equation*

$$\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix} x(n+2) + \begin{bmatrix} \hat{B}_{n,1} \\ \hat{C}_{n+1,2} \\ 0 \end{bmatrix} x(n+1) + \begin{bmatrix} \hat{C}_{n,1} \\ 0 \\ 0 \end{bmatrix} x(n) = \begin{bmatrix} \hat{g}_1(n) \\ \hat{g}_2(n+1) \\ \hat{g}_3(n+2) \end{bmatrix}, \quad (25)$$

212 where the matrix  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}$  is invertible for all  $n \geq n_0$ .

213 **Remark 3.14.** Unlike in [2, 12, 16], we do not change the variable  $x$ . This trick  
214 permits us to simplify significantly the condensed forms in these references. We  
215 emphasize that as in (11), we only require five constant rank conditions within  
216 one step of index reduction, instead of seven as in [16]. Therefore, this trick  
217 will enlarge the domain of application for SiDEs (and also for DAEs, in the  
218 continuous-time case). This trick is also useful for the control analysis of the  
219 descriptor system (1), as will be seen later.

220 **Remark 3.15.** i) Within one loop of Algorithm 1, for each  $n$ , we have used 4  
221 SVDs to remove the hidden redundancies in two matrix pairs. The total cost  
222 depends on the problems itself, i. e., depending on sizes of the matrix pairs  
223 which applied SVDs. Nevertheless, it does not exceed  $\mathcal{O}(m^2 d^2)$ .

224 ii) Unfortunately, since  $Z_n^{(3)}, Z_n^{(4)}, Z_n^{(5)}$  are not orthogonal, in general Algorithm  
225 1 could not be stably implemented. For the numerical solution to the IVP  
226 (2)-(3), we will consider a suitable numerical scheme in Section 5.

## 227 4. Regularization of second order descriptor systems

228 Based on the index reduction procedure for SiDEs in Section 3, in this sec-  
229 tion we construct the strangeness-index concept for the descriptor system (1).  
230 The solvability analysis for first order descriptor systems with variable coef-  
231 ficients have been carefully discussed in [3, 10, 18]. Nevertheless, for second  
232 order descriptor systems, this problem has been rarely considered. We refer the  
233 interested readers to [12, 22] for continuous-time systems.

234 It is well known, that in regularization procedures of continuous-time sys-  
235 tems, one should avoid differentiating equations that involve an input function,  
236 due to the fact that it may not be differentiable. Here, we will also keep this  
237 spirit, and hence, will not shift any equation that involve an input function,  
238 since it may destroy the causality of the considered system, as in Example 2.1.  
239 Instead of it, we will also incorporate proportional state and first order feedback  
240 within each index reduction step of the regularization procedure, as will be seen  
241 later. In the following lemma, we give the condensed form for system (1).



**Lemma 4.1.** Consider the descriptor system (1). Then, there exist two point-wise nonsingular matrix sequences  $\{U_n\}_{n \geq n_0}$ ,  $\{V_n\}_{n \geq n_0}$  such that the following identities hold.

$$(U_n [A_n \ B_n \ C_n], U_n D_n V_n) = \left( \left[ \begin{array}{ccc|ccc} A_{n,1} & B_{n,1} & C_{n,1} & D_{n,1} & 0 & 0 \\ 0 & B_{n,2} & C_{n,2} & 0 & 0 & 0 \\ 0 & 0 & C_{n,3} & 0 & 0 & 0 \\ \hline 0 & B_{n,4} & C_{n,4} & 0 & \Sigma_{\varphi,1} & 0 \\ 0 & 0 & C_{n,5} & 0 & 0 & \Sigma_{\varphi,0} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right], \begin{array}{c} r_{2,n} \\ r_{1,n} \\ r_{0,n} \\ \varphi_{1,n} \\ \varphi_{0,n} \\ v_n \end{array} \right) \text{ for all } n \geq n_0. \quad (26)$$

Here sizes of the block rows are  $r_{2,n}$ ,  $r_{1,n}$ ,  $r_{0,n}$ ,  $\varphi_{1,n}$ ,  $\varphi_{0,n}$ ,  $v_n$ , the matrices  $A_{n,1}$ ,  $B_{n,2}$ ,  $B_{n,4}$ ,  $C_{n,3}$  are of full row rank and the matrices  $\Sigma_{\varphi,1}$ ,  $\Sigma_{\varphi,0}$  are nonsingular and diagonal.

*Proof.* First we apply Lemma 2.7 to four matrices  $A_n$ ,  $B_n$ ,  $C_n$  and  $D_n$  to obtain the matrix  $U_n$  that satisfies (6). Decompose the matrix  $\begin{bmatrix} \check{D}_4 \\ \check{D}_5 \end{bmatrix}$  via one SVD, we then obtain the block  $\begin{bmatrix} 0 & \Sigma_{\varphi,1} & 0 \\ 0 & 0 & \Sigma_{\varphi,0} \end{bmatrix}$ . Finally, we use Gaussian elimination to cancel out all non-zero matrices on the two columns of  $\check{D}$  that contain  $\Sigma_{\varphi,1}$  and  $\Sigma_{\varphi,0}$ , and hence, we obtain the desired form (26).  $\square$

In order to build an index reduction procedure for (1), we also need the following assumption.

**Assumption 4.2.** Assume that the local characteristic invariants  $r_{2,n}$ ,  $r_{1,n}$ ,  $r_{0,n}$ ,  $\varphi_{1,n}$ ,  $\varphi_{0,n}$ ,  $v_n$ , become global, i.e., they are constant for all  $n \geq n_0$ .

Making use of Lemma 4.1, we can transform the descriptor system (1) to the following system

$$\begin{array}{c} r_2 \\ r_1 \\ r_0 \\ \varphi_1 \\ \varphi_0 \\ v \end{array} \begin{array}{ccc|ccc} A_{n,1} & B_{n,1} & C_{n,1} & D_{n,1} & 0 & 0 \\ 0 & B_{n,2} & C_{n,2} & 0 & 0 & 0 \\ 0 & 0 & C_{n,3} & 0 & 0 & 0 \\ \hline 0 & B_{n,4} & C_{n,4} & 0 & \Sigma_{\varphi,1} & 0 \\ 0 & 0 & C_{n,5} & 0 & 0 & \Sigma_{\varphi,0} \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} + \begin{array}{ccc|ccc} D_{n,1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & \Sigma_{\varphi,1} & 0 & 0 & 0 & 0 \\ 0 & 0 & \Sigma_{\varphi,0} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \begin{bmatrix} v_1(n) \\ v_2(n) \\ v_3(n) \end{bmatrix} = \tilde{f}(n), \quad (27)$$

where  $u(n) = V_n v(n)$ ,  $v(n) := \begin{bmatrix} v_1(n) \\ v_2(n) \\ v_3(n) \end{bmatrix}$ ,  $\tilde{f}(n) := U_n f(n)$ , for all  $n \geq n_0$ .

Moreover, we notice that the third and fourth block rows, whose sizes are  $\varphi_1$  and  $\varphi_0$ , are related to the feedback regularization of (1), as shown in the following proposition.

**Proposition 4.3.** *i) Assume that for each  $n \geq n_0$ , the matrix  $\begin{bmatrix} A_{n,1} \\ B_{n+1,2} \\ C_{n+2,3} \end{bmatrix}$  is of full row rank. Then, there exist two matrices sequences  $\{F_{n,1}\}_{n \geq n_0}$ ,  $\{F_{n,0}\}_{n \geq n_0}$  which take values in  $\mathbb{R}^{\varphi_1, d}$  and  $\mathbb{R}^{\varphi_0, d}$ , respectively, such that the following matrix has full row rank for all  $n \geq n_0$*

$$\begin{bmatrix} A_{n,1} \\ B_{n+1,2} \\ C_{n+2,3} \\ \hline B_{n+1,4} + \begin{bmatrix} 0 & \Sigma_{\varphi,1} & 0 \end{bmatrix} F_{n+1,1} \\ C_{n+2,5} + \begin{bmatrix} 0 & 0 & \Sigma_{\varphi,0} \end{bmatrix} F_{n+2,0} \end{bmatrix}.$$

*ii) Consequently, if the upper part of (27) is strangeness-free then there exists a first order feedback of the form*

$$v(n) = F_{n,1}x(n+1) + F_{n,0}x(n), \text{ for all } n \geq n_0, \quad (28)$$

*such that the closed loop system*

$$A_n x(n+2) + (B_n + D_n F_{n,1}) x(n+1) + (C_n + D_n F_{n,0}) x(n) = f(n),$$

259 *is strangeness-free.*

260 *Proof.* Since the part ii) is a direct consequence of part i), we only need to prove

261 i). The part i) is directly followed by applying Lemma 2.8 for  $P = \begin{bmatrix} A_{n,1} \\ B_{n+1,2} \\ C_{n+2,3} \end{bmatrix}$ ,  
 262  $Q = \begin{bmatrix} 0 & \Sigma_{\varphi,1} & 0 \\ 0 & 0 & \Sigma_{\varphi,0} \end{bmatrix}$  and  $G = \begin{bmatrix} B_{n+1,4} \\ C_{n+2,5} \end{bmatrix}$ .  $\square$

263 From Proposition 4.3, we see that we only need to remove the hidden redun-  
 264 dancies in the upper part of (27) as follows. By performing one index reduction  
 265 step for the upper part of (27), as in Section 3, we obtain the following lemma.

**Lemma 4.4.** *Assume that the upper part of the descriptor system (27) is not strangeness-free. Then, for each input sequence  $\{v(n)\}_{n \geq n_0}$ , it has exactly the same solution set as the following system*

$$\begin{array}{c} \tilde{r}_2 \\ \tilde{r}_1 \\ \tilde{r}_0 \\ \varphi_1 \\ \varphi_0 \\ \tilde{v} \end{array} \begin{array}{c|ccc} \tilde{A}_{n,1} & \tilde{B}_{n,1} & \tilde{C}_{n,1} \\ \hline 0 & \tilde{B}_{n,2} & \tilde{C}_{n,2} \\ 0 & 0 & \tilde{C}_{n,3} \\ \hline 0 & B_{n,4} & C_{n,4} \\ 0 & 0 & C_{n,5} \\ 0 & 0 & 0 \end{array} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} + \begin{array}{c|ccc} \tilde{D}_{n,1} & 0 & 0 \\ \hline 0 & 0 & 0 \\ 0 & 0 & 0 \\ \hline 0 & \Sigma_{\varphi,1} & 0 \\ 0 & 0 & \Sigma_{\varphi,0} \\ 0 & 0 & 0 \end{array} \begin{bmatrix} v_1(n) \\ v_2(n) \\ v_3(n) \end{bmatrix} = \tilde{f}(n), \quad (29)$$

266 *for all  $n \geq n_0$ . Here  $\tilde{r}_2 = r_2 - s_2$ ,  $\tilde{r}_1 = r_1 + s_2 - s_1$ ,  $\tilde{r}_0 = r_0 + s_1$ ,  $\tilde{v} \geq v$ , for*  
 267 *some  $s_2 > 0$ ,  $s_1 > 0$ . Furthermore, both pairs  $\left(\tilde{A}_{n,1}, \begin{bmatrix} \tilde{B}_{n,2} \\ \tilde{C}_{n,3} \end{bmatrix}\right)$  and  $\left(\tilde{B}_{n,2}, \tilde{C}_{n,3}\right)$*   
 268 *have no hidden redundancy.*

269 *Proof.* System (29) is directly obtained by applying Lemma 3.9 to the upper  
 270 part of (27). To keep the brevity of this paper, we will omit the details here.  $\square$

271 Similar to the observation made in Section 3, here we also see that an *index*  
 272 *reduction step*, which passes system (27) to the new form (29) has reduced the  
 273 upper rank  $r^u$  by at least  $s_2 + s_1$ . Continue in this way until  $s_2 = s_1 = 0$ , finally  
 274 we obtain a strangeness-free descriptor system in the next theorem.

**Theorem 4.5.** *Consider the descriptor system (1). Furthermore, assume that Assumption 4.2 is fulfilled whenever needed. Then, for each fixed input sequence  $\{u(n)\}_{n \geq n_0}$ , system (1) has the same solution set as the so-called strangeness-free descriptor system*

$$\begin{array}{c} \hat{r}_2 \\ \hat{r}_1 \\ \hat{r}_0 \\ \hat{\varphi}_1 \\ \hat{\varphi}_0 \\ \hat{v} \end{array} \begin{array}{c} \left[ \begin{array}{ccc} \hat{A}_{n,1} & \hat{B}_{n,1} & \hat{C}_{n,1} \\ 0 & \hat{B}_{n,2} & \hat{C}_{n,2} \\ 0 & 0 & \hat{C}_{n,3} \end{array} \right] \begin{array}{c} x(n+2) \\ x(n+1) \\ x(n) \end{array} + \begin{array}{c} \hat{D}_{n,1} \\ 0 \\ 0 \\ \hat{D}_{n,4} \\ \hat{D}_{n,5} \\ 0 \end{array} u(n) = \begin{array}{c} \left[ \begin{array}{c} \hat{f}_1(n) \\ \hat{f}_2(n) \\ \hat{f}_3(n) \\ \hat{f}_4(n) \\ \hat{f}_5(n) \\ \hat{f}_6(n) \end{array} \right], \text{ for all } n \geq n_0, \end{array}$$

(30)

275 where the matrices  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}, \begin{bmatrix} \hat{D}_{n,4} \\ \hat{D}_{n,5} \end{bmatrix}$  have full row rank for all  $n \geq n_0$ .

*Proof.* By repeating index reduction steps until the upper rank  $r^u$  stop decreasing, we obtain the system

$$\begin{array}{c} \hat{r}_2 \\ \hat{r}_1 \\ \hat{r}_0 \\ \hat{\varphi}_1 \\ \hat{\varphi}_0 \\ \hat{v} \end{array} \begin{array}{c} \left[ \begin{array}{ccc} \hat{A}_{n,1} & \hat{B}_{n,1} & \hat{C}_{n,1} \\ 0 & \hat{B}_{n,2} & \hat{C}_{n,2} \\ 0 & 0 & \hat{C}_{n,3} \end{array} \right] \begin{array}{c} x(n+2) \\ x(n+1) \\ x(n) \end{array} + \begin{array}{c} \left[ \begin{array}{ccc} \hat{D}_{n,11} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right] \\ \begin{array}{ccc} 0 & \Sigma_{\hat{\varphi}_1} & 0 \\ 0 & 0 & \Sigma_{\hat{\varphi}_0} \\ 0 & 0 & 0 \end{array} \end{array} v(n) = \begin{array}{c} \left[ \begin{array}{c} \hat{f}_1(n) \\ \hat{f}_2(n) \\ \hat{f}_3(n) \\ \hat{f}_4(n) \\ \hat{f}_5(n) \\ \hat{f}_6(n) \end{array} \right], \end{array}$$

for all  $n \geq n_0$ , where the matrix  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}$  has full row rank for all  $n \geq n_0$ . Here

the new input sequence  $\{v(n)\}_{n \geq n_0}$  satisfies  $u(n) = V_n v(n)$ ,  $V_n$  is nonsingular for all  $n \geq n_0$ . Transform back  $v(n) = V_n^{-1} u(n)$ , and set

$$\begin{array}{c} \hat{D}_{n,1} \\ 0 \\ 0 \\ \hat{D}_{n,4} \\ \hat{D}_{n,5} \\ 0 \end{array} := \begin{array}{c} \left[ \begin{array}{ccc} \hat{D}_{n,11} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right] \\ \begin{array}{ccc} 0 & \Sigma_{\hat{\varphi}_1} & 0 \\ 0 & 0 & \Sigma_{\hat{\varphi}_0} \\ 0 & 0 & 0 \end{array} \end{array} V_n^{-1},$$

276 we obtain exactly the strangeness-free descriptor system (30).  $\square$

277 As a direct corollary of Theorem 4.5, we obtain the existence and uniqueness  
278 of a solution to the closed-loop system via feedback as follows.

279 **Corollary 4.6.** *Under the conditions of Theorem 4.5, the following statements*  
280 *hold true.*

- 281 *i) There exists a first order feedback of the form (28) such that the closed-loop*  
282 *system is solvable if and only if either  $\hat{v} = 0$  or  $\hat{f}_6(n) = 0$  for all  $n \geq n_0$ .*
- 283 *ii) Furthermore, the solution to the corresponding IVP (of the closed-loop sys-*  
284 *tem) is unique if and only if in addition,  $d = m - \hat{v}$ .*

285 **Remark 4.7.** It should be noted that, in analogous to SiDEs, each index re-  
286 duction step of the descriptor system (1) also makes use of Lemma 3.9, where  
287 the matrices  $Z_n^{(i)}$ ,  $i = 3, 4, 5$ , may not be orthogonal. Furthermore, in Lemma  
288 4.1, two matrices  $U_n$ ,  $V_n$  are only nonsingular but not orthogonal. Therefore,  
289 in general, the strangeness-free formulation (30) could not be stably computed.  
290 For the numerical treatment of second-order DAEs, in [22] a different approach  
291 was developed. We will modify it for SiDEs/descriptor systems in the next  
292 section.

293 **Remark 4.8.** Another interesting method in the study of descriptor systems  
294 is the *behavior approach*, where we do not distinguish the state  $x$  and an input  
295  $u$  but combine them in one *behavior vector*. Then, (1) will become a SiDE of  
296 this behavior variable, and hence, we can apply the results in Section 3 for this  
297 system. Nevertheless, due to the reinterpretation of variables, this approach  
298 may alter the strangeness-free form (30). To keep the brevity of this research,  
299 we will not present the details here. For the interested readers, we refer to  
300 [10, 18, 19] for the case of first order DAEs, and [22] for the case of second order  
301 DAEs.

## 302 5. Difference arrays associated with second-order SiDEs/descriptor 303 systems

304 As have shown in two previous sections, to analyze the theoretical solvability  
305 of the SiDE (2) or of the descriptor system (1), first one needs to bring it to a  
306 strangeness-free formulation. Nevertheless, this task is not always doable, for  
307 example when Assumptions 3.4, 4.2 are violated at some index reduction steps.  
308 These difficulties have also been observed for continuous-time systems of both  
309 first and higher orders, and they have been addressed in [10, 22]. The basic  
310 idea, thanks to Campbell [4], while considering DAEs, is to differentiate a given  
311 system a number of times and put every one of them, including the original one,  
312 into a so-called *inflated system*. Then, the strangeness-free formulation will be  
313 determined by appropriate selection of equations inside this inflated system. In  
314 this section we will examine this approach to the descriptor system (1). The  
315 analysis for SiDEs of the form (2) can be obtained by simply setting  $D_n$  to be  
316  $0_{m,p}$  for all  $n$ . We further assume the following condition.

317 **Assumption 5.1.** Consider the descriptor system (1). Assume that there exists  
 318 a first order feedback of the form (28) such that the corresponding IVP of the  
 319 closed-loop system is uniquely solvable.

320 Notice that, in case of the SiDE (2), Assumption 5.1 means that the IVP  
 321 (2)-(3) is uniquely solvable. Now let us introduce the *difference-inflated system*  
 322 of level  $\ell \in \mathbb{N}$  as follows.

$$\begin{aligned} A_n x(n+2) + B_n x(n+1) + C_n x(n) + D_n u(n) &= f(n), \\ A_{n+1} x(n+3) + B_{n+1} x(n+2) + C_{n+1} x(n+1) + D_{n+1} u(n+1) &= f(n+1), \\ &\dots \\ A_{n+\ell} x(n+\ell+2) + B_{n+\ell} x(n+\ell+1) + C_{n+\ell} x(n+\ell) + D_{n+\ell} u(n+\ell) &= f(n+\ell). \end{aligned}$$

We rewrite this system as

$$\begin{aligned} &\underbrace{\begin{bmatrix} C_n & B_n & A_n & & & \\ & C_{n+1} & B_{n+1} & A_{n+1} & & \\ & & \ddots & \ddots & \ddots & \\ & & & \ddots & \ddots & \ddots \\ & & & & C_{n+\ell} & B_{n+\ell} & A_{n+\ell} \end{bmatrix}}_{=:\mathcal{M}} \underbrace{\begin{bmatrix} x(n) \\ x(n+1) \\ x(n+2) \\ \vdots \\ x(n+\ell) \end{bmatrix}}_{=:\mathcal{X}} + \\ &+ \underbrace{\begin{bmatrix} D_n & & & & \\ & D_{n+1} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & D_{n+\ell} \end{bmatrix}}_{=:\mathcal{N}} \underbrace{\begin{bmatrix} u(n) \\ u(n+1) \\ \vdots \\ u(n+\ell) \end{bmatrix}}_{=:\mathcal{U}} = \underbrace{\begin{bmatrix} f(n) \\ f(n+1) \\ \vdots \\ f(n+\ell) \end{bmatrix}}_{=:\mathcal{G}}, \text{ for all } n \geq n_0. \quad (31) \end{aligned}$$

323 **Definition 5.2.** Suppose that the descriptor system (1) satisfies Assumption  
 324 5.1. Let  $\ell$  be the minimum number such that by using elementary matrix's  
 325 row operations, a strangeness-free descriptor system of the form (30) can be  
 326 extracted from (31). Then, the so-called *shift-index* of (1), denoted by  $\nu$ , is set  
 327 by  $\ell/2$  if  $\ell$  is even and by  $(\ell + 1)/2$  otherwise.

328 We give the relation between this shift-index  $\nu$  and the strangeness-index  $\mu$   
 329 in the following proposition.

330 **Proposition 5.3.** Suppose that the descriptor system (1) satisfies Assumption  
 331 5.1. If the strangeness-index  $\mu$  is well-defined, then so is the shift-index  $\nu$ .  
 332 Furthermore, we have that  $\nu \leq \mu$ .

333 *Proof.* The claim is straight forward, since every reformulation step performed  
 334 in Algorithm 1 is a consequence of an inflated system (31) with  $\ell = 2\mu$  or  
 335  $2\mu - 1$ .  $\square$

336 **Remark 5.4.** As will be seen later in Example 5.8, for second order SiDEs, the  
 337 shift index can be strictly smaller than the strangeness index.

**Remark 5.5.** Restricted to the case of first order SiDEs (i.e.,  $A_n = 0$  for all  $n \geq n_0$ ), the strangeness-index  $\mu$  defined in this paper is equal to the forward strangeness-index proposed by Brüll, [2]. For second order system, our strangeness-index is analogous to the one for continuous-time systems proposed by Mehrmann and Shi ([16]), and by Wunderlich ([22]). We, however, emphasize that the canonical forms constructed in this research is simpler and more convenient from the theoretical viewpoint. Besides that, similar to the case of continuous-time systems, the strangeness index  $\mu$  only gives an upper bound for the number of shift-forward operator that have been used, in order to achieve the strangeness-free form (21). For further details, see Remark 17, [16].

Assume that  $\nu$  is already known, now we construct an algorithm to select the strangeness-free descriptor system (30) from the inflated system (31). For notational convenience, we will follow the Matlab language, [14]. Consider the following spaces and matrices

$$\begin{aligned} \mathcal{W} &:= [\mathcal{M}(:, 3n+1 : \text{end}) \quad \mathcal{N}(:, n+1 : \text{end})], \\ U_1 &\text{ basis of } \ker(\mathcal{W}^T), \text{ and } U_{1,\perp} \text{ basis of } \text{range}(\mathcal{W}). \end{aligned} \quad (32)$$

Due to Lemma 2.6 we have that  $U_1^T \mathcal{W} = 0$  and  $U_{1,\perp}^T \mathcal{W}$  has full row rank. Furthermore, the matrix  $\begin{bmatrix} U_1^T \\ U_{1,\perp}^T \end{bmatrix}$  is nonsingular, and hence system (31) is equivalent to the coupled system below.

$$U_1^T \mathcal{M}(:, 1 : 3n) \begin{bmatrix} x(n) \\ x(n+1) \\ x(n+2) \end{bmatrix} + U_1^T \mathcal{N}(:, 1 : n) u(n) = U_1^T \mathcal{G}, \quad (33)$$

$$U_{1,\perp}^T \mathcal{W} \begin{bmatrix} x(n+3) \\ \vdots \\ x(n+\nu) \\ u(n+1) \\ \vdots \\ u(n+\nu) \end{bmatrix} + U_{1,\perp}^T [\mathcal{M}(:, 1 : 3n) \quad \mathcal{N}(:, 1 : n)] \begin{bmatrix} x(n) \\ x(n+1) \\ x(n+2) \\ u(n) \end{bmatrix} = U_{1,\perp}^T \mathcal{G}. \quad (34)$$

Notice that due to the full row rank property of  $U_{1,\perp}^T \mathcal{W}$ , (34) plays no role in the determination of the strangeness-free descriptor system (30). Thus, (30) is a consequence of (33). In the following proposition we show that system (33) is not affected by left equivalence transformation.

**Proposition 5.6.** *Consider two left equivalent systems. Then, at the same level  $\ell$ , their difference-inflated systems of the form (31) are also left equivalent. Consequently, system (33) is not affected by left equivalence transformation.*

*Proof.* Let us assume that (1) is left equivalent to the SiDE

$$\tilde{A}_n x(n+2) + \tilde{B}_n x(n+1) + \tilde{C}_n x(n) + \tilde{D}_n u(n) = \tilde{f}(n), \text{ for all } n \geq n_0. \quad (35)$$

Thus, there exists a pointwise nonsingular matrix sequence  $\{P_n\}_{n \geq n_0}$  such that

$$\begin{bmatrix} \tilde{A}_n & \tilde{B}_n & \tilde{C}_n & \tilde{D}_n \end{bmatrix} = P_n \begin{bmatrix} A_n & B_n & C_n & D_n \end{bmatrix} \text{ and } \tilde{f}(n) = P_n f(n), \text{ for all } n \geq n_0.$$

Therefore, the difference-inflated system of level  $\ell$  for system (35) takes the form

$$\tilde{\mathcal{M}}\mathcal{X} + \tilde{\mathcal{N}}\mathcal{U} = \tilde{\mathcal{G}}, \quad (36)$$

where the matrix coefficients are

$$\tilde{\mathcal{M}} = \text{diag}(P_n, \dots, P_{n+\ell}) \mathcal{M}, \quad \tilde{\mathcal{N}} = \text{diag}(P_n, \dots, P_{n+\ell}) \mathcal{N}, \quad \tilde{\mathcal{G}} = \text{diag}(P_n, \dots, P_{n+\ell}) \mathcal{G}.$$

355 This follows that two systems (31) and (36) are left equivalent, which finishes  
356 the proof.  $\square$

For notational convenience, let us rewrite system (33) as

$$\left[ \begin{array}{ccc|c} \check{A} & \check{B} & \check{C} & \check{D} \end{array} \right] \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \\ u(n) \end{bmatrix} = \check{G}.$$

Scale this system with the matrix  $\check{U}$  obtained in Lemma 2.7, we have

$$\left[ \begin{array}{ccc|c} \check{A}_1 & \check{B}_1 & \check{C}_1 & \check{D}_1 \\ 0 & \check{B}_2 & \check{C}_2 & 0 \\ 0 & 0 & \check{C}_3 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & \check{B}_4 & \check{C}_4 & \check{D}_4 \\ 0 & 0 & \check{C}_5 & \check{D}_5 \end{array} \right] \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \\ u(n) \end{bmatrix} = \begin{bmatrix} \check{G}_1 \\ \check{G}_2 \\ \check{G}_3 \\ 0 \\ \hline \check{G}_4 \\ \check{G}_5 \end{bmatrix}. \quad (37)$$

357 Here the matrices  $\check{A}_1, \check{B}_2, \check{B}_4, \check{C}_3$ , and  $\begin{bmatrix} \check{D}_4 \\ \check{D}_5 \end{bmatrix}$  have full row rank. Notice that  
358 the presence of the 0 block on the right hand side vector is due to the existence  
359 of a solution (Assumption 5.1). In the following theorem we will answer the  
360 question how to derive the strangeness-free formulation (30) from (37).

**Theorem 5.7.** *Assume that the shift index  $\nu$  of the descriptor system (1) is well-defined. Furthermore, suppose that (1) satisfies Assumption 5.1. Then, any solution to the descriptor system (1) is also a solution to the following system*

$$\begin{array}{c} \hat{r}_2 \\ \hat{r}_1 \\ \hat{r}_0 \\ \hat{\varphi}_1 \\ \hat{\varphi}_0 \end{array} \begin{bmatrix} \hat{A}_{n,1} & \hat{B}_{n,1} & \hat{C}_{n,1} \\ 0 & \hat{B}_{n,2} & \hat{C}_{n,2} \\ 0 & 0 & \hat{C}_{n,3} \\ \hline 0 & \hat{B}_{n,5} & \hat{C}_{n,5} \\ 0 & 0 & \hat{C}_{n,6} \end{bmatrix} \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \end{bmatrix} + \begin{bmatrix} \hat{D}_{n,1} \\ 0 \\ 0 \\ \hline \hat{D}_{n,4} \\ \hat{D}_{n,5} \end{bmatrix} u(n) = \begin{bmatrix} \hat{G}_{n,1} \\ \hat{G}_{n,2} \\ \hat{G}_{n,3} \\ \hline \hat{G}_{n,4} \\ \hat{G}_{n,5} \end{bmatrix}, \text{ for all } n \geq n_0, \quad (38)$$

where the matrices  $\begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix}, \begin{bmatrix} \hat{D}_{n,4} \\ \hat{D}_{n,5} \end{bmatrix}$  have full row rank for all  $n \geq n_0$ . Furthermore, we have that  $\sum_{i=0}^2 \hat{r}_i + \sum_{i=0}^1 \hat{\varphi}_i = d$ , or equivalently,

$$\text{rank} \left( \begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix} \right) + \text{rank} \left( \begin{bmatrix} \hat{D}_{n,4} \\ \hat{D}_{n,5} \end{bmatrix} \right) = d. \quad (39)$$

*Proof.* First we will extract the first two block row equations of system (38) from (37), by suitably removing the existence hidden redundancy. Applying Lemma 2.6 consecutively for two following matrix pairs  $(\check{B}_2, \check{C}_3), \left( \check{A}_1, \begin{bmatrix} \check{B}_2 \\ \check{C}_3 \end{bmatrix} \right)$ , we obtain two orthogonal matrices  $\begin{bmatrix} S_n^{(i)} \\ Z_n^{(i)} \end{bmatrix} \in \mathbb{R}^{r_i, r_i}, i = 1, 2$  such that both pairs  $(S_n^{(1)} \check{B}_2, \check{C}_3), \left( S_n^{(2)} \check{A}_1, \begin{bmatrix} \check{B}_2 \\ \check{C}_3 \end{bmatrix} \right)$  have no hidden redundancy. Scale the first and second block row equations of (37) with  $S_n^{(2)}$  and  $S_n^{(1)}$  respectively, we obtain

$$\left[ \begin{array}{ccc|c} S_n^{(2)} \check{A}_1 & S_n^{(2)} \check{B}_1 & S_n^{(2)} \check{C}_1 & S_n^{(2)} \check{D}_1 \\ 0 & S_n^{(1)} \check{B}_2 & S_n^{(1)} \check{C}_2 & 0 \end{array} \right] \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \\ u(n) \end{bmatrix} = \begin{bmatrix} S_n^{(2)} \check{G}_1 \\ S_n^{(1)} \check{G}_2 \end{bmatrix}.$$

Combining these equations with the third, fifth and sixth block equations of (37), we obtain the system

$$\left[ \begin{array}{ccc|c} S_n^{(2)} \check{A}_1 & S_n^{(2)} \check{B}_1 & S_n^{(2)} \check{C}_1 & S_n^{(2)} \check{D}_1 \\ 0 & S_n^{(1)} \check{B}_2 & S_n^{(1)} \check{C}_2 & 0 \\ 0 & 0 & \check{C}_3 & 0 \\ \hline 0 & \check{B}_4 & \check{C}_4 & \check{D}_4 \\ 0 & 0 & \check{C}_5 & \check{D}_5 \end{array} \right] \begin{bmatrix} x(n+2) \\ x(n+1) \\ x(n) \\ u(n) \end{bmatrix} = \begin{bmatrix} S_n^{(2)} \check{G}_1 \\ S_n^{(1)} \check{G}_2 \\ \check{G}_3 \\ \check{G}_4 \\ \check{G}_5 \end{bmatrix}. \quad (40)$$

361 which is exactly our desired system (38). Moreover, due to Lemma 2.3, the  
 362 matrix  $\begin{bmatrix} S_n^{(2)} \check{A}_1 \\ S_n^{(1)} \check{B}_2 \\ \check{C}_3 \end{bmatrix}$  has full row rank. Finally, the identity (39) holds true due to  
 363 Assumption 5.1. □

364 We summarize our result in the following algorithm.



---

**Algorithm 2** Strangeness-free formulation for SiDEs using difference arrays

---

**Input:** The SiDE (1).

**Output:** The strangeness-free descriptor system (38) and the minimal number of shifts  $\ell$ .

- 1: Set  $\ell := 0$ .
  - 2: Construct the difference-inflated system of level  $\ell$ , and rewrite it in the form (31).
  - 3: Find  $U_1$  as in (32) and construct system (33).
  - 4: Find  $\check{U}$  as in Lemma 2.7 and construct system (37).
  - 5: Find the matrices  $S_n^{(1)}, S_n^{(2)}$  in the process used to remove the hidden redundancies in two matrix pairs  $(\check{B}_2, \check{C}_3), \left(\check{A}_1, \begin{bmatrix} \check{B}_2 \\ \check{C}_3 \end{bmatrix}\right)$ , respectively.
  - 6: Construct the system (40).
  - 7: **if**  $\text{rank} \begin{bmatrix} \hat{A}_{n,1} \\ \hat{B}_{n+1,2} \\ \hat{C}_{n+2,3} \end{bmatrix} + \text{rank} \begin{bmatrix} \hat{D}_{n,4} \\ \hat{D}_{n,5} \end{bmatrix} = d$  **then** STOP.
  - 8: **else** set  $\ell := \ell + 1$  and go to 2
  - 9: **end if**
- 

365 In order to illustrate Algorithm 2, we consider two following examples.

**Example 5.8.** Let us revisit system (22) for the case  $\alpha = 0$ . In this system,  $D_n = 0$  for all  $n \geq 0$ . For  $\ell = 2$ , the inflated system (31) reads

$$\left[ \begin{array}{ccc|cc} C_n & B_n & A_n & 0 & 0 \\ 0 & C_{n+1} & B_{n+1} & A_{n+1} & 0 \\ 0 & 0 & C_{n+2} & B_{n+2} & A_{n+2} \end{array} \right] \begin{bmatrix} x(n) \\ x(n+1) \\ x(n+2) \\ x(n+3) \\ x(n+4) \end{bmatrix} = \begin{bmatrix} f(n) \\ f(n+1) \\ f(n+2) \end{bmatrix} \quad (41)$$

Let  $U_1$  be the basis of  $\text{kernel}(\mathcal{W}^T)$ , where  $\mathcal{W} = \begin{bmatrix} 0 & 0 \\ A_{n+1} & 0 \\ B_{n+2} & A_{n+2} \end{bmatrix}$ . We then compute system (33) by scaling (41) with  $U_1^T$ . The resulting system reads

$$U_1^T \begin{bmatrix} C_n & B_n & A_n \\ 0 & C_{n+1} & B_{n+1} \\ 0 & 0 & C_{n+2} \end{bmatrix} \begin{bmatrix} x(n) \\ x(n+1) \\ x(n+2) \end{bmatrix} = U_1^T \begin{bmatrix} f(n) \\ f(n+1) \\ f(n+2) \end{bmatrix}. \quad (42)$$

366 Finally, by performing Steps 6 to 10 we can extract the strangeness-free form  
 367 (24) from (42). Thus, we conclude that the shift index is  $\nu = 1$ , which is the  
 368 same as the shift index in the case  $\alpha \neq 0$ . We recall Example 3.11, in which it  
 369 is shown that the strangeness indices in the two cases are different.

**Example 5.9.** A singular system of second-order differential equations, which describes a three link robot arm [8], is given by

$$\begin{bmatrix} M_0 & 0 \\ 0 & 0 \end{bmatrix} \ddot{x}(t) + \begin{bmatrix} G_0 & 0 \\ 0 & 0 \end{bmatrix} \dot{x}(t) + \begin{bmatrix} K_0 & H_0^T \\ H_0 & 0 \end{bmatrix} x(t) = \begin{bmatrix} B_0 \\ 0 \end{bmatrix} u(t).$$

Here  $M_0$  represents the nonsingular mass matrix,  $G_0$  the coefficient matrix associated with damping, centrifugal, gravity, and Coriolis forces,  $K_0$  the stiffness matrix, and  $H_0$  the constraint. A simple discretized version of this system takes the form

$$\begin{aligned} & \begin{bmatrix} M_0 & 0 \\ 0 & 0 \end{bmatrix} \frac{x(n+2) - 2x(n+1) + x(n)}{h^2} + \begin{bmatrix} G_0 & 0 \\ 0 & 0 \end{bmatrix} \frac{x(n+2) - x(n)}{2h} \\ & + \begin{bmatrix} K_0 & H_0^T \\ H_0 & 0 \end{bmatrix} x(n+1) = \begin{bmatrix} B_0 \\ 0 \end{bmatrix} u(n+1). \end{aligned}$$

where  $h$  is the discretized stepsize.

As a simple example, let us take  $M_0 = G_0 = K_0 = H_0 = B_0 = 1$ ,  $h = 0.01$ . Then, Algorithm 2 terminates after two steps and hence, the shift index is  $\nu = 1$  for all  $n \geq n_0$ . Furthermore, we notice that no matter central, forward or backward difference is chosen for discretizing the derivative  $\dot{x}(t)$ , the shift index remains unchanged  $\nu = 1$ . Of course, the resulting strangeness-free descriptor systems are different.

## 6. Conclusion

By using the algebraic approach, we have analyzed the solvability of second order SiDEs/descriptor systems, based on derived condensed forms constructed under certain constant rank assumptions. In comparison to well-known results [16, 21], we have reduced the number of constant rank conditions in every index reduction step from seven to five. This would enlarge the domain of application for SiDEs (and also for DAEs). However, requiring constant rank assumptions in the discrete-time case seems less nature than in the continuous-time case. To overcome this limitation, we also consider the difference-array method, which is numerically applicable. The theory together with the two algorithms presented in this paper can be extended without difficulty to arbitrarily high order SiDEs/descriptor systems. We also notice that the backward time case ( $n \leq n_0$ ) can be directly extended from the forward time case, as it has been done in [2]. The analysis of two-way case, which happens while considering boundary value problems for SiDEs, is under our on-going research. Besides that, the condensed forms presented in this research also motivate further study on the staircase form for second order systems, which hopefully can generalize classical results for first order systems, e.g. [20].

## References

- [1] R. Agarwal. *Difference Equations and Inequalities: Theory, Methods, and Applications*. Chapman & Hall/CRC Pure and Applied Mathematics. CRC Press, 2000.
- [2] T. Brüll. Existence and uniqueness of solutions of linear variable coefficient discrete-time descriptor systems. *Linear Algebra Appl.*, 431(1-2):247–265, 2009.
- [3] R. Byers, P. Kunkel, and V. Mehrmann. Regularization of linear descriptor systems with variable coefficients. *SIAM J. Cont.*, 35:117–133, 1997.
- [4] S. L. Campbell. Comment on controlling generalized state-space (descriptor) systems. *Internat. J. Control*, 46:2229–2230, 1987.
- [5] S. Elaydi. *An Introduction to Difference Equations*. Undergraduate Texts in Mathematics. Springer New York, 2013.
- [6] G. H. Golub and C. F. Van Loan. *Matrix Computations*. The Johns Hopkins University Press, Baltimore, MD, 3rd edition, 1996.
- [7] P. Ha and V. Mehrmann. Analysis and reformulation of linear delay differential-algebraic equations. *Electr. J. Lin. Alg.*, 23:703–730, 2012.
- [8] M. Hou. A three-link planar manipulator model. Sicherheitstechnische Regelungs- und Meßtechnik, Bergische Universität–GH Wuppertal, Germany, May 1994.
- [9] W. Kelley and A. Peterson. *Difference Equations: An Introduction with Applications*. Harcourt/Academic Press, 2001.
- [10] P. Kunkel, V. Mehrmann, and W. Rath. Analysis and numerical solution of control problems in descriptor form. *Math. Control, Signals, Sys.*, 14:29–61, 2001.
- [11] V.H. Linh, N.T.T. Nga, and D.D. Thuan. Exponential stability and robust stability for linear time-varying singular systems of second order difference equations. *SIAM J. Matr. Anal. Appl.*, 39(1):204–233, 2018.
- [12] P. Losse and V. Mehrmann. Controllability and observability of second order descriptor systems. *SIAM J. Cont. Optim.*, 47(3):1351–1379, 2008.
- [13] D. Luenberger. *Introduction to dynamic systems: theory, models, and applications*. Wiley, 1979.
- [14] The MathWorks, Inc., Natick, MA. *MATLAB Version 8.3.0.532 (R2014a)*, 2014.
- [15] V. Mehrmann. Index concepts for differential-algebraic equations. *ENCYCLOPEDIA APPLIED MATHEMATICS*, 2014.

- 431 [16] V. Mehrmann and C. Shi. Transformation of high order linear differential-  
432 algebraic systems to first order. *Numer. Alg.*, 42:281–307, 2006.
- 433 [17] V. Mehrmann and D.D. Thuan. Stability analysis of implicit difference  
434 equations under restricted perturbations. *SIAM J. Matr. Anal. Appl.*,  
435 36(1):178–202, 2015.
- 436 [18] W. Rath. Derivative and proportional state feedback for linear descriptor  
437 systems with variable coefficients. *Lin. Alg. Appl.*, 260:273–310, 1997.
- 438 [19] W. Rath. *Feedback Design and Regularization for Linear Descriptor Sys-*  
439 *tems with Variable Coefficients*. Dissertation, TU Chemnitz, Chemnitz,  
440 Germany, 1997.
- 441 [20] E. Sontag. *Mathematical Control Theory: Deterministic Finite Dimen-*  
442 *sional Systems*. Texts in Applied Mathematics. Springer New York, 2013.
- 443 [21] L. Wunderlich. Numerical treatment of second order differential-algebraic  
444 systems. In *Proc. Appl. Math. and Mech. (GAMM 2006, Berlin, March*  
445 *27-31, 2006)*, volume 6 (1), pages 775–776, 2006.
- 446 [22] L. Wunderlich. *Analysis and Numerical Solution of Structured and Switched*  
447 *Differential-Algebraic Systems*. Dissertation, Institut für Mathematik, TU  
448 Berlin, Berlin, Germany, 2008.