

Relative controllability of delay differential systems with impulses and linear parts defined by permutable matrices

Zhongli You¹ | JinRong Wang^{1,2}  | Donal O'Regan³ | Yong Zhou^{4,5} 

¹Department of Mathematics, School of Mathematics and Statistics, Guizhou University, Guiyang, P.R. China

²School of Mathematical Sciences, Qufu Normal University, Qufu, P.R. China

³School of Mathematics, Statistics and Applied Mathematics, National University of Ireland, Galway, Ireland

⁴Department of Mathematics, Xiangtan University, Xiangtan, P.R. China

⁵Nonlinear Analysis and Applied Mathematics (NAAM) Research Group, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

Correspondence

JinRong Wang, Department of Mathematics, School of Mathematics and Statistics, Guizhou University, Guiyang, Guizhou 550025, P.R. China.
Email: wjr9668@126.com

Communicated by: C. Cuevas

Funding information

National Natural Science Foundation of China, Grant/Award Number: 11661016; Training Object of High Level and Innovative Talents of Guizhou Province, Grant/Award Number: ([2016]4006); Science and Technology Program of Guizhou Province, Grant/Award Number: ([2017]5788-10); Major Research Project of Innovative Group in Guizhou Education Department, Grant/Award Number: ([2018]012); Foundation of Postgraduate of Guizhou Province, Grant/Award Number: (YJSCXJH[2018]035)

MSC Classification: 34A37; 93B05

This paper investigates the relative controllability of delay differential systems with linear impulses and linear parts defined by permutable matrices. We use the impulsive delay Grammian matrix to discuss the relatively controllability of impulsive linear delay controlled systems and we use the Krasnoselskii's fixed point theorem to discuss the relatively controllability of impulsive semilinear delay controlled systems. Finally, two examples are presented to illustrate our theoretical results.

KEYWORDS

impulsive delay differential equations, impulsive delay Grammian matrix, relative controllability

1 | INTRODUCTION

Mathematical control theory is an important area of research and classical differential controlled systems in finite and infinite dimensional spaces have been discussed in the literature. However, very little is known for delay differential controlled systems and impulsive delay controlled systems.

Khusainov and Shuklin¹ and Diblík and Khusainov² introduced delayed exponential matrix functions and these were used to discuss formulas for solutions to linear delay systems with permutable matrices, stability, inequality, and control problems (see³⁻²³ and the references therein). Motivated by,^{1,2} You and Wang^{24,25} extended the single and multiple delayed exponential matrix function in¹ to the impulsive case and used it to discuss the representation and stability of solutions to linear delay systems with linear impulses.

In this paper, we consider the relative controllability of the following impulsive semilinear delay differential systems

$$\begin{cases} v'(t) = Av(t) + Bv(t - \vartheta) + f(t, v(t)) + Du(t), & t \in \Omega, t \notin \mathcal{T}, \vartheta > 0, \\ \Delta v(t_i) := v(t_i^+) - v(t_i^-) = C_i v(t_i), & t_i \in \mathcal{T} = \{t_1, t_2, \dots, t_k\}, \\ v(t) = \psi(t), & -\vartheta \leq t \leq 0, \end{cases} \quad (1)$$

where $A, B, C_i, D \in \mathbb{R}^{n \times n}$, $AB = BA$, $AC_i = C_i A$, $BC_i = C_i B$ for each $i \in \{1, 2, \dots, k\}$, $v \in C^1([-\vartheta, 0], \mathbb{R}^n) \cup [PC(\Omega, \mathbb{R}^n) \cap (\cup_{i=0}^k C^1(t_i, t_{i+1}))]$ (here $t_0 = 0$ and $t_{k+1} = \tau_1$), $\psi \in C^1_{\vartheta} := C^1([-\vartheta, 0], \mathbb{R}^n)$, $f \in C(\Omega \times \mathbb{R}^n, \mathbb{R}^n)$, $\Omega := [0, \tau_1]$, $\tau_1 > 0$, $0 < t_1 < t_2 < \dots < t_k < \tau_1$, and the control function $u(\cdot)$ takes values from $L^2(\Omega, \mathbb{R}^n)$. Set $v(t_i^+) = \lim_{\epsilon \rightarrow 0^+} v(t_i + \epsilon)$ and $v(t_i^-) = v(t_i)$ represent respectively the right and left limits of $v(t)$ at $t = t_i$.

First, we investigate the relative controllability of the linear impulsive delay controlled system as follows:

$$\begin{cases} v'(t) = Av(t) + Bv(t - \vartheta) + Du(t), & t \in \Omega, t \notin \mathcal{T}, \vartheta > 0, \\ \Delta v(t_i) = C_i v(t_i), & t_i \in \mathcal{T}, \\ v(t) = \psi(t), & -\vartheta \leq t \leq 0, \end{cases} \quad (2)$$

using the impulsive delay Cauchy matrix introduced in.²⁴ Next, we construct a suitable control function for (1), which means that we give a condition necessary and sufficient for $u \in L^2(\Omega, \mathbb{R}^n)$ to lead the solution of (1) to v_1 at the time τ_1 . We apply the Krasnoselskii's fixed point theorem to show that (1) is also relatively controllable under suitable conditions.

The rest of this paper is organized as follows. In Section 2, we give some notations, concepts, and important lemmas. In Section 3, we establish relative controllability results for (1) and (2) respectively. Examples are given to illustrate our main results in the final section.

2 | PRELIMINARIES

Let \mathbb{R}^n be the n -dimensional Euclidian space with the vector norm $\|\cdot\|$ and $\mathbb{R}^{n \times n}$ be the $n \times n$ matrix space with real valued elements. For $v \in \mathbb{R}^n$ and $A \in \mathbb{R}^{n \times n}$, we introduce the vector infinite-norm $\|v\| = \max_{1 \leq i \leq n} |v_i|$ and the matrix infinite-norm

$\|A\| = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|$ respectively, where v_i and a_{ij} are the elements of the vector v and matrix A . Denote by $C(\Omega, \mathbb{R}^n)$ the

Banach space of vector-valued bounded continuous functions from $\Omega \rightarrow \mathbb{R}^n$ and let $C^1(\Omega, \mathbb{R}^n) = \{v \in C(\Omega, \mathbb{R}^n) : v' \in C(\Omega, \mathbb{R}^n)\}$. Let $PC(\Omega, \mathbb{R}^n) := \{v : \Omega \rightarrow \mathbb{R}^n : v \in C((t_i, t_{i+1}], \mathbb{R}^n), i = 0, 1, \dots, k, \text{ and there exists } v(t_i^+), v(t_i^-) \text{ with } v(t_i^-) = v(t_i), i = 1, \dots, k\}$. Denote by PC , the space $PC(\Omega, \mathbb{R}^n)$ of vector-valued piece-wise continuous functions from $\Omega \rightarrow \mathbb{R}^n$ endowed with the norm $\|v\|_{PC} = \sup_{t \in \Omega} \|v(t)\|$ where $\|\cdot\|$ is the norm on \mathbb{R}^n . In addition, $\|\psi\|_{PC} = \sup_{t \in [-\vartheta, 0]} \|\psi(t)\|$. Let Y_1, Y_2 be two Banach spaces, and let $L_b(Y_1, Y_2)$ denote the space of all bounded linear operators from Y_1 to Y_2 . Now $L^p(\Omega, Y_2)$ denotes the Banach space of functions $y : \Omega \rightarrow Y_2$ which are Bochner integrable with norm $\|y\|_{L^p(\Omega, Y_2)}$ (here $1 < p < \infty$).

We recall the delayed matrix exponential $e_{\vartheta}^{\Lambda} : \mathbb{R} \rightarrow \mathbb{R}^{n \times n}$ given by as follows:

$$e_{\vartheta}^{\Lambda t} = \begin{cases} \Theta, & t < -\vartheta, \vartheta > 0, \\ E, & -\vartheta \leq t < 0, \\ E + \Lambda t + \Lambda^2 \frac{(t-\vartheta)^2}{2} + \dots + \Lambda^z \frac{(t-(z-1)\vartheta)^z}{z!}, & (z-1)\vartheta \leq t < z\vartheta, z = 1, 2, \dots, \end{cases} \quad (3)$$

where $\Lambda \in \mathbb{R}^{n \times n}$, and Θ and E are the zero and identity matrices, respectively. From,^{1, Lemma 4}

$$\frac{d}{dt} e_{\vartheta}^{\Lambda t} = \Lambda e_{\vartheta}^{\Lambda(t-\vartheta)}, \quad (4)$$

for all $t \in \mathbb{R}$.

Let $V(\cdot, \cdot) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^{n \times n}$ (see²⁴) be given by

$$V(t, s) = e^{A(t-s)} X(t, s + \vartheta), \quad t > s, \quad (5)$$

and

$$X(t, s) = e_g^{\check{B}(t-s)} + \sum_{s-\vartheta < t_j \leq t} C_j e_g^{\check{B}(t-\vartheta-t_j)} X(t_j, s), \quad \check{B} = e^{-A\vartheta} B. \quad (6)$$

For any solution $v \in C^1([-\vartheta, 0], \mathbb{R}^n) \cup [PC(\Omega, \mathbb{R}^n) \cap (\cup_{i=0}^k C^1(t_i, t_{i+1}))]$ (here $t_0 = 0$ and $t_{k+1} = \tau_1$) of (1), from²⁴, Corollary 3.3 we obtain the following:

$$\begin{aligned} v(t) &= V(t, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(t, s)[\psi'(s) - A\psi(s)]ds \\ &\quad + \int_0^t V(t, s)[f(s, v(s)) + Du(s)]ds. \end{aligned} \quad (7)$$

The following results are important and we keep the details of the proof for the reader.

Lemma 2.1. *For all $t \geq 0$, we have $\|e_g^{\Lambda t}\| \leq e^{\|\Lambda\|t}$.*

Proof. From (3), without loss of generality, for $(z-1)\vartheta \leq t < z\vartheta, z = 1, 2, \dots$, we have the following

$$\begin{aligned} \|e_g^{\Lambda t}\| &= \left\| \sum_{n=0}^z \Lambda^n \frac{(t - (n-1)\vartheta)^n}{n!} \right\| \\ &\leq \sum_{n=0}^z \|\Lambda\|^n \frac{t^n}{n!} \\ &\leq \sum_{n=0}^{\infty} \|\Lambda\|^n \frac{t^n}{n!} = e^{\|\Lambda\|t}. \end{aligned}$$

The result is proved. \square

Lemma 2.2. *For any $t > s$, we have the following:*

$$\|X(t, s)\| \leq \left(\prod_{s-\vartheta < t_j \leq t} (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{\|\check{B}\|(t-s)}, \quad (8)$$

and

$$\|V(t, s)\| \leq \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{-\|\check{B}\|\vartheta} e^{a(t-s)}, \quad (9)$$

where $a = \|A\| + \|\check{B}\|$.

Proof. Without loss of generality, we suppose that $t_m \leq s - \vartheta < t_{m+1}$ and $t_{m+n} \leq t < t_{m+n+1}, m, n = 0, 1, 2, \dots$. To complete our proof we use mathematical induction.

a. For $n = 0$, from Lemma 2.1 via (6),

$$\|X(t, s)\| \leq \|e_g^{\check{B}(t-s)}\| \leq e^{\|\check{B}\|(t-s)}.$$

b. For $n = 1$, from Lemmas 2.1 via (6), we have

$$\begin{aligned} \|X(t, s)\| &\leq \|e_g^{\check{B}(t-s)} + C_{m+1} e_g^{\check{B}(t-\vartheta-t_{m+1})} X(t_{m+1}, s)\| \\ &\leq e^{\|\check{B}\|(t-s)} + \|C_{m+1}\| e^{\|\check{B}\|(t-\vartheta-t_{m+1})} e^{\|\check{B}\|(t_{m+1}-s)} \\ &= (1 + \|C_{m+1}\| e^{-\|\check{B}\|\vartheta}) e^{\|\check{B}\|(t-s)}. \end{aligned}$$

c. For $n = z$, suppose that

$$\|X(t, s)\| \leq \left(\prod_{j=m+1}^{m+z} (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{\|\check{B}\|(t-s)} = \left(\prod_{s-\vartheta < t_j \leq t} (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{\|\check{B}\|(t-s)}.$$

For $n = z + 1$, from Lemmas 2.1 via (6) again,

$$\begin{aligned}
 \|X(t, s)\| &\leq \left(\prod_{j=m+1}^{m+z} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{\|\tilde{B}\|(t-s)} \\
 &\quad + \|C_{m+z+1}\| e^{\|\tilde{B}\|(t-\vartheta-t_{m+z+1})} \left(\prod_{j=m+1}^{m+z} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{\|\tilde{B}\|(t_{m+z+1}-s)} \\
 &= \left(\prod_{j=m+1}^{m+z+1} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{\|\tilde{B}\|(t-s)} \\
 &= \left(\prod_{s-\vartheta < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{\|\tilde{B}\|(t-s)}.
 \end{aligned}$$

From the mathematical induction principle, we obtain (8).

Finally, using (5) via (8) and $\|e^{At}\| \leq e^{\|A\|t}$, one obtains (9) immediately. \square

Definition 2.3. (see²⁶, Definition 4). System (1) is called relatively controllable, if for an arbitrary initial vector function $\psi \in C^1([-\vartheta, 0], \mathbb{R}^n)$, final state of the vector $v_1 \in \mathbb{R}^n$ at time τ_1 , there exists a control $u \in L^2(\Omega, \mathbb{R}^n)$ such that the system (1) has a solution $v \in C^1([-\vartheta, 0], \mathbb{R}^n) \cup [PC(\Omega, \mathbb{R}^n) \cap (\cup_{i=0}^k C^1(t_i, t_{i+1}))]$ (here $t_0 = 0$ and $t_{k+1} = \tau_1$) that satisfies $v(\tau_1) = v_1$.

Lemma 2.4. (Krasnoselskii's fixed point theorem, see²⁷). Let B be a bounded closed and convex subset of Banach space X and let F_1, F_2 be maps of B into X such that $F_1x + F_2y \in B$ for every $x, y \in B$. If F_1 is a contraction and F_2 is compact and continuous, then the equation $F_1x + F_2x = x$ has a solution on B .

Theorem 2.5. (PC-type Ascoli–Arzela theorem, see²⁸, Theorem 2.1). Let $\mathcal{Q} \subset PC(\Omega, X)$ where X is a Banach space. Then \mathcal{Q} is a relatively compact subset of $PC(\Omega, X)$ if, (a) \mathcal{Q} is uniformly bounded subset of $PC(\Omega, X)$; (b) \mathcal{Q} is equicontinuous in (t_i, t_{i+1}) , $i = 0, 1, 2, \dots, k$ (here $t_0 = 0$ and $t_{k+1} = \tau_1$); and (c) $\mathcal{Q}(t) = \{v(t) | v \in \mathcal{Q}, t \in \Omega \setminus \mathcal{T}\}$, $\mathcal{Q}(t_i^+) = \{v(t_i^+) | v \in \mathcal{Q}\}$ and $\mathcal{Q}(t_i^-) = \{v(t_i^-) | v \in \mathcal{Q}\}$ are relatively compact subsets of X .

3 | RELATIVE CONTROLLABILITY

3.1 | Linear systems

We consider an impulsive delay Grammian matrix, an extension of the classical Grammian matrix for linear differential systems, as follows:

$$W_\vartheta[0, \tau_1] = \int_0^{\tau_1} V(\tau_1, s) D D^T V^T(\tau_1, s) ds. \quad (10)$$

Theorem 3.1. System (2) is relatively controllable, if and only if $W_\vartheta[0, \tau_1]$ is nonsingular.

Proof. First, we establish sufficiency. Since $W_\vartheta[0, \tau_1]$ is nonsingular, its inverse $W_\vartheta^{-1}[0, \tau_1]$ is well-defined. For any final state $v_1 \in \mathbb{R}^n$ one can select a control function as follows:

$$u(t) = D^T V^T(\tau_1, t) W_\vartheta^{-1}[0, \tau_1] \eta,$$

where

$$\eta = v_1 - V(\tau_1, -\vartheta) \psi(-\vartheta) - \int_{-\vartheta}^0 V(\tau_1, s) [\psi'(s) - A \psi(s)] ds.$$

Then

$$\begin{aligned}
 v(\tau_1) &= V(\tau_1, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(\tau_1, s)[\psi'(s) - A\psi(s)]ds \\
 &\quad + \int_0^{\tau_1} V(\tau_1, s)Du(s)ds \\
 &= V(\tau_1, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(\tau_1, s)[\psi'(s) - A\psi(s)]ds \\
 &\quad + \int_0^{\tau_1} V(\tau_1, s)DD^TV^T(\tau_1, s)W_{\vartheta}^{-1}[0, \tau_1]\eta ds \\
 &= v_1.
 \end{aligned}$$

We argue by contradiction to prove our necessity result. Assume $W_{\vartheta}[0, \tau_1]$ is singular, ie, there exists at least one nonzero state $\tilde{v} \in \mathbb{R}^n$ such that,

$$\tilde{v}^T W_{\vartheta}[0, \tau_1] \tilde{v} = 0.$$

One obtains the following

$$\begin{aligned}
 0 &= \tilde{v}^T W_{\vartheta}[0, \tau_1] \tilde{v} \\
 &= \int_0^{\tau_1} \tilde{v}^T V(\tau_1, s)DD^TV^T(\tau_1, s)\tilde{v}ds \\
 &= \int_0^{\tau_1} \|\tilde{v}^T V(\tau_1, s)D\|^2 ds,
 \end{aligned}$$

which implies that

$$\tilde{v}^T V(\tau_1, s)D = \underbrace{(0, \dots, 0)}_n := \mathbf{0}^T, \quad \forall s \in \Omega.$$

Since system (2) is relatively controllable, according to Definition 2.3, there exists a control $u_1(t)$ that drives the initial state to zero at τ_1 ie,

$$v(\tau_1) = V(\tau_1, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(\tau_1, s)[\psi'(s) - A\psi(s)]ds + \int_0^{\tau_1} V(\tau_1, s)Du_1(s)ds = \mathbf{0}. \quad (11)$$

Similarly, there also exists a control $u_2(t)$ that drives the initial state to the (nonzero) state \tilde{v} at τ_1 , ie,

$$\begin{aligned}
 v(\tau_1) &= V(\tau_1, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(\tau_1, s)[\psi'(s) - A\psi(s)]ds \\
 &\quad + \int_0^{\tau_1} V(\tau_1, s)Du_2(s)ds \\
 &= \tilde{v}.
 \end{aligned} \quad (12)$$

Then from (11) and (12), we have

$$\tilde{v} = \int_0^{\tau_1} V(\tau_1, s)D[u_2(s) - u_1(s)]ds. \quad (13)$$

Multiplying both sides of (13) by \tilde{v}^T , we obtain

$$\tilde{v}^T \tilde{v} = \int_0^{\tau_1} \tilde{v}^T V(\tau_1, s)D[u_2(s) - u_1(s)]ds = 0.$$

Thus $\tilde{v} = \mathbf{0}$, which conflicts with \tilde{v} being nonzero. Thus, the impulsive delay Grammian matrix $W_{\vartheta}[0, \tau_1]$ is nonsingular. The proof is finished. \square

3.2 | Semilinear systems

We assume the following:

(H_1): The operator $W : L^2(\Omega, \mathbb{R}^n) \rightarrow \mathbb{R}^n$ defined by

$$Wu = \int_0^{\tau_1} V(\tau_1, s)Du(s)ds,$$

has an inverse operator W^{-1} which takes values in $L^2(\Omega, \mathbb{R}^n)/\ker W$. Then we set

$$M = \|W^{-1}\|_{L_b(\mathbb{R}^n, L^2(\Omega, \mathbb{R}^n)/\ker W)}.$$

From²⁹, Remark 3.3 we know

$$M = \sqrt{\|W_\vartheta[0, \tau_1]^{-1}\|}. \quad (14)$$

(H_2): The function $f : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and there exists a constant $q > 1$ and $L_f(\cdot) \in L^q(\Omega, \mathbb{R}^+)$ such that

$$\|f(t, \nu) - f(t, \mu)\| \leq L_f(t)\|\nu - \mu\|, \quad \nu, \mu \in \mathbb{R}^n.$$

Theorem 3.2. Suppose that (H_1) and (H_2) are satisfied. Then system (1) is relatively controllable provided that

$$c \left[1 + \frac{d\|D\|M}{a}(e^{a\tau_1} - 1) \right] < 1, \quad (15)$$

where $d = \left(\prod_{j=1}^k (1 + \|C_j\|e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta}$ and $c = \left(\prod_{j=1}^k (1 + \|C_j\|e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \left[\frac{1}{ap}(e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)}, \frac{1}{p} + \frac{1}{q} = 1, \quad p, q > 1.$

Proof. Let $\tilde{v} \in \mathbb{R}^n$ be the final state. Using hypothesis (H_1) for arbitrary $\nu(\cdot) \in \mathcal{PC}$, we define the control function $u_\nu(t)$ by

$$\begin{aligned} u_\nu(t) = & W^{-1}(\nu_1 - V(\tau_1, -\vartheta)\psi(-\vartheta) - \int_{-\vartheta}^0 V(\tau_1, s)[\psi'(s) - A\psi(s)]ds \\ & - \int_0^{\tau_1} V(\tau_1, s)f(s, \nu(s))ds)(t), \quad t \in \Omega. \end{aligned} \quad (16)$$

We show that, using this control, the operator $\mathcal{P} : \mathcal{PC} \rightarrow \mathcal{PC}$ defined by

$$\begin{aligned} (\mathcal{P}\nu)(t) = & V(t, -\vartheta)\psi(-\vartheta) + \int_{-\vartheta}^0 V(t, s)[\psi'(s) - A\psi(s)]ds \\ & + \int_0^t V(t, s)f(s, \nu(s))ds + \int_0^t V(t, s)Du_\nu(s)ds, \end{aligned}$$

has a fixed point ν , which is a mild solution of (1).

We check that $(\mathcal{P}\nu)(\tau_1) = \nu_1$, which means that u_ν steers the system (1) from $(\mathcal{P}\nu)(0)$ to ν_1 in finite time τ_1 . This implies system (1) is relatively controllable on Ω .

For each positive number r , let $\mathcal{B}_r = \{\nu \in \mathcal{PC} : \|\nu\|_{\mathcal{PC}} \leq r\}$ (a bounded, closed and convex set of \mathcal{PC}). Set $N = \sup_{t \in \Omega} \|f(t, 0)\|$.

We divide the proof into several steps.

Step 1. We claim that there exists a positive number r such that $\mathcal{F}(\mathcal{B}_r) \subseteq \mathcal{B}_r$.

From (H_2) and Hölder's inequality, we obtain that

$$\begin{aligned} \int_0^t e^{a(t-s)} L_f(s) ds & \leq \left(\int_0^t e^{ap(t-s)} ds \right)^{\frac{1}{p}} \left(\int_0^t L_f^q(s) ds \right)^{\frac{1}{q}} \\ & \leq \left[\frac{1}{ap}(e^{apt} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)}, \end{aligned}$$

and

$$\int_0^t e^{a(t-s)} \|f(s, 0)\| ds \leq N \int_0^t e^{a(t-s)} ds = \frac{N}{a}(e^{at} - 1).$$

From (16), using (H_1) and (H_2) , we have the following

$$\begin{aligned}
\|u_v(t)\| &\leq \|W^{-1}\|_{L_b(\mathbb{R}^n, L^2(\Omega, \mathbb{R}^n)/\ker W)} (\|v_1\| + \|V(\tau_1, -\vartheta)\| \|\psi(-\vartheta)\| \\
&\quad + \int_{-\vartheta}^0 \|V(\tau_1, s)\| \|\psi'(s) - A\psi(s)\| ds \\
&\quad + \int_0^{\tau_1} \|V(\tau_1, s)\| \|f(s, v(s))\| ds) \\
&\leq M \left(\|v_1\| + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(\tau_1+\vartheta)} \|\psi(-\vartheta)\| \right. \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_{-\vartheta}^0 e^{a(\tau_1-s)} \|\psi'(s) - A\psi(s)\| ds \\
&\quad + \left. \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_0^{\tau_1} e^{a(\tau_1-s)} (\|f(s, v(s)) - f(s, 0)\| + \|f(s, 0)\|) ds \right) \\
&\leq M \left(\|v_1\| + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(\tau_1+\vartheta)} \|\psi(-\vartheta)\| \right. \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_{-\vartheta}^0 e^{a(\tau_1-s)} \|\psi'(s) - A\psi(s)\| ds \\
&\quad + \left. \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_0^{\tau_1} e^{a(\tau_1-s)} (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \right) \\
&\leq M \left(\|v_1\| + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(\tau_1+\vartheta)} \|\psi(-\vartheta)\| \right. \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_{-\vartheta}^0 e^{a(\tau_1-s)} \|\psi'(s) - A\psi(s)\| ds \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \left[\frac{1}{ap} (e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} \|v\|_{PC} \\
&\quad + \left. \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \frac{N}{a} (e^{a\tau_1} - 1) \right) \\
&\leq M \|v_1\| + Mb + Mc \|v\|_{PC},
\end{aligned}$$

where

$$\begin{aligned}
b &= \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(\tau_1+\vartheta)} \|\psi(-\vartheta)\| \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_{-\vartheta}^0 e^{a(\tau_1-s)} \|\psi'(s) - A\psi(s)\| ds \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \frac{N}{a} (e^{a\tau_1} - 1).
\end{aligned}$$

From (H_1) and (H_2) we have the following

$$\begin{aligned}
\|(\mathcal{F}_1 v)(t)\| &\leq \|V(t, -\vartheta)\| \|\psi(-\vartheta)\| + \int_{-\vartheta}^0 \|V(t, s)\| \|\psi'(s) - A\psi(s)\| ds \\
&+ \int_0^t \|V(t, s)\| \|f(s, v(s))\| ds + \int_0^t \|V(t, s)\| \|D\| \|u_v(s)\| ds \\
&\leq \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t+\vartheta)} \|\psi(-\vartheta)\| \\
&+ \int_{-\vartheta}^0 \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} \|\psi'(s) - A\psi(s)\| ds \\
&+ \int_0^t \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
&+ \int_0^t \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} \|D\| [M\|v_1\| + Mb + Mc\|v\|_{PC}] ds \\
&\leq b + c\|v\|_{PC} + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \|D\| \\
&\quad \times [M\|v_1\| + Mb + Mc\|v\|_{PC}] \int_0^t e^{a(t-s)} ds \\
&\leq b \left[1 + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \frac{\|D\|M}{a} (e^{at} - 1) \right] \\
&\quad + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \frac{\|D\|M}{a} (e^{at} - 1) \|v_1\| \\
&\quad + c \left[1 + \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \frac{\|D\|M}{a} (e^{at} - 1) \right] \|v\|_{PC} \\
&\leq b \left[1 + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \right] + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \|v_1\| \\
&\quad + c \left[1 + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \right] r = r,
\end{aligned}$$

where

$$r = \frac{b \left[1 + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \right] + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \|v_1\|}{1 - c \left[1 + \frac{d\|D\|M}{a} (e^{a\tau_1} - 1) \right]}.$$

Hence, we obtain $\mathcal{F}(\mathcal{B}_r) \subseteq \mathcal{B}_r$ for such an r .

Now, we define operators \mathcal{F}_1 and \mathcal{F}_2 on \mathcal{B}_r as follows

$$(\mathcal{F}_1 v)(t) = V(t, -\vartheta) \psi(-\vartheta) + \int_{-\vartheta}^0 V(t, s) [\psi'(s) - A\psi(s)] ds + \int_0^t V(t, s) Du_v(s) ds,$$

and

$$(\mathcal{F}_2 v)(t) = \int_0^t V(t, s) f(s, v(s)) ds,$$

for $t \in \Omega$, respectively.

Step 2. We claim that \mathcal{F}_1 is a contraction mapping.

Let $v, \gamma \in \mathcal{B}_r$. From (H_1) and (H_2) , for each $\epsilon \in \Omega$, we have the following

$$\begin{aligned}
 \|u_v(t) - u_\gamma(t)\| &\leq M \int_0^{\tau_1} \|V(\tau_1, s)\| \|f(s, v(s)) - f(s, \gamma(s))\| ds \\
 &\leq M \int_0^{\tau_1} \left(\prod_{s < t_j \leq \tau_1} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(\tau_1-s)} L_f(s) (\|v(s) - \gamma(s)\|) ds \\
 &\leq M \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \left[\frac{1}{ap} (e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} \|v - \gamma\|_{PC} \\
 &\leq Mc \|v - \gamma\|_{PC}.
 \end{aligned}$$

Thus,

$$\begin{aligned}
 &\|(\mathcal{F}_1 v)(t) - (\mathcal{F}_1 \gamma)(t)\| \\
 &\leq \int_0^t \|V(t, s)\| \|D\| \|u_v(s) - u_\gamma(s)\| ds \\
 &\leq \int_0^t \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} ds \|D\| Mc \|v - \gamma\|_{PC} \\
 &\leq \frac{d\|D\|Mc}{a} (e^{a\tau_1} - 1) \|v - \gamma\|_{PC},
 \end{aligned}$$

so we obtain

$$\|\mathcal{F}_1 v - \mathcal{F}_1 \gamma\|_{PC} \leq P \|v - \gamma\|_{PC},$$

where $P = \frac{d\|D\|Mc}{a} (e^{a\tau_1} - 1)$. From (15), we have $P < 1$, so \mathcal{F}_1 is a contraction.

Step 3. We claim that $\mathcal{F}_2 : \mathcal{B}_r \rightarrow \mathcal{PC}$ is a compact and continuous operator.

Let $v_n \in \mathcal{B}_r$ with $v_n \rightarrow v$ in \mathcal{B}_r . Using (H_2) , we have $f(s, v_n(s)) \rightarrow f(s, v(s))$ in \mathcal{PC} and thus using the Lebesgue dominated convergence theorem we have

$$\begin{aligned}
 &\|(\mathcal{F}_2 v_n)(t) - (\mathcal{F}_2 v)(t)\| \\
 &\leq \int_0^t \|V(t, s)\| \|f(s, v_n(s)) - f(s, v(s))\| ds \\
 &\leq \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} \int_0^t e^{a(t-s)} \|f(s, v_n(s)) - f(s, v(s))\| ds \\
 &\rightarrow 0 \text{ as } n \rightarrow \infty,
 \end{aligned}$$

which implies that \mathcal{F}_2 is continuous on \mathcal{B}_r .

To check the compactness of $\mathcal{F}_2 : \mathcal{B}_r \rightarrow \mathcal{PC}$, we prove that $\mathcal{F}_2(\mathcal{B}_r)$ is equicontinuous and uniformly bounded. In fact, for any $v \in \mathcal{B}_r$, $t_z < t \leq t + h \leq t_{z+1}$, $z = 0, 1, \dots, k$,

$$\begin{aligned}
 &(\mathcal{F}_2 v)(t+h) - (\mathcal{F}_2 v)(t) \\
 &= \int_0^{t+h} V(t+h, s) f(s, v(s)) ds - \int_0^t V(t, s) f(s, v(s)) ds \\
 &= \int_t^{t+h} V(t+h, s) f(s, v(s)) ds \\
 &\quad + \int_0^t e^{A(t+h-s)} (X(t+h, s+\vartheta) - X(t, s+\vartheta)) f(s, v(s)) ds \\
 &\quad + \int_0^t (e^{A(t+h-s)} - e^{A(t-s)}) X(t, s+\vartheta) f(s, v(s)) ds.
 \end{aligned}$$

Let

$$\begin{aligned}
 S_1 &= \int_t^{t+h} V(t+h, s) f(s, v(s)) ds, \\
 S_2 &= \int_0^t e^{A(t+h-s)} (X(t+h, s+\vartheta) - X(t, s+\vartheta)) f(s, v(s)) ds, \\
 S_3 &= \int_0^t (e^{A(t+h-s)} - e^{A(t-s)}) X(t, s+\vartheta) f(s, v(s)) ds \\
 &= \int_0^t (e^{Ah} - E) V(t, s) f(s, v(s)) ds.
 \end{aligned}$$

From above, we see that

$$\|(\mathcal{F}_2 v)(t+h) - (\mathcal{F}_2 v)(t)\| \leq \|S_1\| + \|S_2\| + \|S_3\|.$$

Now, we only need to check $\|S_i\| \rightarrow 0$ as $h \rightarrow 0$, $i = 1, 2, 3$. Clearly,

$$\begin{aligned}
 \|S_1\| &\leq \int_t^{t+h} \|V(t+h, s)\| \|f(s, v(s))\| ds \\
 &\leq \int_t^{t+h} \left(\prod_{s < t_j \leq t+h} (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{-\|\check{B}\|\vartheta} e^{a(t+h-s)} (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
 &\leq e^{-\|\check{B}\|\vartheta} \left[\frac{1}{ap} (e^{ap} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} \|v\|_{PC} + e^{-\|\check{B}\|\vartheta} \frac{N}{a} (e^{ah} - 1) \\
 &\rightarrow 0 \text{ as } h \rightarrow 0, \\
 \|S_2\| &\leq \int_0^t \|e^{A(t+h-s)}\| \|X(t+h, s+\vartheta) - X(t, s+\vartheta)\| \|f(s, v(s))\| ds \\
 &\leq e^{\|A\|\tau_1} \int_0^t \left\| e_{\vartheta}^{\check{B}(t+h-\vartheta-s)} - e_{\vartheta}^{\check{B}(t-\vartheta-s)} + \sum_{s < t_j \leq t+h} C_j e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} X(t_j, s+\vartheta) \right. \\
 &\quad \left. - \sum_{s < t_j \leq t} C_j e_{\vartheta}^{\check{B}(t-\vartheta-t_j)} X(t_j, s+\vartheta) \right\| (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
 &\leq e^{\|A\|\tau_1} \int_0^t \left\| e_{\vartheta}^{\check{B}(t+h-\vartheta-s)} - e_{\vartheta}^{\check{B}(t-\vartheta-s)} \right. \\
 &\quad \left. + \sum_{s < t_j \leq t} C_j (e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} - e_{\vartheta}^{\check{B}(t-\vartheta-t_j)}) X(t_j, s+\vartheta) \right\| (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
 &\leq e^{\|A\|\tau_1} \int_0^t \|e_{\vartheta}^{\check{B}(t+h-\vartheta-s)} - e_{\vartheta}^{\check{B}(t-\vartheta-s)}\| (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
 &\quad + \sum_{j=1}^k e^{\|A\|\tau_1} \int_0^t \|C_j (e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} - e_{\vartheta}^{\check{B}(t-\vartheta-t_j)}) X(t_j, s+\vartheta)\| L_f(s) \|v(s)\| ds \\
 &\quad + \sum_{j=1}^k e^{\|A\|\tau_1} \int_0^t \|C_j (e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} - e_{\vartheta}^{\check{B}(t-\vartheta-t_j)}) X(t_j, s+\vartheta)\| \|f(s, 0)\| ds \\
 &\leq e^{\|A\|\tau_1} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} r \left(\int_0^t \|e_{\vartheta}^{\check{B}(t+h-\vartheta-s)} - e_{\vartheta}^{\check{B}(t-\vartheta-s)}\|^p ds \right)^{\frac{1}{p}} \\
 &\quad + e^{\|A\|\tau_1} N \int_0^t \|e_{\vartheta}^{\check{B}(t+h-\vartheta-s)} - e_{\vartheta}^{\check{B}(t-\vartheta-s)}\| ds \\
 &\quad + \sum_{j=1}^k e^{\|A\|\tau_1} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} r \left(\int_0^t \|C_j (e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} - e_{\vartheta}^{\check{B}(t-\vartheta-t_j)}) X(t_j, s+\vartheta)\|^p ds \right)^{\frac{1}{p}} \\
 &\quad + \sum_{j=1}^k e^{\|A\|\tau_1} N \int_0^t \|C_j (e_{\vartheta}^{\check{B}(t+h-\vartheta-t_j)} - e_{\vartheta}^{\check{B}(t-\vartheta-t_j)}) X(t_j, s+\vartheta)\| ds \\
 &\rightarrow 0 \text{ as } h \rightarrow 0,
 \end{aligned}$$

and

$$\begin{aligned}
\|S_3\| &\leq \int_0^t \|e^{Ah} - E\| \|V(t, s)\| \|f(s, v(s))\| ds \\
&\leq \|e^{Ah} - E\| \int_0^t \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
&\leq \|e^{Ah} - E\| d \left(\left[\frac{1}{ap} (e^{apt} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} r + \frac{N}{a} (e^{at} - 1) \right) \\
&\rightarrow 0 \text{ as } h \rightarrow 0.
\end{aligned}$$

As a result, we immediately obtain that

$$\|(\mathcal{F}_2 v)(t+h) - (\mathcal{F}_2 v)(t)\| \rightarrow 0 \text{ as } h \rightarrow 0,$$

for all $v \in \mathcal{B}_r$. Therefore, $\mathcal{F}_2(\mathcal{B}_r)$ is equicontinuous in \mathcal{PC} .

Next, repeating the above computations, we have the following

$$\begin{aligned}
\|(\mathcal{F}_2 v)(t)\| &\leq \int_0^t \|V(t, s)\| \|f(s, v(s))\| ds \\
&\leq \int_0^t \left(\prod_{s < t_j \leq t} (1 + \|C_j\| e^{-\|\tilde{B}\|\vartheta}) \right) e^{-\|\tilde{B}\|\vartheta} e^{a(t-s)} (L_f(s) \|v(s)\| + \|f(s, 0)\|) ds \\
&\leq d \left[\frac{1}{ap} (e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} r + \frac{dN}{a} (e^{a\tau_1} - 1).
\end{aligned}$$

Hence $\mathcal{F}_2(\mathcal{B}_r)$ is uniformly bounded. From Theorem 2.5, $\mathcal{F}_2(\mathcal{B}_r)$ is relatively compact in \mathcal{PC} . Thus, $\mathcal{F}_2 : \mathcal{B}_r \rightarrow \mathcal{PC}$ is a compact and continuous operator.

Hence, using the Krasnoselskii's fixed point theorem, \mathcal{F} has a fixed point v on \mathcal{B}_r . Obviously, v is a solution of the system (1) satisfying $v(\tau_1) = v_1$. The boundary condition $v(t) = \psi(t)$, $-\vartheta \leq t \leq 0$ holds from (7). The proof is complete. \square

Remark 3.3. Now, we make a comparison with.²⁹ Let $C_i = \Theta$, $i = 1, 2, \dots, k$, and then system (1) reduces to.^{29, (3)} For relative controllability results for the linear system, one has $V(\cdot, \cdot) = e^{A(\cdot-\cdot)} e_{\vartheta}^{\tilde{B}(-\vartheta-\cdot)}$ and the Grammian matrix (10) which is the same as in.^{29, (10)} For the nonlinear system, set $C_i = \Theta$, $a = N$, $c = e^{-\|\tilde{B}\|\vartheta} M_2$, $d = e^{-\|\tilde{B}\|\vartheta}$, and then (15) becomes

$$e^{-\|\tilde{B}\|\vartheta} M_2 \left[1 + \frac{e^{-\|\tilde{B}\|\vartheta} \|D\| M}{N} (e^{N\tau_1} - 1) \right] < 1. \quad (17)$$

Now, (17) with $M_2[1 + \frac{M}{N}(e^{N\tau_1} - 1)\|C\|] < 1$,^{29, (23)} and one see that the factor (17) improves^{29, (23)} since $e^{-\|\tilde{B}\|\vartheta}$ is in (17).

4 | EXAMPLES

Example 4.1. Set $\tau_1 = 1$. Consider the following semilinear impulsive delay differential controlled system

$$\begin{cases} v'(t) = Av(t) + Bv(t-0.5) + f(t, v(t)) + Du(t), & t \in \Omega := [0, 1], t \notin \mathcal{T}, \\ \Delta v(t_i) = C_i v(t_i), & t_i \in \mathcal{T} := \{0.5\}, \\ \psi(t) = (0.2, 0.3)^T, & -0.5 \leq t \leq 0, \end{cases} \quad (18)$$

where we set $k = 1$, $\vartheta = 0.5$, $\mathcal{T} = \{0.5\}$ and

$$\begin{aligned} A &= \begin{pmatrix} 0.2 & 0 \\ 0 & 0.2 \end{pmatrix}, \quad B = \begin{pmatrix} 0.2 & 0.1 \\ 0 & 0.3 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \\ C_i &= \begin{pmatrix} 0.3 & 0.1 \\ 0 & 0.4 \end{pmatrix}, \quad i = 1, \quad f(t, v(t)) = \begin{pmatrix} 0.2tv_1(t) \\ 0.3tv_2(t) \end{pmatrix}. \end{aligned}$$

Clearly $\|A\| = 0.2$, $\|\check{B}\| = 0.2715$, $\|D\| = 1$ and $\|C_i\| = 0.4 (i = 1)$. Also,

$$\begin{aligned} AB &= \begin{pmatrix} 0.04 & 0.02 \\ 0 & 0.06 \end{pmatrix} = BA, \quad AC_i = \begin{pmatrix} 0.06 & 0.02 \\ 0 & 0.08 \end{pmatrix} = C_i A, \\ BC_i &= \begin{pmatrix} 0.06 & 0.06 \\ 0 & 0.12 \end{pmatrix} = C_i B, \quad i = 1, \\ \check{B} &= e^{-A\vartheta} B = \begin{pmatrix} 1.8221 & 0.3644 \\ 0 & 1.4577 \end{pmatrix}. \end{aligned}$$

Clearly $a = \|A\| + \|\check{B}\| = 0.4715$.

Now, we use (14) to obtain M . For this purpose, we need to obtain $W_\vartheta[0, \tau_1]$ and then $W_\vartheta[0, \tau_1]^{-1}$. The delay Grammian matrix has the following explicit form as follows

$$\begin{aligned} W_\vartheta[0, \tau_1] &= \int_0^{\tau_1} V(\tau_1, s) D D^T V^T(\tau_1, s) ds \\ &= \int_0^1 e^{A(1-s)} X(1, s + 0.5) X^T(1, s + 0.5) e^{A^T(1-s)} ds \\ &= W_1 + W_2, \end{aligned}$$

where

$$W_1 = \int_0^{0.5} e^{A(1-s)} [E + \check{B}(0.5 - s) + C_1] \times [E + \check{B}^T(0.5 - s) + C_1^T] e^{A^T(1-s)} ds,$$

and

$$W_2 = \int_{0.5}^1 e^{A(1-s)} E^2 e^{A^T(1-s)} ds.$$

Note

$$W_1 = \begin{pmatrix} 1.2371 & 0.1230 \\ 0.1230 & 1.4622 \end{pmatrix}, \quad W_2 = \begin{pmatrix} 0.5535 & 0 \\ 0 & 0.5535 \end{pmatrix}.$$

Therefore, we obtain

$$W_\vartheta[0, 1] = \begin{pmatrix} 1.7906 & 0.1230 \\ 0.1230 & 2.0157 \end{pmatrix}, \quad W_\vartheta[0, 1]^{-1} = \begin{pmatrix} 0.5608 & -0.0342 \\ -0.0342 & 0.4982 \end{pmatrix},$$

and

$$M = \sqrt{\|W_\vartheta[0, 1]\|} = \sqrt{0.5266} = 0.7257.$$

For any $v, \mu \in \mathbb{R}^n$,

$$\begin{aligned} \|f(t, v) - f(t, \mu)\| &= \max\{0.2t\|v_1 - \mu_1\|, 0.3t\|v_2 - \mu_2\|\} \\ &\leq 0.3t \max\{\|v_1 - \mu_1\|, \|v_2 - \mu_2\|\} \\ &= 0.3t\|v - \mu\|. \end{aligned}$$

Now, we set $L_f(t) = 0.3t$, $L_f \in L^q(\Omega, \mathbb{R}^+)$ with $p = q = 2$. Note $\|L_f\|_{L^2(\Omega, \mathbb{R}^+)} = (\int_0^1 (0.3s)^2 ds)^{\frac{1}{2}} = 0.1732$,

$$\begin{aligned} d &= \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{-\|\check{B}\|\vartheta} \\ &= (1 + \|C_1\| e^{-0.5\|\check{B}\|})^2 e^{-0.5\|\check{B}\|} = 1.3681, \\ c &= \left(\prod_{j=1}^k (1 + \|C_j\| e^{-\|\check{B}\|\vartheta}) \right) e^{-\|\check{B}\|\vartheta} \left[\frac{1}{ap} (e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} \\ &= d \left[\frac{1}{ap} (e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} = 0.3055, \end{aligned}$$

and

$$c \left[1 + \frac{d\|D\|M}{a}(e^{a\tau_1} - 1) \right] = 0.3055 \left[1 + \frac{1.3681 \times 0.7257}{0.4715}(e^{0.4715} - 1) \right] = 0.6930 < 1.$$

Thus all the conditions of Theorem 3.2 are satisfied, so (18) is relatively controllable on $[0, 1]$.

Example 4.2. In Example 4.1, we keep the other parameters unchanged except f and let

$$f(t, v(t)) = \begin{pmatrix} 0.3t^2 v_1(t) \sin t \\ 0.4t v_2(t) \cos 2t \end{pmatrix}. \quad (19)$$

For any $t \in \Omega$ and $v, \mu \in \mathbb{R}^n$,

$$\begin{aligned} \|f(t, v) - f(t, \mu)\| &= \max\{0.3t^2\|(v_1 - \mu_1) \sin t\|, 0.4t\|(v_2 - \mu_2) \cos 2t\|\} \\ &\leq 0.4t \max\{\|v_1 - \mu_1\|, \|v_2 - \mu_2\|\} \\ &= 0.4t\|v - \mu\|. \end{aligned}$$

Set $L_f(t) = 0.4t$, $L_f \in L^q(\Omega, \mathbb{R}^+)$ with $p = q = 2$. Note

$$\begin{aligned} \|L_f\|_{L^2(\Omega, \mathbb{R}^+)} &= \left(\int_0^1 (0.4s)^2 ds \right)^{\frac{1}{2}} = 0.2309, \\ c &= d \left[\frac{1}{ap}(e^{ap\tau_1} - 1) \right]^{\frac{1}{p}} \|L_f\|_{L^q(\Omega, \mathbb{R}^+)} = 0.4073, \end{aligned}$$

and

$$c \left[1 + \frac{d\|D\|M}{a}(e^{a\tau_1} - 1) \right] = 0.4073 \left[1 + \frac{1.3681 \times 0.7257}{0.4715}(e^{0.4715} - 1) \right] = 0.9239 < 1.$$

From Theorem 3.2, we know (18) with (19) is relatively controllable on $[0, 1]$.

Example 4.3. Consider the relative controllability of system (18) (with $f(t, v(t)) \equiv \mathbf{0}$) on Ω , where A, B, C_i, D are defined in Example 4.1.

According to Theorem 3.1, we know that system (18) is relative controllability when $f(t, v(t)) = \mathbf{0}$. Further, one can get

$$\eta = v_1 - V(1, -0.5)\psi(-0.5) - \int_{-0.5}^0 V(1, s)[\psi'(s) - A\psi(s)]ds = v_1 - \begin{pmatrix} -0.0444 \\ -0.0630 \end{pmatrix}.$$

By using the form of the control $u(t)$, we arrive at

$$\begin{aligned} u(t) &= D^T V^T(\tau_1, t) W_\theta^{-1}[0, \tau_1] \eta \\ &= X^T(1, t + 0.5) e^{A^T(1-t)} W_\theta^{-1}[0, 1] \eta \\ &= \begin{cases} [E + \check{B}^T(0.5 - t) + C_1^T] e^{A^T(1-t)} W_\theta^{-1}[0, 1] \eta, & 0 \leq t < 0.5, \\ E e^{A^T(1-t)} W_\theta^{-1}[0, 1] \eta, & 0.5 \leq t < 1, \end{cases} \end{aligned}$$

where \check{B} and $W_\theta^{-1}[0, 1]$ are given in Example 4.1.

5 | CONCLUSIONS

The main contribution of this paper is to introduce a relative controllability method for impulsive delay controlled systems with linear parts defined by permutable matrices. We use the impulsive delayed Cauchy matrix to construct the impulsive delay Grammian matrix. We give a sufficient and necessary condition to examine when a linear delay controlled system is relatively controllable. Also, we apply a fixed point method to establish a relative controllability result for impulsive semilinear delay controlled systems. The study on controllability of delay differential systems with impulses has potential for future research to multiple delay problems or to problems with a nonlinear term f depending on delayed arguments.

ACKNOWLEDGEMENTS

The authors are grateful to the referees for their careful reading of the manuscript and their valuable comments. We also thank the editor.

This work was supported by the National Natural Science Foundation of China (11661016), Training Object of High Level and Innovative Talents of Guizhou Province ([2016]4006), Science and Technology Program of Guizhou Province ([2017]5788-10), Major Research Project of Innovative Group in Guizhou Education Department ([2018]012), and Foundation of Postgraduate of Guizhou Province (YJSCXJH[2018]035).

ORCID

JinRong Wang  <http://orcid.org/0000-0002-6642-1946>

Yong Zhou  <http://orcid.org/0000-0002-4099-8077>

REFERENCES

1. Ya. Khusainov D, Shuklin GV. Linear autonomous time-delay system with permutation matrices solving. *Stud Univ Žilina*. 2003;17:101-108.
2. Diblík J, Ya. Khusainov D. Representation of solutions of discrete delayed system $x(k+1) = Ax(k) + Bx(k-m) + f(k)$ with commutative matrices. *J Math Anal Appl*. 2006;318:63-76.
3. Ya. Khusainov D, Shuklin GV. Relative controllability in systems with pure delay. *Int J Appl Math*. 2005;2:210-221.
4. Medveď M, Pospíšil M, Škripková L. Stability and the nonexistence of blowing-up solutions of nonlinear delay systems with linear parts defined by permutable matrices. *Nonlinear Anal*. 2011;74:3903-3911.
5. Medveď M, Pospíšil M. Sufficient conditions for the asymptotic stability of nonlinear multidelay differential equations with linear parts defined by pairwise permutable matrices. *Nonlinear Anal*. 2012;75:3348-3363.
6. Diblík J, Fečkan M, Pospíšil M. Representation of a solution of the Cauchy problem for an oscillating system with two delays and permutable matrices. *Ukrainian Math J*. 2013;65:58-69.
7. Diblík J, Ya. Khusainov D, Răužičková M. Controllability of linear discrete systems with constant coefficients and pure delay. *SIAM J Control Optim*. 2008;47:1140-1149.
8. Diblík J, Fečkan M, Pospíšil M. On the new control functions for linear discrete delay systems. *SIAM J Control Optim*. 2014;52:1745-1760.
9. Diblík J, Morávková B. Discrete matrix delayed exponential for two delays and its property. *Adv Diff Equ*. 2013;2013:1-18.
10. Diblík J, Morávková B. Representation of the solutions of linear discrete systems with constant coefficients and two delays. *Abstr Appl Anal*. 2014;2014:1-19.
11. Diblík J, Ya. Khusainov D, Baštinec J, Sirenko AS. Exponential stability of linear discrete systems with constant coefficients and single delay. *Appl Math Lett*. 2016;51:68-73.
12. Boichuk A, Diblík J, Khusainov D, Răužičková M. Fredholm's boundary-value problems for differential systems with a single delay. *Nonlinear Anal*. 2010;72:2251-2258.
13. Pospíšil M. Representation and stability of solutions of systems of functional differential equations with multiple delays. *Electron J Qual Theory Differ Equ*. 2012;54:1-30.
14. Pospíšil M. Representation of solutions of delayed difference equations with linear parts given by pairwise permutable matrices via Z-transform. *Appl Math Comput*. 2017;294:180-194.
15. Luo Z, Wei W, Wang J. On the finite time stability of nonlinear delay differential equations. *Nonlinear Dyn*. 2017;89:713-722.
16. Liang C, Wang J, O'Regan D. Controllability of nonlinear delay oscillating systems. *Electron J Qual Theory Differ Equ*. 2017;2017(47):1-18.
17. Li M, Wang J. Exploring delayed Mittag-Leffler type matrix functions to study finite time stability of fractional delay differential equations. *Appl Math Comput*. 2018;324:254-265.
18. Cao X, Wang J. Finite-time stability of a class of oscillating systems with two delays. *Math Meth Appl Sci*. 2018;41:4943-4954.
19. Li L, Meng F, Ju P. Some new integral inequalities and their applications in studying the stability of nonlinear integro differential equations with time delay. *J Math Anal Appl*. 2011;377:853-862.
20. Meng Q, Feng F, Zheng B. Gronwall–Bellman type nonlinear delay integral inequalities on times scales. *J Math Anal Appl*. 2011;382:772-784.
21. Yuan Z, Yuan X, Meng F, Zhang H. Some new delay integral inequalities and their applications. *Appl Math Comput*. 2009;208:231-237.
22. Bai Y, Guo C. New results on stability and boundedness of third order nonlinear delay differential equations. *Dyn Sys Appl*. 2013;22:95-104.
23. Guo Y. Mean square global asymptotic stability of stochastic recurrent neural networks with distributed delays. *Appl Math Comput*. 2009;215:791-795.
24. You Z, Wang J. Stability of impulsive delay differential equations. *J App Math Comput*. 2018;56:253-268.
25. You Z, Wang J. On the exponential stability of nonlinear delay systems with impulses. *IMA J Math Control Inform*. 2018;35:773-803.
26. Khusainov DY, Shuklin GV. Relative controllability in systems with pure delay. *Int J Appl Math*. 2005;2:210-221.

27. Krasnoselskii M. *Topological Methods in the Theory of Nonlinear Integral Equations*. New York: Pergamon Press; 1964.
28. Wei W, Xiang X, Peng Y. Nonlinear impulsive integro-differential equation of mixed type and optimal controls. *Optimization*. 2006;55:141-156.
29. Wang J, Luo Z, Fečkan M. Relative controllability of semilinear delay differential systems with linear parts defined by permutable matrices. *European J Control*. 2017;38:39-46.

How to cite this article: You Z, Wang J, O'Regan D, Zhou Y. Relative controllability of delay differential systems with impulses and linear parts defined by permutable matrices. *Math Meth Appl Sci*. 2018;1-15. <https://doi.org/10.1002/mma.5400>