

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350173364>

Lyapunov-based stability of delayed linear differential algebraic systems

Article in *Applied Mathematics Letters* · August 2021

DOI: 10.1016/j.aml.2021.107185

CITATIONS

0

READS

26

2 authors, including:




Ann Al Sawoor

Université de Bretagne Occidentale

17 PUBLICATIONS 53 CITATIONS

SEE PROFILE



Lyapunov-based stability of delayed linear differential algebraic systems



Ann Al Sawoor, Miloud Sadkane^{*}

Univ Brest, CNRS - UMR 6205, Laboratoire de Mathématiques de Bretagne Atlantique. 6, Av. Le Gorgeu. 29238 Brest Cedex 3, France

ARTICLE INFO

Article history:

Received 24 December 2020

Received in revised form 3 March 2021

Accepted 3 March 2021

Available online xxxx

Keywords:

Linear differential algebraic systems

Time delay

Lyapunov stability

Decay rate

ABSTRACT

The Lyapunov stability theorem for linear systems is extended to linear delay-differential algebraic systems of index one. In particular, bounds on the decay of the solution are established in terms of solvability of a certain Lyapunov-type matrix equation.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

It is known that if A is a stable matrix (i.e., all of its eigenvalues have negative real parts), then the linear ordinary differential system

$$\dot{x}(t) = Ax(t), \quad t \geq 0 \quad (1)$$

is asymptotically stable, that is, $x(t) \rightarrow 0$ as $t \rightarrow +\infty$.

The Lyapunov theory can be used to inform about the quality of stability and to measure the decay rate of the solution (see, e.g., [1]). Indeed, under the assumption that A is stable, the Lyapunov matrix equation

$$C = -(A^T H + H A) \quad (2)$$

has the unique solution $H = \int_0^\infty e^{tA^T} C e^{tA} dt = H^T > 0$ (i.e., symmetric positive definite) for all matrices $C = C^T > 0$. Conversely, if Eq. (2) is satisfied with some matrices $C = C^T > 0$ and $H = H^T > 0$, then A is stable and, using the quadratic function

$$v(x(t)) = x(t)^T H x(t), \quad (3)$$

^{*} Corresponding author.

E-mail addresses: Ann.AlSawoorMaskoni@univ-brest.fr (A.A. Sawoor), miloud.sadkane@univ-brest.fr (M. Sadkane).

we obtain the bound (see, e.g., [2])

$$\|x(t)\| \leq \sqrt{\|H\| \|H^{-1}\|} e^{-\frac{\lambda_{\min}(C)}{2\|H\|}t} \|x(0)\|, \quad t \geq 0, \quad (4)$$

where $\|\cdot\|$ denotes the 2-norm for vectors and matrices and $\lambda_{\min}(C)$ denotes the smallest eigenvalue of C . See also [3] for a comparison with other bounds. Although this bound is not sharp, it measures the decay rate of the solution (because $\lambda_{\min}(C) > 0$) by involving quantities based on the computation of eigenvalues of symmetric (positive definite) matrices for which reliable algorithms are available [4]. Moreover, the norm of H can be used in preference to the spectral criterion “ $\operatorname{Re}(\lambda) < 0$ for all eigenvalues of A ” for assessing the stability of A : the larger the norm of H , the less stable the matrix A . For related issues, see, e.g., [5–7].

Our goal in this note is to study to what extent a bound similar to (4) holds for linear differential algebraic systems with time delay, of the form

$$\begin{cases} E\dot{x}(t) = Ax(t) + Bx(t - \tau), & t > 0, \\ x(t) = \psi(t), & -\tau \leq t \leq 0, \end{cases} \quad (5)$$

where E, A, B are real $n \times n$ matrices with E singular and τ is a fixed positive delay.

Time delay systems arise generally in applications where a transport phenomenon occurs. A wide range of examples can be found in [8,9]. Solution theory for (5) is established for example in [10–14] and the references therein.

Stability analysis of systems of type (5) has been investigated by many authors; for example, in [15,16] the analysis is based on a careful choice of a Lyapunov–Krasovskii function (a generalization of (3)), and sufficient conditions for stability are given in terms of linear matrix inequalities. In [17], the stability is based on the computation of eigenvalues of certain matrix pencils. In [18], sufficient conditions for asymptotic stability are formulated with the help of the characteristic equation of the system. In [19], a spectrum-based approach is developed for the stability analysis and stabilization of systems described by delay differential algebraic equations. In [20], necessary and sufficient conditions for exponential stability of classes of systems of the form (5) are derived using the roots of the associated characteristic equation. In the present note, using a Lyapunov–Krasovskii type function, bounds on the decay of the solution, analogous to (4), are established in terms of solvability of a certain Lyapunov-type matrix equation.

2. Lyapunov-Based stability analysis

We assume throughout this note that the pencil $\lambda E - A$ is regular (i.e., there exists λ such that $\det(\lambda E - A) \neq 0$) and of index 1 (this is the index of the nilpotent matrix in the Weierstraß canonical form of $\lambda E - A$, see, e.g., [21]). Since the matrices E and A are real, the real Weierstraß canonical form can be used to decompose E and A as

$$E = W \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} T, \quad A = W \begin{pmatrix} J & 0 \\ 0 & I_{n-r} \end{pmatrix} T, \quad (6)$$

where W and T are real nonsingular matrices, the symbol I_k denotes the identity matrix of order k , the matrix J is in real Jordan form (see, e.g., [22, Section 3.4]) and corresponds to the finite eigenvalues of the pencil $\lambda E - A$ and $r = \operatorname{rank}(E)$. Alternatively, the quasi-Weierstraß form or a block-diagonalization via the real generalized Schur form may also be used (see [23,24]).

The spectral projection onto the right deflating subspace of $\lambda E - A$ corresponding to the finite eigenvalues is given by

$$\mathcal{P} = \frac{1}{2\pi i} \oint_{\Gamma} (\lambda E - A)^{-1} E \, d\lambda, \quad (7)$$

where Γ is a closed Jordan curve surrounding the finite eigenvalues. Using (6), we obtain

$$\mathcal{P} = T^{-1} \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} T. \quad (8)$$

As a first step, a bound on $\mathcal{P}x(t)$ analogous to (4) is obtained in Theorem 2.1. Then, Theorem 2.2 extends the bounds to $x(t)$. In these theorems, the assumption that the pencil has index 1 plays an essential role (see the function $v(t, y(t))$ in the proof of Theorem 2.1 and the property (16)).

From (6) and (8) the system (5) can be written

$$\begin{cases} \hat{E}\dot{y}(t) = \hat{A}y(t) + \hat{B}y(t - \tau), & t > 0, \\ y(t) = \hat{\psi}(t), & -\tau \leq t \leq 0, \end{cases} \quad (9)$$

where

$$\begin{cases} \hat{E} = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}, \quad \hat{A} = \begin{pmatrix} J & 0 \\ 0 & I_{n-r} \end{pmatrix}, \quad \hat{B} = W^{-1}BT^{-1}, \\ y(t) = Tx(t), \quad \hat{\psi}(t) = T\psi(t). \end{cases} \quad (10)$$

With this notation, we obtain the following result.

Theorem 2.1. Assume that there exists

$$H = \begin{pmatrix} H_{11} & 0 \\ 0 & H_{22} \end{pmatrix} \quad \text{with } H_{11} = H_{11}^T > 0, \quad H_{22} + H_{22}^T + \frac{1}{\tau}I_{n-r} < 0$$

such that

$$C = - \begin{pmatrix} \hat{A}^T H + H^T \hat{A} + \frac{1}{\tau}I_n & H^T \hat{B} \\ \hat{B}^T H & -\frac{1}{2\tau}I_n \end{pmatrix} = C^T > 0.$$

Then

- (i) the pencil $\lambda E - A$ is stable (i.e., the eigenvalues of J lie in the open left-half plane).
- (ii) The following bound holds for $t \geq 0$

$$\|\mathcal{P}x(t)\| \leq \alpha e^{-\frac{1}{2} \frac{\beta}{\|H_{11}\|} t},$$

where

$$\begin{aligned} \alpha &= \|T\| \|T^{-1}\| \sqrt{\|H_{11}^{-1}\| (\|H_{11}\| + \ln 2)} \max_{-\tau \leq \nu \leq 0} \|\mathcal{P}x(\nu)\|, \\ \beta &= \min \left(\lambda_{\min}(C), \frac{\|H_{11}\|}{2\tau} \right). \end{aligned}$$

Proof.

- (i) Let (λ, x_1) be an eigenpair of J and $\hat{x} = \begin{pmatrix} x_1 \\ 0 \end{pmatrix}$ and $x = \begin{pmatrix} \hat{x} \\ 0 \end{pmatrix}$ be respectively of size n and $2n$. Then, denoting by x^* the conjugate transpose of x , we obtain

$$-x^* C x = 2x_1^* H_{11} x_1 \operatorname{Re} \lambda + \frac{\|x_1\|^2}{\tau}$$

which shows that $\operatorname{Re}(\lambda) < 0$.

- (ii) Consider the Lyapunov–Krasovskii type function

$$v(t, y(t)) = y^T(t) \hat{E} H y(t) + \int_{t-\tau}^t \frac{\|y(\nu)\|^2}{t - \nu + \tau} d\nu, \quad t \geq 0.$$

If we set $y(t) = \begin{pmatrix} y_1(t) \\ y_2(t) \end{pmatrix}$, then $y^T(t)\hat{E}Hy(t) = y_1^T(t)H_{11}y_1(t) \geq 0$ and

$$\begin{aligned} \dot{v}(t, y(t)) &= y^T(t)H^T\hat{E}\dot{y}(t) + (\hat{E}\dot{y}(t))^THy(t) + \frac{\|y(t)\|^2}{\tau} - \frac{\|y(t-\tau)\|^2}{2\tau} \\ &\quad - \int_{t-\tau}^t \frac{\|y(\nu)\|^2}{(t-\nu+\tau)^2} d\nu \\ &= y(t)^TH^T \left(\hat{A}y(t) + \hat{B}y(t-\tau) \right) + \left(\hat{A}y(t) + \hat{B}y(t-\tau) \right)^T Hy(t) \\ &\quad + \frac{\|y(t)\|^2}{\tau} - \frac{\|y(t-\tau)\|^2}{2\tau} - \int_{t-\tau}^t \frac{\|y(\nu)\|^2}{(t-\nu+\tau)^2} d\nu \\ &= - \begin{pmatrix} y(t) \\ y(t-\tau) \end{pmatrix}^T C \begin{pmatrix} y(t) \\ y(t-\tau) \end{pmatrix} - \int_{t-\tau}^t \frac{\|y(\nu)\|^2}{(t-\nu+\tau)^2} d\nu \leq 0. \end{aligned} \quad (11)$$

Note that

$$\begin{aligned} \begin{pmatrix} y(t) \\ y(t-\tau) \end{pmatrix}^T C \begin{pmatrix} y(t) \\ y(t-\tau) \end{pmatrix} &\geq \lambda_{\min}(C) \left\| \begin{pmatrix} y(t) \\ y(t-\tau) \end{pmatrix} \right\|^2 \geq \lambda_{\min}(C) \|y(t)\|^2 \\ &\geq \frac{\lambda_{\min}(C)}{\|H_{11}\|} y^T(t)\hat{E}Hy(t) \end{aligned} \quad (12)$$

$$t - \tau \leq \nu \leq t \Rightarrow (t - \nu + \tau)^2 \leq 2\tau(t - \nu + \tau). \quad (13)$$

Using (11), (12) and (13) we obtain

$$\dot{v}(t, y(t)) \leq -\frac{\beta}{\|H_{11}\|} v(t, y(t)), \quad \beta = \min \left(\lambda_{\min}(C), \frac{\|H_{11}\|}{2\tau} \right).$$

Therefore

$$v(t, y(t)) \leq e^{-\frac{\beta}{\|H_{11}\|}t} v(0, y(0)).$$

Since

$$\begin{aligned} v(t, y(t)) &\geq y^T(t)\hat{E}Hy(t) = y_1^T(t)H_{11}y_1(t) \\ &\geq \lambda_{\min}(H_{11}) \|y_1(t)\|^2 = \|H_{11}^{-1}\|^{-1} \|y_1(t)\|^2, \\ v(0, y(0)) &= y^T(0)\hat{E}Hy(0) + \int_{-\tau}^0 \frac{\|y(\nu)\|^2}{-\nu+\tau} d\nu \\ &\leq \|\hat{E}H\| \|y(0)\|^2 + \max_{-\tau \leq \nu \leq 0} \|y(\nu)\|^2 \ln 2 \\ &\leq (\|H_{11}\| + \ln 2) \max_{-\tau \leq \nu \leq 0} \|y(\nu)\|^2, \end{aligned}$$

we deduce that

$$\|y_1(t)\|^2 \leq \|H_{11}^{-1}\| (\|H_{11}\| + \ln 2) \max_{-\tau \leq \nu \leq 0} \|y(\nu)\|^2 e^{-\frac{\beta}{\|H_{11}\|}t}.$$

The proof terminates by noticing that

$$\|y_1(t)\| = \|T\mathcal{P}x(t)\| \text{ and } \|T^{-1}\|^{-1} \|\mathcal{P}x(t)\| \leq \|T\mathcal{P}x(t)\| \leq \|T\| \|\mathcal{P}x(t)\|.$$

□

Remarks.

- In [Theorem 2.1](#), the connection between H and C is the analogue of Eq. (2). In particular, if $E = I$, $B = 0$, then the bound in (ii) reduces to (4).
- Similar to the ordinary case, the bound on $\|\mathcal{P}x(t)\|$ depends on the solvability of the Lyapunov-type equation involving C and H .

- In the expression of α , the coefficient $\ln 2$ results from our simple choice of the function v . Of course, more sophisticated choices can be considered.
- The factor α depends essentially on the condition number of T , which is an indicator of the quality of the spectral projection (8), and on $\|H_{11}\| = \|\hat{E}H\|$, which is an indicator of the stability of J and therefore of the pencil $\lambda E - A$. A large value of α can result in a transient growth of $\mathcal{P}x(t)$. The factor β depends on $\lambda_{\min}(C)$ and $\|H_{11}\|$ and is responsible for the decay and asymptotic behaviour of $\mathcal{P}x(t)$.
- The assumption of the theorem results in constraints mainly on the matrix B . Indeed, by considering the Schur complement, the positive definiteness of C is equivalent to

$$\hat{A}^T H + H^T \hat{A} + \frac{1}{\tau} I_n < 0, \quad (14)$$

$$\hat{B}^T H \left(\hat{A}^T H + H^T \hat{A} + \frac{1}{\tau} I_n \right)^{-1} H^T \hat{B} + \frac{1}{2\tau} I_n > 0. \quad (15)$$

If we assume that the pencil $\lambda E - A$ is stable, then the condition (14) is easily satisfied. Indeed, by considering, for example, the matrix $K = \frac{c}{\tau} I_r$ of size $r \times r$ with $c > 1$, the Lyapunov equation $J^T H_{11} + H_{11} J + K = 0$ has a unique solution $H_{11} = H_{11}^T > 0$. Then

$$\hat{A}^T H + H^T \hat{A} + \frac{1}{\tau} I_n = \begin{pmatrix} \frac{1-c}{\tau} I_r & 0 \\ 0 & H_{22} + H_{22}^T + \frac{1}{\tau} I_{n-r} \end{pmatrix} < 0.$$

In particular, the choice $H_{22} = -\frac{c}{2\tau} I_{n-r}$ leads to $\hat{A}^T H + H^T \hat{A} + \frac{1}{\tau} I_n = -\frac{c-1}{\tau} I_n < 0$.

The condition (15) can be written $(W^{-1} B T^{-1})^T H H^T (W^{-1} B T^{-1}) - \frac{c-1}{2\tau^2} I_n < 0$ and will be satisfied if $\|B\| \leq (\|H\| \|T^{-1}\| \|W^{-1}\|)^{-1} \left(\frac{c-1}{2\tau^2} \right)^{\frac{1}{2}}$.

For example if $n = 4$, $A = \begin{pmatrix} J & 0 \\ 0 & I_2 \end{pmatrix}$, $J = \begin{pmatrix} -3 & 1 \\ 0 & -2 \end{pmatrix}$, $E = \begin{pmatrix} I_2 & 0 \\ 0 & 0 \end{pmatrix}$, $\tau = 1$, then the equation $J^T H_{11} + H_{11} J + 2I_2 = 0$ has the unique solution $H_{11} = \begin{pmatrix} \frac{1}{15} & \frac{1}{15} \\ \frac{1}{15} & \frac{8}{15} \end{pmatrix}$. Taking $H_{22} = -I_2$ and letting $H = \begin{pmatrix} H_{11} & 0 \\ 0 & H_{22} \end{pmatrix}$ leads to $A^T H + H A^T + \frac{1}{\tau} I_4 = -I_4$ and hence the condition (14) is satisfied. The condition (15) will be satisfied for any matrix B such $-2B^T H^2 B + I_4$ is positive definite, and, in particular if $\|B\| \leq \frac{1}{\sqrt{2}\|H\|} = 0.70711$.

Under the assumption of Theorem 2.1, the pencil $\lambda E - A$ is stable and hence the matrix A is nonsingular. Multiplying equation (5) on the left by $(I_n - \mathcal{P})A^{-1}$ and noting that $(I_n - \mathcal{P})A^{-1}E = 0$, we obtain

$$0 = (I_n - \mathcal{P})x(t) + (I_n - \mathcal{P})A^{-1}Bx(t - \tau), \quad t > 0, \quad (16)$$

which will be used to derive a bound analogous to (4).

Theorem 2.2. Under the assumption of Theorem 2.1, let $t = k\tau + \mu$, $0 \leq \mu \leq \tau$, $k \geq 0$. Then

$$\|x(t)\| \leq \alpha \sum_{j=0}^k \left(\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} \right)^j e^{-\frac{\beta}{2\|H_{11}\|}t} + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|,$$

where α and β are defined in Theorem 2.1.

Proof. For $0 \leq t \leq \tau$, the equality (16) gives

$$\|(I_n - \mathcal{P})x(t)\| \leq \|(I_n - \mathcal{P})A^{-1}B\| \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \quad (17)$$

and for $t = \tau + \mu$, $0 \leq \mu \leq \tau$, it gives

$$\|(I_n - \mathcal{P})x(t)\| \leq \|(I_n - \mathcal{P})A^{-1}B\| [\|\mathcal{P}x(t - \tau)\| + \|(I_n - \mathcal{P})x(t - \tau)\|].$$

From [Theorem 2.1](#) and the inequality [\(17\)](#) we obtain

$$\|\mathcal{P}x(t-\tau)\| \leq \alpha e^{-\frac{\beta}{2\|H_{11}\|}\mu}, \quad \|(I_n - \mathcal{P})x(t-\tau)\| \leq \|(I_n - \mathcal{P})A^{-1}B\| \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|.$$

Therefore

$$\|(I_n - \mathcal{P})x(t)\| \leq \alpha \|(I_n - \mathcal{P})A^{-1}B\| e^{-\frac{\beta}{2\|H_{11}\|}\mu} + \|(I_n - \mathcal{P})A^{-1}B\|^2 \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|. \quad (18)$$

Continuing this way we easily obtain for $t = k\tau + \mu$, $0 \leq \mu \leq \tau$ a generalization of [\(17\)](#) and [\(18\)](#) as follows

$$\|(I_n - \mathcal{P})x(t)\| \leq \alpha \sum_{j=1}^k \|(I_n - \mathcal{P})A^{-1}B\|^j e^{-\frac{\beta}{2\|H_{11}\|}(t-j\tau)} + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|.$$

Hence

$$\begin{aligned} \|x(t)\| &\leq \|\mathcal{P}x(t)\| + \|(I_n - \mathcal{P})x(t)\| \\ &\leq \alpha \sum_{j=0}^k \|(I_n - \mathcal{P})A^{-1}B\|^j e^{-\frac{\beta}{2\|H_{11}\|}(t-j\tau)} + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \\ &= \alpha \sum_{j=0}^k \left(\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} \right)^j e^{-\frac{\beta}{2\|H_{11}\|}t} + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|. \end{aligned}$$

□

Corollary 1. Under the assumptions of [Theorem 2.1](#), let $t = k\tau + \mu$, $0 \leq \mu \leq \tau$, $k \geq 0$.

(i) If $\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} < 1$, then

$$\|x(t)\| \leq \left(\frac{\alpha}{1 - \|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}}} + \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \right) e^{-\frac{\beta}{2\|H_{11}\|}t}.$$

(ii) If $\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} = 1$, then

$$\|x(t)\| \leq \left(\alpha \left(\frac{t}{\tau} + 1 \right) + \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \right) e^{-\frac{\beta}{2\|H_{11}\|}t}.$$

(iii) If $1 < \|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}}$ and $\|(I_n - \mathcal{P})A^{-1}B\| \leq 1$, then

$$\|x(t)\| \leq \left(\frac{\alpha e^{\frac{\beta\tau}{2\|H_{11}\|}}}{\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} - 1} + \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \right) e^{-\ln\left(\|(I_n - \mathcal{P})A^{-1}B\|^{-\frac{1}{\tau}}\right)t}.$$

Proof. For the three cases we use [Theorem 2.2](#).

(i) If $\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} < 1$, then

$$\|x(t)\| \leq \alpha \sum_{j \geq 0} (\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}})^j e^{-\frac{\beta}{2\|H_{11}\|}t} + e^{-\frac{\beta\tau(k+1)}{2\|H_{11}\|}} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|$$

and the desired inequality follows from $(k+1)\tau \geq t$.

(ii) If $\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} = 1$, then

$$\|x(t)\| \leq \alpha(k+1) e^{-\frac{\beta}{2\|H_{11}\|}t} + e^{-\frac{\beta\tau(k+1)}{2\|H_{11}\|}} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\|$$

and the desired inequality follows from $t \leq (k+1)\tau \leq t + \tau$.

(iii) If $1 < \|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}}$ and $\|(I_n - \mathcal{P})A^{-1}B\| \leq 1$, then

$$\begin{aligned} \|x(t)\| &\leq \alpha \sum_{j=0}^k (\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}})^{j-k} (\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}})^k e^{-\frac{\beta}{2\|H_{11}\|}t} \\ &\quad + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \\ &\leq \alpha \sum_{j \geq 0} \left((\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}})^{-1} \right)^j (\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}})^k e^{-\frac{\beta}{2\|H_{11}\|}t} \\ &\quad + \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \\ &\leq \left[\frac{\alpha e^{\frac{\beta\tau}{2\|H_{11}\|}}}{\|(I_n - \mathcal{P})A^{-1}B\| e^{\frac{\beta\tau}{2\|H_{11}\|}} - 1} + \max_{-\tau \leq \nu \leq 0} \|x(\nu)\| \right] \|(I_n - \mathcal{P})A^{-1}B\|^{k+1} \end{aligned}$$

and the desired inequality follows again from $t \leq (k+1)\tau$. \square

As a consequence we have the following

Corollary 2. Under the assumption of [Theorem 2.1](#), the solution of (5) is asymptotically stable if $\|(I_n - \mathcal{P})A^{-1}B\| < 1$ and stable if $\|(I_n - \mathcal{P})A^{-1}B\| = 1$.

In particular, when $E = I$ and hence $\mathcal{P} = I_n$, [Corollary 2](#) implies that the solution of (5) is asymptotically stable, see also [\[25, Proposition 5.3\]](#).

Acknowledgements

The authors would like to thank the reviewers for useful remarks and suggestions.

References

- [1] Z. Gajić, M.T.J. Qureshi, Lyapunov Matrix Equation in System Stability and Control, Dover, Mineola, N.Y., 2008.
- [2] S.K. Godunov, Modern Aspects of Linear Algebra, in: Translations of Mathematical Monographs, vol. 175, American Math. Society, Providence, RI, 1998.
- [3] K. Veselić, Bounds for exponentially stable semigroups, Linear Algebra Appl. 358 (2003) 309–333.
- [4] G.H. Golub, C.F. Van Loan, Matrix Computations, third ed., The Johns Hopkins Univ. Press, Baltimore, MD, 1996.
- [5] M. Sadkane, R.B. Sidje, Efficient computation of the spectral projections of regular matrix pairs, J. Comput. Appl. Math. 298 (2016) 72–81.
- [6] L.N. Trefethen, M. Embree, Spectra and Pseudospectra: The Behavior of Nonnormal Matrices and Operators, Princeton University Press, Princeton, 2005.
- [7] C.F. Van Loan, How near is a stable matrix to an unstable matrix, Contemp. Math. 47 (1985) 465–477.
- [8] E. Fridman, Introduction to Time-Delay Systems, Analysis and Control, Springer International Publishing, 2014.
- [9] W. Michiels, S.-I. Niculescu, Stability and Stabilization of Time-Delay Systems: An Eigenvalue-Based Approach, in: Advances in Design and Control, vol. 12, SIAM Publications, Philadelphia, 2007.
- [10] S.L. Campbell, Nonregular 2D descriptor delay systems, IMA J. Math. Cont. Inf. 12 (1995) 57–67.
- [11] P. Ha, Analysis and Numerical Solutions of Delay Differential-Algebraic Equations (Dissertation), Technische Universität Berlin, 2015, <http://dx.doi.org/10.14279/depositionce-4385>.
- [12] B. Unger, Discontinuity propagation in delay differential-algebraic equations, Electron. J. Linear Algebr. 34 (2018) 582–601.

- [13] S. Trenn, B. Unger, Delay regularity of differential-algebraic equations, in: Proc. 58th IEEE Conf. Decision Control, Nice, France, S., 2019, pp. 989–994.
- [14] B. Unger, Well-Posedness and Realization Theory for Delay Differential-Algebraic Equations (Dissertation), Technische Universität Berlin, 2020, <http://dx.doi.org/10.14279/depositonce-10707>.
- [15] E. Fridman, New Lyapunov–Krasovskii functionals for stability of linear retarded and neutral type systems, Systems Control Lett. 43 (2001) 309–319.
- [16] E. Fridman, Stability of linear descriptor systems with delay: A Lyapunov-based approach, J. Math. Anal. Appl. 273 (2002) 24–44.
- [17] S.-I. Niculescu, P. Fu, J. Chen, On the stability of linear delay-differential algebraic systems: Exact conditions via matrix pencil solutions, in: Proc. 45th IEEE Conf. Decision Control, 2006.
- [18] S.L. Campbell, V.H. Linh, Stability criteria for differential-algebraic equations with multiple delays and their numerical solutions, Appl. Math. Comput. 208 (2009) 397–415.
- [19] W. Michiels, Spectrum-based stability analysis and stabilisation of systems described by delay differential algebraic equations, IET Control Theory Appl. 5 (S) (2011) 1829–1842.
- [20] N.H. Du, V.H. Linh, V. Mehrmann, D.D. Thuan, Stability and robust stability of linear time-invariant delay differential-algebraic equations, SIAM J. Matrix Anal. Appl. 34 (2013) 1631–1654.
- [21] G.W. Stewart, J.-G. Sun, Matrix Perturbation Theory, Academic Press, San Diego, California, 1990.
- [22] R.A. Horn, C.R. Johnson, Matrix Analysis, second ed., Cambridge University Press, New York, 2013.
- [23] T. Berger, A. Ilchmann, S. Trenn, The quasi-Weierstraß form for regular matrix pencils, Linear Algebra Appl. 436 (2012) 4052–4069.
- [24] B. Kågström, P. Poromaa, Computing eigenspaces with specified eigenvalues of a regular matrix pair (A, B) and condition estimation: theory, algorithms and software, Numer. Algorithms 12 (1996) 369–407.
- [25] K. Gu, V.L. Kharitonov, J. Chen, Stability of Time-Delay Systems, Birkhäuser, Boston, MA, 2003.